Title: Intrinsic growth rules of patients infected, dead and recovered with 2019

novel coronavirus in mainland China

Chuanliang Han[1△], Yimeng Liu[2,3,4△], Saini Yang[2,3,4*]

[1]State Key Laboratory of Cognitive Neuroscience and Learning & IDG/McGovern

Institute for Brain Research, Beijing Normal University, Beijing 100875, China

[2]Key Laboratory of Environmental Change and Natural Disaster, Ministry of

Education, Beijing Normal University, Beijing 100875, China

[3]State Key Laboratory of Earth Surface Processes and Resource Ecology, Beijing

Normal University, Beijing 100875, China

[4]Academy of Disaster Reduction and Emergency Management Ministry of

Emergency Management and Ministry of Education, Faculty of Geographical Science,

Beijing Normal University, Beijing 100875, China

[△]C.L.H and Y.M.L contributed equally to this work

*Correspondence authors:

Saini Yang,

Key Laboratory of Environmental Change and Natural Disaster, Ministry of

Education & State Key Laboratory of Earth Surface Processes and Resource Ecology

& Academy of Disaster Reduction and Emergency Management Ministry of

Emergency Management and Ministry of Education, Faculty of Geographical Science,

Beijing Normal University, Beijing 100875, China

Email: yangsaini@bnu.edu.cn

Abstract

An outbreak of a novel coronavirus (SARS-CoV-2)-infected pneumonia (COVID-19)

was first diagnosed in Wuhan, China, in December 2019 and then spread rapidly to

other regions. We collected the time series data of the cumulative number of

confirmed infected, dead, and cured cases from the health commissions in 31

provinces in mainland China. A descriptive model in a logistic form was formulated to

infer the intrinsic epidemic rules of COVID-19, which illustrates robustness spatially

and temporally. Our model is robust (R^2>0.95) to depict the intrinsic growth rule for

the cumulative number of confirmed infected, dead, and cured cases in 31 provinces

in mainland China. Furthermore, we compared the intrinsic epidemic rules of

COVID-19 in Hubei with that of severe acute respiratory syndrome (SARS) in

Beijing, which was obtained from the Ministry of Public Health of China in 2003. We

found that the infected case is the earliest to be saturated and has the lowest

semi-saturation period compared with deaths and cured cases for both COVID-19 and

SARS. All the three types of SARS cases are later to saturate and have longer

semi-saturation period than that of COVID-19. Despite the virus caused SARS

(SARS-CoV) and the virus caused COVID-19 (SARS-CoV-2) are homologous, the

duration of the outbreak would be shorter for COVID-19.

Introduction

In December 2019, a novel coronavirus-infected pneumonia causing human infection was first identified in Wuhan, a city of Hubei province located in central China with a population of 11 million people[1,2]. The coronavirus was firstly named as 2019-nCoV on January 12, 2020 and may have a probable bat origin[3]. Human to human transmission through 2019-nCoV has been confirmed among closed contacts since the middle of December 2019[4,5]. As of February 29, 2020, infections of 2019-nCoV have been identified in all Chinese provincial administrative regions and in dozens of countries around the world[6-10]. There have been a total of 79824 confirmed cases of 2019-nCoV infections in mainland China, including 2870 deaths and 41625 cases cured[11]. So far, the coronavirus has been renamed as COVID-19 on February 12, 2020 officially. Different from a previous famous beta coronavirus in 2003, severe acute respiratory syndrome coronavirus (SARS-CoV)[12-14] that has a mortality rates of 10%[15], that of COVID-19 is 4.13% in Hubei province, and 0.84% in rest of mainland China[11] up to February 29, 2020.

Here, we used a descriptive model to fit data on the 2019-nCoV cases (infection, death, cure) from January 20, 2020 to February 29, 2020, aiming to capture the epidemiologic characteristics and dynamics of COVID-19 in 31 provinces in mainland China. Furthermore, we compared these characteristics with SARS in 2003 using the same model structure in order to validate this model and understand the differences among these two diseases.

Results

Intrinsic growth rules of COVID-19 infections in mainland China

Hubei province contributed a majority of infections to this statistical numbers (Fig

1A). Based on the cumulative infected case data of COVID-19, we firstly established

a descriptive model with three parameters for each of the 31 provinces in mainland

China (Fig 1B & Fig 2B, See Methods). Fitting goodness of models was expressed by

the R-square, which ranged from 0.95 to 0.999, indicating that the models we

established fitted the data well. The panel B of Figure 1 illustrates examples of time

series data of infected cases and corresponding fitting curves in five provinces

(provinces of Xinjiang, Beijing, Guangdong, Zhejiang and Heilongjiang) and that of

all provinces can be found in Supplementary Figure S1. It is clear that different

province varied in patterns of increase of infected cases (Fig 1B).

In our model, parameter k is an important parameter to depict the intrinsic growth rule

of cumulative number of infected patients (Equation (8)). We assume that the growth

pattern follows a sigmoid form, and the parameter k here denotes how rapid the

number of infected cases get to saturate. If k is large, the S form of the curve would be

sharp, and the curve will rise up very quickly to be saturated to the maximum number.

While if k is only slightly larger than 0, the curve would show a slow and smooth

increase from 0 to the maximum.

To further illustrate the spatial distribution of the infected cases' growth pattern, we

exerted further analyses of the fitted infected case curves of 30 provinces. We

excluded Tibet in this analysis because the number of confirmed infected case did not

increase since Jan 29, 2020. The only one confirmed case in Tibet has been cured on

Feb 12. The spatial distribution of parameter k is shown in Fig 2A, illustrating

provinces varied in the increase patterns of cumulative infected cases. Fitted curves of

provinces such as Jilin and Qinghai has larger value of k, which means that the

number of infections peaked earlier in these regions than in other provinces. By

contrast, provinces like Hainan, Hebei, Ningxia, has smaller value of k than other

provinces, suggesting the infections of these provinces would continue increasing

when infections of other provinces reach saturation (Fig 2B). Fig 2C shows the spatial

distribution of predicted maximum number of infected cases (parameter A) in each

province of mainland China. With the data up to Feb. 29th, our model shows that the

maximum number of infected COVID-19 cases in mainland China would be 80119

(95% confidence interval: [79133, 81105]). Among all the provinces in mainland

China, the Hubei province is predicted to have largest number of infected COVID-19

cases, which would be 66750 (95% confidence interval: [65939,67561]).

Our model could not only well explain the data in various geographic space, but also

provided robust results with different length of time series (Fig 2D-F). We used our

model to fit the data by Feb 19 and Feb 29 respectively, and the parameters estimated

by the two dataset do not have significant difference (for k, t=0.09, p=0.93; for t_0,

t=0.04, p=0.97; for A, t=0.02, p=0.98; two-tailed test). This model's spatial and

temporal robustness provides us confidence in further analyses.

Comparison of intrinsic epidemic rules between COVID-19 in Wuhan and SARS

in Beijing (2003)

To compare the intrinsic growth rule of infected cases, dead cases and cured cases

between COVID-19 and SARS, we applied the descriptive model to fit the data of

SARS in Beijing 2003 (Fig 3A). Since the time series data of SARS was full phase of

the outbreak, the parameters estimated by our model was solid (R-square>0.95). We

found the k value of fitted curves of infected SARS cases in Beijing was 0.18, which

was much lower than that of COVID-19 ($k = 0.25$) in Hubei Province (Fig 3 ABC).

Furthermore, we found that all 30 provinces' k values were significantly larger than

0.18 (t=10.71, p<0.0001, right-tailed test) (Fig 3D). We also noted the t_0 of fitted

curves of infected SARS cases in Beijing was larger than that of fitted curves of

infected COVID-19 cases in all provinces (Fig 3E). Moreover, the increasing rate (k)

was negatively correlated (Pearson, r = -0.44, p = 0.0125) to semi-saturation period

(Fig 3F). We concluded that at the provincial level, the COVID-19 spread more

quickly than the SARS did. The infections of COVID-19 increased more rapid and

would reach saturation earlier than SARS according to our model.

Intrinsic rules of infection, death and recovery of COVID-19 and SARS2003

The outbreak of SARS in Beijing was the most serious among all the provincial

administrative regions, and the outbreak of COVID-19 was most serious in Hubei

province. Here we used our descriptive model to fit the dead and cured cases' curves

for SARS in Beijing and COVID-19 in Hubei province (Fig 4 A-C, F-H, R-square≥

0.95, for other provinces in mainland China, see Fig S2, S3), and compared the

growth features of three types cases of SARS and COVID-19 by two parameters, k

and t_0 (Fig 4 DEIJ). We found that k values of the three types cases of SARS were

lower than that of COVID-19 (Fig 4 DE), and semi-saturation period for them in

SARS was higher than COVID-19's, which suggested the duration of COVID-19

would be shorter than that of SARS. For both SARS and COVID-19, their infected

curves have the lowest semi-saturation period (Fig 4 EJ), while cured curves have the

highest which is in keeping with the common sense that the cure is always later than

infection and death. For both SARS in Beijing and COVID-19 in Hubei, the k values

of death case curves were lowest compared with curves of infected cases and cured

cases. For both SARS and COVID-19, the k value of infected case curve was larger

than that of cured curve.

Discussion

We modeled the epidemiologic characteristics of 2019-nCoV for 31 provinces in

mainland China based on the data from Jan 10 to Feb 29 during the outbreak (Hubei

province reported laboratory confirmed 2019-nCoV cases since Jan 10, while other

provinces in mainland China reported laboratory confirmed 2019-nCoV cases since

Jan 20). Furthermore, we compared these characteristics with SARS in 2003 using the

same model framework.

Compared with dynamical models such as SIR (Susceptible - Infectious - Recovered)

or SEIR (Susceptible - Exposed - Infectious - Recovered), our model is relatively

simple but robust. We used a differential equation in logistic form to describe the

dynamics of the infected, dead and healed cases caused by COVID-19 respectively

(See Methods). The increasing of the population no matter in which type of case

(infection, death or recovery) has multiple reasons. When the outbreak of COVID-19

originated in Wuhan, it did not cause too much concern at the beginning so that it

leads to many infections and starts to export potential exposed cases to other

provinces. But as the successive closure of the cities in Hubei province, and isolation

of the community in all the provinces in mainland China, the virus spread tendency

was weakened. In our model, we also considered these measures taken by the

government. We rewrote Equation (1) to Equation (9).

$$\frac{dI(t)}{dt} = \alpha I(t) - rI^2(t) \tag{9}$$

where α is growth rate of the number of infections; *I(t)* is cumulative number of

infections at time *t*; r is a parameter to reflect the effect of measures aiming at

controlling the virus spread. Here we decompose $I(t)$ into two parts. The first part

$\alpha I(t)$ represents the rapid infections growth in the early phase of the outbreak

without any measures for epidemic control. The second part $rI^2(t)$ illustrates the

slower infections growth after taking effective isolation measures for epidemic control.

For death cases, this idea still holds. But things are different for cured case. In contrast,

at the early phase of the outbreak, the number of cured cases increase slowly. But

once there are effective treatments, the number of cured cases will rise up quickly.

The growth of cured cases would again be slow since the number of patients is limited.

All these facts were considered in our highly summarized model (Equation (8)) in a

macroscopic scale. Each fitted model has a R-square value greater than 0.95 to the

original data set, so that we could further discuss some intrinsic mechanisms for the

growth rules of infection, death and recovery through the parameters estimated from

this model (See Methods).

The value of k from Equation (8) is a critical parameter. k represents how rapid the

number of infected cases get to the saturation. We found the variability of k values

among fitted curves of infected COVID-19 cases in 30 provinces (exclude Tibet) (Fig

2 AB) is large (Ranging from 0.19 to 0.38). It is clear to show the real situation in

each province and this guides us to be more careful against the epidemic. Further we

compared this characteristics between COVID-19 and SARS in 2003 in typical

provinces (Fig 3 AB). We noted that k values of fitted COVID-19 infection curves in

30 provinces of mainland China (exclude Tibet) are larger than that of SARS in

Beijing 2003 (Fig 3 C), which suggests the time cost of this epidemic will be shorter

compared with SARS in 2003. This result may relate to several aspects. Firstly, it may

due to the basic spread property of the virus. Besides, the isolation policy of all the

provinces in mainland China was more strictly enforced than it was in 2003.

Moreover, the improved clinical skills today may also helpful for shortening the

duration of epidemic.

Furthermore, we analyzed semi-saturation period (Fig 4 EK) (t_0 in Equation (8)) and

make comparison in three types of cases (infection, death and recovery) (Fig 4 A-C

G-I). k values for the fitted curves of the three types of cases caused by COVID-19

were larger than that of SARS, while t_0 values were the opposite. This again suggests

that the epidemic of COVID-19 will be quicker to pass away than SARS. For both

SARS and COVID-19, fitted curves of infections have lowest semi-saturation period

(Fig 4 EK), while fitted curves of cured cases have the highest semi-saturation period,

which follows the common sense that the cure was always later than infection and

death since it costs time to get on the research of effect plans to cure the patients. For

both SARS and COVID-19, the k value of infected case curve was larger than that of

cured curve, while that of death case is the lowest.

Methods

Sources of data

We obtained the time series data of 2019-nCoV cases from the National Health

Commission of China, and the provincial Health Commission of 31 provincial

administrative regions in mainland China (January 10 to February 29, 2020). The data

used in this study include the number of confirmed 2019-nCoV cases, the cumulative

number of suspected, dead, and cured cases. The 200 infected cases found in jail of

Shandong Province on Feb 20 were not counted in. All cases were laboratory

confirmed following the case definition by national health commission of China[16].

The basic test procedure has been described in detail in previous work[4,17].

The severe acute respiratory syndrome (SARS) in 2003 was another fatal coronavirus

epidemics over the last two decades, we collected the data (the cumulative number of

suspected, dead, and cured cases) of SARS during March 1 to August 16, 2003, in

Beijing from the National Health and Family Planning Commission of PRC.

Epidemic curve modeling

We assumed the number of susceptible people per unit time was proportional to the

number of uninfected persons, and isolation or closure of the city could affect the

spread effect of infected to susceptible people to a certain extent. Previous studies

have showed that daily cumulative number of patients could be explained by logistic

function[18,19]. Based on the above assumption and existing findings, we modeled the

epidemic information of virus as logistic form (Equation (1))

239
$$\begin{cases} \frac{dI(t)}{dt} = \alpha I(t)(1 - \frac{I(t)}{N_1}) \\ I(0) = I_0 \end{cases} \tag{1}$$

240 where N_1 is maximum number of cumulative infections, $I(t)$ is cumulative number of

241 patients at time t, α is the incidence growth rate, and I_0 is the number of infected

242 cases at the initial time.

243 Analytical solution of Equation (1) could be written in the form of Equation (2)

244
$$I(t) = \frac{N_1}{1 + C_1 e^{-\alpha t}} \tag{2}$$

245
$$C_1 = \frac{N_1}{I_0} - 1 \tag{3}$$

246 Similarly, we could write the analytical solutions for the dead and cured cases by

247 Equation (4) and Equation (6).

248
$$D(t) = \frac{N_2}{1 + C_2 e^{-\beta t}} \tag{4}$$

249
$$C_2 = (\frac{N_2}{D_0} - 1) \tag{5}$$

250
$$R(t) = \frac{N_3}{1 + C_3 e^{-\gamma t}} \tag{6}$$

251
$$C_3 = (\frac{N_3}{R_0} - 1) \tag{7}$$

252 Where β, γ is the growth rate of dead cases and the growth rate of cured cases,

253 respectively. N_2 and N_3 is maximum number of cumulative dead and cured cases. D_0

254 and R_0 is the number of dead and cured cases at initial time.

255 Here we simplified Equation (2),(4),(6) and unified them as Equation (8)

256
$$N(t) = \frac{A}{1 + e^{-k(t - t_0)}} \tag{8}$$

where $N(t)$ is the general form of $I(t)$, $D(t)$ or $R(t)$, A denotes N_1, N_2, or N_3, and k

represents α, β, or γ, indicating the intrinsic rule for growth of cases. t_0 represents

the semi-saturation period, illustrating the inflection point of the sigmoid curve.

The data processing and modeling were performed on MATLAB (The MathWorks)

with custom scripts. The nonlinear least square (NLS) algorithm was adopted for data

fitting and parameter estimation. We used the MATLAB function "nlinfit" to

minimize the sum of squared differences between the data points and the fitted values.

Equation (8) was used in this work as the descriptive model to depict the intrinsic

growth rule for infected, dead and cured cases.

References:

Zhu, N. *et al.* A Novel Coronavirus from Patients with Pneumonia in China, 2019. *N Engl J Med* **382**, 727-733, doi:10.1056/NEJMoa2001017 (2020).

Hui, D. S. *et al.* The continuing 2019-nCoV epidemic threat of novel coronaviruses to global health - The latest 2019 novel coronavirus outbreak in Wuhan, China. *Int J Infect Dis* **91**, 264-266, doi:10.1016/j.ijid.2020.01.009 (2020).

Zhou, P. *et al.* A pneumonia outbreak associated with a new coronavirus of probable bat origin. *Nature*, doi:10.1038/s41586-020-2012-7 (2020).

Li, Q. *et al.* Early Transmission Dynamics in Wuhan, China, of Novel Coronavirus-Infected Pneumonia. *N Engl J Med*, doi:10.1056/NEJMoa2001316 (2020).

Wang, D. *et al.* Clinical Characteristics of 138 Hospitalized Patients With 2019 Novel Coronavirus-Infected Pneumonia in Wuhan, China. *JAMA*, doi:10.1001/jama.2020.1585 (2020).

Holshue, M. L. *et al.* First Case of 2019 Novel Coronavirus in the United States. *N Engl J Med*, doi:10.1056/NEJMoa2001191 (2020).

Ki, M. & nCo, V. T. Epidemiologic characteristics of early cases with 2019 novel coronavirus (2019-nCoV) disease in Republic of Korea. *Epidemiol Health*, e2020007, doi:10.4178/epih.e2020007 (2020).

Shigemura, J., Ursano, R. J., Morganstein, J. C., Kurosawa, M. & Benedek, D. M. Public responses to the novel 2019 coronavirus (2019-nCoV) in Japan: Mental health consequences and target populations. *Psychiatry Clin Neurosci*, doi:10.1111/pcn.12988 (2020).

Munster, V. J., Koopmans, M., van Doremalen, N., van Riel, D. & de Wit, E. A Novel Coronavirus Emerging in China - Key Questions for Impact Assessment. *N Engl J Med* **382**, 692-694, doi:10.1056/NEJMp2000929 (2020).

Nishiura, H. *et al.* The Rate of Underascertainment of Novel Coronavirus (2019-nCoV) Infection: Estimation Using Japanese Passengers Data on Evacuation Flights. *J Clin Med* **9**, doi:10.3390/jcm9020419 (2020).

Prevention, C. C. f. D. C. a. *The latest situation of COVID-19*, <http://www.nhc.gov.cn/xcs/yqtb/202003/9d462194284840ad96ce75eb8e4c8039.shtml> (2020).

Drosten, C. *et al.* Identification of a novel coronavirus in patients with severe acute respiratory syndrome. *New Engl J Med* **348**, 1967-1976, doi:DOI 10.1056/NEJMoa030747 (2003).

Ksiazek, T. G. *et al.* A novel coronavirus associated with severe acute respiratory syndrome. *N Engl J Med* **348**, 1953-1966, doi:10.1056/NEJMoa030781 (2003).

Kuiken, T. *et al.* Newly discovered coronavirus as the primary cause of severe acute respiratory syndrome. *Lancet* **362**, 263-270, doi:Doi

10.1016/S0140-6736(03)13967-0 (2003).

15   WHO. *Summary of probable SARS cases with onset of illness from 1*
November   2002   to   31   July   2003,
<https://www.who.int/csr/sars/country/table2004_04_21/en/> (2003).

16   Prevention, C. C. f. D. C. a. *Prevention and control measures of COVID-19*,
<http://www.nhc.gov.cn/jkj/s3577/202002/a5d6f7b8c48c451c87dba14889b30
147/files/3514cb996ae24e2faf65953b4ecd0df4.pdf> (2020).

17   Huang, C. *et al.* Clinical features of patients infected with 2019 novel
coronavirus   in   Wuhan,   China.   *Lancet*   **395**,   497-506,
doi:10.1016/S0140-6736(20)30183-5 (2020).

18   Huang, D., Guan, P. & Zhou, B. Fitness of morbidity and discussion of
epidemic characteristics of SARS based on logistic models. *Chinese Journal*
of Public Health **19**, 1-2 (2003).

19   Wang, Y. & Liu, x. The compound Logistic model used to describe epidemic
situation dynamics of SARS in Beijing. *Journal of China Jiliang University*
16, 159-162 (2005).

ADDITIONAL INFORMATION

Acknowledgements:
This study is sponsored by the National Key Research and Development Program of
China (2018YFC1508903), the National Natural Science Foundation of China
(41621061) and the support of International Center for Collaborative Research on
Disaster Risk Reduction (ICCR-DRR).

Contributions：
C.L.H, Y.M.L and S.N.Y designed the research. C.L.H and Y.M.L performed the
research. C.L.H and Y.M.L. analyzed the data. Y.M.L, C.L.H and S.N.Y wrote the
paper.

Corresponding author:
Correspondence to Yang Saini.
Email: yangsaini@bnu.edu.cn

Competing interests
The authors declare no competing interests.

Figure 1 Spatial distribution
of the number of COVID-19
infected cases in mainland
China.
Panel A is the spatial distribution of
the number of COVID-19 infected
cases of provinces in mainland
China.
Panel B takes five provinces
(Xinjiang, Beijing, Guangdong,
Zhejiang and Heilongjiang) as
example to show the time series of infected cases and corresponding fitted curve. The
horizontal axis is the x_{th} day after Jan 1, 2020. And the vertical axis denotes the
number of infected cases in corresponding province. The black dots in each panel are
raw data of the cumulative infected cases. The red line is the fitted curve by our
descriptive model. The dashed red lines show the 95% confidence interval of the
fitted curves.
370
371

Fig 2 Spatial distribution of model parameters and robustness of the model

Panel A demonstrates the spatial distribution of parameter *k* of provinces in mainland China (exclude Tibet, same for B-F).

Panel B shows the fitted curves of infected cases in 30 provinces (exclude Tibet). The vertical axis is the number of infected cases (normalized by dividing the maximum number of infected cases in each province). The color of each province is consistent with Panel A.

Panel C is the spatial distribution of the predicted maximum number (parameter A in our model) of COVID-19 infected cases of provinces in mainland China.

Panel D-F show the difference of parameters estimated from long (by the end of Feb 29) and short (by the end of Feb 29) time series data, each dot denotes for one province.

Fig 3 Comparison of intrinsic epidemic rules between COVID-19 in Wuhan and SARS in Beijing (2003)

Panel A shows the raw data and fitted curve of SARS cases in Beijing. The horizontal axis is the x_{th} day after March 1, 2003. The vertical axis denotes the cumulative number of SARS cases in Beijing. The black dots are the raw data of the SARS cases. The red line is the fitted curve by our descriptive model. The dashed red lines are the 95% confidence interval of the fitted curves.

Panel B shows the raw data and fitted curve of COVID-19 cases in Hubei province. The horizontal axis

is the x_{th} day after Jan 1, 2020. The vertical axis denotes the cumulative number of
COVID-19 cases in Hubei. The black dots are raw data of the COVID-19 cases. The
red line is the fitted curve by our descriptive model. The dashed red lines are the 95%
confidence interval of the fitted curves.
Pannel C shows the fitted curves of infected SARS cases (gray curve) and COVID-19
(black curve), which are normalized by dividing the maximum number of infected
cases in each province.
Panel D illustrates the *k* value (increasing rate of infected cases) of fitted curves of
infected COVID-19 cases in each province in mainland China, where Hubei is
represented by the solid red square (the same in D and E). *k* value of fitted curves of
infected SARS cases in Beijing is represented by the solid red triangle (the same in D
and E).
Panel E illustrates the t_0 (semi-saturation period) of fitted curves of infected
COVID-19 cases in each province in mainland China, as well as t_0 of fitted curves of
infected SARS cases in Beijing.
The scatter plot of *k* and t_0 is shown in Panel F, indicating significant negative
correlation   between   *k*   and t_0.

Fig 4 Intrinsic rules of infection, death and recovery of COVID-19 and SARS2003

Panel A-C show the raw data and fitted curve of SARS cases in Beijing. Their horizontal axises are the x_{th} day after Mar 1, 2003. The black dots in Panel A, B and C are raw data of the SARS infected cases, dead cases and cured cases, respectively. The red lines are the fitted curves by our descriptive model. The dashed red lines are the 95% confidence interval of the fitted curves.

Panel D and Panel E respectively shows the k value and the t_0 of fitted curves of three types cases caused by SARS (the circle, triangle and square denotes infected, dead and cured cases, respectively, and the same in Panel I and J). The error bars show the 95% confidence interval of the parameters.

Panel F-H show the raw data and fitted curve of COVID-19 cases in Hubei. Their horizontal axises are the x_{th} day after Jan 1, 2020. The black dots in Panel F, G and H are raw data of the COVID-19 infected cases, dead cases and cured cases, respectively. The red lines are the fitted curves by our descriptive model. The dashed red lines are the 95% confidence interval of the fitted curves.

Panel I and Panel J respectively shows the k value and the t_0 of fitted curves of three types cases caused by COVID-19, which is same with D and E.

451
452 **Supplement Figure S1 Intrinsic growth rules of patients infected with 2019 novel**
453 **coronavirus in all provinces of mainland China**
454 The black dots in each panel are raw data of the cumulative infected cases. The red
455 line is the fitted curve by our descriptive model.
456

457
458 **Supplement Figure S2 Intrinsic growth rules of patients dead with 2019 novel**
459 **coronavirus in all provinces of mainland China (Excluded 0 death case**
460 **provinces)**
461 The black dots in each panel are raw data of the cumulative dead cases. The red line is
462 the fitted curve by our descriptive model.

Supplement Figure S3 Intrinsic growth rules of patients cured with 2019 novel coronavirus in all provinces of mainland China

466 The black dots in each panel are raw data of the cumulative cured cases. The red line
467 is the fitted curve by our descriptive model.

471