The long-term impact on self-reported health, function and comorbidities from lower limb

apophysitis: Protocol of a cross-sectional study

Kasper Krommes[1], Kristian Thorborg[1], Lasse Christensen[2], Per Hölmich[1]

[1]Sports Orthopedic Research Center – Copenhagen, Orthopedic Department, Amager-Hvidovre,

Copenhagen University hospital

[2]Department of Sports Science and Clinical Biomechanics, University of Southern Denmark,

Odense, Denmark

Contact to corresponding author: Email: kasper.krommes@regionh.dk, twitter: @Krommes

Registration identifier: ClinicalTrials.gov: NCT04313621

Conflict of interest: The authors have no potential conflicts of interest to declare.

Author contributions: KK conceptualized and planned the study, and drafted the protocol. KK, LC,

KT, and PT revised and approved the protocol.

Revision history

Version	Date	Amendment
1.1	24-APR-2020	Added identifier for ethical and data approval, and pre-registration. Wording of title changed. Added details to analysis plan. Added secondary objective. Attached statistical code (stata) and link to the survey in Danish.

Abstract

Lower limb apophysitis cause long-term pain, decrease in function, and can reduce or completely

hinder participation in sports and physical activity. Yet, there is little knowledge on the long-term

consequences for health. Our objective with this investigation is to capture self-reported health-

status for all adults diagnosed with lower limb apophysitis in the period of 1977 to 2020 in Danish

secondary care and compare these data with normative values for the background population. We

are therefore conducting a national cross-sectional study based on data from the Danish National

Patient Registry. In this protocol we describe the planned methods.

Background, rationale, and review of the literature

Being physically active during adult life is key for health and prevention of disease, and carries

additional benefits during adolescence such as improved academic abilities and cognitive

function.[1–4] Besides somatic advantages, participating in sports fosters meaningful social

networks, lowers the risk of criminal activity, drinking, and substance abuse.[5–7] The levels of

physical activity declines during early adolescence, and less than 20% are currently meeting

recommendations for moderate to vigorous physical activity or sports participation.[8]

As adolescence is a period of increased autonomy, behaviors established during this period could

potentially last into adulthood.[9,10] Several barriers exist for participation in physical activity and

sports during adolescence, such as the risk of injury and pain during activity. In line with this, lower

limb pain is the most frequent cause for seeking primary care during adolescence[11], and up to half

of sports-active adolescents regularly take pain medication for injury-related pain.[12] In addition,

almost a third of adolescents quitting their sport, reports injuries or pain as the main reason

Adolescents are specifically susceptible to growth-related overuse injuries in the knee and heel.[13–

17] For early adolescence, when growth velocity peaks (ages 10-16),[18] the most common growth-

related injuries are apophysitis in the lower limb; *Morbus Severs* in the heel, and *Morbus Sinding*

Larsen Johansson and *Morbus Osgood Schlatter* in the knee.[19] All three entities cause long-term

pain, decrease in function, and can reduce or completely hinder participation in sports and physical

activity.[20–25] The most common, *Osgood Schlatter*, affects 1 in 4 active adolescents.[15,26–29] Yet

there is little knowledge on the long-term consequences for health, with only a small case-series on

younger participants[24] and in college-aged students.[30]

Objective

To review the long-term consequences of having had a lower limb apophysitis in a larger sample,

we aim to capture self-reported health-status for all adults in Denmark having been diagnosed with

such from 1977 to 2020 and compare these data with normative values for the background

population. Secondly, to examine if self-reported historical apophysitis symptoms (duration,

symptom severity, restrictions in participation) are related to current symptoms or health by

comparing subgroups.

Research question

What is the health status of adults with a history of lower limb apophysitis?

Methods

Design

We wish to conduct a national cross-sectional study based on self-reported data from patients

registered with a relevant diagnosis code in the Danish National Patient Registry (LPR). The study

is approved by the regional research ethics committee (Region Hovedstaden, H-20016972) and

the regional data review board (P-2020-433). The study is registered at ClinicalTrials.gov

(NCT04313621), and the protocol is published as pre-print on medRxiv (DOI:

10.1101/2020.03.01.20029660). On the first page of the survey presented to the participants, there

will be information about possible risks, study-participant rights, and aspects of data-sharing and

data-protection, where they will be asked to consent. If participants wish to redraw their answers

from the survey, they can do so. A contact email for the project coordinator (KK) is provided in the

invitation message. The study protocol is based on relevant items from the STROBE checklist[31] for

reporting of cross-sectional studies and the SPIRIT checklist for trial protocols.[32] The study is

descriptive and exploratory, and no pre-determined hypotheses are therefore being tested; instead,

it aims to generate potential future hypotheses to be tested in longitudinal designs.

Recruitment and eligibility

All participants aged 18-55 having received diagnosis codes pertaining to lower limb apophysitis

(Severs, Sinding-Larsen Johansson, Osgood Schlatter) in the years 1977-2020 in Denmark will be

eligible to participate. All eligible participants will be contacted and invited to participate by filling

out the electronic survey online. Sample size will be of convenience with no pre-determined cut-off

for the smallest or largest sample of interest. Rather, the final sample of respondents will be based

on however many have been designated the diagnosis codes and responds to the survey.

Measures

The full survey in Danish is freely available (doi.org/10.6084/m9.figshare.14504958.v1). Outcomes

and their design is based on relevant sports- and knee-related measures, and the 'National Health

Profile 2017' (Sundhedsprofilen 2017), where 3% (180.000) of the Danish population is surveyed

on a range of health and disease questions.[33] The following outcomes are ranged from the most

important to the least important regarding this study's objective.

<u>Self-rated health</u>

Measured on the SF-12 (Short-form 12 item) health survey and specific questions which computes

a Mental Component Summary scale (MCS-12) and a Physical Component Summary scale (PCS-12).[34] From a scale from 0-100 where higher scores indicate better health. The PCS-12 will be used in this study. In the National Health Profile "Bad physical health" are defined as a score above 35.37 on the PCS, and 'Bad physical health" as a score above 35.76 on MCS.

Knee-related pain and symptoms, and physical activity

We will ask about the current extent of physical activity[35] and knee-related pain and symptoms (KOOS).[36] The subscale dimensions used in this context will be KOOS-pain, KOOS-symptoms, and KOOS Sport/rec. Physical Activity Scale 2 (PAS-2) is used to capture daily and weekly physical activity.[37,38]

Other diseases

To describe this population and their rate of other diseases, participants will be asked if they have any other diseases in accordance with conditions from the National Health Profile.[33]

Apophysitis-related history

We will ask questions related to the duration and severity of their apophysitis.

Sleep problems, pain, and mental health

These factors are related to prognosis or severity for many other well-researched musculoskeletal conditions.[33,39]

Data collection and sharing

All data will be directly entered into REDCap (Research Electronic Data Capture),[40] a logged secure system designed for capturing sensitive non-commercial research data that is hosted at Hvidovre Hospital. REDCap contains options for valid values, range checks, data validation, branching, scheduling, and stop-rules to increase data quality.

Invitations to the questionnaire will be posted to participants digital government-issued inbox (e-boks) used for confidential correspondence (e.g. doctors' appointments), using their social security number (CPR number) obtained from the NPR. From pilot-testing, we expect the mean duration of filling out the questionnaire will be in the range of 10-15 minutes. If participants have questions or comments, there will be a contact-email posted in the survey-invitation.

The full dataset, excluding all personal identifiers, will be posted alongside the pre-print when the

results have been disseminated.

Analyses

The flow of participants from the potential total pool, to invitations, final respondents, and missing

data, will be described in detail and visualized in a consort-style flowchart. As this is an exploratory

single-group study with no pre-determined hypothesis and only self-assessed outcomes, no

blinding will be performed. However, we are committed to a pre-specified analysis strategy as

outlined below, and have compiled the statistical code for this before the completion of data

collection (Supplemental file 2: 'Stata code.pdf') All statistical analyses will be performed using

Stata version 16.1 (StataCorp, USA). Residuals of continuous variables will be visually inspected

for normality using a quartile-quartile plot. If the assumption of normal distribution of data is not

met, we will transform variables with appropriate methods for analytical use. The primary analysis

will be descriptive and reported in means and 95% confidence intervals, and if non-normally

distributed, reported as medians and inter-quartile ranges. The between-group difference in sub-

groups will be compared using an independent-samples t-test on PCS-12 and KOOS scores for

the continuous variables and will be reported with standardized effect sized (Cohens d) and

assessed as trivial ($d < 0.2$), small ($d \geq 0.2$), medium ($d \geq 0.5$) and large ($d \geq 0.8$).[41] The between-

group difference in diagnosis groups (Severs, Sinding-Larsen Johansson, Osgood Schlatter) will

be compared using a one-way analysis of variance (ANOVA) followed by the tukey post-hoc test to

analyze the difference between the groups (Severs, Sinding-Larsen Johansson, Osgood Schlatter)

on PCS-12 and KOOS scores. An alpha level of <5% is considered statistically significant. For the

dichotomous variables, the prevalence of knee and heel-related conditions is compared between

subgroups using unadjusted logistic regression. Similarly, outcomes are compared between strata

based on diagnosis, also using unadjusted logistic regression. The association between subgroups

are reported using odds ratios and a designated a priori defined level of magnitude. We have

outlined an explanation of this pre-determined relationship of odds ratio with magnitude (table 1),

ranging from a 'very large' magnitude of a reduction or increase in odds to a 'negligible' difference

with inspiration from GRADE Handbook (GRADing the quality of Evidence and the strength of

recommendations).[42]

Table 1. Odds ratio magnitude inspired from GRADE Handbook		
Odds ratio	Corresponding to	Magnitude
<0.2	> 5-fold reduction	Very large
0.21-0.50	> 100% reduction	Large
0.51-0.75	> 25% reduction	Moderate
0.76-0.90	> 10% reduction	Small
0.91-1.10	< 10% increase/reduction	Negligible difference
1.10-1.24	> 10% increase	Small
1.25-1.99	> 25% increase	Moderate
2.0-4.99	> 100% increase	Large
>5.0	> 5-fold increase	Very large

If the contingency tables needed for calculating odds ratios should contain null-responses and

therefore rendering the analysis impossible, we will post hoc explore other analysis strategies and

label these analysis as post hoc.

The *a priori* subgroups of interest are:

1. Participants reporting short (<6 months) vs. long (>6 months) apophysitis symptom duration.

2. Participants reporting significant limitation to sport and physical activity during their

apophysitis vs. those that were not significantly affected ('very' or 'totally' limited in sports

participation or physical activity during their apophysitis) pa participation during apo).

3. Participants who currently have knee pain or symptoms from the same general area vs.

those who currently do not.

4. Participants who report having met WHO recommendations (150 or 75 min of moderate-to-

vigorous physical activity weekyl)for physical activity in their adult life vs. does that report

having been less physically active.

5. Participants that report currently having a large bony prominence thought to originate from

their apophysitis (only Osgood Schlatter patients) vs. those that does not

6. Participants that report severe symptoms during their apophysitis vs. those who only report

having experienced light or moderate symptoms (based on pain intensity and restriction in

physical activity and sport).

If relevant sub-groups should emerge *post hoc*, they will be denoted as such. We will also compare

data from this sample with normative values for the background population.

Limitations

As this study only can include patients attending hospitals that are part of secondary care or

specialized care, the results will potentially only be relevant for severe cases, as mild cases likely

are managed either in their sports club, by their parents, by their general practitioner or other

primary care health professionals with direct access. Nevertheless, the patients in most need of

management and a more solid evidence-base will be the sample in question.

References

1. Henriksson P, Shiroma EJ, Henriksson H, et al. Fit for life? Low cardiorespiratory fitness in adolescence is associated with a higher burden of future disability. *Br J Sports Med*. Published online August 14, 2020.
2. Thompson WR, Sallis R, Joy E, Jaworski CA, Stuhr RM, Trilk JL. Exercise Is Medicine. *American Journal of Lifestyle Medicine*. 2020;14(5):511-523.
3. Donnelly JE, Hillman CH, Castelli D, et al. Physical Activity, Fitness, Cognitive Function, and Academic Achievement in Children: A Systematic Review. *Medicine & Science in Sports & Exercise*. 2016;48(6):1197.
4. von Rosen P, Olofsson O, Väsbom S, Heijne A. Correlates of health in adolescent elite athletes and adolescents: A cross-sectional study of 1016 adolescents. *Eur J Sport Sci*. 2019;19(5):707-716.
5. Werner EE, Smith RS. *Vulnerable but Invincible: A Longitudinal Study of Resilient Children and Youth*. 3 edition. Adams Bannister Cox Pubs; 1989.
6. Christensen KO, Lindstad J, Boje-Kovacs B, Sigurd F. Unge, sport og kriminalitet Undersøgelse af sportsklubbers kriminalitetsforebyggende potentiale i udsatte boligområder. :41.
7. Sigfusson J. Evidence Based Primary Prevention The Icelandic Model. :83.
8. Krommes K. Physical Activity & Sport for Health – What does it benefit? Published online February 17, 2020.
9. Nelson MC, Story M, Larson NI, Neumark-Sztainer D, Lytle LA. Emerging adulthood and college-aged youth: an overlooked age for weight-related behavior change. *Obesity (Silver Spring)*. 2008;16(10):2205-2211.
10. Rindfuss RR. The young adult years: diversity, structural change, and fertility. *Demography*. 1991;28(4):493-512.
11. Prathivadi Bhayankaram N, Lacey RJ, Barnett LA, Jordan KP, Dunn KM. Musculoskeletal consultations from childhood to adulthood: a longitudinal study. *J Public Health (Oxf)*.
12. Sari D, Thorlund JB, MIkkelsen U, Møller M. Pain medication use in Danish youth handball: a cross-sectional study of 466 youth handball players. In: ; 2020.
13. Alexander CJ. Effect of growth rate on the strength of the growth plate-shaft junction. *Skeletal Radiol*. 1976;1(2):67-76.
14. van der Sluis A, Elferink-Gemser MT, Coelho-e-Silva MJ, Nijboer JA, Brink MS, Visscher C. Sport injuries aligned to peak height velocity in talented pubertal soccer players. *Int J Sports Med*. 2014;35(4):351-355.
15. de Lucena GL, dos Santos Gomes C, Guerra RO. Prevalence and associated factors of Osgood-Schlatter syndrome in a population-based sample of Brazilian adolescents. *Am J Sports Med*. 2011;39(2):415-420.

16. Fuglkjær S, Hartvigsen J, Wedderkopp N, et al. Musculoskeletal extremity pain in Danish
school children – how often and for how long? The CHAMPS study-DK. *BMC Musculoskelet*
Disord. 2017;18.
17. Fuglkjær S, Dissing KB, Hestbæk L. Prevalence and incidence of musculoskeletal extremity
complaints in children and adolescents. A systematic review. *BMC Musculoskeletal Disorders*.
2017;18(1):418.
18. Hawkins D, Metheny J. Overuse injuries in youth sports: biomechanical considerations. *Med*
Sci Sports Exerc. 2001;33(10):1701-1707.
19. Arnold A, Thigpen CA, Beattie PF, Kissenberth MJ, Shanley E. Overuse Physeal Injuries in
Youth Athletes. *Sports Health*. 2017;9(2):139-147.
20. Rathleff MS, Winiarski L, Krommes K, et al. Pain, Sports Participation, and Physical Function
in 10-14 Year Olds With Patellofemoral Pain and Osgood Schlatter: A Matched Cross-
Sectional Study of 252 Adolescents. *J Orthop Sports Phys Ther*. Published online January 6,
2020:1-26.
21. Valasek AE, Young JA, Huang L, Singichetti B, Yang J. Age and Sex Differences in Overuse
Injuries Presenting to Pediatric Sports Medicine Clinics. *Clin Pediatr (Phila)*. 2019;58(7):770-
777.
22. Duong MM, Nicholson AD, Li SQ, Gilmore A, Cooperman DR, Liu RW. Relationship Between
Sever Disease and Skeletal Maturity. *J Pediatr Orthop*. 2020;40(2):93-96.
23. López-Alameda S, Alonso-Benavente A, López-Ruiz de Salazar A, Miragaya-López P, Alonso-
Del Olmo JA, González-Herranz P. [Sinding-Larsen-Johansson disease: analysis of the
associated factors]. *Rev Esp Cir Ortop Traumatol*. 2012;56(5):354-360.
24. Guldhammer C, Rathleff MS, Jensen HP, Holden S. Long-term Prognosis and Impact of
Osgood-Schlatter Disease 4 Years After Diagnosis: A Retrospective Study. *Orthopaedic*
Journal of Sports Medicine. 2019;7(10):2325967119878136.
25. Holden S, Krommes K, Olesen JL, et al. Prognosis of Osgood Schlatter- poorer than
anticipated? A prospective cohort study with 24-month follow-up. *(In Review)*. Published online
2020.
26. Price RJ, Hawkins RD, Hulse MA, Hodson A. The Football Association medical research
programme: an audit of injuries in academy youth football. *Br J Sports Med*. 2004;38(4):466-
471.
27. Junge T, Runge L, Juul-Kristensen B, Wedderkopp N. Risk Factors for Knee Injuries in
Children 8 to 15 Years: The CHAMPS Study DK. *Medicine & Science in Sports & Exercise*.
2016;48(4):655–662.
28. Trifonov Rexen C, Andersen LB, ErsbØll AK, Jespersen E, Franz C, Wedderkopp N. Injuries in
Children with Extra Physical Education in Primary Schools. *Medicine & Science in Sports &*
Exercise. 2014;46(4):745–752.
29. Materne O, Chamari K, Farooq A, et al. Association of Skeletal Maturity and Injury Risk in Elite
Youth Soccer Players: A 4-Season Prospective Study With Survival Analysis. *Orthopaedic*
Journal of Sports Medicine. 2021;9(3):2325967121999113.
30. Ross MD, Villard D. Disability levels of college-aged men with a history of Osgood-Schlatter
disease. *J Strength Cond Res*. 2003;17(4):659-663.
31. von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP. The
Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement:
guidelines for reporting observational studies. *Journal of Clinical Epidemiology*.
2008;61(4):344-349.
32. Chan A-W, Tetzlaff JM, Altman DG, et al. SPIRIT 2013 statement: defining standard protocol
items for clinical trials. *Ann Intern Med*. 2013;158(3):200-207.
33. Danskernes Sundhed - Den Nationale Sundhedsprofil 2017.
34. Ware J, Kosinski M, Keller SD. A 12-Item Short-Form Health Survey: construction of scales
and preliminary tests of reliability and validity. *Med Care*. 1996;34(3):220-233.

35. Danquah I. *Validering Af Spørgsmål Om Fysisk Aktivitet Og Siddetid Ida Høgstedt Danquah,*
Sofie Smedegaard Skov Bo Rytkjær Callesen, Ida Voss Christina Bjørk Petersen, Janne
Schurmann Tolstrup – Til Brug for National Monitorering. Statens Institut for Folkesundhed;
2016.
36. Roos EM, Roos HP, Lohmander LS, Ekdahl C, Beynnon BD. Knee Injury and Osteoarthritis
Outcome Score (KOOS)--development of a self-administered outcome measure. *J Orthop*
Sports Phys Ther. 1998;28(2):88-96.
37. Pedersen ESL, Mortensen LH, Brage S, Bjerregaard AL, Aadahl M. Criterion validity of the
Physical Activity Scale (PAS2) in Danish adults. *Scand J Public Health*. 2018;46(7):726-734.
38. Construct validity of a revised Physical Activity Scale and testing by cognitive interviewing -
Lise G. Andersen, Mogens Groenvold, Torben Jørgensen, Mette Aadahl, 2010.
39. Vaegter HB, Handberg G, Kent P. Brief Psychological Screening Questions Can be Useful for
Ruling Out Psychological Conditions in Patients With Chronic Pain. *Clin J Pain*.
2018;34(2):113-121.
40. Harris PA, Taylor R, Thielke R, Payne J, Gonzalez N, Conde JG. Research Electronic Data
Capture (REDCap) - A metadata-driven methodology and workflow process for providing
translational research informatics support. *J Biomed Inform*. 2009;42(2):377-381.
41. *Statistical Power Analysis for the Behavioral Sciences*. Routledge; 2013.
42. GRADE handbook.
306