

1 Management of lower limb apophysitis: Protocol of a cross-sectional study on
2 long-term impact on health status

3 *(Pre-print – preliminary protocol without peer-review)*

4
5 Kasper Krommes¹, Kristian Thorborg¹, Per Hölmich¹

6
7 Sports Orthopedic Research Center – Copenhagen, Orthopedic Department, Amager-
8 Hvidovre, Copenhagen university hospital

9
10 February 29th, 2020. Version 1.0. Doi: xxx (identifier pending)

11
12 Contact to corresponding author: Email: kasper.krommes@regionh.dk, twitter: @Krommes

13 Registration identifier: ClinicalTrials.gov: xxx (identifier pending)

14 Conflict of interest: The authors have no potential conflicts of interest to declare.

15 Author contributions: KK conceptualized and planned the study, and drafted the protocol.

16 KK, PH and KT revised and approved the protocol.

17
18 **Abstract**

19
20 Lower limb apophysitis cause long-term pain, decrease in function, and can reduce or
21 completely hinder participation in sports and physical activity, yet there is little knowledge
22 on the long-term consequences for health. Our objective with this investigation is to
23 capture self-reported health-status for all adults having been diagnosed with lower limb
24 apophysitis in the period of 1980 to 2020 and compared these data with normative values
25 for the background population. We are therefore conducting a national cross-sectional
26 study based on data from the Danish National Patient Registry (NPR). In this protocol we
27 describe, as detailed as possible, the planned methods.


28

29 Background, rationale, and review of the literature

30

31 Physical activity during the adult life is crucial for health and prevention of disease but has
 32 additional benefits during adolescence (fig. 1). During adolescence the levels of physical
 33 activity declines with only 1 in 6 meeting the recommendations for moderate to vigorous
 34 physical activity or sports participation (fig. 1).

35



36

37

38

39 Given that adolescence is a period of increasing autonomy, positive behaviors established
 40 during this period could potentially last into later adulthood.^{1,2} There are several barriers to
 41 participation in physical activity and sports during adolescence, one such is the risk of
 42 injury and pain during these activities. During adolescence, lower limb pain is the most
 43 frequent cause for seeking primary MSK care³ and up to 50% of sports active adolescents
 44 regularly consume painkillers for injury-related pain.⁴

45

46 Adolescents are specifically susceptible to growth-related overuse injuries in the knee and

47 heel.⁵⁻⁹ For early adolescence, when growth velocity peaks (ages 10-16),¹⁰ the most
48 common growth related injuries is apophysitis in the lower limb; *Severs* in the heel, and
49 *Sinding Larsen Johansson* and *Osgood Schlatter* in the knee.¹¹ All three entities cause
50 long-term pain, decrease in function, and can reduce or completely hinder participation in
51 sports and physical activity.¹²⁻¹⁷ The most common, *Osgood Schlatter*, affects 1 in 5
52 active adolescents.⁷ Yet there is little knowledge on the long-term consequences for
53 health, with only a small case-series on younger participants¹⁶ and one in college-aged
54 students.¹⁸

55

56 Objective

57 To review the long-term consequences of having had a lower limb apophysitis in a large
58 sample, we aim to capture self-reported health-status for all adults having been diagnosed
59 with such in the period of 1980 to 2020.

60

61 Research question

62 What is the health status of adults with a history of lower limb apophysitis?

63

64

65 Methods

66

67 Design

68 We wish to conduct a national cross-sectional study based on data from the Danish
69 National Patient Registry (NPR). The study is reported to the regional research ethics
70 committee (Region Hovedstaden) but has not yet received approval (identifier pending,
71 submitted February 29th, 2020). In addition, it's been reported to the local data authority
72 (Region Hovedstadens Videnscenter for Dataanmeldelser), but has not yet received
73 approval (identifier pending, submitted February 29th, 2020). The study is being registered
74 at ClinicalTrials.gov (identifier pending, submitted February 29th, 2020), and the protocol is
75 published here at medRxiv (submitted March 1st, 2020). Participants will be thoroughly
76 informed about possible risks, study-participant rights, and aspects of data-sharing and
77 data-protection, and asked to consent in the first page of the survey. If patients wish to
78 redraw their answers to the survey, they can do so. A contact email for the project

79 coordinator (KK) is provided in the invitation message. The study protocol is based on
80 relevant items from the STROBE checklist¹⁹ for reporting of cross-sectional studies and
81 the SPIRIT checklist for trial protocols.²⁰ The study is descriptive and exploratory, and no
82 pre-determined hypothesis are being tested, rather it aims to generate future hypothesis to
83 be tested in longitudinal designs.

84

85 Recruitment and eligibility

86 All patients aged 18-50 having received diagnostics codes pertaining to lower limb
87 apophysitis (Severs, Sinding-Larsen Johansson, Osgood Schlatter) in the years 1980-
88 2020 in Denmark will be eligible to participate. Diseased participants will be censored. All
89 eligible patients will be contacted and invited to participate by filling out the electrical
90 survey online. Sample size will be of convenience with no pre-determined cut-off for
91 smallest or largest sample of interest, rather, the final sample of respondents will be based
92 on however many have been designated the diagnostic codes and responds to the survey.

93

94 Outcomes

95

96 Primary outcome: Physical and mental health

97 Measured on the SF-12 (Short-form 12 item) health survey.²¹

98

99 Other diagnostic codes

100 To describe this population and their rate of other diseases, all other designated diagnostic
101 codes from the NPR will be captured and reported.

102

103 Sleep problems, mental health, BMI and self-efficacy

104 These factors are related to prognosis or severity for many other well-researched
105 musculoskeletal conditions. We will utilize validated questions²² and questionnaires
106 (PSEQ-2).^{23(p2)}

107

108 Knee-related pain and symptoms, and physical activity

109 We will ask to current extent, satisfaction, and level of physical activity and sport (Tegner
110 scale).²⁴ In addition, we will ask to knee-related pain and symptoms (KOOS)²⁵ and other
111 types of pain.

112

113 Apophysitis-related history

114 We will ask questions related to duration, severity and treatment of their apophysitis.

115

116 Data collection and sharing

117 All data will be directly entered into REDCap (Research Electronic Data Capture),²⁶ a
118 logged secure system designed for capturing sensitive non-commercial research data that
119 is hosted at Hvidovre Hospital. REDCap contains options for valid values, range checks,
120 data validation, branching, scheduling, and stop-rules to increase data quality.

121

122 Invitations to the questionnaire will be posted to participants digital government-issued
123 inbox (e-boks) used for confidential correspondence (e.g. doctors' appointments), using
124 their social security number (CPR number) obtained from the NPR. We expect the mean
125 duration of a reply will be in the range of 8-10 minutes. If participants have questions or
126 comments, there will be a contact-email posted in the survey-invitation.

127

128 The full dataset, excluding all personal identifiers, will be posted alongside the pre-print
129 when the results have been disseminated.

130

131 Analyses

132 The flow of participants from the potential total pool, to invitations, final respondents, and
133 missing data, will be described in detail. As this is a single-group study with no pre-
134 determined hypothesis, no blinding will be performed. Analyses will be descriptive and
135 reported in means and 95% confidence intervals, and if non-normally distributed, reported
136 as medians and inter-quartile ranges. We will compare sub-groups of interest observing
137 overlaps of confidence intervals. The *a priori* subgroups of interest are:

- 138 1. Patients reporting short (<6 months) vs. long (>1 year) apophysitis symptom duration.
- 139 2. Patients reporting significant limitation to sport and physical activity during their
140 apophysitis vs. those that was not significantly affected.
- 141 3. Patients that currently have knee pain or symptoms from the same general area vs.
142 those that currently does not.

- 143 4. Patients who continued moderate or high levels of physical activity after cessation of
144 apophysitis symptoms vs. does that have been less than moderate physically active
145 since cessation of their apophysitis symptoms.
- 146 5. Patients that report currently having a large bony prominence thought to originate
147 from their apophysite (Only Osgood Schlatter patients) vs. those that does not
- 148 6. Patients that report severe symptoms during their apophysitis vs. does that only
149 report have experienced light or moderate symptoms (based on pain intensity and
150 frequency of pain-flares, and restriction in physical activity and sport).
- 151 7. Patients that report having undergone a period of rapid growth during their
152 apophysitis vs. does that does not

153

154 If relevant sub-groups should emerge *post hoc*, they will be denoted as such. We will also
155 compare data from this sample with normative values for the background population.

156

157

158 References

159

- 160 1. Nelson MC, Story M, Larson NI, Neumark-Sztainer D, Lytle LA. Emerging adulthood and college-aged
161 youth: an overlooked age for weight-related behavior change. *Obesity (Silver Spring)*. 2008;16(10):2205-
162 2211.
- 163 2. Rindfuss RR. The young adult years: diversity, structural change, and fertility. *Demography*.
164 1991;28(4):493-512.
- 165 3. Prathivadi Bhayankaram N, Lacey RJ, Barnett LA, Jordan KP, Dunn KM. Musculoskeletal consultations
166 from childhood to adulthood: a longitudinal study. *J Public Health (Oxf)*.
- 167 4. Sari D, Thorlund JB, Mikkelsen U, Møller M. Pain medication use in Danish youth handball: a cross-
168 sectional study of 466 youth handball players. In: Copenhagen; 2020.
- 169 5. Alexander CJ. Effect of growth rate on the strength of the growth plate-shaft junction. *Skeletal Radiol*.
170 1976;1(2):67-76.
- 171 6. van der Sluis A, Elferink-Gemser MT, Coelho-e-Silva MJ, Nijboer JA, Brink MS, Visscher C. Sport injuries
172 aligned to peak height velocity in talented pubertal soccer players. *Int J Sports Med*. 2014;35(4):351-355.
- 173 7. de Lucena GL, dos Santos Gomes C, Guerra RO. Prevalence and associated factors of Osgood-Schlatter
174 syndrome in a population-based sample of Brazilian adolescents. *Am J Sports Med*. 2011;39(2):415-420.
- 175 8. Fuglkjær S, Hartvigsen J, Wedderkopp N, et al. Musculoskeletal extremity pain in Danish school children –
176 how often and for how long? The CHAMPS study-DK. *BMC Musculoskelet Disord*. 2017;18.
- 177 9. Fuglkjær S, Dissing KB, Hestbæk L. Prevalence and incidence of musculoskeletal extremity complaints in
178 children and adolescents. A systematic review. *BMC Musculoskeletal Disorders*. 2017;18(1):418.
- 179 10. Hawkins D, Metheny J. Overuse injuries in youth sports: biomechanical considerations. *Med Sci Sports*
180 *Exerc*. 2001;33(10):1701-1707.

- 181 11. Arnold A, Thigpen CA, Beattie PF, Kissenberth MJ, Shanley E. Overuse Physeal Injuries in Youth Athletes.
182 *Sports Health*. 2017;9(2):139-147.
- 183 12. Rathleff MS, Winiarski L, Krommes K, et al. Pain, Sports Participation, and Physical Function in 10-14 Year
184 Olds With Patellofemoral Pain and Osgood Schlatter: A Matched Cross-Sectional Study of 252
185 Adolescents. *J Orthop Sports Phys Ther*. January 2020:1-26.
- 186 13. Valasek AE, Young JA, Huang L, Singichetti B, Yang J. Age and Sex Differences in Overuse Injuries
187 Presenting to Pediatric Sports Medicine Clinics. *Clin Pediatr (Phila)*. 2019;58(7):770-777.
- 188 14. Duong MM, Nicholson AD, Li SQ, Gilmore A, Cooperman DR, Liu RW. Relationship Between Sever
189 Disease and Skeletal Maturity. *J Pediatr Orthop*. 2020;40(2):93-96.
- 190 15. López-Alameda S, Alonso-Benavente A, López-Ruiz de Salazar A, Miragaya-López P, Alonso-Del Olmo
191 JA, González-Herranz P. [Sinding-Larsen-Johansson disease: analysis of the associated factors]. *Rev Esp*
192 *Cir Ortop Traumatol*. 2012;56(5):354-360.
- 193 16. Gulddammer C, Rathleff MS, Jensen HP, Holden S. Long-term Prognosis and Impact of Osgood-Schlatter
194 Disease 4 Years After Diagnosis: A Retrospective Study. *Orthopaedic Journal of Sports Medicine*.
195 2019;7(10):2325967119878136.
- 196 17. Holden S, Krommes K, Olesen JL, et al. Prognosis of Osgood Schlatter- poorer than anticipated? A
197 prospective cohort study with 24-month follow-up. (*In Review*).
- 198 18. Ross MD, Villard D. Disability levels of college-aged men with a history of Osgood-Schlatter disease. *J*
199 *Strength Cond Res*. 2003;17(4):659-663.
- 200 19. von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP. The Strengthening the
201 Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting
202 observational studies. *Journal of Clinical Epidemiology*. 2008;61(4):344-349.
- 203 20. Chan A-W, Tetzlaff JM, Altman DG, et al. SPIRIT 2013 statement: defining standard protocol items for
204 clinical trials. *Ann Intern Med*. 2013;158(3):200-207.
- 205 21. Ware J, Kosinski M, Keller SD. A 12-Item Short-Form Health Survey: construction of scales and preliminary
206 tests of reliability and validity. *Med Care*. 1996;34(3):220-233.
- 207 22. Vaegter HB, Handberg G, Kent P. Brief Psychological Screening Questions Can be Useful for Ruling Out
208 Psychological Conditions in Patients With Chronic Pain. *Clin J Pain*. 2018;34(2):113-121.
- 209 23. Nicholas MK, McGuire BE, Asghari A. A 2-Item Short Form of the Pain Self-Efficacy Questionnaire:
210 Development and Psychometric Evaluation of PSEQ-2. *The Journal of Pain*. 2015;16(2):153-163.
- 211 24. Briggs KK, Steadman JR, Hay CJ, Hines SL. Lysholm score and Tegner activity level in individuals with
212 normal knees. *Am J Sports Med*. 2009;37(5):898-901.
- 213 25. Roos EM, Roos HP, Lohmander LS, Ekdahl C, Beynon BD. Knee Injury and Osteoarthritis Outcome Score
214 (KOOS)--development of a self-administered outcome measure. *J Orthop Sports Phys Ther*.
215 1998;28(2):88-96.
- 216 26. Harris PA, Taylor R, Thielke R, Payne J, Gonzalez N, Conde JG. Research Electronic Data Capture
217 (REDCap) - A metadata-driven methodology and workflow process for providing translational research
218 informatics support. *J Biomed Inform*. 2009;42(2):377-381.