TITLE
Gaps in Sexually Transmitted Infection Screening among
Men who Have Sex with Men in PrEP Care in the United States

SHORT TITLE
Gaps in STI Screening among MSM in PrEP

AUTHORS
Christina Chandra,[1] Kevin M. Weiss,[1] Colleen F. Kelley,[2] Julia L. Marcus,[3] Samuel M. Jenness[1]

[1] Department of Epidemiology, Rollins School of Public Health, Emory University, Atlanta, GA

[2] Department of Medicine, Division of Infectious Diseases, Emory University School of Medicine, Atlanta, GA

[3] Harvard Medical School and Harvard Pilgrim Health Care Institute, Boston, MA

CORRESPONDENCE
Samuel M. Jenness, Emory University, 1520 Clifton Road, Atlanta, GA 30323. Email:
samuel.m.jenness@emory.edu. Alternate: Christina Chandra (christina.lynne.chandra@emory.edu)

FUNDING
This work was supported by the National Institutes of Health [grant number R21 MH112449].

CONFLICTS OF INTEREST
The authors declare no conflicts of interest.

WORD COUNTS
Abstract: 241 [250 max]
Text: 2975 [3000 max]
References: 40 [40 max]

KEYWORDS
HIV; preexposure prophylaxis; sexually transmitted infections; screening; men who have sex with men

SUMMARY
Nearly half of U.S. men who have sex with men in PrEP care are not receiving consistent bacterial STI screening at sites of sexual exposure, and levels are worse in the Southeast region where the burden of STI is highest.

ABSTRACT

Background   The U.S. Centers for Disease Control and Prevention (CDC) recommends comprehensive sexually transmitted infection (STI) screening every 3–6 months for men who have sex with men (MSM) using HIV preexposure prophylaxis (PrEP). The gaps between these recommendations and clinical practice by region have not been quantified.

Methods   We collected data between 2017 and 2019 on STI screening among MSM across the U.S., stratified by current, prior, and never PrEP use. Poisson regression models with robust error variance were used to model factors, including residence in the Southeast states, associated with consistent exposure site-specific STI screening during PrEP care.

Results   Of 3259 HIV-negative MSM, 19% were currently using PrEP, 6% had used PrEP in the past, and 75% had never used PrEP. Among ever PrEP users, 87%, 78%, 57%, and 64% reported consistent screening for STIs by blood sample, urine sample or urethral swab, rectal swab, or pharyngeal swab, respectively, during PrEP care. Compared to PrEP users in all other regions, PrEP users in the Southeast were significantly less likely to be consistently screened for urogenital [adjusted prevalence ratio (aPR): 0.86, 95% confidence interval (CI): 0.76, 0.98] and rectal STIs (aPR: 0.76, 95% CI: 0.62, 0.93) during PrEP care.

Conclusions   Substantial gaps exist between CDC recommendations for STI screening during PrEP care and current clinical practice, particularly for rectal and pharyngeal exposure sites that can harbor asymptomatic infections and for MSM in Southeast states where the STI burden is substantial.

INTRODUCTION

In the United States (U.S.), cases of the three major reportable bacterial sexually transmitted infections (STIs) — syphilis, gonorrhea, and chlamydia — increased by up to 71% between 2014 and 2018, reaching all-time highs [1]. Rates of new HIV diagnoses have remained stable overall and increased in key subgroups during this time [2]. Men who have sex with men (MSM) are disproportionately affected by both bacterial STIs and HIV, accounting for more than half of new infectious syphilis cases, 40% of gonorrhea cases, and 66% of HIV cases in 2018 despite only representing 4% of the population [1–3].

One hypothesis for the increase in STI cases among MSM has been the growth in use of HIV preexposure prophylaxis (PrEP). In a recent meta-analysis, the pooled STI prevalence among PrEP users within three months of PrEP initiation was 24%, and STI incidence was estimated at 72 per 100 person-years among persistent PrEP users globally [4]. U.S. studies have found that STI prevalence or incidence increased among MSM after initiation of PrEP [5,6]. The difference in STI rates before and after PrEP initiation could be a result of multiple factors, including increased detection of cases from more frequent screening in PrEP care, changes in sexual risk behavior after initiation of PrEP (sometimes called "risk compensation"), and trends in sexual behavior and STI rates that pre-date PrEP [1,7–10]. Many PrEP users are at elevated risk of STIs simply because PrEP is recommended for MSM with a history of inconsistent condom use and multiple partners [11].

The U.S. Centers for Disease Control and Prevention (CDC) recommends comprehensive STI screening every 3–6 months for MSM using PrEP [11]. This includes blood tests to screen for syphilis, and urine samples and pharyngeal and rectal swabs for gonorrhea and chlamydia infections. Prior modeling research suggested that while STI diagnoses may initially increase with expanded PrEP use due to variations of risk compensation, routine screening and treatment of STIs during PrEP care could prevent over 40% of gonorrhea and chlamydia cases among MSM over ten years [12]. These projections assumed complete adherence to PrEP-related STI screening and treatment recommendations; secondary model scenarios projected that STI incidence could actually increase if fewer than 50% of PrEP users were consistently screened and treated for STIs.

Quantifying any gaps between PrEP STI screening recommendations and clinical practice is therefore critical to understanding the role of PrEP care in STI control. Several recent studies have evaluated this in different regions, generally finding much lower rates of STI screening than recommended [7,13–16]. While these studies have provided important evidence regarding the gaps in PrEP care, there are still no national-level studies of MSM that have compared these gaps across geographic regions. Geographic heterogeneity in clinical practice is important to understand because of

the regional differences in infrastructure for PrEP services (e.g., the ability to assess kidney function and serological confirmation of Hepatitis B status for PrEP eligibility) and the spatial clustering of STIs in the Southeast region [17,18]. Differences in STI screening practices may also reflect state-level factors that may impact PrEP care, including access to health care (e.g., Medicaid expansion) and other social determinants of health that may vary by state (e.g., housing stability, employment) [19,20]. By examining these differences, we may better understand which regions might benefit the most from interventions to improve comprehensive PrEP care that includes recommended STI screening.

In this study, we compared STI screening among MSM by PrEP use history in a national study of MSM in the U.S. Among PrEP users, we then evaluated factors associated with consistent STI screening at PrEP care visits, including residence in the Southeast and recent sexual exposure at anatomical sites. We hypothesized that there would be strong geographic differences in STI screening consistency among PrEP users even after controlling for demographic and behavioral factors.

METHODS

Study Design. ARTnet was a cross-sectional web-based study of MSM in the U.S. conducted between 2017 and 2019 [21]. The study recruited participants through a parent national American Men's Internet Survey (AMIS) [22]. ARTnet eligibility criteria included male sex at birth, current male identity, lifetime history of sexual activity with another man, and age between 15 and 65 years. The study surveyed participants about recent sexual behaviors, HIV and STI screening, and egocentric network data [23]. We restricted the participants in this analysis to PrEP-eligible MSM, defined as MSM who had ever had an HIV test and who self-reported as HIV-negative. The Emory University Institutional Review Board reviewed and approved the study.

Measures. We quantified participant-reported data on HIV and STI screening from blood samples, urine sample/urethral swabs, and rectal and pharyngeal swabs in the past 12 months overall and at PrEP care visits. Participants who had ever used PrEP were asked: "At these PrEP (i.e., Truvada) check-up visits, how often did you get tested for HIV and sexually transmitted diseases (STD)? STDs could include gonorrhea, chlamydia, and syphilis." Participants could answer "Always", "Sometimes", "Rarely", or "Never" for HIV screening or STI screening site/method: throat (pharyngeal swab), anus (rectal swab), genital/penis (urine sample or urethral swab), or blood sample. We assumed blood-based screening was used to screen for syphilis. For statistical models, we dichotomized the outcomes to consistent screening ("Always" or "Sometimes") versus inconsistent screening ("Rarely" or "Never").

We then investigated correlates of consistent STI screening. We considered age group, race/ethnicity, current health insurance, annual household income, recent sexual exposure at anatomical site, and geographic region of residence. Using data on the five most recent reported partners in the past year, respondents were classified as having been exposed at the urethra (any insertive anal or insertive oral intercourse), rectum (any receptive anal intercourse), or pharynx (any receptive oral intercourse). For geography, we considered Census division and state by matching reported residential ZIP Codes against Census data [21]. Our primary exposure of interest was residence in 12 Southeast states as defined by the U.S. Bureau of Economic Analysis: Alabama, Arkansas, Florida, Georgia, Kentucky, Louisiana, Mississippi, North Carolina, South Carolina, Tennessee, Virginia, and West Virginia [24].

Statistical Analysis. Our primary research objective was to quantify the association between residence in Southeast states and consistent screening for STIs among MSM who were currently using or had previously used PrEP. We first estimated the bivariable prevalence ratios between our hypothesized correlates — age, race/ethnicity, current health insurance, annual household income, recent sexual exposure at anatomical site, and geography — and consistent site-specific STI screening using Poisson regression with robust error variance. In multivariable Poisson regression models, we estimated the prevalence ratios between our primary exposure (residence in a Southeast state) and consistent STI screening adjusted for four variables that we hypothesized would confound this relationship: age, race/ethnicity, annual household income, and sexual exposure at anatomical site. We did not control for variables that we considered mediators of the associations, such as health insurance that may vary by state as a result of Medicaid expansion or other state-level factors like HIV prevention funding that may impact STI screening for PrEP users. All analyses were conducted in R 3.6.1 [25]. Analysis scripts are provided in a GitHub repository (https://github.com/EpiModel/PrEP-STI-Test) and the primary data are available upon request.

RESULTS

After excluding PrEP-ineligible participants, 3259 MSM were included in this analysis. Of those, 631 (19%) were currently using PrEP at the time of the study, 178 (6%) had previously used PrEP, and 2450 (75%) had never used PrEP. **Table 1** presents the demographics and HIV/STI screening history by PrEP use history. Overall, most participants were non-Hispanic white (74%) and more than half were between 15 and 34 years old (51%). Overall, 23% of the participants lived in Southeast states. PrEP users were generally underrepresented in the Southeast, with 25% of MSM who had never used PrEP residing in the Southeast compared to 19% of current PrEP users and prior PrEP users. Four in five (82%) current PrEP

users reported having any private health insurance compared with 72% of prior PrEP users and 75% of never-PrEP users. Only 3% of current PrEP users were uninsured compared with 10% of prior PrEP users and 9% of those who had never used PrEP. A higher proportion of current PrEP users reported screening for a bacterial STI in the past year (89%), compared to prior PrEP users (71%) and participants who had never used PrEP (44%). Similarly, 98% of current PrEP users were screened for HIV in the past year compared to 80% of prior PrEP users and 65% of those who had never used PrEP.

Among MSM who had ever used PrEP, 84% of current users reported attending PrEP care visits at least every three months compared to 71% of prior PrEP users (**Table 2**). Overall, 91% of MSM who had ever used PrEP were always screened for HIV at their PrEP care visits. Greater proportions of current PrEP users, compared to prior PrEP users, reported always being screened for STIs by throat swab (44% versus 35%), rectal swab (37% versus 32%), blood (for syphilis) (70% versus 57%), and urine sample/urethral swab (59% versus 50%) at these visits. In contrast, among all MSM who had ever used PrEP, 35% had never provided a rectal swab, 28% had never provided a throat swab, and 17% had never provided a urine sample or urethral swab for STI screening during PrEP care visits.

A higher percentage of MSM who had ever used PrEP reported consistent screening for syphilis (87%) and screening for urogenital STIs (78%) (**Table 3**) compared to pharyngeal STI screening (64%) and rectal STI screening (57%) (**Table 4**). Older MSM had the lowest levels of consistent screening for urogenital STIs (80% versus 71%), pharyngeal STIs (70% versus 60%), and rectal STIs (67% versus 53%) (**Tables 3 and 4**). Reporting of consistent STI screening ranged widely by geography. Consistent screening was less common in Southeast states for urogenital STI screening (66% versus 80% in all other states), pharyngeal STI screening (55% versus 66%), and rectal STI screening (44% versus 60%). Exposure at the anatomical site was associated with consistent screening for urogenital STIs (78% versus 68% among those unexposed), pharyngeal STIs (64% versus 63%), rectal STIs (60% versus 44%).

Among MSM who had ever used PrEP, age, race/ethnicity, health insurance, annual household income, Census division, and Southeast states were not significantly associated with syphilis screening at PrEP care visits in bivariable Poisson regression analysis (**Table 3**). For urogenital, rectal, and pharyngeal STIs, however, there was a general trend of younger MSM reporting consistent screening compared to older MSM at PrEP care visits (**Tables 3 and 4**). In these unadjusted analyses, MSM in Southeast states were less likely to report consistent urogenital STI screening [prevalence ratio (PR) = 0.83; 95% confidence interval (CI) = 0.73, 0.94], rectal STI screening (PR = 0.73; 95% CI = 0.59, 0.89), and pharyngeal STI screening (PR = 0.83; 95% CI = 0.71, 0.97) during PrEP care (**Table 4**). MSM who

reported exposure at the rectal site were 35% more likely to report consistent rectal STI screening at PrEP care visits (PR = 1.35; 95% CI = 1.11, 1.65). There was no bivariable association between sexual exposure at anatomical site and consistent STI screening for urogenital and pharyngeal STIs.

In adjusted models, MSM in Southeast states were less likely to report consistent STI screening compared to MSM in other regions (**Table 5**). MSM in the Southeast had 14% lower prevalence of consistent urogenital STI screening [adjusted prevalence ratio (aPR) = 0.86; 95% CI = 0.76, 0.98] and 24% lower prevalence of consistent rectal STI screening (aPR = 0.76; 95% CI = 0.62, 0.93) at PrEP care. MSM residing in the Southeast were also less likely to receive consistent pharyngeal STI screening (aPR = 0.87; 95% CI = 0.74, 1.03). There was no association between Southeast residence and consistent syphilis screening (aPR = 1.02; 95% CI = 0.95, 1.10).

DISCUSSION

In this national study of MSM in the United States, consistency of STI screening at PrEP care visits was lower than recommended, especially for rectal and pharyngeal infections that are mostly asymptomatic. Our findings also highlight the regional variation in gaps between recommendations and PrEP clinical practice overall, and raise concerns about whether comprehensive PrEP care as currently practiced would be effective for STI control, particularly in the Southeast where the burden of bacterial STIs is highest [1,18].

This study contributes to the growing literature on patterns of STI screening within PrEP care. With the exception of one national study [14], research has been limited to individual clinics or health systems, often with limited sample sizes [7,13,15,16]. These studies analyze a variety of data streams, including electronic health records, health insurance claims, or laboratory screening [15,16,26]. However, many different measures have been used as primary outcomes: screening at last PrEP care visit [13]; STI or rectal STI screening in the prior 12 months [7]; or proportion screened for STIs within 6 months of PrEP initiation [14]. Our study, in contrast, measured STI screening *across* PrEP care visits that may better reflect ongoing consistency of clinical practice.

Despite these different study designs and outcomes, the research base has now consistently shown suboptimal STI screening among PrEP users, especially for extragenital screening [13–16]. Similar to a study conducted in Baltimore, we found that among all site-specific screening, rectal swabs were reported least often [16]. A study in New York City, in contrast, found that pharyngeal STI screening was the least reported (at the last PrEP care visit) [13]. These results are also consistent with findings of infrequent extragenital (rectal of pharyngeal) STI screening among MSM *outside* of PrEP care [27–29]. Infrequent

extragenital screening is a public health concern because of the higher prevalence of asymptomatic chlamydia or gonorrhea infections that may remain undetected [30]. Asymptomatic STIs at these sites also lead to increased risk of HIV infection among partners in the sexual network not using PrEP [23,31].

Several factors have been previously found to be associated with more recent or comprehensive STI screening: younger age, white race, college education, reporting exposure at anatomical sites that may trigger screening, and previous syphilis diagnoses [7,13,15,16]. Our study additionally found that residence in the Southeast was associated with inconsistent screening for urogenital, rectal, and pharyngeal STIs, even after adjusting for potential confounders. These results have implications for regional improvements of comprehensive health services in PrEP care. A recent study found that 96% of current or potential PrEP providers were somewhat or completely comfortable with screening for STIs, suggesting that providers are well-equipped to provide comprehensive STI care and that other barriers exist to STI screening and treatment [32]. However, the cost of ancillary services associated with PrEP care such as STI screening may be a barrier, as providers have noted that high cost of medication and health insurance coverage may hinder the real-world effectiveness of PrEP [33]. Moreover, 7 of the 12 Southeast states (in our definition) have not expanded Medicaid under the Affordable Care Act, potentially limiting the ability of MSM to obtain adequate coverage for consistent STI screening [34]. Gaps in comprehensive PrEP care may exacerbate the large racial and socioeconomic disparities in HIV and STI incidence in the region [35]. Wider implementation of alternative STI screening approaches, such as self-collected rectal swabs or home screening kits, may also improve STI screening coverage for MSM using PrEP [36–38].

Limitations. This study has several limitations. First, this web-based sample does not necessarily represent the larger population of MSM with respect to age, race/ethnicity, health insurance, annual household income, and geographic residential location. Most participants were white, had higher income levels, had some form of private health insurance, and lived in the mid-Atlantic/Northeast or the West Coast. Compared to venue-based sampling, web-based sampling of MSM may yield a sample with more non-Hispanic white men and those in higher income groups [39]. Second, stratifying participant data by the covariates of interest yielded small sample sizes in some categories (reflected in wide confidence intervals for some estimates). Third, we relied on self-reported data on the exposures and outcomes, and there may be misclassification of screening frequency due to not understanding what specific STIs they are being tested. Fourth, the data were also limited in classifying anatomical site exposure since the ARTnet study collected data on a maximum of five partners; this may not align with their PrEP visits. Fifth, we assumed that prior and current PrEP users accessed PrEP in their reported residential ZIP code, but it

is possible that prior PrEP users accessed PrEP in a state where they previously lived or that current PrEP users access PrEP in a neighboring state. Sixth, we may be overestimating consistency of syphilis screening, because participants may misinterpret blood draws for HIV screening or other tests at PrEP care visits as inclusive of STI screening. Finally, the STI screening frequency measure (always, sometimes, rarely, or never) allows for an approximate but not perfect estimate of adherence to CDC recommendations for PrEP care, particularly because "sometimes" can represent a wide range of screening frequencies as interpreted by respondents. For this reason, we used the term "consistent" rather than "as recommended."

Conclusions. Data from this large national study of MSM improved the understanding of geographic differences in comprehensive health services for MSM using PrEP. Although a higher proportion of PrEP users reported recent STI screening compared to non-PrEP users, barriers exist to complete adherence to CDC PrEP guidelines of comprehensive screening for STIs at all exposure sites among MSM. This was especially true in Southeast states where the STI burden is high and where the federal government has prioritized funding HIV prevention programs through the Ending the HIV Epidemic initiative [40]. As PrEP is scaled up in the Southeast through this national initiative, resources to support STI screening and treatment as part of PrEP care should simultaneously be strengthened. Otherwise, suboptimal STI screening in PrEP care may lead to increases overall STI incidence among MSM.

REFERENCES

1. Centers for Disease Control and Prevention. Sexually Transmitted Disease Surveillance 2018. US Dep Health Hum Serv **2019**; Available at: https://www.cdc.gov/std/stats.

2. Centers for Disease Control and Prevention. HIV Surveillance Report, 2018 (Preliminary). Centers for Disease Control and Prevention, 2019. Available at: http://www.cdc.gov/hiv/library/reports/hiv-surveillance.html.

3. Purcell DW, Johnson CH, Lansky A, et al. Estimating the Population Size of Men Who Have Sex with Men in the United States to Obtain HIV and Syphilis Rates. Open AIDS J **2012**; 6:98–107.

4. Ong JJ, Baggaley RC, Wi TE, et al. Global Epidemiologic Characteristics of Sexually Transmitted Infections Among Individuals Using Preexposure Prophylaxis for the Prevention of HIV Infection: A Systematic Review and Meta-analysis. JAMA Netw Open **2019**; 2:e1917134–e1917134.

5. MONTAÑO MA, DOMBROWSKI JC, DASGUPTA S, et al. Changes in Sexual Behavior and STI Diagnoses among MSM Initiating PrEP in a Clinic Setting. AIDS Behav **2019**; 23:548–555.

6. Marcus JL, Hurley LB, Hare CB, et al. Preexposure Prophylaxis for HIV Prevention in a Large Integrated Health Care System: Adherence, Renal Safety, and Discontinuation. J Acquir Immune Defic Syndr 1999 **2016**; 73:540–546.

7. Menza TW, Lipira L, Bhattarai A, Ramirez FJ, Orellana ER. Self-Reported Screening for Rectal Sexually Transmitted Infections Among Men Who Have Sex With Men: Sex Transm Dis **2019**; 46:683–688.

8. Volk JE, Marcus JL, Phengrasamy T, et al. No New HIV Infections With Increasing Use of HIV Preexposure Prophylaxis in a Clinical Practice Setting. Clin Infect Dis Off Publ Infect Dis Soc Am **2015**; 61:1601–1603.

9. Paz-Bailey G, Mendoza MCB, Finlayson T, et al. Trends in condom use among MSM in the United States: the role of antiretroviral therapy and seroadaptive strategies. AIDS Lond Engl **2016**; 30:1985–1990.

10. Harawa NT, Holloway IW, Leibowitz A, et al. Serious concerns regarding a meta-analysis of preexposure prophylaxis use and STI acquisition: AIDS **2017**; 31:739–740.

11. Centers for Disease Control and Prevention. Preexposure Prophylaxis for the Prevention of HIV Infection in the United States – 2017 Update: a Clinical Practice Guideline. 2018. Available at: https://www.cdc.gov/hiv/pdf/risk/prep/cdc-hiv-prep-guidelines-2017.pdf.

12. Jenness SM, Weiss KM, Goodreau SM, et al. Incidence of Gonorrhea and Chlamydia Following Human Immunodeficiency Virus Preexposure Prophylaxis Among Men Who Have Sex With Men: A Modeling Study. Clin Infect Dis **2017**; 65:712–718.

13. Parsons JT, John SA, Whitfield THF, Cienfuegos-Szalay J, Grov C. Human Immunodeficiency Virus/Sexually Transmitted Infection Counseling and Testing Services Received by Gay and Bisexual Men Using Preexposure Prophylaxis at Their Last PrEP Care Visit. Sex Transm Dis **2018**; 45:798–802.

14. Huang YA, Tao G, Samandari T, Hoover KW. Laboratory Testing of a Cohort of Commercially Insured Users of HIV Preexposure Prophylaxis in the United States, 2011–2015. J Infect Dis **2018**; 217:617–621.

15. Spinelli MA, Scott HM, Vittinghoff E, et al. Provider Adherence to Pre-exposure Prophylaxis Monitoring Guidelines in a Large Primary Care Network. Open Forum Infect Dis **2018**; 5.

16. Schumacher C, Wu L, Chandran A, et al. STI Screening among Gay, Bisexual and Other Men who Have Sex with Men Prescribed PrEP in Baltimore City, Maryland. Clin Infect Dis **2019**;

17. Sullivan PS, Mena L, Elopre L, Siegler AJ. Implementation Strategies to Increase PrEP Uptake in the South. Curr HIV/AIDS Rep **2019**; 16:259–269.

18. Marotta P. Assessing Spatial Relationships between Race, Inequality, Crime, and Gonorrhea and Chlamydia in the United States. J Urban Health Bull N Y Acad Med **2017**; 94:683–698.

19. Patel RR, Mena L, Nunn A, et al. Impact of insurance coverage on utilization of pre-exposure prophylaxis for HIV prevention. PLoS ONE **2017**; 12. Available at: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5448799/. Accessed 4 March 2020.

20. Philbin MM, Parker CM, Parker RG, Wilson PA, Garcia J, Hirsch JS. The Promise of Pre-Exposure Prophylaxis for Black Men Who Have Sex with Men: An Ecological Approach to Attitudes, Beliefs, and Barriers. AIDS Patient Care STDs **2016**; 30:282–290.

21. Weiss KM, Goodreau SM, Morris M, et al. Egocentric sexual networks of men who have sex with men in the United States: Results from the ARTnet study. Epidemics **2020**; 30:100386.

22. Zlotorzynska M, Sullivan P, Sanchez T. The Annual American Men's Internet Survey of Behaviors of Men Who Have Sex With Men in the United States: 2016 Key Indicators Report. JMIR Public Health Surveill **2019**; 5:e11313.

23. Jenness SM, Weiss KM, Prasad P, Zlotorzynska M, Sanchez T. Bacterial Sexually Transmitted Infection Screening Rates by Symptomatic Status Among Men Who Have Sex With Men in the United States: A Hierarchical Bayesian Analysis. Sex Transm Dis **2019**; 46:25–30.

24. BEA Regions. Available at: https://apps.bea.gov/regional/docs/regions.cfm.

25. R Core Team. R: A Language and Environment for Statistical Computing. Vienna, Austria: R Foundation for Statistical Computing, 2019. Available at: https://www.R-project.org/.

26. Hevey MA, Walsh JL, Petroll AE. PrEP Continuation, HIV and STI Testing Rates, and Delivery of Preventive Care in a Clinic-Based Cohort. AIDS Educ Prev Off Publ Int Soc AIDS Educ **2018**; 30:393–405.

27. de Voux A, Bernstein KT, Kirkcaldy RD, Zlotorzynska M, Sanchez T. Self-Reported Extragenital Chlamydia and Gonorrhea Testing in the Past 12 Months Among Men Who Have Sex with Men in the United States-American Men's Internet Survey, 2017. Sex Transm Dis **2019**; 46:563–570.

28. Barbee L, Dhanireddy S, Tat S, Marrazzo J. Barriers to Bacterial Sexually Transmitted Infection Testing of HIV-Infected Men Who Have Sex With Men Engaged in HIV Primary Care. Sex Transm Dis **2015**; 42:590–594.

29. Patton ME, Kidd S, Llata E, et al. Extragenital gonorrhea and chlamydia testing and infection among men who have sex with men–STD Surveillance Network, United States, 2010-2012. Clin Infect Dis **2014**; 58:1564–70.

30. Kent CK, Chaw JK, Wong W, et al. Prevalence of rectal, urethral, and pharyngeal chlamydia and gonorrhea detected in 2 clinical settings among men who have sex with men: San Francisco, California, 2003. Clin Infect Dis Off Publ Infect Dis Soc Am **2005**; 41:67–74.

31. Bernstein KT, Marcus JL, Nieri G, Philip SS, Klausner JD. Rectal gonorrhea and chlamydia reinfection is associated with increased risk of HIV seroconversion. J Acquir Immune Defic Syndr 1999 **2010**; 53:537–543.

32. Petroll AE, Walsh JL, Owczarzak JL, McAuliffe TL, Bogart LM, Kelly JA. PrEP Awareness, Familiarity, Comfort, and Prescribing Experience among US Primary Care Providers and HIV Specialists. AIDS Behav **2017**; 21:1256–1267.

33. Krakower D, Ware N, Mitty JA, Maloney K, Mayer KH. HIV Providers' Perceived Barriers and Facilitators to Implementing Pre-Exposure Prophylaxis in Care Settings: A Qualitative Study. AIDS Behav **2014**; 18:1712–1721.

34. Kaiser Family Foundation. Status of State Medicaid Expansion Decisions: Interactive Map. Henry J Kais. Fam. Found. 2020; Available at: https://www.kff.org/medicaid/issue-brief/status-of-state-medicaid-expansion-decisions-interactive-map/. Accessed 4 March 2020.

35. Goodreau SM, Rosenberg ES, Jenness SM, et al. Sources of racial disparities in HIV prevalence in men who have sex with men in Atlanta, GA, USA: a modelling study. Lancet HIV **2017**; 4:e311–e320.

36. Moncada J, Schachter J, Liska S, Shayevich C, Klausner JD. Evaluation of Self-Collected Glans and Rectal Swabs from Men Who Have Sex with Men for Detection of Chlamydia trachomatis and Neisseria gonorrhoeae by Use of Nucleic Acid Amplification Tests. J Clin Microbiol **2009**; 47:1657–1662.

37. Park I, Schapiro J, Hurley L, et al. LB 2.59 Improving std screening in hiv care through implementation of self-collected extragenital swabs. Sex Transm Infect **2017**; 93:A92–A92.

38. Fisher M, Wayal S, Smith H, et al. Home Sampling for Sexually Transmitted Infections and HIV in Men Who Have Sex with Men: A Prospective Observational Study. PLoS ONE **2015**; 10:e0120810.

39. Chen Y-T, Bowles K, An Q, et al. Surveillance Among Men Who have Sex with Men in the United States: A Comparison of Web-Based and Venue-Based Samples. AIDS Behav **2018**; 22:2104–2112.

40. Fauci AS, Redfield RR, Sigounas G, Weahkee MD, Giroir BP. Ending the HIV Epidemic: A Plan for the United States. JAMA **2019**; 321:844–845.

TABLES

Table 1. Characteristics of PrEP-Eligible ARTnet Participants, Stratified by PrEP Use History

	Total[a]		Current PrEP Use		Prior PrEP Use		Never Used PrEP	
	N or Mean	%[b] or SD	N or Mean	% or SD	N or Mean	% or SD	N or Mean	% or SD
Total Sample	3259	100.0	631	100.0	178	100.0	2450	100.0
Race/Ethnicity								
Black (Non-Hispanic)	134	4.1	28	4.4	4	2.3	102	4.2
Hispanic	422	12.9	81	12.8	23	12.9	318	13.0
Other (Non-Hispanic)	283	8.7	53	8.4	11	6.2	219	8.9
White (Non-Hispanic)	2420	74.3	469	74.3	140	78.7	1811	73.9
Age	37.3	13.6	39.5	12.4	34.1	11.5	36.9	13.9
15–24	713	21.9	71	11.3	36	20.2	606	24.7
25–34	947	29.1	196	31.1	76	42.7	675	27.6
35–44	518	15.9	122	19.3	26	14.6	370	16.3
45–54	573	17.6	149	23.6	24	13.5	400	16.3
55–65	508	15.6	93	14.7	16	9.0	399	16.3
Residence in Census Division[c]								
Pacific	562	17.2	155	24.6	31	17.4	376	15.3
Mountain	281	8.6	30	4.8	21	11.8	230	9.4
West North Central	187	5.7	32	5.1	13	7.3	142	5.8
East North Central	466	14.3	92	14.6	25	14.0	349	14.2
West South Central	323	9.9	61	9.7	11	6.2	251	10.2
East South Central	132	4.1	19	3.0	6	3.4	107	4.4
South Atlantic	701	21.5	119	18.9	35	19.7	547	22.3
Middle Atlantic	442	13.6	92	14.6	20	11.2	330	13.5
New England	165	5.1	31	4.9	16	9.0	118	4.8
Residence in Southeast[b]								
Yes	760	23.3	121	19.2	33	18.5	606	24.7
No	2499	76.7	510	80.8	145	81.5	1844	75.3
Health Insurance								
Private	2441	76.4	516	82.2	128	71.9	1797	75.2
Public	514	16.1	94	15.0	33	18.5	387	16.2
None	242	7.6	18	2.9	17	9.6	207	8.7
Annual Household Income								
$0 to $19,999	372	12.3	54	9.3	18	10.7	300	13.2
$20,000 to $39,999	574	19.1	93	16.1	34	20.2	447	19.7
$40,000 to $74,999	861	28.6	153	26.4	56	33.3	652	28.8
$75,000 or more	1206	40.0	279	48.2	60	35.7	867	38.3
Screened for HIV in the Past 12 Months								
Yes	2106	71.9	561	98.2	129	79.6	1416	64.5
No	822	28.1	10	1.8	33	20.4	779	35.5
Screened for an STI in Past 12 Months								
Yes	1677	54.4	547	89.1	122	70.9	1008	43.8
No	1408	45.6	67	10.9	50	28.1	1291	56.2

Abbreviations: PrEP, preexposure prophylaxis; SD, standard deviation; USD, U.S. Dollars; STI, sexually transmitted infection

[a] Totals may not add up to sample total due to missing data for health insurance, annual household income, and screening for

[b] Column percents

[c] https://www2.census.gov/geo/pdfs/maps-data/maps/reference/us_regdiv.pdf

[d] Residence in Alabama, Arkansas, Florida, Georgia, Kentucky, Louisiana, Mississippi, North Carolina, South Carolina, Tennessee, Virginia, and West Virginia

Table 2. Frequency of PrEP Care Visits and HIV/STI Screening at PrEP Care Visits among MSM with Current or Prior PrEP Use

	Total[a]		Current PrEP Use		Prior PrEP Use	
	N	%	N	%	N	%
Total Sample	809	100.0	631	100.0	178	100.0
Frequency of PrEP Care Visits						
Every Month	23	2.9	13	2.1	10	5.7
Every 3 Months	629	78.3	514	82.0	115	65.3
Every 6 Months	83	10.3	75	12.0	8	4.5
Every 9 Months	3	0.4	1	0.2	2	1.1
Every 12 Months	8	1.0	8	1.3	0	0.0
Returned Only Once	21	2.6	3	0.5	18	10.2
Returned at Some Other Interval of Time	11	1.4	6	1.0	5	2.8
Did Not Return Regularly	25	3.1	7	1.1	18	10.2
Frequency of HIV screening at PrEP Care Visits		0				
Always	685	91.1	566	92.5	119	85.0
Sometimes	50	6.6	40	6.5	10	7.1
Rarely	7	0.9	3	0.5	4	2.9
Never	10	1.3	3	0.5	7	5.0
Frequency of Throat Swab at PrEP Care Visits						
Always	316	42.5	267	44.1	49	35.3
Sometimes	161	21.6	134	22.1	27	19.4
Rarely	56	7.5	44	7.3	12	8.6
Never	211	28.4	160	26.4	51	36.7
Frequency of Rectal Swab at PrEP Care Visits						
Always	270	36.2	226	37.2	44	31.7
Sometimes	155	20.8	128	21.1	27	19.4
Rarely	58	7.8	50	8.2	8	5.8
Never	263	35.3	203	33.4	60	43.2
Frequency of Blood Sample at PrEP Care Visits						
Always	492	67.4	415	69.9	77	56.6
Sometimes	143	19.6	114	19.2	29	21.3
Rarely	32	4.4	21	3.5	11	8.1
Never	63	8.6	44	7.4	19	14.0
Frequency of Urine Sample or Swab at PrEP Care Visits						
Always	425	57.3	356	59.0	69	49.6
Sometimes	150	20.2	128	21.2	22	15.8
Rarely	42	5.7	33	5.5	9	6.5
Never	125	16.8	86	14.3	39	28.1

Abbreviations: STI, sexually transmitted infection; PrEP, preexposure prophylaxis

[a] Totals may not add up to sample total due to missing data

Table 3. Bivariable Associations of Consistent Syphilis and Urogenital STI Screening with Demographic and Behavioral Factors among MSM Who Have Ever Used PrEP

	Syphilis Screening[a] (N=730)						Urogenital STI Screening (N=742)					
	Consistent[b]		Inconsistent				Consistent		Inconsistent			
	N	%	N	%	PR	95% CI	N	%	N	%	PR	95% CI
Total Sample	635	87.0	95	13.0			575	77.5	167	22.5		
Age												
15-24	75	83.3	15	16.7	1.00	-	73	80.2	18	19.8	1.00	-
25-34	215	87.8	30	12.2	1.05	0.95, 1.17	201	80.1	50	19.9	1.00	0.89, 1.12
35-44	123	90.4	13	9.6	1.09	0.97, 1.21	104	75.4	34	24.6	0.94	0.82, 1.08
45-54	140	85.9	23	14.1	1.03	0.92, 1.15	127	77.9	36	22.1	0.97	0.85, 1.11
55-65	82	85.4	14	14.6	1.02	0.91, 1.16	70	70.7	29	29.3	0.88	0.75, 1.04
Race/Ethnicity												
Black (Non-Hispanic)	30	93.8	2	6.3	1.00	-	28	87.5	4	12.5	1.00	-
Hispanic	84	91.3	8	8.7	0.97	0.87, 1.09	78	84.8	14	15.2	0.97	0.83, 1.13
Other (Non-Hispanic)	53	91.4	5	8.6	0.97	0.87, 1.10	45	77.6	13	22.4	0.89	0.73, 1.07
White (Non-Hispanic)	468	85.4	80	14.6	0.91	0.83, 1.00	424	75.7	136	24.3	0.87	0.75, 0.99
Health Insurance												
None	28	84.8	5	15.2	1.00	-	23	69.7	10	30.3	1.00	-
Public	98	87.5	14	12.5	1.03	0.88, 1.21	91	79.8	23	20.2	1.15	0.90, 1.46
Private	507	87.1	75	12.9	1.03	0.89, 1.19	459	77.5	133	22.5	1.11	0.88, 1.40
Annual Household Income												
$0 to $19,999	54	83.1	11	16.9	1.00	-	50	76.9	15	23.1	1.00	-
$20,000 to $39,999	86	80.4	21	19.6	0.97	0.84, 1.12	87	79.1	23	20.9	1.03	0.87, 1.21
$40,000 to $74,999	165	87.3	24	12.7	1.05	0.93, 1.19	147	75.4	48	24.6	0.98	0.84, 1.14
$75,000 or more	279	89.1	34	10.9	1.07	0.96, 1.21	244	77.2	72	22.8	1.00	0.87, 1.16
Residence in Census Division												
New England	36	92.3	3	7.7	1.00	-	33	80.5	8	19.5	1.00	-
Middle Atlantic	89	85.6	15	14.4	0.93	0.82, 1.05	86	79.6	22	20.4	0.99	0.83, 1.18
East North Central	89	84.0	17	16.0	0.91	0.80, 1.03	83	77.6	24	22.4	0.96	0.80, 1.16
West North Central	29	74.4	10	25.6	0.81	0.66, 0.99	24	60.0	16	40.0	0.75	0.56, 1.00
South Atlantic	119	86.9	18	13.1	0.94	0.84, 1.05	93	67.9	44	32.1	0.84	0.70, 1.02
East South Central	18	85.7	3	14.3	0.93	0.76, 1.13	13	61.9	8	38.1	0.77	0.53, 1.11
West South Central	62	89.9	7	10.1	0.97	0.86, 1.10	57	82.6	12	17.4	1.03	0.85, 1.24
Mountain	34	79.1	9	20.9	0.86	0.72, 1.02	32	74.4	11	25.6	0.92	0.73, 1.17
Pacific	159	92.4	13	7.6	1.00	0.91, 1.11	154	87.5	22	12.5	1.09	0.93, 1.28
Residence in Southeast[c]												
No	517	87.2	76	12.8	1.00	-	484	80.0	121	20.0	1.00	-
Yes	118	86.1	19	13.9	0.99	0.92, 1.06	91	66.4	46	33.6	0.83	0.73, 0.94
Exposure at Anatomical Site[d]												
No							15	68.2	7	31.8	1.00	-
Yes							560	77.8	160	22.2	1.14	0.86, 1.52

Abbreviations: STI, sexually transmitted infection; PrEP, preexposure prophylaxis; PR, prevalence ratio; CI, confidence interval

[a] Participants who provided blood samples for STI screening were considered to be screened for syphilis

[b] Consistent screening defined as always or sometimes (versus rarely or never) being screened at PrEP care visits compared

[c] Alabama, Arkansas, Florida, Georgia, Kentucky, Louisiana, Mississippi, North Carolina, South Carolina, Tennessee, Virginia, and W Virginia

[d] Defined as any insertive anal intercourse or receiving oral sex for urogenital STI screening

Table 4. Bivariable Associations of Consistent Rectal and Pharyngeal STI Screening with Demographic and Behavioral Factors among MSM Who Have Ever Used PrEP

| | Rectal STI Screening (N=746) | | | | | | Pharyngeal STI Screening (N=744) | | | | | |
| | Consistent | | Inconsistent | | | | Consistent | | Inconsistent | | | |
	N	%	N	%	PR	95% CI	N	%	N	%	PR	95% CI
Total Sample	425	57.0	321	43.0			477	64.1	267	35.9		
Age												
15-24	61	67.0	30	33.0	1.00	-	63	70.0	27	30.0	1.00	-
25-34	146	57.9	106	42.1	0.86	0.72, 1.03	163	64.9	88	35.1	0.93	0.79 1.09
35-44	74	53.6	64	46.4	0.80	0.65, 0.99	89	64.5	49	35.5	0.92	0.77 1.11
45-54	91	55.2	74	44.8	0.82	0.67, 1.00	102	61.8	63	38.2	0.88	0.74 1.06
55-65	53	53.0	47	47.0	0.79	0.63, 1.00	60	60.0	40	40.0	0.86	0.70 1.06
Race/Ethnicity												
Black (Non-Hispanic)	16	50.0	16	50.0	1.00	-	19	59.4	13	40.6	1.00	-
Hispanic	57	62.0	35	38.0	1.24	0.85, 1.82	61	66.3	31	33.7	1.12	0.81, 1.54
Other (Non-Hispanic)	39	67.2	19	32.8	1.34	0.91, 1.99	43	74.1	15	25.9	1.25	0.90, 1.73
White (Non-Hispanic)	313	55.5	251	44.5	1.11	0.78, 1.58	354	63.0	208	37.0	1.06	0.79, 1.42
Health Insurance[a]												
None	19	57.6	14	42.4	1.00	-	22	66.7	11	33.3	1.00	-
Public	58	50.0	58	50.0	0.87	0.62, 1.23	74	63.8	42	36.2	0.96	0.73, 1.26
Private	347	58.4	247	41.6	1.01	0.75, 1.37	380	64.2	212	35.8	0.96	0.75, 1.23
Annual Household Income												
$0 to $19,999	35	53.0	31	47.0	1.00	-	44	66.7	22	33.3	1.00	-
$20,000 to $39,999	59	53.6	51	46.4	1.01	0.76, 1.35	62	56.4	48	43.6	0.85	0.67, 1.07
$40,000 to $74,999	109	55.9	86	44.1	1.05	0.81, 1.37	125	64.4	69	35.6	0.97	0.79, 1.18
$75,000 or more	187	58.8	131	41.2	1.11	0.87, 1.42	206	65.0	111	35.0	0.97	0.81, 1.18
Residence in Census Division												
New England	29	70.7	12	29.3	1.00	-	30	73.2	11	26.8	1.00	-
Middle Atlantic	65	60.2	43	39.8	0.85	0.66, 1.09	74	69.2	33	30.8	0.95	0.76, 1.18
East North Central	57	52.8	51	47.2	0.75	0.57, 0.97	59	54.6	49	45.4	0.75	0.58, 0.96
West North Central	21	52.5	19	47.5	0.74	0.52, 1.06	24	61.5	15	38.5	0.84	0.62, 1.15
South Atlantic	65	46.1	76	53.9	0.65	0.50, 0.85	79	56.0	62	44.0	0.77	0.60, 0.97
East South Central	7	33.3	14	66.7	0.47	0.25, 0.89	9	42.9	12	57.1	0.59	0.35, 0.99
West South Central	40	58.0	29	42.0	0.82	0.62, 1.09	47	68.1	22	31.9	0.93	0.73, 1.19
Mountain	19	45.2	23	54.8	0.64	0.43, 0.94	20	47.6	22	52.4	0.65	0.45, 0.94
Pacific	122	69.3	54	30.7	0.98	0.79, 1.22	135	76.7	41	23.3	1.05	0.86, 1.28
Residence in Southeast[b]												
No	364	60.1	242	39.9	1.00	-	400	66.2	204	33.8	1.00	-
Yes	61	43.6	79	56.4	0.73	0.59, 0.89	77	55.0	63	45.0	0.83	0.71, 0.97
Exposure at Anatomical Site[c]												
No	61	44.2	77	55.8	1.00	-	15	62.5	9	37.5	1.00	-
Yes	364	59.9	244	40.1	1.35	1.11, 1.65	462	64.2	258	35.8	1.03	0.75, 1.41

Abbreviations: STI, sexually transmitted infection; PrEP, preexposure prophylaxis; PR, prevalence ratio; CI, confidence interval
[a] Consistent screening defined as always or sometimes (versus rarely or never) being screened at PrEP care visits compared
[b] Alabama, Arkansas, Florida, Georgia, Kentucky, Louisiana, Mississippi, North Carolina, South Carolina, Tennessee, Virginia, and W Virginia
[c] Defined as any receptive anal intercourse for rectal STI screening and any performance of oral sex on a partner for pharyngeal STI screening

Table 5. Multivariable Associations of Consistent[a] STI Screening with Residence in the Southeast U.S.[b] Among MSM who Have Ever Used PrEP

	Adjusted PR[c]	95% CI
Syphilis Screening[d]		
Residence Outside Southeast	1.00	-
Residence in Southeast	1.02	0.95, 1.10
Urogenital STI Screening		
Residence Outside Southeast	1.00	-
Residence in Southeast	0.86	0.76, 0.98
Rectal STI Screening		
Residence Outside Southeast	1.00	-
Residence in Southeast	0.76	0.62, 0.93
Pharyngeal STI Screening		
Residence Outside Southeast	1.00	-
Residence in Southeast	0.87	0.74, 1.03

Abbreviations: STI, sexually transmitted infection; PrEP, preexposure prophylaxis; PR, prevalence ratio; CI, confidence interval

[a] Consistent screening defined as always or sometimes (versus rarely or never) being screened at PrEP care visits compared

[b] Residence in Alabama, Arkansas, Florida, Georgia, Kentucky, Louisiana, Mississippi, North Carolina, South Carolina, Tennessee, Virginia, and West Virginia

[c] Adjusted for age, race/ethnicity, annual household income, and sexual exposure at anatomical site

[d] Not adjusted for sexual exposure at anatomical site