Transmission potential of COVID-19 in Iran

Authors: Kamalich Muniz-Rodriguez[1], MPH, Isaac Chun-Hai Fung[1]*, PhD, Shayesteh R. Ferdosi, PhD,

Sylvia K. Ofori, MPH, Yiseul Lee, MPH, Amna Tariq, MPH, Gerardo Chowell*, PhD

Affiliations: Jiann-Ping Hsu College of Public Health, Georgia Southern University (I. C.-H. Fung, K.

Muniz-Rodriguez, S. K. Ofori), Brain Tumor Unit, The Translational Genomics Research Institute (S. R.

Ferdosi), School of Public Health, Georgia State University (Y. Lee, A. Tariq, G. Chowell)

[1] These two individuals serve as co-first authors.

Email addresses: K. Muniz-Rodriguez (km11200@georgiasouthern.edu), I. C. H. Fung

(cfung@georgiasouthern.edu), S. R. Ferdosi (shayesteh01@gmail.com), S. K. Ofori

(so01935@georgiasouthern.edu), Y. Lee (ylee97@student.gsu.edu), A. Tariq (atariq1@student.gsu.edu),

G. Chowell (gchowell@gsu.edu)

*Correspondence should be addressed to: I. C. H. Fung, Department of Biostatistics, Epidemiology and

Environmental Health Sciences, Jiann-Ping Hsu College of Public Health, Georgia Southern University.

Telephone: 912-478-5079. Email: cfung@georgiasouthern.edu; G. Chowell, Department of Population

Health Sciences, School of Public Health, Georgia State University. Telephone: 404-413-9322. Fax: 404-

413-2344. Email: gchowell@gsu.edu.

Research Letter

Figure, n = 1

Technical Appendix, n = 1, with 6 supplementary tables and 8 supplementary figures

Abstract: n=50; Word limit = 50

We estimated the reproduction number of 2020 Iranian COVID-19 epidemic using two different methods:

R_0 was estimated at 4.4 (95% CI, 3.9, 4.9) (generalized growth model) and 3.50 (1.28, 8.14) (epidemic

doubling time) (February 19 - March 1) while the effective R was estimated at 1.55 (1.06, 2.57) (March 6-

19).

Keywords: Communicable Diseases; Coronavirus; COVID-19; Epidemiology; Infections; Iran.

Main text

Main text

Iran has been experiencing a devastating epidemic of COVID-19 in 2020 (1). The virus appears

to be spreading rapidly with multiple cases reported among the elite (2). One model estimated that 18,300

(95% CI, 3,770 to 53,470) cases would have had occurred by February 24 (3). The Iranian authorities

have adopted social distancing measures to slow disease transmission (4). As the epidemic continues, our

results call for efforts to minimize COVID-19-related morbidity and mortality.

To confront the epidemic, the transmission potential of COVID-19 is a useful metric to guide the

outbreak response efforts including the scope and intensity of interventions. During the early transmission

phase, the basic reproduction number, R_0, represents such a measurement quantifying the average number

of secondary cases that primary cases generate in a completely susceptible population in the absence of

interventions or behavioral changes. $R_0>1$ indicates the possibility of sustained transmission, whereas

$R_0<1$ implies that transmission chains cannot sustain epidemic growth. As the epidemic continues its

course, the effective reproduction number, R_e, offers a time-dependent record of the average number of

secondary cases per case as the number of susceptible individuals gets depleted and control interventions

takes effect. Herein we employed two different methods to quantify the reproduction number using the

curve of reported COVID-19 cases in Iran and its five regions (Table S1).

Using a Wikipedia entry as a starting point, Farsi-speaking coauthor SFR double-checked the data

of daily reported new cases, from February 19 through March 19, 2020 (the day before the Iranian New

Year), using Iranian official press releases and other credible news sources, and corrected the data as per

the official data (Figure S1, Tables S2 and S3). Incident cases for the 5 regions were missing for two

days: March 2-3, 2020, which were not included in our analysis. We noted that the reported national

number of new cases did not match the sum of the new cases reported in Iran's five regions on March 5,

2020; we decided to treat each time series as independent and used the data as reported. Using Method 1,

we estimated R_0 using data from February 19 through March 1, 2020. Using Method 2, we estimated R_0

from the early transmission phase (February 19 through March 1, 2020) and R_e based on the growth rate

estimated from March 6 through 19, 2020 during which the epidemic slowed down likely reflecting the

impact of social distancing measures.

Two methods were utilized to estimate the reproduction number. The first method utilized a

generalized growth model (GGM) (5) with the growth rate and its scaling factor to characterize the daily

reported incidence, followed by simulation of the calibrated GGM, using a discretized probability

distribution of the serial interval and assuming a Poisson error structure. See Technical Appendix for

detailed description of Method 1.

The second method for estimating the reproduction number is based on the calculation of the

epidemic's doubling times, which correspond to the times when the cumulative incidence doubles. We

estimated the epidemic doubling time using the curve of cumulative daily reported cases. To quantify

parameter uncertainty, we used parametric bootstrapping with a Poisson error structure around the

number of new reported cases in order to derive 95% confidence intervals of our estimates (6-8).

Assuming exponential growth, the epidemic growth rate is equal to ln(2)/doubling time. Assuming that

the pre-infectious and infectious periods follow an exponential distribution, R_0 is approximately equal to 1

+ growth rate × serial interval. See Technical Appendix and Equation 4.14, Table 4.1 in Vynnycky and

White (9).

In both methods, the serial interval was assumed to follow a gamma distribution (mean: 4.41

days; standard deviation: 3.17 days) (10, 11). MATLAB version R2019b and R version 3.6.2 were used

for data analyses and creating figures. It was determined *a priori* that $\alpha = 0.05$.

Using Method 1, we estimated an R_0 of 4.4 (3.9, 4.9) for the COVID-19 epidemic in Iran. We

estimated a growth rate of 0.65 (95% CI, 0.56, 0.75) and a scaling parameter of 0.96 (95% CI, 0.93, 1)

(Table S4). The latter indicated a near-exponential growth of the epidemic (**Figure**). Using Method 2, we

found that from February 19 through March 1, the cumulative incidence of confirmed cases in Iran had

doubled 8 times. The estimated epidemic doubling time was 1.20 (95% CI, 1.05, 1.45) days and the

corresponding R_0 estimate was 3.50 (95% CI, 1.28, 8.14). From March 6 through March 19, the

cumulative incidence of confirmed cases had doubled 1 time, with a doubling time of 5.46 (5.29, 5.65)

days. The corresponding effective R estimate was 1.55 (95% CI, 1.06, 2.57) (Table S5, Figures S7, S8).

Our results are robust, as they are consistent with the Iranian COVID-19 R_0 estimates of 4.7 and 4.86

generated by Ahmadi et al. and Sahafizadeh and Sartoli (12, 13), respectively, but are higher than the R_0

of 2.72 estimated by Ghaffarzadegan & Rahmandad (14).

Our study has limitations. First, our analysis is based on the number of daily reported cases

whereas it would be ideal to analyze case counts by date of symptoms onset, which were not available.

Second, case counts could be underreported due to underdiagnosis, given subclinical or asymptomatic

cases, or limited testing capacity to test mild cases. The rapid increase in case counts might represent a

belated realization of the severity of the epidemic and a rapid process of catching up with testing many

suspected cases. If the reporting ratio remains constant over the study period, and given the near-

exponential growth of the epidemic's trajectory, our estimates would remain reliable; however, this is a

strong assumption. Third, while data are not stratified according to imported and local cases, we assumed

that they were infected locally, as it is likely that transmission has been ongoing in Iran for some time (3).

In conclusion, we used two different methods to compute the R_0 of COVID-19 epidemic in Iran.

Our mean estimate was at 4.4 (95% CI, 3.9, 4.9) for Method 1, and at 3.50 (95% CI, 1.28, 8.14) for

Method 2, which means that on average around four susceptible individuals were infected by one

infectious individual during the early growth phase. When social distancing interventions are in place, the

epidemic slows. The doubling time of the COVID-19 epidemic in Iran increased from 1.20 (95% CI,

1.05, 1.45) to 5.46 (95% CI, 5.29, 5.65) days, and the effective R dropped to 1.55 (95% CI, 1.06, 2.57).

While the COVID-19 epidemic in Iran has now slowed down substantially, the situation in Iran is still

dire with the average daily number of ~2290 new reported cases over 14 days from March 20 through

April 2, 2020. Tightening social distancing interventions is needed to bring the epidemic under control in

Iran.

Co-first authors' biographies

103 Kamalich Muniz-Rodriguez, a doctoral student in epidemiology, and Dr. Isaac Chun-Hai Fung, an

104 associate professor of epidemiology, are at Jiann-Ping Hsu College of Public Health, Georgia Southern

105 University. Their research interests include infectious disease epidemiology, digital health and disaster

106 emergency responses.

Acknowledgement

108 GC acknowledges support from NSF grant 1414374 as part of the joint NSF-NIH-USDA Ecology and

109 Evolution of Infectious Diseases program. ICHF acknowledges salary support from the National Center

110 for Emerging and Zoonotic Infectious Diseases, Centers for Disease Control and Prevention

111 (19IPA1908208). This article is not part of ICHF's CDC-sponsored projects.

Conflict of interest statement

113 We do not have any conflict of interest to declare.

Disclaimer

115 This article does not represent the official positions of the Centers for Disease Control and Prevention, the

116 National Institutes of Health, or the United States Government.

Ethics statement

118 The Georgia Southern University Institutional Review Board has made a non-human subjects

119 determination for our project (H20364), under G8 exemption category.

References

1.   Wood G. Coronavirus could break Iranian Society. The Atlantic. . 2020 [cited February 29,
2020]; Available from: https://www.theatlantic.com/ideas/archive/2020/02/iran-cannot-handle-
coronavirus/607150/

2.   Fassihi F, Gladstone R. Iran Vice President Is One of 7 Officials to Contract Coronavirus. 2020
[cited March 1, 2020]; Available from:
https://www.nytimes.com/2020/02/27/world/middleeast/coronavirus-iran-vice-
president.html?action=click&module=RelatedLinks&pgtype=Article
3.   Tuite AR, Bogoch II, Sherbo R, Watts A, Fisman D, Khan K. Estimation of COVID-2019 burden
and potential for international dissemination of infection from Iran. medRxiv.
2020;2020.02.24.20027375v1.
4.   Fassihi F, Gladstone R. Confusion and Contradiction Reign in Iran's Coronavirus Reporting. New
York Times. 2020 [cited March 1, 2020]; Available from:
https://www.nytimes.com/2020/02/28/world/middleeast/coronavirus-iran-confusion.html
5.   Viboud C, Simonsen L, Chowell G. A generalized-growth model to characterize the early
ascending phase of infectious disease outbreaks. Epidemics. 2016 2016/06/01/;15:27-37.
6.   Banks HT, Hu S, Thompson WC. Modeling and inverse problems in the presence of uncertainty:
CRC Press; 2014.
7.   Chowell G, Ammon CE, Hengartner NW, Hyman JM. Transmission dynamics of the great
influenza pandemic of 1918 in Geneva, Switzerland: Assessing the effects of hypothetical interventions. J
Theor Biol. 2006 Jul 21;241(2):193-204.
8.   Chowell G, Shim E, Brauer F, Diaz-Duenas P, Hyman JM, Castillo-Chavez C. Modelling the
transmission dynamics of acute haemorrhagic conjunctivitis: application to the 2003 outbreak in Mexico.
Stat Med. 2006 Jun 15;25(11):1840-57.
9.   Vynnycky E, White RG. An Introduction to Infectious Disease Modelling. Oxford: Oxford
University Press; 2010.
10.   Nishiura H, Linton NM, Akhmetzhanov AR. Serial interval of novel coronavirus (2019-nCoV)
infections. medRxiv. 2020:2020.02.03.20019497.
11.   You C, Deng Y, Hu W, Sun J, Lin Q, Zhou F, et al. Estimation of the Time-Varying
Reproduction Number of COVID-19 Outbreak in China. medRxiv. 2020:2020.02.08.20021253.
12.   Ahmadi A, Fadaei Y, Shirani M, Rahmani F. Modeling and Forecasting Trend of COVID-19
Epidemic in Iran. medRxiv. 2020:2020.03.17.20037671.
13.   Sahafizadeh E, Sartoli S. Estimating the reproduction number of COVID-19 in Iran using
epidemic modeling. medRXiv. 2020:2020.03.20.20038422.
14.   Ghaffarzadegan N, Rahmandad H. Simulation-based Estimation of the Spread of COVID-19 in
Iran. medRXiv. 2020:2020.03.22.20040956.

Figure. The mean basic reproduction number of COVID-19 epidemic in Iran, with 95% confidence

interval. Estimates for growth rate, r, and the scaling of the growth rate parameter, p, are also provided.

The plot in the lower panel depicts the fit of the Generalized Growth Model (Method 1) to the Iranian

data assuming Poisson error structure as of March 1, 2020.