Isolation and contact tracing can tip the scale to containment of

COVID-19 in populations with social distancing

Mirjam E. Kretzschmar[1#], Ganna Rozhnova[1], Michiel van Boven[1]

[1] Julius Center for Health Sciences and Primary Care, University Medical Center Utrecht,

Utrecht University, Utrecht, the Netherlands

corresponding author:

Mirjam E. Kretzschmar

Julius Center for Health Sciences and Primary Care

University Medical Center Utrecht, Utrecht

Email: m.e.e.kretzschmar@umcutrecht.nl

Phone: +31 88 75 687 61

Abstract

Background: Novel coronavirus (SARS-CoV-2) has extended its range of transmission in all parts of the world, with substantial variation in rates of transmission and severity of associated disease. Many countries have implemented social distancing as a measure to control further spread.

Methods: We evaluate whether and under which conditions containment or slowing down COVID-19 epidemics are possible by isolation and contact tracing in settings with various levels of social distancing. We use a stochastic transmission model in which every person generates novel infections according to a probability distribution that is affected by the incubation period distribution (time from infection to symptoms), distribution of the latent period (time from infection to onset of infectiousness), and overall transmissibility. The model distinguishes between close contacts (e.g., within a household) and other contacts in the population. Social distancing affects the number of contacts outside but not within the household.

Findings: The proportion of asymptomatic or unascertained cases has a strong impact on the controllability of the disease. If the proportion of asymptomatic infections is larger than 30%, contact tracing and isolation cannot achieve containment for an R_0 of 2.5. Achieving containment by social distancing requires a reduction of numbers of non-household contacts by around 90%. Depending on the realized level of contact reduction, tracing and isolation of only household contacts, or of household and non-household contacts are necessary to reduce the effective reproduction number to below 1. A combination of social

distancing with isolation and contact tracing leads to synergistic effects that increase the

prospect of containment.

Interpretation: Isolation and contact tracing can be an effective means to slow down

epidemics, but only if the majority of cases are ascertained. In a situation with social

distancing, contact tracing can act synergistically and tip the scale towards containment, and

can therefore be a tool for controlling COVID-19 epidemics as part of an exit strategy from

current lockdown measures.

Funding: This research was partly funded by ZonMw project number 91216062.

Research in context

Evidence before this study

As of 8 April 2020, the novel coronavirus (SARS-CoV-2) has spread to more than 170

countries and has caused near 90,000 deaths of COVID-19 worldwide. In the absence of

effective medicines and vaccines, the preventive measures are limited to social distancing,

isolation of confirmed and suspected cases, and identification and quarantining of their

contacts. Evidence suggests that a substantial portion of transmission may occur before the

onset of symptoms and before cases can be isolated, and that many cases remain

unascertained. This has potentially important implications for the prospect of containment

by combinations of these measures.

Added value of this study

Using a stochastic transmission model armed with current best estimates of epidemiological

parameters, we evaluated under which conditions containment could be achieved with

combinations of social distancing, isolation and contact tracing. We investigated the level of

social distancing needed for containment, and how an additional implementation of

isolation and contact tracing may likely help to in reducing the effective reproduction

number to below 1, the critical threshold. We analyzed what proportion of household and

non-household contacts need to be isolated effectively to achieve containment depending

on the level of social distancing in the population. We estimated the impact of combinations

of these measures on epidemic growth rate and doubling time for the number of infections.

We find that under realistic assumptions on the level of social distancing, additional isolation

and contact tracing are needed for stopping the epidemic. Whether quarantining only

household contacts is sufficient, depends on levels of social distancing and timeliness of

tracing and isolation.

Implications of all the available evidence

Our analyses based on best understanding of the epidemiology of COVID-19, highlight that

if social distancing is not complete, isolation and contact tracing at least of household

contacts can help to delay and lower the epidemic peak. High levels of timely contact

tracing of household and non-household contacts may be sufficient to control the epidemic.

Introduction

As of early April 2020, the number of infections of the novel coronavirus (SARS-CoV-2) is still

increasing at an exponential rate in many countries, while the virus is expanding its range to

all parts of the world. There are no registered effective medicines, treatment options are

mainly supportive, and there are no vaccines available, limiting preventive measures mainly

to social distancing combined with isolation of infected persons and those that have high

likelihood of being infected, for instance because they have been traced as contacts of

infected persons [1, 2]. However, recently it was questioned by high-ranked officials of the

World Health Organization (WHO) whether social distancing or lockdowns are sufficient to

stop the SARS-CoV-2 outbreak (https://www.reuters.com/article/us-health-coronavirus-

who-ryan/lockdowns-not-enough-to-defeat-coronavirus-whos-ryan-idUSKBN2190FM).

Possibly, additional measures are needed, for example active tracing of contacts in

combination with isolation of infected contacts. Also, such measures will be important in the

context of exit strategies, i.e. once social distancing measures are reduced or lifted, as has

been suggested recently [3]. It is unclear, how effective such combinations of interventions

can be in populations with social distancing in place.

To what extent local containment or local slowing down of an epidemic by isolation and

contact tracing is successful depends on the fraction of infections that remain asymptomatic

or have mild disease, and on the infectiousness before the onset of symptoms [4, 5]. It is

known that a high probability of asymptomatic infection, a high proportion of transmission

occurring before the onset of symptoms, a long delay between case finding and isolation,

and high overall transmissibility all factor in negatively in the likelihood that an outbreak can

be contained [6-9]. For SARS-CoV-2, evidence indicates that a high fraction of infected

persons is infectious before they show symptoms (up to 50%), that a substantial fraction of

infections may be asymptomatic or show only mild symptoms (up to 80%), and that the

epidemic doubling time in the absence of interventions is approximately one week or

perhaps even less [5, 10-16]. On the other hand, it is also reported that with intensive

contact tracing it may be possible to trace the majority (>80%) of secondary infections [17].

Here we provide model-based analyses of the impact of isolation and contact tracing in a

setting with various levels of social distancing measures, using varying levels of the

effectiveness and timeliness of contact tracing. It is important to consider the impact of each

of these interventions in isolation but also in combination, as it is well known that each

intervention that reduces transmission in the population (i.e. reduces the reproduction

number) is expected to increase the effectiveness of additional interventions in a synergistic

manner [18]. We show how those measures impact on the cumulative numbers of infecteds

during an outbreak and compare with recent outbreak data from various countries [19].

We focus on conditions that make containment of an early epidemic possible, but also on

the impact of isolation and contact tracing when containment is not possible, or in an exit

phase from social distancing. In a population with social distancing in place, we investigate

what the additional impact of isolation and contact tracing can be for reaching containment

or slowing down the spread. We report effective reproduction number, the (exponential)

rate of increase, and the doubling time of the epidemic for scenarios with various

combinations of interventions. Considering that the capacity of healthcare systems is

limited, it is important to assess which interventions are most effective in slowing down the

rate of increase of healthcare demand during an ongoing outbreak. As it is likely that, on the

one hand, isolation and contact tracing will be more effective in close contact settings with

well-defined contacts (household, workplace) than in the community (commuting, public

spaces), while, on the other hand, the potential impact of household interventions on the

epidemic could be smaller, we stratify the analyses by transmission setting (henceforth

called household and non-household) [8]. As many of the relevant epidemiological and

intervention parameters are still quite uncertain, or may be variable in different settings, we

focus throughout on a systematic analysis of the relation between key parameters for

timeliness and completeness of contact tracing and main outcomes such as the effective

reproduction number and the epidemic growth rate.

Methods

We modify a model that was developed earlier for similar aims in another context [8]. The stochastic model describes an epidemic in its early phase as a branching process. Starting from a small set of initially infected individuals, the model calculates the numbers of latently infected persons, infectious persons, and persons that are diagnosed and isolated in time steps of one day. Latent infection, infectivity during the infectious period, and daily contact rates are quantified using distributions taken from the literature (Table 1). We distinguish between household contacts and non-household contacts, which differ in the risk of infection and the delay and effectiveness of tracing and isolation. Intervention effectiveness is determined by the daily probability of being diagnosed during the infectious period (Table 2). Furthermore, intervention effectiveness depends on the delays in tracing household and non-household contacts, respectively, and the proportions of contacts can be found and isolated. We assume that isolation is perfect, i.e. that isolated persons cannot transmit any longer. The model is described by a set of difference equations, and allows for explicit computation of the basic reproduction number R_0 and the effective reproduction number under interventions R_e. The model was coded in Mathematica 12.1.

Natural history

We assume that the latent period lasts between 1 and 3 days. Individuals then become infectious for at most 10 days. Infectivity is high at the beginning of the infectious period and decays to low levels during these 10 days (Figure S1a). The probability of symptoms onset increases during the first 4 days of the infectious period, thereby influencing the daily probability of diagnosis during the infectious period (see below).

An infectious individual makes contacts with household members and persons outside the

household. We model the daily number of household contacts with a Poisson distribution,

and the numbers of non-household contacts with a negative binomial distribution (Table 1),

with parameters based on the average household size in the Netherlands, and numbers of

contacts observed in a contact study in the Netherlands (FigureS1b) [20]. With the chosen

parameters, the mean number of contacts per day is 13.2 (SD 8.5).

On each day of the infectious period, an individual makes a number of contacts according to

the contact distribution. This number is reduced by a factor describing the probability that

the contact person has already been infected during earlier contacts with the index person.

More precisely, the number of contacts is reduced by a factor f_k per day k of the infectious

period, describing the probability that a contact has already been infected on previous days

of the infectious period of the index case. The probability of transmission upon contact with

a susceptible household contact is given by the distribution in Figure S1a. As contacts with

persons outside the household are often less close, we assume that the transmission

probability for these contacts is lower by factor 0.25. Figure S1c shows the percentage of

onward transmissions per day of the infectious period, e.g. more than half of all onward

transmissions occur during the first 3 days of the infectious period.

Social distancing

Social distancing can be self-imposed, if people decide to reduce their social contacts during

the outbreak, and it can be government-imposed by closing schools, workplaces, and other

venues of social gatherings. Here we assume that when social distancing is applied,

household contacts remain unchanged, but the mean number of non-household contacts is

reduced. This is implemented by a reduction factor in the mean of the negative binomial

distribution describing non-household contact numbers. The reduction factor for social

distancing was varied between 0% and 95%. In scenarios with social distancing we assumed

that only 60% of cases are symptomatic or can be ascertained.

184   Diagnosis, contact tracing, and isolation

An infectious person becomes symptomatic with a given probability per day of the infectious

period. For SARS-CoV-2 the probability of developing symptoms is high in the first few days

of the infectious period. We assume that if an infected and infectious person has not

developed symptoms by day 6, the probability that he/she will still do so is very small. The

probability of developing symptoms determines whether he/she will be diagnosed and

isolated. The total probability of developing symptoms determines the fraction that remains

asymptomatic or otherwise undiagnosed, i.e. if the total probability of developing symptoms

is smaller than 1, a proportion of the infected persons will remain undiagnosed and can

transmit throughout their infectious period. With the assumed distributions, on average

about half of all potential onward transmissions will have occurred before an infected

person is diagnosed and isolated. The diagnosis can be delayed, which in the model is

implemented by setting the diagnosis probability to zero for the number of days of delay.

This delay describes the time between symptom onset until a symptomatic person visits a

GP or hospital and gets diagnosed.

If an individual is diagnosed, contacts will be traced, and in case they are infected will be

diagnosed and isolated. Tracing goes back in time to trace all contacts during the infectious

period of the index case. There may be a delay before contacts are found and diagnosed,

and only a fraction of all contacts may be found. These parameters, tracing delay and tracing

coverage, may be different for household and non-household contacts. We assume that all

traced infected persons are immediately isolated and cannot transmit any further.

Therefore, the only individuals who will continue transmitting are those who are not found

by tracing and are not yet diagnosed.

Baseline scenario

For assessing the effectiveness of contact tracing and isolation, we use a best case scenario,

where all parameters are set to optimistic values. We assume that when a case is diagnosed,

he/she will immediately be isolated and this will stop onward transmission completely.

Furthermore, we assume that all contacts will be traced, and if found infected will be

isolated immediately. We assume that it takes 1 day to find and isolate both household and

non-household contacts. The rationale for using these overly optimistic assumptions as a

baseline is that we want to investigate the maximum contribution contact tracing can

provide for achieving containment. In sensitivity analyses we investigated for various control

parameters at which point of diverging from the baseline parameters control of the

outbreak will be lost (Supplementary Information).

Output variables

We compute cumulative numbers of cases for a time period of 50 days after the epidemic

has reached more than 100 cases. We compare doubling time of the epidemic with only

isolation of diagnosed cases to situations with various levels of social distancing, isolation

and contact tracing. We assume that all diagnosed cases are isolated upon diagnosis and

cannot transmit further. We plot the epidemic from the time the cumulative number of

infected persons is above 100. We compare these trajectories to similar trajectories for

China, Germany, Italy, Japan, Turkey, and the US using data downloaded from

ourworldindata.org [19]. We show what the additional contribution of isolation and contact

tracing can be in the attempt to bend the curve towards longer doubling times.

The model allows an explicit calculation of the basic reproduction number R_0 and the

effective reproduction number R_e [8]. R_0 is defined as the number of secondary cases an

index case generates on average in a susceptible population, and R_e is the number of

secondary infections per case when an intervention is in place. R_0 is determined by daily

transmission probabilities and numbers of contacts. The effective reproduction number is in

addition determined by the level of social distancing, diagnosis probabilities, tracing delays,

and tracing coverages per day of the infectious period. We can therefore investigate how R_e

depends on R_0 and on the intervention parameters.

We are interested in the critical tracing coverage, i.e. what proportion of non-household

contacts needs to be found and isolated to control the outbreak, for populations with

various levels of social distancing. Furthermore, we study the epidemic growth rate (or

epidemic doubling time) without and with contact tracing and isolation and various levels of

social distancing. In sensitivity analyses, we study how these quantities depend on the delay

to diagnosis of cases and on the delay in contact tracing. For example, we assume that

household contacts can be traced with a high coverage and without delay, but that tracing

of non-household contact may take longer and be less complete.

Based on the distribution of the latent and infectious periods and infectivity, we calculate

the exponential growth rate and doubling time under various assumptions on the

intervention parameters. This gives additional information for situations where the outbreak

is not controllable, because intervention measures will lower the growth rate and increase

the epidemic doubling time.

We investigate how controllability of the outbreak depends on the fraction of infections that

develop symptoms and therefore vary this percentage between 0% and 100%. We then

considered combinations of interventions and their impact on effective reproduction

number, growth rate, and doubling time of the epidemic. We varied levels of social

distancing, and coverage of tracing of household and non-household contacts. In our

analysis for different levels of social distancing we assumed that 60% of infected persons

develop symptoms.

Role of the funding source

The funders of the study had no role in study design, data collection, data analysis, data

interpretation, writing of the manuscript, or the decision to submit for publication. All

authors had full access to all the data in the study and were responsible for the decision to

submit the manuscript for publication.

Results

Basic and effective reproduction numbers

In the baseline scenario without interventions we calibrate the transmission probability such

that the basic reproduction number equals R_0 = 2.5. In that case, 39% of transmissions take

place among household contacts. The basic reproduction number of household contacts is

0.97, and that of non-household contacts 1.53. Hence, if all non-household transmissions

could be prevented, the outbreak would be just under the control limit. In the baseline

scenario without interventions the exponential growth rate is 0.19 per day and the doubling

time is 3.6 days, which agrees with published estimates [13]. For sensitivity analysis of the

relation between R_0 and R_e for various combinations of parameters, see the Supplementary

Information.

Outbreak doubling times

Isolation of diagnosed cases alone reduces the doubling time to almost 4 days. If social

distancing is implemented 28 days after the first case was diagnosed, and the number of

non-household contacts is reduced by 70% [21], the doubling time will increase to around 14

days (Figure 1a). The figure also shows that it takes a week before the impact on the

epidemic growth starts to be visible. If higher levels of social distancing can be achieved, the

doubling times can be increased more (Figure 1b), but even for 90% reduction of contacts

the curve has not been flattened. In Figure 1c the impact of the timing of social distancing

measures on the further course of the outbreak is shown. The figure demonstrates how

every week delay in implementing these measures leads to an order of magnitude higher

numbers of cases, thus emphasizing the importance of fast implementation of social

distancing measures. Finally, in Figure 4d we show a scenario where social distancing is

implemented at day 28 after diagnosing the first case and reduces non-household contacts

by 70%. Then at day 49, also contact tracing is implemented. It is assumed that all infected

household contacts and 50% of the non-household contacts of a newly diagnosed case are

traced and isolated. This additional measure increases the doubling time to … days. We

contrast this scenario with data from epidemics in a number of countries that have

implemented social distancing at various levels and time points.

Fraction of non-household contacts needed to trace and isolate

297   If there is a diagnosis delay, the question arises how successful contact tracing has to be to

298   keep the outbreak under control. We therefore compute the minimum fraction of non-

299   household contacts that need to be traced and isolated (henceforth termed "critical tracing

300   coverage") to bring R_e to below 1 (for examples with various assumptions on timeliness of

301   contact tracing see Supplementary Information).

Impact of asymptomatic cases

Not being diagnosed can be a consequence of not developing symptoms, having only mild

symptoms, or any other reason why infected persons might not be identified by health care.

We subsume these possible reasons for cases not being ascertained under the term

"asymptomatic". With increasing proportion of asymptomatic cases, the possibility of

controlling the outbreak with contact tracing and isolation quickly fades. This is shown in

Figure 2, where we plotted the critical tracing coverage for non-household contacts for

several values of R_0 as a function of the fraction of symptomatic cases (i.e. the fraction of

those who will eventually develop symptoms during their entire infectious period).

Household contacts are assumed to be always traced and isolated. The figure shows that for

$R_0 = 2.5$ control is not possible with isolation and contact tracing, if less than 60% of all

infected persons develop symptoms or are otherwise not detected by the health care

system, even if all other parameters are at the most optimistic values. Other interventions

measures such as social distancing are then needed for containment of the outbreak.

Social distancing and contact tracing

Social distancing in theory can reduce the effective reproduction number to below 1, but

only if non-household contacts are almost reduced to zero. In reality this is not possible,

because there are many non-household contacts which cannot be avoided to keep essential

processes in society going. Therefore social distancing will not be complete and may not be

sufficient for containment. Additional tracing and isolation of household contacts may then

be the additional effort needed to achieve containment. In figure 3a we show for various

levels of social distancing, expressed as proportion reduction of non-household contacts,

what fraction of household- or additionally non-household contacts need to be traced and

isolated to reduce the effective reproduction number. Similarly, figure 3b shows the

increase of the doubling time that can be achieved with these contact tracing measures

depending on level of social distancing. Figure 4a shows for various values of R_0, what are

the critical tracing coverages of household contacts only in order to reach an effective

reproduction number of 1, which is the threshold to containment. If tracing household

contacts is not sufficient, also non-household contacts have to be traced and isolated, for

which the critical coverages are shown in figure 4b. For example, if social distancing leads to

a reduction of 50% of non-household contacts, additional tracing and isolating household

contacts can further reduce the effective reproduction number, but cannot achieve

containment. If then also 20% of non-household contacts can be traced and isolated, the

transition to containment can be made.

Discussion

Our analyses show that rapid diagnosis and isolation of infections based on COVID-19

disease alone cannot control outbreaks of SARS-CoV-2, but that the addition of tracing and

isolation of traced cases could in theory be successful depending on the value of R_0

(Figure 2). In practice, however, the potential for containment will be seriously jeopardized

by various delays and imperfections in the tracing process. Especially delays in diagnosis and

isolation, and the existence of asymptomatic and mild infections that contribute to onward

transmission will make an outbreak uncontrollable. As evidence is mounting that the

proportion of asymptomatic and mild cases is large and leads to substantial numbers of

unascertained cases [22], most countries have now switched to strategies of social

distancing or complete lockdowns. Such measures have proven effective earlier during the

2009 influenza pandemic [23, 24]. However, social distancing can never be complete, as

healthcare workers and doctors have to continue their work, but also personnel of

supermarkets, public transport employees, and others need to have contact outside their

households. Therefore, transmission chains may remain. We find that in a situation where

60% of cases are ascertained, social distancing of non-household contacts fails to contain

the epidemic even if contacts outside the household are reduced by 80%. In that case,

combining the social distancing with tracing and isolation of household contacts may suffice

to bring the balance towards containment. If social distancing is less severe, more intensive

contact tracing and also tracing of non-household contacts is needed (Figures 3 and 4). If

social distancing reduces non-household contacts only by 50%, tracing and isolation also of

non-household contacts is needed for containment. If this is not possible, for example due

to constraints of the public health system, tracing and isolation of household contacts can at

least substantially increase the doubling time of the epidemic (Figure 3b).

In the recent report by Ferguson and colleagues (https://www.imperial.ac.uk/mrc-global-

infectious-disease-analysis/news--wuhan-coronavirus/) it is assumed that non-household

contacts would be reduced by 75% by social distancing [25], and a contact study in the UK

shows that numbers of contacts are reduced by around 73% [21]. Using these estimates, our

model predicts that additional tracing and isolation of infected household contacts would

suffice as an additional measure to achieve containment. Possibly, this could be achieved by

using digital tracing applications as implemented in South Korea, where it has apparently

helped to bring the epidemic under control [3].

Even though the pandemic situation shows that COVID-19 epidemics cannot be contained by

contact tracing and rapid isolation alone, this does not render contact tracing useless. On

the contrary, contact tracing and isolation when used in addition to social distancing, may be

the tool needed to make this mix of strategies successful. Our analyses show that isolation

and contact tracing when combined with social distancing can contribute to reducing the

growth rate and increasing the doubling time of epidemics, thereby buying time, spreading

the number of severe cases out over a longer period of time, and potentially also reducing

the total number of infections (the "final size") [26]. This will lower peak healthcare demand,

alleviate the stress on healthcare systems, and contribute to reducing the burden of disease.

Our analyses of contact tracing add to an earlier study by a more systematic analysis of the

relation between key parameters (transmissibility, fraction asymptomatic, fraction of

contacts traced, diagnosis delays), and by incorporating household versus non-household

contacts [27]. Household contacts are at a higher risk of becoming infected than non-

household contacts as persons in a household will usually have repeated contacts. On the

other hand, our analyses show that household infections contribute less to onward

transmission than non-household infections simply because the numbers of household

contacts are much lower than numbers of other contacts. As a consequence, the

effectiveness in isolating non-household contacts is key for a successful contact tracing

strategy.

A strength of the model is that quantitative information about distributions of the latent and

infectious periods, and the infectivity per day of the infectious period can be incorporated

easily and detailed, such that if new and better data on those quantities emerge, the

analysis can be updated quickly. In particular, the model can incorporate non-standard

distributions based on empirical data (e.g. viral load measurements to quantify

infectiousness per day), should they become available.

A limitation of the analyses presented here is that they apply to a situation in which the

epidemic is described by a branching process and is growing exponentially. This also applies

to another modelling using a (one-type) branching process.[27] Ultimately, as the number of

persons who are or have been infected increases, the number of persons that are still

susceptible will start to dwindle, and epidemic growth will ultimately come to a halt. Hence,

strictly speaking our results apply to the early stages of an epidemic. In fact, even when the

number of infected persons is still relatively small in the early stage of an epidemic it is

possible that exponential growth is not observed, for instance due to local depletion of

susceptible persons in combination with clustering in contact patterns, spatial effects, and

inhomogeneous mixing [28]. However, estimates of the effective reproduction number are

independent of the dynamics and give information about the ability of an intervention to

stop of slow down epidemic spread. Also, at present the epidemics in many countries are

still growing exponentially [19].

In conclusion, our results show that in populations where social distancing is implemented,

isolation and contact tracing can play an essential role in gaining control of the COVID-19

epidemic. On their own, none of these strategies are able to contain COVID-19 for realistic

parameter settings, but in a combined strategy they can just tip the balance towards

containment. These insights provide guidance for policy makers, who will have to decide

when and how to release severe lockdown or social distancing measures, and whether

additional contact tracing and isolation is then a feasible alternative keep a resurging

epidemic at bay.

Contributors

MEK, MvB, and GR conceived the study. MK designed and programmed the model, and

produced output. All authors interpreted the results, contributed to writing the manuscript,

and approved the final version for submission.

Declaration of interests

431   We declare no competing interests.

Tables

Table 1: Disease and transmission related parameters.

Parameter	Distribution/values	Reference
Latent period	1-3 days	**[29]**
Infectious period	10 days	**[29]**
Incubation period	Weibull(3.0, 7.2) mean: 6.4 (day) sd: 2.3 (day)	[10]
Number of household contacts	Poisson(2.15) mean: 2.15 (day^{-1})	https://www.statista.com/statistics/521777/netherlands-average-household-size-by-number-of-residents/
Number of non-household contacts	NB(2, 0.15)[*] mean: 11.3 (day^{-1}) sd: 8.7 (day^{-1})	[20]
Reduction factor of transmissibility for non-household contacts	0.25	[30]
Transmission probability per contact		calibrated such that R_0=2.5

440   Table 2: Parameters related to diagnosis and contact tracing.

Parameter	Value/range	Reference
Delay until diagnosis of index case	Baseline: 1 day; varied from 1 to 10 days	[13] [31]
Time needed to find household contacts	Baseline: 1 day	
Time needed to find other contacts	Baseline: 1 day; varied from 1 to 7 days	[32] [33]
Percentage of household contacts isolated	Baseline: 100%; not varied in this analysis	
Proportion of other contacts isolated	Varied from 0% to 100%	
Percentage symptomatic (also interpretable as ascertainment rate)	Baseline: 100%; varied from 1.5% to 100%	
Percentage reduction contact rate outside the household	Baseline: 0%; varied from 0% to 95%	

References

1. Bernard Stoecklin S, Rolland P, Silue Y, Mailles A, Campese C, Simondon A, et al. First cases of coronavirus disease 2019 (COVID-19) in France: surveillance, investigations and control measures, January 2020. Euro Surveill. 2020;25(6). Epub 2020/02/20. doi: 10.2807/1560-7917.ES.2020.25.6.2000094. PubMed PMID: 32070465.

2. Anderson RM, Heesterbeek H, Klinkenberg D, Hollingsworth TD. How will country-based mitigation measures influence the course of the COVID-19 epidemic? The Lancet. 2020;395(10228):931-4. doi: 10.1016/S0140-6736(20)30567-5.

3. Ferretti L, Wymant C, Kendall M, Zhao L, Nurtay A, Abeler-Dörner L, et al. Quantifying SARS-CoV-2 transmission suggests epidemic control with digital contact tracing. Science. 2020:eabb6936. doi: 10.1126/science.abb6936.

4. Lauer SA, Grantz KH, Bi Q, Jones FK, Zheng Q, Meredith HR, et al. The Incubation Period of Coronavirus Disease 2019 (COVID-19) From Publicly Reported Confirmed Cases: Estimation and Application. Ann Intern Med. 2020. Epub 2020/03/10. doi: 10.7326/M20-0504. PubMed PMID: 32150748.

5. Cheng H-Y, Jian S-W, Liu D-P, Ng T-C, Huang W-T, Lin H-H. High transmissibility of COVID-19 near symptom onset. medRxiv. 2020:2020.03.18.20034561. doi: 10.1101/2020.03.18.20034561.

6. Fraser C, Riley S, Anderson RM, Ferguson NM. Factors that make an infectious disease outbreak controllable. Proc Natl Acad Sci U S A. 2004;101(16):6146-51. Epub 2004/04/09. doi: 10.1073/pnas.0307506101. PubMed PMID: 15071187; PubMed Central PMCID: PMCPMC395937.

7. Klinkenberg D, Fraser C, Heesterbeek H. The effectiveness of contact tracing in emerging epidemics. PLoS One. 2006;1:e12. Epub 2006/12/22. doi: 10.1371/journal.pone.0000012. PubMed PMID: 17183638; PubMed Central PMCID: PMCPMC1762362.

8. Kretzschmar M, van den Hof S, Wallinga J, van Wijngaarden J. Ring vaccination and smallpox control. Emerg Infect Dis. 2004;10(5):832-41. Epub 2004/06/18. doi: 10.3201/eid1005.030419. PubMed PMID: 15200816; PubMed Central PMCID: PMCPMC3323203.

9. Muller J, Kretzschmar M, Dietz K. Contact tracing in stochastic and deterministic epidemic models. Math Biosci. 2000;164(1):39-64. Epub 2000/03/08. doi: 10.1016/s0025-5564(99)00061-9. PubMed PMID: 10704637.

10. Backer JA, Klinkenberg D, Wallinga J. Incubation period of 2019 novel coronavirus (2019-nCoV) infections among travellers from Wuhan, China, 20-28 January 2020. Euro Surveill. 2020;25(5). Epub 2020/02/13. doi: 10.2807/1560-7917.ES.2020.25.5.2000062. PubMed PMID: 32046819.

11. Chen N, Zhou M, Dong X, Qu J, Gong F, Han Y, et al. Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. Lancet. 2020;395(10223):507-13. Epub 2020/02/03. doi: 10.1016/S0140-6736(20)30211-7. PubMed PMID: 32007143.

12. Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. Lancet. 2020;395(10223):497-506. Epub 2020/01/28. doi: 10.1016/S0140-6736(20)30183-5. PubMed PMID: 31986264.

13. Li Q, Guan X, Wu P, Wang X, Zhou L, Tong Y, et al. Early Transmission Dynamics in Wuhan, China, of Novel Coronavirus-Infected Pneumonia. N Engl J Med. 2020. Epub 2020/01/30. doi: 10.1056/NEJMoa2001316. PubMed PMID: 31995857.

14. Riou J, Althaus CL. Pattern of early human-to-human transmission of Wuhan 2019 novel coronavirus (2019-nCoV), December 2019 to January 2020. Euro Surveill. 2020;25(4). Epub 2020/02/06. doi: 10.2807/1560-7917.ES.2020.25.4.2000058. PubMed PMID: 32019669; PubMed Central PMCID: PMCPMC7001239.

15. Rothe C, Schunk M, Sothmann P, Bretzel G, Froeschl G, Wallrauch C, et al. Transmission of 2019-nCoV Infection from an Asymptomatic Contact in Germany. N Engl J Med. 2020. Epub 2020/02/01. doi: 10.1056/NEJMc2001468. PubMed PMID: 32003551.

16. Tang B, Wang X, Li Q, Bragazzi NL, Tang S, Xiao Y, et al. Estimation of the Transmission Risk of the
2019-nCoV and Its Implication for Public Health Interventions. J Clin Med. 2020;9(2). Epub
2020/02/13. doi: 10.3390/jcm9020462. PubMed PMID: 32046137.
17. Keeling MJ, Hollingsworth TD, Read JM. The Efficacy of Contact Tracing for the Containment of
the 2019 Novel Coronavirus (COVID-19). medRxiv. 2020:2020.02.14.20023036. doi:
10.1101/2020.02.14.20023036.
18. Dodd PJ, White PJ, Garnett GP. Notions of synergy for combinations of interventions against
infectious diseases in heterogeneously mixing populations. Mathematical biosciences.
2010;227(2):94-104. Epub 2010/07/01. doi: 10.1016/j.mbs.2010.06.004. PubMed PMID:
20600157.
19. Our world in data. 2020 [1 April 2020]. Available from: https://ourworldindata.org/coronavirus
20. Mossong J, Hens N, Jit M, Beutels P, Auranen K, Mikolajczyk R, et al. Social contacts and mixing
patterns relevant to the spread of infectious diseases. PLoS Med. 2008;5(3):e74. Epub
2008/03/28. doi: 10.1371/journal.pmed.0050074. PubMed PMID: 18366252; PubMed Central
PMCID: PMCPMC2270306.
21. Jarvis CI, van Zandvoort K, Gimma A, Prem K, Klepac P, Rubin GJ, et al. Quantifying the impact of
physical distance measures on the transmissionof COVID-19 in the UK. CMMID Repository2020.
22. Li R, Pei S, Chen B, Song Y, Zhang T, Yang W, et al. Substantial undocumented infection facilitates
the rapid dissemination of novel coronavirus (SARS-CoV2). Science. 2020:eabb3221. doi:
10.1126/science.abb3221.
23. Cauchemez S, Ferguson NM, Wachtel C, Tegnell A, Saour G, Duncan B, et al. Closure of schools
during an influenza pandemic. Lancet Infect Dis. 2009;9(8):473-81. Epub 2009/07/25. doi:
10.1016/S1473-3099(09)70176-8. PubMed PMID: 19628172.
24. Te Beest DE, Birrell PJ, Wallinga J, De Angelis D, van Boven M. Joint modelling of serological and
hospitalization data reveals that high levels of pre-existing immunity and school holidays shaped
the influenza A pandemic of 2009 in the Netherlands. J R Soc Interface. 2015;12(103). Epub
2014/12/30. doi: 10.1098/rsif.2014.1244. PubMed PMID: 25540241; PubMed Central PMCID:
PMCPMC4305427.
25. Mahase E. Covid-19: UK starts social distancing after new model points to 260 000 potential
deaths. BMJ. 2020;368:m1089. doi: 10.1136/bmj.m1089.
26. Diekmann O, Heesterbeek H, Britton T. Mathematical Tools for Understanding Infectious Disease
Dynamics: Princeton University Press; 2013.
27. Hellewell J, Abbott S, Gimma A, Bosse NI, Jarvis CI, Russell TW, et al. Feasibility of controlling
COVID-19 outbreaks by isolation of cases and contacts. Lancet Glob Health. 2020. Epub
2020/03/03. doi: 10.1016/S2214-109X(20)30074-7. PubMed PMID: 32119825.
28. Chowell G, Viboud C, Simonsen L, Moghadas SM. Characterizing the reproduction number of
epidemics with early subexponential growth dynamics. J R Soc Interface. 2016;13(123). Epub
2016/10/07. doi: 10.1098/rsif.2016.0659. PubMed PMID: 27707909; PubMed Central PMCID:
PMCPMC5095223.
29. Bi Q, Wu Y, Mei S, Ye C, Zou X, Zhang Z, et al. Epidemiology and Transmission of COVID-19 in
Shenzhen China: Analysis of 391 cases and 1,286 of their close contacts. medRxiv.
2020:2020.03.03.20028423. doi: 10.1101/2020.03.03.20028423.
30. Chan JF, Yuan S, Kok KH, To KK, Chu H, Yang J, et al. A familial cluster of pneumonia associated
with the 2019 novel coronavirus indicating person-to-person transmission: a study of a family
cluster. Lancet. 2020;395(10223):514-23. Epub 2020/01/28. doi: 10.1016/S0140-6736(20)30154-
9. PubMed PMID: 31986261.
31. Donnelly CA, Ghani AC, Leung GM, Hedley AJ, Fraser C, Riley S, et al. Epidemiological
determinants of spread of causal agent of severe acute respiratory syndrome in Hong Kong.
Lancet. 2003;361(9371):1761-6. Epub 2003/06/05. doi: 10.1016/S0140-6736(03)13410-1.
PubMed PMID: 12781533.
32. Baker MG, Thornley CN, Mills C, Roberts S, Perera S, Peters J, et al. Transmission of pandemic
A/H1N1 2009 influenza on passenger aircraft: retrospective cohort study. BMJ. 2010;340:c2424.

Epub 2010/05/25. doi: 10.1136/bmj.c2424. PubMed PMID: 20495017; PubMed Central PMCID:
PMCPMC2874661.
33.Swaan CM, Appels R, Kretzschmar ME, van Steenbergen JE. Timeliness of contact tracing among
flight passengers for influenza A/H1N1 2009. BMC Infect Dis. 2011;11:355. Epub 2011/12/30. doi:
10.1186/1471-2334-11-355. PubMed PMID: 22204494; PubMed Central PMCID:
PMCPMC3265549.

Figure legends

Figure 1: The cumulative number of cases in an outbreak shown from the time when the number of cases exceeds 100. In (A) the mean and standard deviation of 20 simulation runs is shown for a situation where up to day 28 after the first case was diagnosed, only diagnosis and case isolation is implemented. At day 28 after diagnosis of the first case (arrow) social distancing was implemented, which reduced the number of non-household contacts by 70%. The green lines show the doubling times without intervention (3.6 days), with isolation of diagnosed cases (3.9 days), and with social distancing (13.7 days). In (B) simulations are shown with varying levels of social distancing (contact reductions of 0%, 30%, 50%, 70%, 80%, 90%; colors from brown to red; gray lines are doubling times). In (C) social distancing is implemented at the level of 70% reduction at different times after the diagnosis of the first case (2, 3, 4, 5, and 6 weeks, see arrows). In (D) a scenario is shown (red dotted line) with implementation of social distancing at a contact reduction of 70% (red arrow) at day 28; additional contact tracing and isolation and quarantine of contacts starts at day 49 (orange arrow) after diagnosis of the first case. We contrast this scenario with data from outbreaks in several countries that have implemented social distancing at various levels and times during the outbreak.

Figure 2: The critical tracing coverage needed for control of the outbreak for varying percentages of asymptomatic infections and values of R_0 between 1.5 and 3.5. If more than 40% of cases escape diagnosis because they are asymptomatic or have only mild infections, for R0 = 2.5 the outbreak is not controllable even with our optimistic baseline values for the intervention parameters.

Figure 3: The additional impact of isolation and contact tracing in a population where social

distancing is practiced on (A) the exponential growth rate and (B) the doubling time of the

epidemic. The additional impact of tracing and isolating household contacts only is shown in

the curves from green to yellow (tracing coverage varying from 0% to 100%). If 100% of

household contacts are traced and isolated, additional benefit is gained with tracing also

non-household contact with increasing tracing coverage shown colored blue to pink (varying

from 0% to 100%). In these computations we assumed that 60% of cases are ascertained

and R_0=2.5.

Figure 4: The critical tracing coverages required for containment in a population with social

distancing. In (A) the critical tracing coverage is shown if only household contacts are traced

and isolated. In (B) the tracing of household contacts is assumed to be 100% and the curves

show the additional tracing of non-household contacts needed to achieve containment.

Again 60% of cases are assumed to be ascertained and tracing coverages are shown for

various values of R_0.

Figure 1

605   Figure 2

610   Figure 3

614   Figure 4