Recommendations for standardized management of CML patients in the core epidemic area of COVID-19

——Multi-center survey results in Hubei Province, China

Summary

Background

Since late December 2019, the outbreak of the novel coronavirus disease, COVID-19, that began in Wuhan, has become endemic in China and more than 100 countries and regions in the world. There is no report about the prevalence of COVID-19 in CML patients untill now. We aimed to describe the clinical course, outcomes of CML patients with COVID-19 and prevalence of COVID-19 in CML patients.

Methods

In this multi-center survey, cross-sectional survey, observational study, the clinical data of CML patients with COVID-19 in each center were collected. Simultaneously, an online survey was conducted for information about the CML patients under the management at each center by asking the CML patients to complete a questionnaire,from February 15, 2020 to February 21, 2020. The questionnaire includes demographic data, place of residence, smoking status, CML diagnosis and treatment, comorbidities, combined medications, epidemiological history, symptoms(fever, cough, shortness of breath, etc) during the epidemic. Additional clinical data was collected on respondents suspected or confirmed to have COVID-19. We described and analyzed the prevalence of COVID-19 in CML patients, and focus on the clinical characteristics and outcomes of COVID-19 patients. Data were compared between the CML patients with optimal response and those with non-optimal response. The primary outcome was prevalence of COVID-19 in CML patients, as of Feb 21, 2020. Secondary outcomes included the history of epidemiology of CML patients, the clinical characteristics and outcomes of CML patients with COVID-19 .

Findings

Of 392 respondents, 223(56.9%) were males, and 240(61.2%) were 50 years or younger. Only 10 patients took drugs irregularly due to the influence of the epidemic because of traffic control, pharmacies unable to operate normally, etc. In the history of epidemiology, there were 4 patients with definite contact with COVID-19, of which 3 were remote contact and 1 was close contact. 12 respondents had fever, cough or shortness of breath during the epidemic, 1 case (common type) was confirmed with COVID-19 and cured after treatment. 1 patient was clinically diagnosed and succumbed. 1 of 299 (0.3%) patients with an optimal response was diagnosed with COVID-19. Of the 50 patients who failed to respond to CML treatment or had a poor response, 1 patient (2%) had a clinical diagnosis of COVID-19.

Interpretation

While the 392 CML respondents required regular referrals to hospitals, they did not have much contact with COVID-19 patients during the outbreak. Patients who failed to achieved an optimal response to CML therapy appear more likely to have a symptomatic infection with SARS-CoV-2. Older patients with comorbidities are at increased risk of death.

Funding This work was supported by grants from the National Natural Science Foundation of China(NSFC)(81873440&81700142).

Introduction

Since late December 2019, the outbreak of the 2019 novel coronavirus disease (COVID-19) in Wuhan, China, has been endemic in China and more than 100 countries and regions in the world.[1-7] By March 1, 2020, China had reported a total of 80,026 confirmed cases of novel coronavirus disease, with 2,912 deaths. In Hubei province which is still the core area of the COVID-19 epidemic in China there were 67,103 cases, accounting for 83.9% of total Chinese cases. The number of confirmed cases in South Korea, Iran, Italy, France, Japan and other countries continues to rise.

Some studies have suggest that cancer patients are more susceptible to infection with SARS-CoV-2 than healthy people and have a worse prognosis because their immune systems are suppressed by the effects of the tumors and anti-cancer treatment[8]. But this view is controversial[9, 10].

Chronic myelogenous leukemia (CML) is a neoplastic disease of hematopoietic stem cells that has an annual incidence of 0.4-1.75/100,000 [11-15] and accounts for 15% of adult leukemia. Since imatinib and subsequent generations of newer tyrosine kinase inhibitors (TKIs) have been used with CML, the prognosis of CML patients has improved significantly. In fact, CML patients are currently approaching normal life expectancy. However, as the survival rates of CML patients improves, the prevalence of CML is increasing. The prevalence of CML in France is estimated to reach 18 / 100,000 in 2018 and 24 / 100,000 in 2030[16]. We sought to determine whether CML patients have been as susceptible to SARS-CoV-2 as other cancer patients during the current epidemic, and what their prognosis has been. Standardized management of this large number of patients has become a new challenge for our hematologists. In order to optimize the care of CML patients as the pandemic spreads, it is therefore essential to understand the characteristics of SARS-CoV-2 infections in CML patients and the experience of their management in Hubei province.

Methods:

Study design and participants

When the outbreak started, the 29 centers of the Hubei Anti-Cancer Association Chronic Myeloid Leukemia Standardized Management Collaboration Group adopted comprehensive management measures for CML patients. Various kinds of online consultation, online smart hospitals and online courses were made available to provide patients with information while they remained at home. Individualized treatment schemes were proposed according to the clinical situation of each patient. It was recommended that all patients, especially those with stable conditions, refrain from leaving their homes, and even avoid going to hospitals. Volunteers were organized to deliver drugs to CML patients in their homes in order to ensure their normal treatment would not be interrupted. Patients with COVID-19-related symptoms were promptly diagnosed and treated accordingly.

All patients were diagnosed with 2019 novel coronavirus disease, according to WHO interim guidance[17] or the 5th version of the novel coronavirus infection pneumonia diagnosis and treatment (The National health commission, People's Republic of China)[18] . Laboratory confirmation of SARS-CoV-2 infection was performed by the local health authority. Identification

of COVID-19 patients was achieved by reviewing and analysing admission logs and histories from all available electronic medical records and patient care resources. Efficacy of CML was assessed according to the 2020 ELN guidelines[19]. The investigation was approved by the ethics committee of Union hospital, Tongji Medical college, Huazhong University of science and Technology.

Data collection

From February 15, 2020 to February 21, 2020, the clinical data of CML patients with COVID-19 in each center were collected. Simultaneously, an online survey was conducted for information about the CML patients under the management at each center by asking the CML patients to complete a questionnaire. A total of 413 questionnaires were collected, of which 392 were valid, but 21 of these were excluded because the respondents did not live in Hubei province during the outbreak. The questionnaire asked about general demographic characteristics, place of residence, smoking status, CML diagnosis and treatment, comorbidities, combined medications, epidemiological history, and symptoms of fever, cough, shortness of breath, etc. during the epidemic. Additional clinical data was collected on respondents suspected or confirmed to have COVID-19. Any missing or uncertain records were collected and clarified through direct communication with involved health-care providers and their families.

Outcomes

The primary outcome was prevalence of COVID-19 in CML patients, as of Feb 21, 2020. Secondary outcomes included the history of epidemiology of CML patients, the clinical characteristics and outcomes of CML patients with COVID-19.

Statistical analysis

The aim of this study is to report prevalence of COVID-19 in CML patients. COVID-19 infection in CML patients with different efficacy was compared. We also report the clinical courses and clinical outcomes of the COVID-19 patients. There were, therefore, no formal hypotheses being implemented to drive the sample size calculation and we included the maximum number of patients who met the inclusion criteria.

Results

A total of 413 questionnaires were collected, of which 392 were valid. Among them, 21

questionnaires were excluded because the respondents did not live in Hubei province during the outbreak. The following is the general situation of the respondents. There were 223 males, accounting for 56.9%, and 61.2% of the respondents were 50 years or younger. The age distribution chart is shown in Table 1. The other details are shown in Table 2. The epidemiological history is shown in Table 3. Specific data of clinical diagnosis and confirmed patients are shown in table 4.

Table 1(the age distribution chart)

General demographic characteristics	Count
Gender	
Male	223
Female	169
Age(years)	
Less than 18 years of age	16
18-35	82
36-50	142
51-65	121

66-75	24
More than 75 years old	7
Diagnose CML time	
Less than 3 months	9
3-12 months	56
1-3 years	136
3-10 years	139
More than 10 years	52
Current TKI status	
Dasatinib	82
Imatinib	259
Nilotinib	37
Else	14
Efficacy evaluation of CML	
Optimum response	299
Warnings	20
Failure	30
Not known	43
Comorbidities	
Hypertension	26
Other Malignancies	10
Kidney diseases	9
Diabetes	14
Else	53
None	280
Combined medications	
Corticosteroids	10

Antihypertensive drugs （Angiotensin-Converting Enzyme Inhibitors）	32
Other anti-tumor drugs	3
Else	58
None	289
Smoking history	
Not have	309
Have	83
Have you forgotten to take medicine in the past 2 weeks	
No	315
Yes	77 （ 10 patients took drugs irregularly due to the influence of the epidemic）

Table 2（general demographic characteristics）

The epidemiological history	Count
Residence	
Wuhan	82
Hubei province (except wuhan)	310
contact with infected person	
Yes	4（3 were in remote contact and 1 was in close contact）
No	330
not known	58
Cluster onset	
yes	1 （3 people confirmed of COVID-19 in her family. This respondent did not live with the COVID-19 patient, the respondent does not have an infection at present）
no	391

Table 3 (the epidemiological histories)

clinical data of the cases			
		Case 1	Case 2
Gender		female	male
age		89 years	47 years
Diagnose CML time		11 years	3 years
Current TKI status		Imatinib (Gleevec)	HQP1351 clinical trials
Efficacy evaluation of CML		failure	Optimal response
Comorbidities		coronary heart disease	None
Combined medications		Angiotensin-Converting Enzyme Inhibitors	None
Residence		Wuhan	Shiyan, Hubei province
The epidemiological history		Yes (exposure to relevant environment)	Yes (exposure to relevant environment)
Other family members affected		No	No
Signs and symptoms	fever	No	Yes
	cough	Yes	Yes
	dyspnea	Yes	Yes
	pharyngalgia	No	Yes
	diarrhea	No	No
Laboratory characteristics	White blood cell count ($\times 10^9$ cells per L)	5.13	4.0
	Lymphocyte count ($\times 10^9$ cells per L)	0.17(Lymphopenia)	0.78(Lymphopenia)

	Elevated ALT (>45 U/L) or AST (>35 U/L)	Yes	No
	Cardiac enzyme	Troponin positive	Normal
	Creatinine	elevate （139umol/l）	Normal
	Confirmatory test done (SARS-CoV-2 quantitative RT-PCR)	No	Yes
Typical pneumonia on the chest CT scan		Yes	No
clinical classification		Critical type	Common type
Outcome		death	Cure

Table 4 (clinical data of the cases)

Discussion

While the 392 CML respondents required regular referrals to hospitals, they did not have much contact with COVID-19 patients during the outbreak, mainly due to the government policy of prevention and control. The survey found that 12 respondents had fever, cough or shortness of breath during the epidemic period, and 9 of them did not go to the hospital because their symptoms were mild or remitted with oral medication. Three people went to the hospital for treatment. After chest CT scan and SARS-CoV-2 quantitative RT-PCR testing, one of these three patients was confirmed to have of COVID-19 infection and was treated and cured. COVID-19 infection was excluded in a second patient. The third patient was clinically diagnosed based on the 5th version of the novel coronavirus infection pneumonia diagnosis and treatment (National Health Commission of the People's Republic of China) that was used at the time. This patient, who was clinically diagnosed and treated as a confirmed case, was an 89-year-old female with coronary heart disease and long-term ACEI treatment. She had been treated with imatinib, but

the response was evaluated as failure. After initially having a cough, the patient quickly developed dyspnea that was not relieved with oxygen therapy at home. She was admitted to the emergency department where she was found to have extreme lymphocytopenia, respiratory failure, renal insufficiency, and myocardial damage. Her condition rapidly deteriorated and she succumbed after three days of hospitalization. This is consistent with previous reports that elderly patients with comorbidities have a poor prognosis and high mortality[20-22], and the degree of lymphopenia is related to the severity of the disease.

Another CML patient with a confirmed COVID-19 infection was a 47-year-old male who had failed CML treatment, had the T315I mutation and was participating in an ongoing clinical trial of an alternative CML treatment (HQP1351). Although he obtained an optimal MMR response, he was not taking TKIs during the outbreak due to side effects.   He presented with cough and fever and was found to be infected with SARS-CoV-2, but a chest CT scan did not show pneumonia and he was cured and discharged after treatment. The specific data of the clinical diagnosis and confirmed patients are shown in table 4.

We then analyzed SARS-CoV-2 infections in patients based on their response to anti-CML therapy (see Table 5 for details), and found that only 1 of 299 (0.3%) patients with an optimal response was diagnosed with COVID-19. Of the 50 patients who failed to respond to CML treatment or had a poor response, 1 patient (2%) had a clinical diagnosis of COVID-19.   Thus patients who failed to achieved an optimal response to CML therapy appear more likely to have a symptomatic infection with SARS-CoV-2.   This tendency requires further study, but there are two possibilities that might explain it. First, an optimal response to TKI treatment may be associated with immune recovery.   CML patients exhibit selective depletion of effector T reg cells (eT reg cells)[23, 24], while TKIs increase the number of natural killer cells (NK), NK-LGL and T-LGLs cells[25], which play a role in regulating immunity.   Second, previous studies have reported that imatinib and other TKI drugs have antiviral activity *in vitro* against Middle East respiratory syndrome coronavirus (MERS-COV) and severe acute respiratory syndrome coronavirus (SARS-COV)[26-28].   It will be interesting to see if TKIs also have activity against anti-SARS-CoV-2.

We analyzed the 82 respondents living in Wuhan and found that there was 1 clinically diagnosed case (1.2%).   Of the Japanese citizens evacuated from Wuhan, 8/565 (1.4%) had a

confirmed infection with COVID-19[29]. Among 310 CML living patients in Hubei province outside of Wuhan, 1 case of infection (0.32%) was diagnosed.

In summary, through the Hubei Anti-Cancer Association Chronic Myeloid Leukemia Standardized Management Collaboration Group, we conducted online consultations, online smart hospitals and online courses to provide information, diagnosis and treatment to patients during the epidemic, and organized volunteers to deliver drugs to CML patients at their homes. The sample size of the study was relatively small and should be expanded to more accurately compare the prevalence of the infection in the different groups. Nevertheless, our findings suggest that standardized management to obtain the best clinical response reduces the risk of SARS-CoV-2 infection and improves the prognosis of those who are infected. CML is a chronic hematologic tumor, and comprehensive measures should be taken to delay and reduce the risk of infection by restricting patient exposure to situations with possible transmission. During the epidemic, special attention should be focused on those CML patients who are older, have co-morbidities and whose response to CML treatment is non-optimal.

	Optimal response（299 persons）	Warnings or failure（50 persons）
Forgotten to take medicine in the past 2 weeks	63	6
Contact with infected person	4	0
Have Comorbidities	82	13
Diagnosed COVID-19	1	1

Table 5 （Data were compared between the CML patients with optimal response and those with non-optimal response）

1. 2019-nCoV acute respiratory disease, Australia: Epidemiology Report 1 (Reporting week 26 January - 1 February 2020). *Communicable diseases intelligence (2018)* 2020; **44**.
2. A Bastola, R Sah, AJ Rodriguez-Morales, et al. The first 2019 novel coronavirus case in Nepal. *The Lancet Infectious diseases* 2020.
3. JY Kim, PG Choe, Y Oh, et al. The First Case of 2019 Novel Coronavirus Pneumonia Imported into Korea from Wuhan, China: Implication for Infection Prevention and Control Measures. *Journal of Korean medical science* 2020; **35**(5): e61.

4.   LT Phan, TV Nguyen, QC Luong, et al. Importation and Human-to-Human Transmission of a Novel Coronavirus in Vietnam. *The New England journal of medicine* 2020.

5.   M Ki, TFF -nCoV. Epidemiologic characteristics of early cases with 2019 novel coronavirus (2019-nCoV) disease in Republic of Korea. *Epidemiology and health* 2020: e2020007.

6.   ML Holshue, C DeBolt, S Lindquist, et al. First Case of 2019 Novel Coronavirus in the United States. *The New England journal of medicine* 2020.

7.   WK Silverstein, L Stroud, GE Cleghorn, JA Leis. First imported case of 2019 novel coronavirus in Canada, presenting as mild pneumonia. *Lancet (London, England)* 2020.

8.   W Liang, W Guan, R Chen, et al. Cancer patients in SARS-CoV-2 infection: a nationwide analysis in China. *The Lancet Oncology* 2020.

9.   H Wang, L Zhang. Risk of COVID-19 for patients with cancer. *The Lancet Oncology* 2020.

10. Y Xia, R Jin, J Zhao, W Li, H Shen. Risk of COVID-19 for cancer patients. *The Lancet Oncology* 2020.

11. Y Chen, H Wang, H Kantarjian, J Cortes. Trends in chronic myeloid leukemia incidence and survival in the United States from 1975 to 2009. *Leukemia & lymphoma* 2013; **54**(7): 1411-7.

12. SJ Harrison, PR Johnson, TL Holyoake. The Scotland Leukaemia Registry audit of incidence, diagnosis and clinical management of new patients with chronic myeloid leukaemia in 1999 and 2000. *Scottish medical journal* 2004; **49**(3): 87-90.

13. CS Chang, K Lee, YH Yang, MT Lin, CN Hsu. Estimation of CML incidence: disagreement between national cancer registry and health claims data system in Taiwan. *Leukemia research* 2011; **35**(5): e53-4.

14. M Höglund, F Sandin, K Hellström, et al. Tyrosine kinase inhibitor usage, treatment outcome, and prognostic scores in CML: report from the population-based Swedish CML registry. *Blood* 2013; **122**(7): 1284-92.

15. M Höglund, F Sandin, B Simonsson. Epidemiology of chronic myeloid leukaemia: an update. *Annals of hematology* 2015: S241-7.

16. M Delord, S Foulon, JM Cayuela, P Rousselot, J Bonastre. The rising prevalence of chronic myeloid leukemia in France. *Leukemia research* 2018; **69**: 94-9.

17.WHO. Clinical management of severe acute respiratory infection when novel coronavirus (nCoV) infection is suspected. Jan 11, 2020. (accessed Feb 8, 2020). *https://www.who.int/publications-detail/clinical-management-of severe-acute-respiratory-infection-when-novel-coronavirus-(ncov)- infection-is-suspected (accessed Feb 8, 2020).*

18.National Health Commission of the People's Republic of China. The 5th version of the novel coronavirus infection pneumonia diagnosis and treatment. *http://www.nhc.gov.cn/yzygj/s7653p/202002/3b09b894ac9b4204a79db5b8912d4440.shtml*

19.A Hochhaus ,M Baccarani ,RT Silver, et al. European LeukemiaNet 2020 recommendations for treating chronic myeloid leukemia. *Leukemia* 2020.

20. [The epidemiological characteristics of an outbreak of 2019 novel coronavirus diseases (COVID-19) in China]. *Zhonghua liu xing bing xue za zhi = Zhonghua liuxingbingxue zazhi* 2020; **41**(2): 145-51.

21. [An update on the epidemiological characteristics of novel coronavirus pneumonia （COVID-19）]. *Zhonghua liu xing bing xue za zhi = Zhonghua liuxingbingxue zazhi* 2020; **41**(2): 139-44.

22. N Chen, M Zhou, X Dong, et al. Epidemiological and clinical characteristics of 99 cases of

2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. *Lancet (London, England)* 2020.

23. A Tanaka, H Nishikawa, S Noguchi, et al. Tyrosine kinase inhibitor imatinib augments tumor immunity by depleting effector regulatory T cells. *The Journal of experimental medicine* 2020; **217**(2).

24. H Ureshino, T Shindo, S Kimura. Role of cancer immunology in chronic myelogenous leukemia. *Leukemia research* 2020; **88**: 106273.

25. N Climent, M Plana. Immunomodulatory Activity of Tyrosine Kinase Inhibitors to Elicit Cytotoxicity Against Cancer and Viral Infection. *Frontiers in pharmacology* 2019; **10**: 1232.

26. CM Coleman, JM Sisk, RM Mingo, EA Nelson, JM White, MB Frieman. Abelson Kinase Inhibitors Are Potent Inhibitors of Severe Acute Respiratory Syndrome Coronavirus and Middle East Respiratory Syndrome Coronavirus Fusion. *Journal of virology* 2016; **90**(19): 8924-33.

27. J Dyall, CM Coleman, BJ Hart, et al. Repurposing of clinically developed drugs for treatment of Middle East respiratory syndrome coronavirus infection. *Antimicrobial agents and chemotherapy* 2014; **58**(8): 4885-93.

28. JM Sisk, MB Frieman, CE Machamer. Coronavirus S protein-induced fusion is blocked prior to hemifusion by Abl kinase inhibitors. *The Journal of general virology* 2018; **99**(5): 619-30.

29. H Nishiura, T Kobayashi, Y Yang, et al. The Rate of Underascertainment of Novel Coronavirus (2019-nCoV) Infection: Estimation Using Japanese Passengers Data on Evacuation Flights. *Journal of clinical medicine* 2020; **9**(2).