

Assessment of menstrual hygiene practice and associated factor among High school female students in Harar Eastern Ethiopia 2019

Astawus Alemayehu Felleke * and Abdi Ahmed Gerada

* Corresponding Author

Full Name: Astawus Alemayehu Felleke

Affiliation: Instructor at Public Health Department at Harar Health Science College, Harar Ethiopia.

Email: astawusalemayehu@gmail.com

Co-author

Full Name: Abdi Ahmed Gerada

Affiliation: Instructor at Public Health Department in Harar Health Science College, Harar Ethiopia.

Email: abcdef356@live.com

Abstract

Background: Menstruation is a visible manifestation of cyclic uterine bleeding as a result of the interaction of different hormones. During menstruation, girls face gender problems. These

are, early marriage, premature childbirth, higher infant mortality and potential vaginal infections resulting in infertility.

Objective: To assess the level of menstrual hygiene practice and associated factors among high school female students in Harar town Eastern, Ethiopia, 2019. A cross-sectional quantitative study were employed from April 02-05/2019. a minimum sample size of 301 Data were collected using a self-administered structured questionnaire.

Result: According to study 168(55.8%) had good practiced the rest had no. students who have no pocket money from family(AOR 0.36:95% CI,0.309,0.989) was 64% less likely poor practiced than students who have permanent pocket money, students who have no educated father(AOR 0.39:95% CI, 0.180,0.872) was 61% less likely poor practiced than who have educated father and students who had not Freely discuss with parents(AOR 0.45:95% CI,0.22,0.903) was 55% less likely poor practiced than who had Freely discuss with parents.

Conclusion: According to this study result, The major female students of high school in Harar town had good knowledge about menstrual hygiene practice. Also, the practice of good menstrual hygiene was more among students who live in the urban 0.27 times than students who live in the rural area, good menstrual hygiene practice was more among students who have permanent pocket money from family 0.36 times than students who have no permanent pocket money from family, good menstrual hygiene practice was more among students who have educated father 0.4 times than students who have no educated father and students who had Freely discuss about menstruation with parents had more good menstrual hygiene practice 0.45 times than who had not Freely discuss about menstruation with parents.

1. Introduction

In 2011 globally, around fifty two percent of the female population is reproductive age. Most of these women and girls menstruate each month for between two and seven days. Girls typically start to menstruate during puberty or adolescence. [1]

MHM is the absorption menstrual blood onto clean material which can be changed in privacy, including the availability of soap and clean water, to clean reusable sanitary materials and body, as well as a suitable place of disposal. [3].

A girl needs to practice a good personal hygiene during their periods and personal hygiene starts from the selection of good sanitary materials, correctly usage, proper disposal, body cleanliness, diet, etc.[4].

MHM practice is hold back by practical, economic, social and cultural factors. At the time of menstruation, adolescent women or girls are faced different challenges related to the management of menstrual hygiene in secondary and preparatory schools. [6]

Globally 20% girls miss from school due to their monthly period, and one in ten will drop out altogether. 39% and 8% of girls use water but no soap for washing their menstrual protection in India and Afghanistan respectively. Girls' participation and performance affected at school during menstruation. In various studies from Africa, the Middle East, and Asia, up to 95% of girls reported missing school because of their periods. [7]

MHP of girls are also different in different countries . In Iran 51% of girls don't take a bath for 8 days after the onset of their period, In Afghanistan 84% of girls never wash their genital areas. In Malawi 30% of girls don't use the latrine during menstruation. Their is 20% girls similarly observed in India communities and 11% and 60% girls only change their menstrual cloths once a day in Ethiopia and India respectively. [8]

In Uganda menstruation is a taboo, embarrassing issue and shameful. It was manifested by girls' avoiding eye contact, nervous laughs and the fact that they often turned their faces towards the floor when talking. [11]

A comparative study done in Udupi Taluk, India showed that only 197 (35.82%) and 83 (33.27%) of rural and urban participants respectively had awareness about menstruation prior to menarche. [12]

According to a cross-sectional study done in India, menstrual hygiene practices were different in 19 years aged girls and above when compared to younger ages. There was significant associations

between type of pads/napkin usage and the age of the participants, Proportion older girls which use disposable pads were highly increased than the young girls, Also between age and place of storage, changing pads at night; amount of pads used per day found to be significantly associated, A higher proportion of older girls practiced changing pads during nights and used more pads per day more than the young girls. [13]

A study conducted in Mehalmeda northern Ethiopia indicate that, all respondents use menstrual sanitary pad, Three hundred forty nine (70.9%) use commercially made sanitary pad, 128(26.01%) homemade sanitary pad and 17(3.4%) use other methods such as underwear and sponge. One hundred forty-six (29.7%) of the participants reuse sanitary. [16]

A study conducted in Northeast Ethiopia indicated that from 455 respondents who had their menarche, 389 (85.49%) of them did not change menstrual soak up at school and the main reasons were absence of separate toilet for female students is 45.50%, fear other students is 39.07%, lack of water sources is 18.77% and shortage of sanitary napkins/pads 15.17%. [17]

A study done in, Northeast Ethiopia, most girls, 478 (86.75%) had heard about menstruation before they had menarche and leading sources of information were their sisters (42.68%), mothers (38.28%), friends(29.50%) and teachers(13.39%), In this study, about 319 (57.89%) of the girls knew correctly that menstrual cycle is a physiologic process and they reported different feelings during their menarche such as feel shame among 195 (42.86%) of girls, 144 (31.65%) and 77 (16.92%)were upset/tensioned and irritated or disgusted respectively. The mean score of the schoolgirls' knowledge of menstruation hygiene management practice is found to be 6.95 ± 2.03 on a scale of 1–12. [17]

A study in Nigeria Amassoma Community, Bayelsa State revealed that 64.2% of the participants were aware of menstruation and their mothers is important source of information, while friends and television also contributed to their information. [20]

According to a study conducted in Wegera, Northwest Ethiopia In general out of 423 study participants one hundred twenty-six (29.8%) of the respondents practiced good menstrual hygiene whereas 297(70.2%) practiced poor menstrual hygiene. [25]

Knowledge of menstrual hygiene was positively associated with practice on menstrual hygiene. This finding was consistent with other studies. A Cross-Sectional Study among Adolescents Female in An Urban Slum of Mumbai, a significant association was observed between having

good/fair knowledge and good practices ($p < 0.01$). Similarly finding there is association between knowledge of menstruation and menstrual hygiene practice ($p < 0.05$). [27]

The objective of this study is to assess menstrual hygiene practice of female high school students and associated factors, to prevent problem which related to poor menstrual hygiene practice. Due to its direct effect on -school absenteeism, increased susceptibility to RTIs, infections of the perineum, increased potential risk of contracting blood-borne diseases such as HIV or HBV and gender discrepancy.

2. Methods and materials

2.1 Study areas and study design

The study was conducted in Harari regional states, Harar town which is located in eastern Ethiopia. The regional administrative Harar town found 516 k/m away from the Addis Ababa city with an elevation of 1830 meters above the sea level. The total population of Harare regional state is estimated to be 232,000. From the total population, 62.6% live in Harar town and the rest 37.4 % live in rural areas. The regional state has six urban and three rural Districts (Woreda) and further the region is subdivided into 19 urban and 17 rural Keble's. The major ethnic groups in the region are Oromo, Amhara and Harari; and the main religions followed are Islam and Ethiopian Orthodox. According to Harari Educational Bureau 2011 E.C (2019 G.C) report, there are 8 governmental secondary schools and 9 non-governmental secondary schools. Among 8 governmental secondary schools, 6 secondary schools are found in an urban area the rest 2 secondary schools found in rural areas. All 8 non-governmental secondary schools are found in urban areas. The total numbers of female and male students in governmental secondary schools are 2510 and 3671 respectively. The total numbers of female and male students in non-governmental secondary schools are 724 and 645 respectively .total female and male students in both secondary schools totally 7550.

Out of 7550 male and female secondary school students, Only 1887 female students learn grades 9 and 10. A school-based cross-sectional study design using a quantitative method was conducted from April 02 – 05/2019.

2.2 Sample size and sampling procedure

Sample size were determined by using single population proportion formula with the assumptions of 95% confidence interval, marginal error of ($d = 5\%$), and by taking 29.8% considering the

proportion of students who practiced good menstrual hygiene from studies done in the country, sample size was 321. Since the source population was less than 10,000 populations the correction formula was used and considering 10 % non-response rate the final sample size becomes 301. The sample size were proportionally allocated accordingly, A simple random and systematic sampling was done to select girls on both governmental and private secondary school girls who were starting menstruating at least one time before data collection.

2.3. Data collection tools and procedures

The data were collected by self-administered technique, using a questionnaire adapted from relevant literature of similar studies. It consists of basic socio-demographic characteristics, source of information or knowledge and awareness about menstruation, menstrual hygiene practice and associated factors and the questionnaire was prepared in the English version and was translated back to Amharic and Afan Oromo (local language) to check its consistency. Respondents those score 6-12 points and 0-5 points under practice were declared as having good and poor practices respectively.

2.4. Method of data processing and analysis

After checking for completeness and consistency of data, data were coded and entered into SPSS version 20 for analysis. Descriptive and analytical statistics were applied. Bivariate analysis was used to examine the association between dependent and independent variables. All variables with $p \leq 0.20$ in the bivariate analysis were fitted into the multivariate logistic regression model to identify factors associated with the menstrual hygienic practice. P value ≤ 0.05 were considered as a level of significance.

2.5. Operational definition

The students' practices were scored by using a scoring system adapted from a past study.

The students' knowledge and practices were scored by using a scoring system adapted from a past study. Each correct response under knowledge was attracted 1(one) point, whereas any wrong or don't know the answer was attracted no (0) mark. Respondents that scored 0-4 points under knowledge were declared as having poor knowledge; whereas those that scored 5-8 were declared as having good knowledge. Similarly, those students that score 6-12 points and 0-5 points under practice were declared as having good and poor practices respectively. As similarly as described by a previous study (25)

2.6. Ethical consideration

Ethical clearance was obtained from Harar health science college institutional Ethical review board Official letters will receive from the department of public health and was submitted to Harari regional state of Education Bureau, the respective directors of the high schools ((SOS, Zoom in, Abadiir secondary school, Hamaresa secondary school, Harar senior school and Bethlehem Catholic school). The purposes of the study will be explained and informed consent was secured from each participant. Confidentiality was maintained throughout the study. Participant’s involvement in the study was on a voluntary basis.

3. Result

3.1. Socio-demographic characteristics of the respondents

Three hundred one female students participated in the study and Majority, 262(87%) and 39(13%) were from public and private schools respectively. The mean age of the study participants was 15.87 years. Among the participants 124 (41.2%) of them were in the age group of 14-15 years, 160(53.2%) of them were in the age group of 16-17 years and the rest 17(5.6%) is \geq 18 years. A majority, 279(92.7%) of the participants had started their menstruation in the age range of 10-14 years. Most, 142 (47.2%) of them were from the Oromo ethnic group, 102(33.9%) from Amara, 19(6.3%) from Tigre, 10(3.3%) from Hararii and 28(9.3%) were from other and also 125 (41.5%) were Orthodox Christians, 135(44.9%) from Muslim, 39(13%) from protestant and 2(0.7%) were from other religion .210 (69.8%) of the participants were from urban areas and the rest from rural. Mothers of schoolgirls 124 (41.2%) were self-employed, 96(31.9%) were housewives, 53 (17.6%) was a governmental employee and 21(7%) were a private employee. Majority, 192(63.8%) of the respondents families earned $>$ 2000 ETB. As shown in the below “Table 1”.

Table 1, Sociodemographic characteristics of female students in Harar town Ethiopia, 2019.

Variables	Category	Number	%
Age	14 - 15 years	124	41.2
	16 - 17 years	160	53.2
	\geq 18 years	17	5.6
Age at menarche	10 -14 years	279	92.7
	15 -19 years	22	7.3
Residence	Urban	210	69.8
	Rural	91	30.2
Religion	Orthodox	125	41.5
	Muslim	135	44.9
	Other	41	13.7
Ethnicity	Amhara	102	33.9
	Oromoo	142	47.2
	Tigre	19	6.3

	Hararii	10	3.3
	other	28	9.3
Grade level	Grade 9	173	57.5
	Grade 10	128	42.5
Father's educational status	Can't read and write	47	15.6
	primary school	50	16.6
	high school	86	28.6
	college diploma	30	10.0
	college/university degree	68	22.6
	other	20	6.6
Mothers' educational status	Can't read and write	89	29.6
	primary school	67	22.3
	high school	77	25.6
	college diploma	30	10.0
	college/university degree	36	12.0
	other	2	0.7
Father's occupation	government employee	103	34.2
	private employee	49	16.3
	self-employee	126	41.9
	other	23	7.6
Mother's occupation	governmental employee	53	17.6
	private employee	21	7.0
	self-employed	124	41.2
	housewife	96	31.9
	other	7	2.3
Family monthly income	<500 birr	20	6.6
	500 - 1000 birr	28	9.3
	1000 - 1500 birr	18	6.0
	1500 - 2000 birr	43	14.3
	> 2000 birr	192	63.8
School type	Public	262	87.0
	Private	39	13.0
Permanent money pocket	Yes	194	64.5
	No	107	35.5

3.2. Source of information about menstrual hygiene practice

Among 301 study participants, 276(91.7%) were heard about menstruation before menarche and The main sources of information about menstruation were, 163(54.2%), 70 (23.3%), 13(4.3%), 15(5%) and 14(5%) from mothers, teacher, health personnel, TV and others respectively. As showed in the above (figs 3)

3.3 Knowledge on menstruation and menstrual hygiene

According to the data obtained from the participants, 184 (61.1%) study participants had good knowledge about menstrual hygiene, 117(38.9%) respondents had poor menstrual hygiene knowledge Majority of the girls 275(91.4%) knew about menstruation before they had menarche. The dominant source of information and advice for the girls were mothers 163(54.2%). The main material known by 275(91.4%) respondents were available in the market. Out of 301 study participants, 264 (87.7 %) of girls knew that the cause of menstruation was a physiological

process, 6 (2 %) of the girls believed that it was a curse from God. More than half, 180 (59.8 %) of the respondents knew the origin of the menstrual blood was from the uterus. 234 (77.7%) knew that the exact duration of normal menses is 2 to 7 days but was the other had no in addition to this only 200(66.4%) of the study participants knew correctly that normal menstrual cycles vary between 20 to 35 days. as shown in the below “Table 2”.

Table 2, Knowledge regarding menstruation and menstrual hygiene practice among female secondary school students in Harar town, 2019.

Respondents characteristics	Frequency	Percent (%)
At what age girl usually get her first period?		
11 - 16 years	299	99.3
> 16 years	2	0.7
How long is the normal menstrual bleeding duration?		
< 2 days	22	7.3
2 - 7 days	234	77.7
> 7 days	28	9.3
I don't know	17	5.6
What is the normal duration of the menstrual cycle?		
< 20 days	50	16.6
20 - 35 days	200	66.4
> 35 days	23	7.6
I don't know	28	9.3
Do you know sanitary pads in the market?		
Yes	275	91.4
No	26	8.6
Cause of menstruation		
Physiological Process	264	87.7
It Caused By Sin	25	8.3
Is Curse Of God	6	2.0
I Don't Know	6	2.0
Source of bleeding during menstruation		
Vagina	84	27.9
Urinary Bladder	16	5.3
Uterus	180	59.8
I Don't Know	21	7.0
Knowledge about Menstrual hygiene practice		
Good knowledge	184	61.1
Poor knowledge	117	38.9

Less than half of the study participants 127(42.2%) responded that menstruation is a secrete issue of whom, 30 (10%) were due to deep-rooted culture in the society, 19 (6.3%) were due to believes

and customs of the society, 66(21.9%) were both reasons and 12 (4%) were due to other reasons. The majority of participant 273(90%) can go to school during menstruation the rest girls cannot go to school during menstruation.

3.4 Menstrual hygiene practice of respondents

According to the data obtained from the participants, 168 (55.8%) of the respondents had good practiced and whereas 133(44.2%) practiced poor menstrual hygiene. All of the study participants use sanitary pads among them 219(72.8%) use disposable sanitary materials, 35(11.6%) use reusable sanitary pads,14(4.7%) use disposable piece of rags and 28(9.3%)use paper/toilet paper/underwear and 5(1.7%) uses other material. Out of 301 participants, 277(92%)girls were washing their genitalia during their menstruation were 24(8%) girls had no wash their genitals during menstruation. Among these females 84(27.9%) wash twice per day,106(35.2%) wash three times per day and 87(28.9%) wash greater than four times per day. Among this girl, 126(41.9%) girls used both soap and water,149(49.5%) girls use only water and 2(0.7%) use other methods. Out of 301 study participants, 296 (98.3%) girls changed their menstrual hygiene materials were 5(1.7%) girls had no change their menstrual hygiene materials. Among this girl who changes their sanitary material, 272(90.4%) girls change their menstrual hygiene material 1-6 times per day and 24(8%) girls change their menstrual hygiene material 7 times per day. 163 (54.2%) throw in the latrine,33(11%) throw used pad in the open field,98(32.6%) throw used material wrap in paper and put in the bin and 7(2.3%) throw in another place. Among 301 study participants, 99 (32.9%) girls store reusable material in drawers,103(34.2%) girls store reusable material in dress cabinet,41(13.6%) girls store reusable material in bathroom and 29(9.6%) girls store reusable material with routine clothe and Only 176(58.5%) respondents take bath during menstruation the rest had no take bath. Among this, 151(50.2%) girls take bath < 2 times per day and the rest 25(8.3%) take bath > 2 times per day.

3.5. Factors associated with menstrual hygiene practice

The students who live in the rural area (AOR 0.269:95% CI, 0.125, 0.577) was 74% less likely poor menstrual hygiene practice than students who live in the urban, students who have no permanent pocket money from family (AOR 0.36:95% CI, 0.309, 0.989) was 64% less likely poor menstrual hygiene practice than students who have permanent pocket money from family, students who have no educated father (AOR 0.39:95% CI, 0.180,0.872) was 61% less likely poor menstrual

hygiene practiced than students who have educated father and students who had not Freely discuss about menstruation with parents (AOR 0.45:95% CI, 0.22,0.903) was 55% less likely poor menstrual hygiene practiced than students who had Freely discuss about menstruation with parents as showed in the below “Table 3”.

Table 3, Bivariate and multivariate logistic regression analysis for Factors associated with menstrual hygiene practice of female students, harar town in 2019.

Independent factors	Menstrual practice		COR(95%CI)	P- value	AOR (95% CI)	P- value
	Good	Poor				
Grade level						
Grade 9	105 (60.7%)	68 (39.3%)	1.5(1.00,2.52)	0.048	1.6(0.94,2.87)	0.081
Grade 10	63 (49.2%)	65 (50.8%)	1	1		
What Types of school do you learn?						
Public	143 (88.3%)	119(45.41%)	0.67(0.33,1.35)	0.266	1.8(0.77,4.18)	0.170
Private	25 (9.54%)	14 (5.34%)	1	0.082		
Do your parents provide Permanent pocket money?						
Yes	124(63.91%)	70 (36.08%)	1	1		
No	44(41.12%)	63 (58.9%)	0.39(0.24,0.64)	0.001	0.36(0.19,0.653)	0.001
Where do you live						
Urban	138 (65.7%)	72 (34.3%)	1	1		
Rural	30 (33%)	61 (67%)	0.25(0.15,0.43)	0.001	0.26(0.12,0.577)	0.001
How old are you when you start menstruation						
10 -14 years	152 (54.5%)	127 (45.5%)	0.44(0.17,1.18)	0.105	0.87(0.27,2.744)	0.820
15 -19 years	16 (72.7%)	6 (27.3%)	1	1		
Family monthly income						
<500 birr	9 (45%)	11(55%)	0.75(0.29,1.89)	0.547	0.79(0.22,2.821)	0.724
500 - 1000 birr	20 (71.4%)	8 (28.6%)	2.30(0.96,5.47)	0.060	1.15(0.39,3.414)	0.792
1000 - 1500 birr	13 (72.2%)	5 (27.8%)	2.39(0.82,6.97)	0.110	1.44(0.413,5.09)	0.563
1500 - 2000 birr	26 (60.5%)	17 (39.5%)	1.4(0.717,2.76)	0.320	1.08(0.480,2.45)	0.845
> 2000 birr	100 (52.1%)	92 (47.9%)	1			
Father's educational status?						
Uneducated	21 (31.3%)	46 (68.6%)	0.27(0.15,0.48)	0.001	0.39(0.18,0.872)	0.021
Educated	147 (62.8%)	87 (37.2%)	1			
Mother's educational status?						
Uneducated	30 (33%)	61 (67%)	0.25(0.15,0.432)	0.001	0.633(0.31,1.28)	0.207
Educated	138 (65.7%)	72 (34.3%)	1			
Heard about menstruation before menarche?						
Yes	158(57.45%)	117(42.55%)	1	1		
No	10 (38.5%)	16 (61.5%)	0.46(0.20,1.057)	0.067	1.38(0.433,4.40)	0.586
Do you know sanitary pad in the market						
Yes	160 (58.2%)	115 (41.8%)	1			
No	8 (30.8%)	18 (69.2%)	0.31(0.134,0.76)	0.010	0.36(0.12,1.114)	0.077
Knowledge about menstrual hygiene practice?						
Good knowledge	113 (61.4%)	71 (38.6%)	1	1		
Poor knowledge	55 (47%)	62 (53%)	0.55(0.34,0.89)	0.015	0.847(0.477,1.5)	0.323

Freely discuss menstruation issues with your parents?

Yes							
No	140 (65.7%)	73 (34.3%)	1	1	0.45(0.22,0.903)	0.025	
	28 (31.8%)	60 (68.2%)	0.24(0.14,0.41)				

Note: 1.00= Reference ; Bold number =p<0.05

4. Discussion

In this study, 184 (61.1%) study participant had good knowledge about menstrual hygiene, 117(38.9%) respondents had poor menstrual hygiene knowledge this is high when compared with the study done in Wegera District was found that 145(34.3%) had good knowledge this difference may be due to poor counseling of teachers.

In this study, 168 (55.8%) of the respondents had good practiced and whereas 133(44.2%) practiced poor menstrual hygiene. This finding almost shows good menstrual practice but the study was done in Wegera District, Northwest Ethiopia In general out of 423 study participants 126 (29.8%) of the respondents practiced good menstrual hygiene whereas 297(70.2%) practiced poor menstrual hygiene. This difference might be due to the reason that students have knowledge about menstrual hygiene practice than Wegera district female high school students.

In this study among 301 study participants, 276(91.7%) were heard about menstruation before menarche A study was done in, Northeast Ethiopia, the majority of the girls, 478 (86.75%) had heard about menstruation before they had menarche this almost similar.

In this study Majority of the girls, 275(91.4%) knew about menstruation before they had menarche was as study was done in India indicated that only 42% of the girls had knowledge about menstruation before their onset of menarche and A similar study conducted in Wegera District was found that 83.2% of the participants had prior knowledge about menstruation before menarche this difference may due to most of the female high school students in our study freely discuss with family about menstruation.

According to this study, all of the study participants (301) use menstrual absorbent was the study conducted in Mehalmeda Amhara regional state of Ethiopia all respondents use a menstrual absorbent, this is also similar.

This study revealed that 198(94.3%) and 77 (84.6%) residency of the urban and rural participants respectively had awareness about menstruation prior to menarche. This finding is higher than a study done in Udipi Taluk, India 83 (33.27%) and 197 (35.82%) of the urban and rural participants had awareness about menstruation prior to menarche respectively This difference

might be due to socio-cultural factors like talking about menstruation might be taboo in Indian mothers so the girls might not have awareness.

In this study showed that Among 301 study participants 276(91.7%) were heard about menstruation before menarche and The main sources of information about menstruation were, 163(54.2%), 70 (23.3%), 13(4.3%), 15(5%) and 14(5%) from mothers, teacher, health personnel, TV and others respectively in similar study conduct in Nigeria Amassoma Community, Bayelsa State revealed that 64.2% of the participants were aware of menstruation and the most important source of information are their mothers, while friends and television also contributed to their information this is almost similar.

This study showed that the practice of good menstrual hygiene was more among students who live in the urban area (AOR 0.269:95% CI, 0.125, 0.577) than students who live in the rural area This finding is similar with the study done in Kano Northwestern Nigeria. It was observed that girls who live in the urban had a good practice (AOR=2.708:95%CI,1.33,3.123) of menstrual hygiene than who live in the rural area.

List of abbreviations

HBV	Hepatitis B Virus
HIV	Human Immunodeficiency virus
MHM	Menstrual hygiene management
MHP	Menstrual hygiene practices
RTI's	Reproductive tract infections

Declarations

Competing interests

There is no competing of interest

Funding

Self sponsor which means there is no other funding source

Author Contributions

Corresponding Author: Investigation, Funding Acquisition, Formal Analysis

Co-author: Investigation, Funding Acquisition

Acknowledgments

First of all, we would like to express thanks to Harar health science college for giving us this opportunity and to thanks our Advisor S/Firehiwot demise for her constructive comments, and support for the accomplishment of this research. Our deepest gratitude also goes to our honored instructors Asefa Tola and Abdu Omar for their commitment and dedication while commenting us on the research and also we want to extend our deep appreciations and thanks for Arif Hussien, Siraj Adem for their, moral and material support during our research thesis work and final we would like to thanks for the followings: regional education office, Harar senior secondary school, Harar Bethlehem primary and secondary school, SOS primary and secondary school, Hamaresa secondary school, Abadir primary and secondary school, Zoom In primary and secondary school, which helped us during our research thesis.

For Supporting Data :

Corresponding Author

astawusalemayehu@gmail.com

Co-author

abcdef356@live.com

Reference

1. Population W, Sheet D. World Population Data Sheet The World At 7 Billion. 2011;1–15.
3. An exploratory study into Menstrual Hygiene Management amongstrural , primary school girls in Uganda : what implications does menstrual related absenteeism have for future interventions ?
4. Lawan UM, Yusuf NW, Musa AB. Menstruation and menstrual hygiene amongst adolescent school girls in Kano, Northwestern Nigeria. *Afr J Reprod Health*. 2010;14(3):201–7.
6. Gultie T, Hailu D, Workineh Y. Age of menarche and knowledge about menstrual hygiene management among adolescent school girls in Amhara province, Ethiopia: implication to health care workers & school teachers. *PLoS One* [Internet]. 2014;9(9):e108644.
7. Sarah House TM. Module six Module six. Co-published byACF,care,IRC,OXFAM,plan,Save the children,UNICEF, USAID,WASH,WaterAid,World vision; 2012. p. 130–50.
8. House S, Mahon T, Cavill S. Menstrual hygiene matters hygiene around the world. First edit. Co-published byACF,care,IRC,OXFAM,plan,Save the children,UNICEF,USAID,WASH,WaterAid ,World vision; 2012. 1-352 p.
11. Boosey R, Prestwich G, Deave T. Menstrual hygiene management amongst schoolgirls in the Rukungiri district of Uganda and the impact on their education: a cross-sectional study. *PanAfr MedJ* [Internet]. 2014;19:1–13.
12. Kamath R, Ghosh D, Lena A, Chandrasekaran V. *Global Journal Of Medicine And Public Health A study on knowledge and practices regarding menstrual hygiene among rural and urban adolescent girls in Udupi Taluk , Manipal , India*. 2013;1–9.
13. Omidvar S, Begum K. Factors influencing hygienic practices during menses among girls from south India- A cross sectional study. *Int J Collab Res Intern Med Public Heal* [Internet]. 2010;2(12):411–23.
16. Gultie TK. Practice of menstrual hygiene and associated factors among female Mehalmeda high school students in Amhara regional state , Ethiopia. 2014;2(3):189–95.
17. Tegegne TK, Sisay MM. Menstrual hygiene management and school absenteeism among female adolescent students in Northeast Ethiopia. *BMC Public Health*. 2014;14(1):1–14.
20. Ayinde MO. Self care practices of menstrual hygiene among adolescents school going girls in Amassoma Community , Bayelsa State. *Int J Nurs Midwifery*. 2013;5(August):99–105.
25. Meseret Abay Fisseha, Yigzaw Kebede, Hedija Yenus Yeshita. Menstrual Hygiene Practice and Associated Factors among Secondary School Girls in Wegera District, Northwest Ethiopia; a Cross-Sectional Study. *Computational Biology and Bioinformatics*. Vol. 5, No. 1, 2017, pp. 6-11. doi: 10.11648/j.cbb.20170501.12
27. Prateek S, Saurabh R. A Cross Sectional Study of Knowledge and Practices about Reproductive Health among Female Adolescents in An Urban Slum of Mumbai. *Journal of Family and Reproductive Health*. December 2011; 5 (4).

