4 **Title**:

DNA methylation differences at birth after conception through assisted reproductive technologies

Elmar W. Tobi[1,2,*], Catarina Almqvist[3,4], Anna Hedman[5], Ellika Andolf[5], Jan Holte[6,7,8], Jan I. Olofsson[9] ,

Håkan Wramsby[10] , Margaretha Wramsby[11], Göran Pershagen[12], Bastiaan T. Heijmans[2], Anastasia N.

Iliadou[3]

1. Periconceptional Epidemiology, Department of Obstetrics and Gynaecology, Erasmus MC, University
Medical Center Rotterdam, Dr. Molewaterplein 50, 3015GE Rotterdam, The Netherlands
2. Molecular Epidemiology, Department of Biomedical Data Sciences, Leiden University Medical Center,
2300RC Leiden, The Netherlands
3. Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, 171 77 Stockholm, Sweden
4. Pediatric Allergy and Pulmonology Unit, Astrid Lindgren Children's Hospital, 171 76 Stockholm, Sweden
5. Department of Clinical Sciences, Danderyd hospital, Karolinska Institutet 182 88 Stockholm Sweden
6. Carl von Linné Clinic, Uppsala, Sweden
7. Department of Women's and Children's Health, Uppsala University, 751 85 Uppsala, Sweden
8. Center for Reproductive Biology in Uppsala, University of Agricultural Sciences and Uppsala University,
Uppsala, Sweden.
9. Department of Women's and Children's Health, Karolinska Institutet, 171 77 Stockholm, Sweden.

10. Livio Fertilitetscentrum Kungsholmen, S:t Görans Sjukhus, 112 81 Stockholm, Sweden

11. Livio Fertilitetscentrum Gärdet Storängsvägen 10, 115 42 Stockholm

12. Institute of Environmental Medicine, Karolinska Institutet, 171 77 Stockholm, Sweden

*corresponding author: Assoc Prof Anastasia Nyman
Department of medical epidemiology and biostatistics,
Karolinska Institutet
Email: Anastasia.nyman@ki.se
Tel:+ 46 708 267874
30

Abstract

Study Question: What is the relationship with assisted reproductive technology (ART) and DNA

methylation (DNAm) patterns in cord-blood and is there any difference between in vitro fertilisation

(IVF) and intra-cytoplasmic sperm injection (ICSI)?

Summary Answer: We identify nineteen CpGs at which DNAm was associated with being conceived via

assisted reproductive technology (ART). No difference was found between IVF and ICSI.

What is known already: Prior studies on either IVF or ICSI show conflicting outcomes as both wide-

spread effects on DNAm and highly localized associations have been reported. No study on both IVF and

ICSI and genome-wide neonatal DNAm has been performed.

Study design, size, duration: We measured 87 infants conceived with in vitro fertilisation (IVF) or intra-

cytoplasmic sperm injection (ICSI) and 70 who were not in a cross-sectional study.

Participants/materials, setting, methods: Participants were from the UppstART study, which were

recruited from fertility and reproductive health clinics, and the Born into Life cohort, which is created

from the larger LifeGene study. We measured DNAm from DNA extracted from cord blood collected at

birth with the Illumina Infinium HumanMethylation450k BeadChip micro-array. Group differences in

DNA methylation at individual CpG dinucleotides (CpGs) were determined using robust linear models

and post-hoc Tukey's tests.

Main results and the role of chance: We found no association with global methylation levels, imprinted

loci and meta-stable epialleles. In contrast, we identify nineteen CpGs at which DNA methylation was

associated with being conceived via ART (effect estimates: 0.5-4.9%, $P_{FDR}<0.05$), but no difference was

found between IVF and ICSI. The associated CpGs map to genes related to brain function/development

or genes connected to the plethora of conditions linked to subfertility, but functional annotation did not

point to likely functional consequences.

Limitations, reasons for caution: We measured DNAm in cord blood and not at later ages and/or other

tissues and findings remain to be replicated in an independent study.

Wider implications of the findings: We find ART is associated with DNA methylation differences in cord

blood, but these differences are limited in number, effect size and with unknown functional

consequences in adult blood. We did not find indications for a disparity between IVF and ICSI.

Study funding/competing interest(s): The authors declare no competing interests. The funders

had no role in the study, study design, analysis and decision to publish.

Keywords
63   DNA methylation; ART; IVF; ICSI; EWAS; cord-blood; epigenetics

Introduction

The usage of assisted reproductive technology (ART) is increasing world-wide and its possible health consequences are a topic of intense study. Evidence is building for an association of ART with long-term health outcomes for the offspring, including autism spectrum disorders (Liu *et al.*, 2017) and cardiovascular health (higher blood pressure, suboptimal cardiac diastolic function and vessel thickness) (Guo *et al.*, 2017). A body of evidence links ART to short term health effects, including low birth weight, placenta associated anomalies (Vermey *et al.*, 2019) and pregnancy complications (Qin *et al.*, 2016), congenital malformations and imprinting disorders (Turkgeldi *et al.*, 2016). However, it is still debated if these associations stem from the application of ART or the underlying infertility leading couples to ART (Luke *et al.*, 2016). Animal studies provided potential molecular mechanisms for these observations by highlighting that the ART procedure may induce changes to epigenetic marks (Morgan *et al.*, 2008; Wang *et al.*, 2010). Epigenetic marks, such as DNA methylation (DNAm), influence the transcription potential of genomic regions (Jaenisch and Bird, 2003). Human studies likewise point towards a link between early development, DNAm and (late-life) phenotypes, but the causality of these associations remains unknown (Tobi *et al.*, 2018).

To date, several studies on DNAm and ART have been performed. Multiple studies have focused on candidate gene regions and/or global methylation levels (Lazaraviciute *et al.*, 2014; Canovas *et al.*, 2017). Genome-wide efforts have focused on samples taken from extra-embryonic lineages (Xu *et al.*, 2017; Choufani *et al.*, 2018) and when focused on material from the infant itself suffered from a small sample size (Melamed *et al.*, 2015) or batch effects correlating with ART status (Estill *et al.*, 2016). Larger studies have focused on specific ART techniques, namely intra-cytoplasmic sperm injection (ICSI) (El Hajj *et al.*, 2017) or in vitro fertilisation (IVF) (Castillo-Fernandez *et al.*, 2017) or IVF and less invasive gamete intra-fallopian transfer (GIFT) coupled with intra-uterine insemination (IUI) (Novakovic *et al.*, 2019). It has been hypothesized that the technique used may matter for the possible consequences on DNAm

patterns (Loke and Craig, 2016). Overall, these studies report conflicting outcomes, with two reporting

wide-spread associations between ART and (cord-)blood DNAm at birth (El Hajj *et al.*, 2017) which

appear to fade with age (Novakovic *et al.*, 2019), while another study reported a DNA methylation

difference at a single genomic locus only (Castillo-Fernandez *et al.*, 2017).

We undertook an epigenome wide association study on ART, comparing DNAm patterns in cord

blood of children conceived via ART (N=87) (Iliadou *et al.*, 2019) with that of children from medically

unassisted conceptions (henceforth "MUC", N=70) (Almqvist *et al.*, 2011). Since we had detailed

information on the ART technique used for 77 of the children conceived via ART we investigated

possible different outcomes for IVF and ICSI on DNAm which has been hypothesized to be important

(Loke and Craig, 2016), but is yet to be tested. In addition, we explored the possible functional

consequences of these methylation differences using external datasets.

Methods

Study Subjects

ART controls: Born into Life

The prospective longitudinal birth cohort study Born into Life was originally created from the larger

LifeGene study (Almqvist *et al.*, 2011). Between the years 2010 and 2012, pregnant women who were

already participating in the LifeGene study and living in Stockholm County were recruited to Born into

Life (Smew *et al.*, 2018). The inclusion criteria were that they had responded to baseline questionnaires

from the LifeGene study, were pregnant, and gave informed consent. They were recruited both before

and after 10-14 gestational weeks, but no later than 26-28 weeks. Originally 107 pregnant women were

included in Born into Life. We only included the 77 women for which cord blood had been successfully

collected in this study.

Women in the Born into Life were asked to complete questionnaires regarding pregnancy,

lifestyle and health at 10-14 and 26-28 gestational weeks. Data on maternal self-reported smoking

during pregnancy and body mass index (BMI), in kg/m2 obtained at the first antenatal care visit in

gestational week week 5-12 were collected from birth records. Data regarding highest attained

educational level, ranging from mandatory secondary school to high school, university or other, were

retrieved from the baseline LifeGene questionnaires. Maternal age at delivery, in years, was calculated

from mothers' date of birth. Parity and data regarding the infants' sex, gestational age in weeks and

mode of delivery, defined as vaginal delivery or Caesarean section, were also collected from the birth

records. Cord blood and placenta samples were obtained from delivery. Cord blood was aspirated 2

minutes after birth by the assisting midwife into a test tube and kept in room temperature awaiting

transportation. The birth records regarding both mother and child at delivery were collected from

Danderyd Hospital, Stockholm.

ART cases: UppstART

The UppstART study has been described in detail elsewhere (Iliadou *et al.*, 2019). Participants of the

UppstART study were recruited from three of the four fertility and reproductive health clinics in

Stockholm (one public, two private) and one private clinic in Uppsala county, which also serves a large

volume of patients from Stockholm. Recruitment took place from September 2011 to December 2013.

IVF treatment(s) of the participants were followed until December 2014 or drop-out/consent withdrawn

(n=4), which ever came first. The participants were asked to answer a web-based baseline questionnaire

within a few days of their clinic visit and prior to their IVF treatment start, which included an extensive

list of questions on sociodemographic, anthropometric and life-style factors. Once the IVF treatment

began and the participants reached the stage of oocyte retrieval, they were asked to respond to a

second online follow-up questionnaire, with a shorter version of the baseline questionnaire, to identify

any changes in lifestyle factors since the initiation of their treatment. Staff at seven delivery units in

Stockholm and Uppsala were recruited to assist in collecting samples from UppStART participants during

delivery. Delivery clinics were provided with a sample collection kit including tubes for collection of

cord-blood for DNA extraction (EDTA). Samples were stored in -20°C freezers at the delivery units until

they were collected by UppStART study staff and deposited into the KI Biobank.

DNA methylation measurements

Genome-wide DNA methylation data were generated using the Illumina Infinium Human Methylation

450K BeadChip (450k array). A total of 500ng of genomic DNA isolated from cord blood was bisulfite

treated using the EZ-96 DNA methylation kit (Zymo Research, Orange County, USA). We used the D-

optimum criterion to assign samples over two 96-well plates and individual 450k arrays, ensuring even

distributions of ART cases and controls, ART methods (IVF or ICSI), gender, gestational age and birth

month across the two 96-well bisulfite plates and each 450k array. The 450K arrays were measured at

ServiceXS (Leiden, The Netherlands). The quality of the generated 450K array data was assessed using

both sample dependent and sample independent quality metrics using the *Bioconductor* package

MethylAid (van Iterson *et al.*, 2014) with default settings. We used the *Bioconductor* package *omicsPrint*

(van Iterson *et al.*, 2018) to check for sample duplications and admixtures and sample sexes were

checked using the X-chromosomal CpGs. One sample was not the correct gender and was deleted. We

used principal component analysis (PCA) and hierarchical clustering on the raw autosomal beta-values

to search for outliers and suspect patterns in the dataset, finding none. The cell proportions of cord

blood were imputed using IDOL (Koestler *et al.*, 2016) on the "FlowSorted.CordBloodCombined.450k"

reference set using the estimateCellCounts2() function in the *FlowSorted.Blood.EPIC* R package(Salas *et*

al., 2018; Gervin *et al.*, 2019). Normalization of the dataset was performed by NOOB background and

color correction in combination with Functional Normalization (Fortin *et al.*, 2014) using four principal

components (eigenvalue > 1). All measurements with <3 beads (0.05% of probes), < 1 intensity value

(0.012% of probes) and a detection p-value >0.01 (0.11% of CpGs) were set as missing. The

measurement success rate per sample was >99%. CpGs with probes that did not map to unique genomic

locations (Chen *et al.*, 2013) or with a < 95% measurement success rate (0.23% of CpGs) were then

removed. We used custom scripts that add on the functions from the *minfi* package (Aryee *et al.*, 2014)

and implements parallelization where possible (https://github.com/molepi/DNAmArray).

Outlier detection was performed by PCA and hierarchical clustering on the autosomal beta-values.

Hierarchical clustering identified one possible outlier. This individual had a very low gestational age (30

weeks), the lowest in the dataset, and this clustering is therefore likely based on biological reasons.

High-quality DNA methylation data were obtained for 157 individuals, 70 MUC from Born into Life, 6 ART

from Born into Life (specific type of ART unknown), 81 ART from UppStART (specific type of ART

unknown for N=4) for a total of 441,836 autosomal CpGs. The datasets generated and analyzed during

the current study are not publicly available due to Swedish privacy and data safety laws but are available

from ANI and CA on reasonable request and after meeting legal requirements.

Measures of global methylation

180 Genome-wide average DNA methylation (GWAM) (Li *et al.*, 2018) was calculated by averaging all beta-

181 value measurements across the autosomes for each individual. The *R* package *REMP* (Zheng *et al.*, 2017)

182 was used to infer DNA methylation at either *ALU* or *LINES*-1 sequences and then the average across all

183 *ALU* or *LINES*-1 elements was calculated for each neonate.

TFBS enrichment

185 We calculated transcription factor binding site enrichments using the R package *PWMEnrich* (Stojnic and

186 Diez, 2018) using binding motifs from motifDB. We calculated position weight matrices using the DNA

187 base background frequencies calculated for the CpGs tested with 25bp flanking sequences. Enrichment

188 was tested relative to this background.

Statistics

We compared unassisted and ART cohort descriptives via Kruskal Wallis (maternal BMI, maternal SES,

years till index pregnancy, CD4T, NK and nucleated red blood cells), t-test (maternal and gestational age,

birthweight, CD8T, Monocytes, granulocytes and B cell proportions) assuming equal or unequal variance

where appropriate and chi-square tests (child sex, parity and maternal smoking). DNA methylation was

always analyzed as beta-value which is reported as % (e.g. x100) in Tables. GWAM, *ALU* and *LINES*-1

methylation levels were compared via t.test (unassisted vs ART) or ANOVA (medically unassisted

fertilization versus IVF versus ICSI).

We used the R package *cate* (Wang and Zhao, 2015) to identify latent variables (e.g. "hidden

variables") influencing the relation between DNA methylation and ART (yes/no) or ART method

(IVF/ICSI/unassisted). Two latent variables were identified ($P<0.001$) for ART and four latent variables for

ART method ($P<0.001$). In both cases the first two latent variables correlated with the proportion of

nucleated red blood cells (nRBC) in blood (rho>0.94, $P<0.001$) and the other two latent variables

correlated with CD4T or B cell proportions in blood and several batch effects (rho>|0.3|, $P<0.001$)

arguing together with directed acyclic graphs (DAG) (*Supplemental text*), for a basic model adjusting for

batch effects and imputed cell proportions.

We performed epigenome-wide association analyses (EWAS) on being conceived by ART or not

using robust linear regression (rlm) from the R *MASS* package with White's estimator for robust

standard errors as implemented in the R package *sandwich* (Zeileis, 2006) (which leads to a model

robust for outlying beta-values and heteroscedasticity). To test for differences in neonates conceived via

IVF or ICSI or without medical assistance we used type II ANOVA with White's estimator for robust

standard errors to account for any heteroscedasticity and unequal variances between groups using the R

car package. We employed heteroscedasticity robust Tukey tests to test for DNA methylation

differences between IVF, ICSI and unassisted conceptions post-hoc. In all instances we used the beta-

value of an individual CpG dinucleotide as the outcome and ART status as the dependent. We adjusted

for the sex of the individual, height of the sample on the 450k array glass slide (continues variable from

1-6), scan batch and bisulfite plate and the imputed cell proportions (CD4T, CD8T, Mono, NK, Bcell,

Gran, nRBC). We ran additional sensitivity analyses by omitting the cell proportions or expanding the

adjustment with minimal adjustment sets coming from DAG analysis with the following models, i: years

till the index pregnancy, maternal age, pre-pregnancy BMI, educational attainment as proxy for socio-

economic status, smoking history ii) gestational length and parity. In addition, we performed look-ups in

other published meta-EWAS of prenatal exposures.

Regions were tested with the same adjustments as defined above in a linear mixed model with a

random effect for individual and a factorial co-variate denoting the specific CpG dinucleotide of each

DNAm measurement in the R *lme4* package using compound symmetry as correlation structure and the

R *lmerTest* package for the Satterthwaite's approximation of the degree of freedom of the fixed effects

within each linear mixed model.

All analyses were performed in R (version 3.6.1) (R core team, 2019). All p-values reported are

two-sided and multiple testing correction is done using FDR (false discovery rate).

Ethical approval

Ethical approval for Born into Life was granted by the Regional Ethics Review Board in Stockholm,

Sweden. Informed consent from both parents was obtained for all study participants. The UppstART

study has been approved by the regional ethics review board at the Karolinska Institutet (Dnr 2011/230-

31/1, Dnr 2011/1427-32, Dnr 2012/131-32, Dnr 2012/792-32, 2013/1700-32, 2014/1956-32, Dnr

2015/1604-32). Women and their partners were approached by the clinic nurse and asked to participate

in the study. To facilitate the process of informed consent, the couple was provided with information

approved by the regional ethical board, both verbally and in written format about the purpose of the

study, methods, possible risks, and that participation was voluntary. Additionally, participants were

informed that they could withdraw from the study at any time with no impact to their medical care. The

requirement for inclusion in the study was an understanding of the Swedish language and exclusion was

the use of donor gametes.

Results

Study subjects

Pre-processing and normalization resulted in DNAm data of 157 infants from cord blood, 87 of which were conceived with the help of ART (Table 1). We had detailed information on the ART technique used for 77 of the 87 newborns conceived by ART and with DNAm data, of which 44 were conceived via IVF and 33 via ICSI. The mothers that conceived with the help of ART were not different in body mass index (BMI, P=0.25) and age (P=0.25), but had a lower socio-economic status (P<0.001) and more were former smokers (P<0.001) than mothers with a MUC. It took on average two years longer for the mothers who conceived through ART to become pregnant than for mothers with MUC (P<0.001). There was no difference between ART and MUC in length of gestation (P=0.16) and parity (P=0.20). The percentage of male newborns in the MUC group was higher (61.4% vs 50.6%), although this difference was not significant (P=0.23).

DNAm is a key determinant of cell identity (Jaenisch and Bird, 2003) and cord blood consists of multiple cell types that influence DNAm (Jones *et al.*, 2007). Seven major cell types, including nucleated red blood cells, were imputed from the genome-wide DNAm data of the newborns (Gervin *et al.*, 2019). In concordance with a prior report on unassisted and ICSI newborns (El Hajj *et al.*, 2017), there were no differences in these seven cell proportions between ART and MUC cord blood samples ($P_{nominal}$>0.13, Supplemental Figure S1).

Global DNAm comparison between ART and unassisted conceptions

First, we investigated genome-wide average DNA methylation (GWAM) (Li *et al.*, 2018) in the cord blood of neonates conceived via IVF or ICSI or MUC. There was no difference between these 3 groups (P=0.89, Supplemental Figure S2). Next, we investigated the genome-wide average DNA methylation of ALU and

263   LINES-1 elements, an often used proxy for global DNA methylation (Zheng *et al.*, 2017), finding no

264   difference between groups (P>0.2, Figure S3).

DNAm comparison between ART and medically unassisted conceptions

We compared DNA methylation at 441,836 autosomal CpGs between 87 ART and 70 MUC neonates

(Supplemental Figure S4). Nineteen CpGs were associated with being conceived via ART in a model

adjusting for sex, batches and cell heterogeneity (P_{FDR}<0.05, Figure S5). The associations extended to

neighboring probes (Figure 1) for the proximal promoter of *AK054845*, a gene of unknown function (five

CpGs $P_{nominal}$<0.05), and the proximal promoter of a smaller variant of *GABRB3* (NM_001191321.2) which

is expressed in the brain (eight CpGs $P_{nominal}$<0.05) and has a neuro-developmental role (Tanaka *et al.*,

2012). Indeed, several CpGs can be linked to genes with roles in brain function or (early) brain

development. *ZNF771* (cg06450634), as a transcription factor (TF), is a driving factor behind a large gene

network in the brain (Maulik *et al.*, 2018) while *CYFIP1* (cg13051607) (Hsiao *et al.*, 2016), *GABRB3*

(cg01251603, cg15066197,cg14859324) (Tanaka *et al.*, 2012) and *SHANK1* (cg14560133) (Sungur *et al.*,

2018) are all autism spectrum disorder candidate genes, a late life phenotype tentatively associated

with ART (Liu *et al.*, 2017).

Omitting the adjustment for cell heterogeneity had little to no effect on the effect estimates for

all nineteen CpGs (Supplemental Table S1) just like additional adjustment for maternal and pregnancy

characteristics (Supplemental Table S2). The nineteen CpGs were not among those previously reported

in other EWASs (FDR corrected P-value of <0.05) with prenatal smoking (Joubert *et al.*, 2016b), folate

use (Joubert *et al.*, 2016a), maternal hypertension and pre-eclampsia (Kazmi *et al.*, 2019), maternal BMI

(Sharp *et al.*, 2017) and birthweight (Küpers *et al.*, 2019). Moreover, a look-up of these previously

reported CpGs in our study did not yield more nominally associated CpGs with ART than may be

expected by chance (P>0.07).

DNA methylation comparison between IVF, ICSI and unassisted conceptions

To investigate any differences between IVF and ICSI we performed post-hoc Tukey's tests for DNA methylation differences between neonates conceived via IVF and ICSI for these nineteen CpGs. No differences were found ($P_{FDR-19\ tests}>0.15$). Next, we extended our analysis to all CpGs by performing an ANOVA analysis to test for differences in DNA methylation across the 441,836 autosomal CpGs between neonates conceived with IVF (N=44) or ICSI (N=33) or MUC (N=70). This yielded five CpGs already identified in the EWAS for ART status and in all cases the difference was between the MUC and both IVF and ICSI, thus finding no evidence for IVF or ICSI specific associations.

Functional annotation

We performed KEGG and Gene Ontology tests (Phipson *et al.*, 2015), finding no enrichment for the genes linked to nineteen CpGs associated with ART, nor when we relaxed the significance threshold to suggestive associations ($P<10^{-5}$, N=37). We did not find significant enrichments of single transcription factor (TF) binding motifs (Stojnic and Diez, 2018), although multiple CpGs overlapped TF binding sites of TFs with a role in early development like *TFAP2C* (Sharma *et al.*, 2016) overlapping cg27266479 mapping to *H6PD* and *RAX* (Bennett *et al.*, 2008) overlapping cg16771467 to *ATP8B1*. Look-up of the nineteen CpGs in reference data (Bonder *et al.*, 2017) uncovered no link between DNAm and gene-expression in adult blood. We measured DNAm in cord blood and average DNAm levels may vary between tissues, but still reflect variation in other tissues (Slieker *et al.*, 2013). There was little to no correlation of methylation at the nineteen CpGs between adult whole blood and tissues from sixteen cadavers (Slieker *et al.*, 2013; Relton *et al.*, 2015), with the notable exception of cg14560133 (*SHANK1*, β = -1.42% (SE=0.3%), $P=2.1\times10^{-6}$) which showed moderate correlation with 5/8 tissues (rho>0.53, P<0.04), Figure 2). The three CpG dinucleotides at *GABRB3* (cg01251603, cg15066197,cg14859324), a gene active in the brain, showed weak to moderate correlation between adult blood and four brain regions (rho=0.39-0.62, P<0.001, N=71-74) in another reference dataset (Hannon *et al.*, 2015).

Imprinted regions and metastable epialleles

Multiple candidate gene studies have looked at DNA methylation or gene-expression differences of

imprinted genes between ART and MUC with mixed results. Therefore, we looked at 374 CpGs on the

450k array known to overlap imprinted differentially methylated regions (DMRs) (Yuen *et al.*, 2011).

There was no overlap with the CpGs showing a (suggestive) association with ART in our study (P<10^{-5},

N=22). These 374 CpGs where distributed over 59 regions. There was no difference in DNAm between

MUC and ART infants when we tested each of these 59 regions (P$_{FDR}$>0.86). Similarly, so-called meta-

stable epialleles (MEs) are hypothesized to be especially sensitive to the early prenatal environment

(Kessler *et al.*, 2018). There were 187 CpGs overlapping MEs, none of which were associated with ART

(P$_{FDR}$>0.06).

Prior studies

We performed a look-up of CpGs and regions associated in prior studies with one form of ART. The sole

region associated with IVF in the meDIP sequencing study from Castillo-Fernandez *et al.* was not covered

by our genome-scale screen with the 450k array (Castillo-Fernandez *et al.*, 2017). Next, El Haij *et al.*

found 4730 CpG dinucleotides associated with ICSI after FDR adjustment (which are unavailable as

Supplemental information) of which two regions, out of five regions selected for validation with

pyrosequencing, were validated in an additional sample (El Hajj *et al.*, 2017). The seven CpGs in these

two regions which cover the proximal promoters of *ATG4C* and *SNORD114-9* were not associated with

ART in our study (P>0.23) nor with ICSI only (post hoc Tukey's test: P>0.40).

Using the 850k array, Novakovic *et al.* identified 2340 CpG dinucleotides at which DNA

methylation in Guthrie card blood was associated with IVF and GIFT coupled with IUI (Novakovic *et al.*,

2019), 1228 of which were covered in our 450k study. Of these, 38 were nominally associated (P<0.05)

and with the same direction of effect in our study. This decreased to 12 loci when we tested for DNA

methylation differences between MUC and IVF only. Novakovic *et al.* also identified three differentially

334   methylated regions where the difference in methylation associated with IVF remained in an adult

335   sample. CpG dinucleotides at *CHRNE* (chr17:4803506–4805392), *PRSS16* (chr6:27185896–27186199)

336   and *TMEM18* (chr2:731073–732037) showed a similar direction of effect in our study as that of

337   Novakovic *et al.* and the association was almost nominally significant for *PRSS16* (cg10279314:

338     $\beta=2.3\%[SE=1.2\%]$, P=0.057, cg09395805: $\beta=3.1[SE=1.6]$,P=0.049, cg07555084: $\beta=2.2[SE=1.2]$,P=0.074).

Discussion

We identified nineteen CpGs at which DNAm was associated with ART and the association extended to neighboring CpG dinucleotides at *AK054845* and *GABRB3*. We found no indication for a disparity between IVF or ICSI on DNAm patterns. The associations were robust to adjustment for cellular heterogeneity and maternal characteristics and did not overlap with those loci associated with other common prenatal exposures. We found no evidence that imprinted loci and meta-stable epialleles are especially sensitive to ART. The nineteen CpGs can be annotated to relevant genes, but the functional implications of variation in DNA methylation at these CpGs remains to be elucidated.

The nineteen CpG dinucleotides could be mapped to fifteen genes. Four of the nineteen were located in a proximal promoter and here the neighboring CpG dinucleotides were also nominally associated with ART. In other cases the associations were either limited to one CpG dinucleotide, which may stem from the sparse coverage of the 450K array across the methylome and in other cases is consistent with the fact that DNAm acts through the altering of the binding potential of a specific TFBS (Bonder *et al.*, 2017). Indeed, none of the CpGs were located in a CpG island, which is consistent with data from other exposures showing that associations between DNAm and prenatal and/or environmental conditions are enriched at regulatory regions like enhancers and CpG poor promoters that show intermediate levels of methylation (Tobi *et al.*, 2014).

Several of the CpGs can be linked to genes with a role in brain function and/or development. *ZNF771* (cg06450634) has been identified as a driving factor behind a large gene network in the brain (Maulik *et al.*, 2018) and *CYFIP1* (cg13051607) (Hsiao *et al.*, 2016), *GABRB3* (cg01251603, cg15066197,cg14859324) (Tanaka *et al.*, 2012) and *SHANK1* (cg14560133) (Sungur *et al.*, 2018) are all autism spectrum disorder candidate genes, a late life phenotype tentatively associated with ART (Liu *et al.*, 2017). In addition, several CpG dinucleotides can be linked to genes connected to the plethora of conditions that can be linked to subfertility. *H6PD* (cg27266479) is a candidate gene for polycystic ovary

syndrome (PCOS) (Martínez-García *et al.*, 2012). *INTS1* (cg03207674) has a crucial role in the developing

blastocyst, as inhibition of *INTS1* function causes growth arrest (Hata and Nakayama, 2007). *NOS3*

(cg00478390) knockout mice are used as an *in vivo* model of (recurrent) embryo loss, as nitric oxide

metabolism plays an important role in implantation (Pallares and Gonzalez-Bulnes, 2010).

The latter may hint that the various medical reasons for ART may underlie the associations,

rather than the ART process itself, which is a key question in the study of ART (Luke *et al.*, 2016). We

employed directed acyclic graphs (DAG) (Krieger and Davey Smith, 2016) and DAG indicated that and

ascertainment of a direct effect is possible with a small minimal adjustment set. There was little to no

effect on the effect estimates from adjustment for various maternal characteristics, gestational age and

the number of years until pregnancy (as a proxy for in-/subfertility). In addition, we did not find any

evidence for different or stronger effects of ICSI on DNAm, as has been hypothesized (Loke and Craig,

2016), and the fertility clinics from which we recruited all used the same culturing media for ICSI and

IVF. The last two items highlighted might argue that not the (reasons for) infertility, but the ART process

itself may explain the associations.

Our analysis did not find wide-spread genome-wide differences like other genome-scale studies

to date (El Hajj *et al.*, 2017; Novakovic *et al.*, 2019), but is consistent with a genome-scale study on only

IVF using immunoprecipitation of methylated DNA with next generation sequencing (Castillo-Fernandez

et al., 2017). We extensively compared individual results, as far as possible, and found little overlap. This

may be due to study power, the different scope of the measurement techniques used (Castillo-

Fernandez *et al.*, 2017) and the different sources of genomic DNA (Guthrie card versus whole cord

blood) (Novakovic *et al.*, 2019). In addition, it is possible that different media was used between

countries or that the fact that only a small proportion of our IVF and ICSI group consisted of embryos

that were frozen and/or cultured extensively in comparison to other studies. The influence of this latter

aspect we were underpowered to test as most ART procedures entailed fresh embryos that were not

cultured extensively *in vitro*.

390   Our study is one of the larger genome-scale DNAm studies on ART to date and we are the first to

study both IVF and ICSI on this scale, nonetheless our study has important limitations to consider. First,

DNAm is one of the drivers of cell identity (Jaenisch and Bird, 2003) and we measured cord blood, which

may be a tissue less relevant in relation to study outcomes although it may still mark processes in

relevant tissues due to mitotic inheritance (Heijmans and Mill, 2012). We used the latest methods to

impute the seven major cell types in cord-blood (Gervin *et al.*, 2019) and in conjuncture with an earlier

report (El Hajj *et al.*, 2017) these imputed cell types did not differ between infants conceived with or

without ART and adjustment did not alter the effect estimates. However, most of these major cell types

consist of subsets of more specialized cells (subsequently showing unique DNAm profiles at increasingly

select loci) and their influence could not be investigated. However, none of the nineteen CpG

dinucleotides is linked to a gene with a role in (auto)immune function or hematopoiesis. Another major

driver of DNAm variation is genetic variation (Bonder *et al.*, 2017) and we did not control for genetic

variation in our analyses. Nine out of nineteen CpG dinucleotides did not have a single nucleotide

polymorphism (SNP) influencing the DNA methylation levels, as determined in a large meta-analysis

(Bonder *et al.*, 2017) and none of these 450k probes had an overlap with a single SNP arguing that

genetic heterogeneity is unlikely to underlie our findings. In addition, the influence of the prenatal

environment on DNAm variation may be independent and additive of genetic variation (Tobi *et al.*,

2012).

408   Also important is to consider the fact that an ART population is inherently different from the

general population (Luke *et al.*, 2016). Despite controlling for possible confounding factors, and cross-

checking our results with other EWAS on prenatal complications/exposures, unresolved confounding

remains a possibility. Our study differs with some of the earlier described in ART studies, which is useful

for triangulation between studies (Lawlor *et al.*, 2017), as our control group has a higher, rather than a

lower, SES as is normal in most of studies on ART.

The data presented showcases modest and specific DNAm differences associated with ART, of

which the functional relevance in adult tissues is unknown. We did not find any difference in DNAm

patterns between IVF and ICSI. Our study found little evidence for the hypothesis that ART, be it IVF or

ICSI, leads to wide-spread disturbances of DNAm patterns nor for the hypothesis that ICSI has a

different/larger relation with DNAm patterns. Our findings warrant cautious interpretation and awaits

confirmation, given the sample size, tissue studied and the unknown functional consequences of the

identified DNAm differences.

Author contributions

Conceptualization: ANI, CA. Methodology: EWT, BTH. Investigation: EWT, BTH, ANI, CA. Formal Analysis:

EWT. Validation: EWT. Resources: ANI, AH, CA, GP, EA, JIO, MW, HW. Data curation: ANI, AH. Writing –

original draft: EWT. Writing – Review & Editing: BTH, ANI, AH, CA, GP, EA, JIO, MW, HW. Visualization:

EWT. Supervision: BTH, ANI. Project administration: ANI, AH. Funding Acquisition: ANI, CA, GP, EA

Acknowledgements

We wish to thank the Biobank at the Karolinska Institutet for professional biobank service.

Competing interests

The authors declare no conflicts of interest

Abbreviations

ART: assisted reproductive technology

IVF: *in vitro* fertilisation

ICSI: intra-cytoplasmic sperm injection

DNAm: DNA methylation

EWAS: epigenome-wide association analysis

CpGs: CpG dinucleotides

IUI: intra-uterine insemination

GIFT: gamete intra-fallopian transfer

MUC: medically unassisted conceptions

450K array: Illumina Infinium Human Methylation 450K BeadChip

PCA: principal component analysis

TFBS: transcription factor binding site

TF: transcription factor

SNP: single nucleotide polymorphism

SES: socio-economic status

BMI: body mass index

Funding

EWT was supported by a VENI grant from the Netherlands Organization for Scientific Research

(91617128) and JPI-H2020 Joint Programming Initiative a Healthy Diet for a Healthy Life (JPI HDHL)

under proposal number 655 (PREcisE Project) through ZonMw (529051023). Financial support was

provided from the European Union's Seventh Framework Program IDEAL (259679), the Swedish

Research Council (K2011-69X-21871-01-6, 2011-3060 and 2018-02640), and the Strategic Research

Program in Epidemiology Young Scholar Awards, Karolinska Institute (to ANI) and through the Swedish

Initiative for Research on Microdata in the Social And Medical Sciences (SIMSAM) framework grant no

340-2013-5867, grants provided by the Stockholm County Council (ALF-projects), the Strategic Research

Program in Epidemiology at Karolinska Institutet, the Swedish Heart-Lung Foundation and Danderyd

University Hospital (Stockholm, Sweden). The funders had no role in study design, data collection,

analysis, decision to publish or preparation of the manuscript.

References

Almqvist C, Adami HO, Franks PW, Groop L, Ingelsson E, Kere J, Lissner L, Litton JE, Maeurer M, Michaëlsson K, *et al.* LifeGene - A large prospective population-based study of global relevance. *Eur J Epidemiol* 2011;**26**:67–77.

Aryee MJ, Jaffe AE, Corrada-Bravo H, Ladd-Acosta C, Feinberg AP, Hansen KD, Irizarry RA. Minfi: a flexible and comprehensive Bioconductor package for the analysis of Infinium DNA methylation microarrays. *Bioinformatics* 2014;**30**:1363–1369. England.

Bennett RL, Blalock WL, Choi E-J, Lee YJ, Zhang Y, Zhou L, Paul Oh S, Stratford May W. RAX is required for fly neuronal development and mouse embryogenesis. *Mech Dev* [Internet] 2008;**125**:777–785.

Bonder MJ, Luijk R, Zhernakova D V, Moed M, Deelen P, Vermaat M, Iterson M van, Dijk F van, Galen M van, Bot J, *et al.* Disease variants alter transcription factor levels and methylation of their binding sites. *Nat Genet* 2017;**49**:131–138. University of Groningen, University Medical Center Groningen, Department of Genetics, Groningen, the Netherlands Molecular Epidemiology Section, Department of Medical Statistics and Bioinformatics, Leiden University Medical Center, Leiden, the Netherlands, United States.

Canovas S, Ross PJ, Kelsey G, Coy P. DNA Methylation in Embryo Development: Epigenetic Impact of ART (Assisted Reproductive Technologies). *BioEssays* 2017;**39**:1–11.

Castillo-Fernandez JE, Loke YJ, Bass-Stringer S, Gao F, Xia Y, Wu H, Lu H, Liu Y, Wang J, Spector TD, *et al.* DNA methylation changes at infertility genes in newborn twins conceived by in vitro fertilisation. *Genome Med* [Internet] 2017;**9**:28. Genome Medicine.

Chen YA, Lemire M, Choufani S, Butcher DT, Grafodatskaya D, Zanke BW, Gallinger S, Hudson TJ,

Weksberg R. Discovery of cross-reactive probes and polymorphic CpGs in the Illumina Infinium

HumanMethylation450 microarray. *Epigenetics* 2013;**8**:203–209. Genetics and Genome Biology

Program, Hospital for Sick Children, Toronto, ON, Canada.

Choufani S, Turinsky AL, Melamed N, Greenblatt E, Brudno M, Bérard A, Fraser WD, Weksberg R, Trasler

J, Monnier P, *et al.* Impact of Assisted Reproduction, Infertility, Sex, and Paternal Factors on the

Placental DNA Methylome. *Hum Mol Genet* [Internet] 2018;Available from:

http://fdslive.oup.com/www.oup.com/pdf/production_in_progress.pdf.

Estill MS, Bolnick JM, Waterland RA, Bolnick AD, Diamond MP, Krawetz SA. Assisted reproductive

technology alters deoxyribonucleic acid methylation profiles in bloodspots of newborn infants.

Fertil Steril [Internet] 2016;**106**:629-639.e10. Elsevier Inc.

Fortin J-P, Labbe A, Lemire M, Zanke BW, Hudson TJ, Fertig EJ, Greenwood CM, Hansen KD. Functional

normalization of 450k methylation array data improves replication in large cancer studies. *Genome*

Biol [Internet] 2014;**15**:503.

Gervin K, Salas LA, Bakulski KM, Zelm MC Van, Koestler DC, Wiencke JK, Duijts L, Moll HA, Kelsey KT,

Kobor MS, *et al.* Systematic evaluation and validation of reference and library selection methods

for deconvolution of cord blood DNA methylation data. *Clin Epigenetics* [Internet] 2019;**11**:125.

Guo XY, Liu XM, Jin L, Wang TT, Ullah K, Sheng JZ, Huang HF. Cardiovascular and metabolic profiles of

offspring conceived by assisted reproductive technologies: a systematic review and meta-analysis.

Fertil Steril [Internet] 2017;**107**:622-631.e5. Elsevier.

Hajj N El, Haertle L, Dittrich M, Denk S, Lehnen H, Hahn T, Schorsch M, Haaf T. DNA methylation

signatures in cord blood of ICSI children. *Hum Reprod* [Internet] 2017;**32**:1761–1769. Oxford

University Press.

Hannon E, Lunnon K, Schalkwyk L, Mill J. Interindividual methylomic variation across blood, cortex, and

cerebellum: implications for epigenetic studies of neurological and neuropsychiatric phenotypes.

Epigenetics [Internet] 2015;**10**:1024–1032.

Hata T, Nakayama M. Targeted disruption of the murine large nuclear KIAA1440/Ints1 protein causes

growth arrest in early blastocyst stage embryos and eventual apoptotic cell death. *Biochim Biophys*

Acta [Internet] 2007;**1773**:1039–1051.

Heijmans BT, Mill J. Commentary: The seven plagues of epigenetic epidemiology. *IntJEpidemiol*

2012;**41**:74–78. Molecular Epidemiology, Leiden University Medical Center, Leiden, The

Netherlands, Netherlands. bas.heijmans@lumc.nl.

Hsiao K, Harony-Nicolas H, Buxbaum JD, Bozdagi-Gunal O, Benson DL. Cyfip1 Regulates Presynaptic

Activity during Development. *J Neurosci* [Internet] 2016;**36**:1564–1576.

Iliadou AN, Öberg AS, Pege J, Rodriguez-Wallberg KA, Olofsson JI, Holte J, Wramsby H, Wramsby M,

Cnattingius S, Cesta CE. The Uppsala-Stockholm Assisted Reproductive Techniques (UppStART)

study. *BMJ Open* 2019;**9**:. BMJ Publishing Group.

Iterson M van, Cats D, Hop P, Heijmans BT. omicsPrint: detection of data linkage errors in multiple omics

studies. In Stegle O, editor. *Bioinformatics* [Internet] 2018;**34**:2142–2143.

Iterson M van, Tobi EW, Slieker RC, Hollander W den, Luijk R, Slagboom PE, Heijmans BT. MethylAid:

visual and interactive quality control of large Illumina 450k datasets. *Bioinformatics* [Internet]

2014;**30**:3435–3437. Department of Molecular Epidemiology, Leiden University Medical Center,

2333 ZC Leiden, the Netherlands Department of Molecular Epidemiology, Leiden University

Medical Center, 2333 ZC Leiden, the Netherlands Department of Molecular Epidemiology, Leiden

Un.

531 Jaenisch R, Bird A. Epigenetic regulation of gene expression: how the genome integrates intrinsic and

532   environmental signals. *Nat Genet* 2003;**33 Suppl**:245–254. United States.

533 Jones SC, Jordan WJ, Meiners SJ, Miller AN, Methven AS. Fungal spore dispersal by the Eastern box turtle

534   (Terrapene carolina carolina). *Am Midl Nat* [Internet] 2007;**157**:121–126.

535 Joubert BR, Dekker HT den, Felix JF, Bohlin J, Ligthart S, Beckett E, Tiemeier H, Meurs JB van, Uitterlinden

536   AG, Hofman A, *et al.* Maternal plasma folate impacts differential DNA methylation in an

537   epigenome-wide meta-analysis of newborns. *Nat Commun* [Internet] 2016a;**7**:10577.

538 Joubert BR, Felix JF, Yousefi P, Bakulski KM, Just AC, Breton C, Reese SE, Markunas CA, Richmond RC, Xu

539   CJ, *et al.* DNA Methylation in Newborns and Maternal Smoking in Pregnancy: Genome-wide

540   Consortium Meta-analysis. *Am J Hum Genet* [Internet] 2016b;**98**:680–696.

541 Kazmi N, Sharp GC, Reese SE, Vehmeijer FO, Lahti J, Page CM, Zhang W, Rifas-Shiman SL, Rezwan FI,

542   Simpkin AJ, *et al.* Hypertensive Disorders of Pregnancy and DNA Methylation in Newborns.

543   *Hypertens (Dallas, Tex 1979)* [Internet] 2019;**74**:375–383.

544 Kessler NJ, Waterland RA, Prentice AM, Silver MJ. Establishment of environmentally sensitive DNA

545   methylation states in the very early human embryo. *Sci Adv* [Internet] 2018;**4**:eaat2624.

546 Koestler DC, Jones MJ, Usset J, Christensen BC, Butler RA, Kobor MS, Wiencke JK, Kelsey KT. Improving

547   cell mixture deconvolution by identifying optimal DNA methylation libraries (IDOL). *BMC*

548   *Bioinformatics* [Internet] 2016;**17**:120.

549 Krieger N, Davey Smith G. The tale wagged by the DAG: broadening the scope of causal inference and

550   explanation for epidemiology. *Int J Epidemiol* [Internet] 2016;**45**:1787–1808. Oxford University

551   Press.

552 Küpers LK, Monnereau C, Sharp GC, Yousefi P, Salas LA, Ghantous A, Page CM, Reese SE, Wilcox AJ,

Czamara D, *et al.* Meta-analysis of epigenome-wide association studies in neonates reveals

widespread differential DNA methylation associated with birthweight. *Nat Commun* 2019;**10**:.

Nature Publishing Group.

Lawlor DA, Tilling K, Davey Smith G. Triangulation in aetiological epidemiology. *Int J Epidemiol* [Internet]

2017;dyw314Available from: https://academic.oup.com/ije/article-

lookup/doi/10.1093/ije/dyw314.

Lazaraviciute G, Kauser M, Bhattacharya S, Haggarty P, Bhattacharya S. A systematic review and meta-

analysis of DNA methylation levels and imprinting disorders in children conceived by IVF/ICSI

compared with children conceived spontaneously. *Hum Reprod Update* [Internet] 2014;**20**:840–

852.

Li S, Wong EM, Dugué P-A, McRae AF, Kim E, Joo J-HE, Nguyen TL, Stone J, Dite GS, Armstrong NJ, *et al.*

Genome-wide average DNA methylation is determined in utero. *Int J Epidemiol* [Internet]

2018;**47**:908–916.

Liu L, Gao J, He X, Cai Y, Wang L, Fan X. Association between assisted reproductive technology and the

risk of autism spectrum disorders in the offspring: a meta-analysis. *Sci Rep* [Internet] 2017;**7**:46207.

Nature Publishing Group.

Loke YJ, Craig JM. Are the effects of IVF on DNA methylation driven by intracytoplasmic sperm injection

and male infertility? *Epigenomics* [Internet] 2016;**8**:881–884. Future Medicine Ltd London, UK.

Luke B, Stern JE, Hornstein MD, Kotelchuck M, Diop H, Cabral H, Declercq ER. Is the wrong question

being asked in infertility research? *J Assist Reprod Genet* [Internet] 2016;**33**:3–8. Springer New York

LLC.

Martínez-García MA, San-Millán JL, Escobar-Morreale HF. The R453Q and D151A polymorphisms of

Hexose-6-Phosphate Dehydrogenase Gene (H6PD) influence the polycystic ovary syndrome (PCOS)

and obesity. *Gene* [Internet] 2012;**497**:38–44.

Maulik U, Sen S, Mallik S, Bandyopadhyay S. Detecting TF-miRNA-gene network based modules for

5hmC and 5mC brain samples: a intra- and inter-species case-study between human and rhesus.

BMC Genet [Internet] 2018;**19**:9.

Melamed N, Choufani S, Wilkins-Haug LE, Koren G, Weksberg R. Comparison of genome-wide and gene-

specific DNA methylation between ART and naturally conceived pregnancies. *Epigenetics* [Internet]

2015;**10**:474–483. Taylor & Francis.

Morgan HD, Jin XL, Li A, Whitelaw E, O'Neill C. The culture of zygotes to the blastocyst stage changes the

postnatal expression of an epigentically labile allele, agouti viable yellow, in mice. *BiolReprod*

2008;**79**:618–623. Royal North Shore Hospital, Disciplines of Medicine and Physiology, University of

Sydney, Sydney, 2065 New South Wales, Australia.

Novakovic B, Lewis S, Halliday J, Kennedy J, Burgner DP, Czajko A, Kim B, Sexton-Oates A, Juonala M,

Hammarberg K, *et al.* Assisted reproductive technologies are associated with limited epigenetic

variation at birth that largely resolves by adulthood. *Nat Commun* [Internet] 2019;**10**:3922.

Springer US.

Pallares P, Gonzalez-Bulnes A. The effect of embryo and maternal genotypes on prolificacy, intrauterine

growth retardation and postnatal development of Nos3-knockout mice. *Reprod Biol* 2010;**10**:241–

248. Elsevier B.V.

Phipson B, Maksimovic J, Oshlack A. missMethyl: an R package for analyzing data from Illumina's

HumanMethylation450 platform. *Bioinformatics* [Internet] 2015;**32**:btv560.

Qin J, Liu X, Sheng X, Wang H, Gao S. Assisted reproductive technology and the risk of pregnancy-related

complications and adverse pregnancy outcomes in singleton pregnancies: A meta-analysis of

cohort studies. *Fertil Steril* 2016;**105**:73-85.e6. Elsevier Inc.

R core team. R: A language and environment for statistical computing. In Team RDC, editor. *R Found Stat*

Comput [Internet] 2019; R Foundation for Statistical ComputingAvailable from: http://www.r-

project.org/.

Relton CL, Gaunt T, McArdle W, Ho K, Duggirala A, Shihab H, Woodward G, Lyttleton O, Evans DM, Reik

W, *et al.* Data Resource Profile: Accessible Resource for Integrated Epigenomic Studies (ARIES).

IntJEpidemiol 2015;**44**:1181–1190. MRC Integrative Epidemiology Unit, and School of Social and

Community Medicine, University of Bristol, Bristol, UK, Institute of Genetic Medicine, Newcastle

University, Newcastle upon Tyne, UK, caroline.relton@bristol.ac.uk MRC Integrative Epidemiology

Un.

Salas LA, Koestler DC, Butler RA, Hansen HM, Wiencke JK, Kelsey KT, Christensen BC. An optimized

library for reference-based deconvolution of whole-blood biospecimens assayed using the Illumina

HumanMethylationEPIC BeadArray. *Genome Biol* [Internet] 2018;**19**:64.

Sharma N, Kubaczka C, Kaiser S, Nettersheim D, Mughal SS, Riesenberg S, Hölzel M, Winterhager E,

Schorle H. *Tpbpa-Cre* -mediated deletion of TFAP2C leads to deregulation of *Cdkn1a* , *Akt1* and the

ERK pathway, causing placental growth arrest. *Development* [Internet] 2016;**143**:787–798.

Sharp GC, Salas LA, Monnereau C, Allard C, Yousefi P, Everson TM, Bohlin J, Xu Z, Huang R-C, Reese SE, *et*

al. Maternal BMI at the start of pregnancy and offspring epigenome-wide DNA methylation:

findings from the pregnancy and childhood epigenetics (PACE) consortium. *Hum Mol Genet*

[Internet] 2017;**26**:4067–4085. Oxford University Press.

Slieker RC, Bos SD, Goeman JJ, Bovée JV, Talens RP, Breggen R van der, Suchiman HED, Lameijer E-W,

Putter H, Akker EB van den, *et al.* Identification and systematic annotation of tissue-specific

differentially methylated regions using the Illumina 450k array. *Epigenetics Chromatin* [Internet]

2013;**6**:26. Molecular Epidemiology, Leiden University Medical Center, Leiden, The Netherlands.

B.T.Heijmans@lumc.nl.

Smew AI, Hedman AM, Chiesa F, Ullemar V, Andolf E, Pershagen G, Almqvist C. Limited association

between markers of stress during pregnancy and fetal growth in 'Born into Life', a new prospective

birth cohort. *Acta Paediatr Int J Paediatr* 2018;**107**:1003–1010.

Stojnic R, Diez D. PWMEnrich: PWM enrichment analysis R package version. 2018;Available from:

https://bioconductor.org/packages/release/bioc/html/PWMEnrich.html.

Sungur AÖ, Schwarting RKW, Wöhr M. Behavioral phenotypes and neurobiological mechanisms in the

Shank1 mouse model for autism spectrum disorder: A translational perspective. *Behav Brain Res*

[Internet] 2018;**352**:46–61. Elsevier B.V.

Tanaka M, DeLorey TM, Delgado-Escueta A, Olsen RW. *GABRB3, Epilepsy, and Neurodevelopment*

[Internet]. *Jasper's Basic Mech Epilepsies* [Internet] 2012;**51**:.

Tobi EW, Goeman JJ, Monajemi R, Gu H, Putter H, Zhang Y, Slieker RC, Stok AP, Thijssen PE, Müller F, *et*

al. DNA methylation signatures link prenatal famine exposure to growth and metabolism. *Nat*

Commun [Internet] 2014;**5**:5592.

Tobi EW, Slagboom PE, Dongen J van, Kremer D, Stein AD, Putter H, Heijmans BT, Lumey LH. Prenatal

famine and genetic variation are independently and additively associated with dna methylation at

regulatory loci within IGF2/H19. *PLoS One* 2012;**7**:.

Tobi EW, Slieker RC, Luijk R, Dekkers KF, Stein AD, Xu KM, Slagboom PE, Zwet EW van, Lumey LH,

Heijmans BT. DNA methylation as a mediator of the association between prenatal adversity and

risk factors for metabolic disease in adulthood. *Sci Adv* [Internet] 2018;**4**:eaao4364. American

Association for the Advancement of Science.

Turkgeldi E, Yagmur H, Seyhan A, Urman B, Ata B. Short and long term outcomes of children conceived

with assisted reproductive technology. *Eur J Obstet Gynecol Reprod Biol* [Internet] 2016;**207**:129–

136. Elsevier.

Vermey BG, Buchanan A, Chambers GM, Kolibianakis EM, Bosdou J, Chapman MG, Venetis CA. Are

singleton pregnancies after assisted reproduction technology (ART) associated with a higher risk of

placental anomalies compared with non-ART singleton pregnancies? A systematic review and

meta-analysis. *BJOG An Int J Obstet Gynaecol* [Internet] 2019;**126**:209–218. Blackwell Publishing

Ltd.

Wang J, Zhao Q. The CATE Package for High Dimensional Factor Analysis and Confounder Adjusted

Multiple Testing. 2015; Comprehensive R Archive Network (CRAN)Available from: https://cran.r-

project.org/package=cate.

Wang Z, Xu L, He F. Embryo vitrification affects the methylation of the H19/Igf2 differentially methylated

domain and the expression of H19 and Igf2. *Fertil Steril* [Internet] 2010;**93**:2729–2733. Elsevier.

Xu N, Barlow GM, Cui J, Wang ET, Lee B, Akhlaghpour M, Kroener L, Williams J, Rotter JI, Chen YI, *et al.*

Comparison of Genome-Wide and Gene-Specific DNA Methylation Profiling in First-Trimester

Chorionic Villi From Pregnancies Conceived With Infertility Treatments. *Reprod Sci* [Internet]

2017;**24**:996–1004.

Yuen RK, Jiang R, Peñaherrera MS, McFadden DE, Robinson WP. Genome-wide mapping of imprinted

differentially methylated regions by DNA methylation profiling of human placentas from triploidies.

Epigenetics Chromatin [Internet] 2011;**4**:10. BioMed Central.

Zeileis A. Object-oriented computation of sandwich estimators. *J Stat Softw* 2006;**16**:1–16. American

Statistical Association.

Zheng Y, Joyce BT, Liu L, Zhang Z, Kibbe WA, Zhang W, Hou L. Prediction of genome-wide DNA

methylation in repetitive elements. *Nucleic Acids Res* [Internet] 2017;**45**:8697–8711.

Tables

Table 1. Cohort characteristics

Variable	MUC[#]	ART	IVF	ICSI
N	70	87	44	33
Frozen (N)[1]	-	10	7	3
Blastocyst (N)[2]	-	8	5 (*3 frozen*)	3 (*2 frozen*)
BMI in kg/m^2 (SD)	22.8 (3.5)	23.4 (3.4)	23.5 (3.3)	23.4 (3.0)
Age in y(SD)	32.4 (3.6)	33.1 (3.6)	33.0 (3.7)	33.0 (3.7)
SES[3] (SD)	3.0 (0.3)	2.6 (0.8)[*]	2.7 (0.7)	2.3 (0.9)
Smoking		[*]		
Never	65	56	27	23
Ever	4	26	17	9
NA	1	5	0	1
Years to index pregnancy				
Mean (SD)	0.2 (0.7)	2.6 (1.3)[*]	2.7 (1.2)	2.4 (1.5)
Min - max	0- 4	0 – 7	1- 7	0-7
Parity				
First	47	62	34	22
Second	21	17	10	7
Third	2	1	0	1
NA	0	7	0	3
N Male children (%)	43 (61.4)	44 (50.6)	22 (50.0)	16 (48.5)
Gestational age in weeks	39.1 (1.6)	39.5 (2.0)	39.2 (2.3)	39.7 (1.6)

1. The number of embryos that were frozen and thawed before placement *in utero*.
2. The number of embryos that were grown *ex utero* to the blastocyst stage before placement *in utero*. Off note, five of these blastocysts were frozen and thawed.
3. Socio-economic status (SES) graded on a 4 level scale based on the highest attained educational level.
[*]Significantly different P<0.001, [#] medically unassisted conception

CpG ID	Location hg19	Nearest gene[1]	distance[1]	Estimate (SE)[2]	P[2]	P_{FDR}
cg27266479	chr1:9294882	H6PD	0	-1.94(0.27)	3.18e-13	8.28E-08
cg04811592	chr3:69834386	MITF	0	0.99(0.21)	1.59e-06	0.039
cg24959663	chr5:10578618	ANKRD33B	0	3.91(0.54)	3.75e-13	8.28E-08
cg22916646	chr5:162672583	-	-	2.18(0.36)	1.22e-09	1.08E-04
cg01500567	chr6:44355777	CDC5L	0	0.48(0.1)	5.46e-07	0.020
cg00478390	chr7:150703765	NOS3	0	-0.92(0.19)	7.37e-07	0.024
cg03207674	chr7:1523569	INTS1	0	0.72(0.14)	1.92e-07	9.43E-03
cg17123384	chr7:83379152	-	-	2.78(0.51)	6.39e-08	3.53E-03
cg19347588	chr10:3868336	KLF6	40862	0.81(0.16)	4.49e-07	0.018
cg07569385	chr13:20766226	GJB2	0	1.53(0.31)	1.24e-06	0.036
cg06485032	chr13:22615064	AK054845	0	-3.47(0.67)	2.59e-07	0.011
cg13051607	chr15:22956714	CYFIP1	0	1.39(0.24)	1.15e-08	7.24 E-04
cg01251603	chr15:26874098	GABRB3	0	-4.89(0.8)	1.18e-09	1.08 E-04
cg15066197	chr15:26874202	GABRB3	0	-4.71(0.79)	2.48e-09	1.83 E-04
cg14859324	chr15:26874363	GABRB3	0	-2.26(0.46)	7.68e-07	0.024
cg06450634	chr16:30430044	ZNF771	0	3.26(0.46)	2.15e-12	3.17E-07
cg08783253	chr17:40996565	AOC2	42	-2.93(0.61)	1.48e-06	0.038
cg16771467	chr18:55315872	ATP8B1	0	0.11(0.02)	1.34e-06	0.037
cg14560133	chr19:51199453	SHANK1	0	-1.42(0.3)	2.07e-06	0.048

679 680 681 682 683 684 685 686 687 688

689 The results of the ART vs MUC EWAS are ordered on genomic location (hg19).

690   1.   Nearest gene within 100kb and the distance to the TSS.

691   2.   Estimate and standard error (SE) and P value, with and without false discovery rate (FDR) correction of the

692   estimate of ART vs unassisted pregnancies in the model: Beta ~ ART (yes/no) + height on micro-array slide

693   + scan batch + bisulfite plate + sex + CD4T + CD8T + Mono + Bcell + Nk + Gran + nRBC.

Figures

 Figure 1. Manhattan plot of EWAS for ART status

 Figure 1. A Manhattan plot showing the −log10 P-values (y-axis) for the association of DNA methylation
 at individual CpG dinucleotides across the 22 autosomal chromosomes (x-axis). CpG dinucleotides 5kb
 up- and downstream of the lead association have been colored in the same color as the lead association.

Figure 2. correlation between adult blood DNA methylation and internal tissues

Figure 2. The spearman correlation plotted for 8 tissues from the body (N=16, P<0.1) and 4 from the
brain (N>71, P<0.01) in colors ranging from bright red (*rho*= -1.0) to dark blue (*rho*=1.0). The light gray
lines along the rows demark three CpG dinucleotides from the *GABRB3* region associated with ART. The
light gray line running along one column denotes the separation of the two tissue reference datasets
used. *"Fat sub": subcutaneous fat, "Left myocard": left myocardium, "Muscle Skel": skeleton muscle,*
"prefr. Cortex": prefrontal cortex, "ent.cortex": entorhinal cortex,"s.t.gyrus": superior temporal gyrus.

717