

1 **Change in outbreak epicenter and its impact on the**
2 **importation risks of COVID-19 progression: a modelling**
3 **study**

4
5 Oyelola A Adegboye^{1*}, Adeshina I Adekunle¹, Anton Pak¹, Ezra Gayawan², Denis HY Leung³,
6 Diana P Rojas⁴, Faiz Elfaki⁵, Emma S McBryde¹, Damon P Eisen¹

7
8 ¹Australian Institute of Tropical Health and Medicine, James Cook University, Townsville,
9 Australia

10 ²Biostatistics and Spatial Statistics Research Group, Department of Statistics, Federal
11 University of Technology, Akure, Nigeria.

12 ³School of Economics, Singapore Management University, Singapore, Singapore

13 ⁴College of Public Health. Medical and Veterinary Sciences, James Cook University,
14 Townsville, Australia

15 ⁵Department of Mathematics, Statistics and Physics, Qatar University, Doha, Qatar

16

17 *Corresponding author:

18 Oyelola A. Adegboye, Email: oyelola.adegboye@jcu.edu.au, Phone: +61 7 4781 5707.

19 Australian Institute of Tropical Health and Medicine, James Cook University, 1 James Cook
20 Drive, Douglas, Queensland, Australia, 4814.

21

22

23 [Abstract](#)

24 The outbreak of Severe Acute Respiratory Syndrome Coronavirus-2 (SARS-CoV-2) that
25 originated in the city of Wuhan, China has now spread to every inhabitable continent, but
26 now the attention has shifted from China to other epicenters, especially Italy. This study
27 explored the influence of spatial proximities and travel patterns from Italy on the further
28 spread of SARS-CoV-2 around the globe. We showed that as the epicenter changes, the
29 dynamics of SARS-CoV-2 spread change to reflect spatial proximities.

30

31

32 As at March 10th 2020, 109 countries and territories have confirmed at least one
33 case of COVID-19 and a total number of 113,702 COVID-19 cases have been confirmed
34 globally – 80,924 in China and 32,778 outside China [1]. Human population movement plays
35 an important role in the spread of infectious diseases. Although China has served as the
36 epicenter of the current SARS-CoV-2 outbreak, real-time phylogenetic analysis [2] suggests
37 that the virus was introduced to most African countries and to Latin America from people
38 traveling from Italy. Migration has remained the major source of concern for the current
39 COVID-19 outbreak and most countries have focused on China as the likely source of any
40 importation. The movement of people between China and sub-Saharan Africa has increased
41 rapidly over the last decade [3, 4], the spread of COVID-19 to the African continent has been
42 more related to the current COVID-19 outbreak in Italy. There has been much attention on
43 the importation of infectious diseases such as Ebola, Tuberculosis, Malaria or viral hepatitis
44 from Africa to Europe [5, 6]. Conversely, there are few reported cases of importing diseases
45 from the European countries to Africa. This, perhaps, makes most African countries to focus
46 their COVID-19 surveillance efforts on travelers from China without much attention paid to
47 the possible importation from other countries.

48 In addition to the phylogenetic analysis, a number of other circumstantial factors
49 support the introduction of the virus to other countries, as it appears more likely that the
50 introduction of COVID-19 to South America occurred from Italy rather than from China.
51 First, since January 23, 2020, China has increased their containment measures and
52 decreased the number of outbound international flights. Second, early febrile airport
53 screening mostly targeted travelers coming from Asia overlooking those from other parts of
54 the world – including travelers from Europe; an approach that turned out to have limited
55 effectiveness [7]. Third, as most of the infections by SARS-CoV-2 are asymptomatic and mild

56 [8-10], it is more likely that infected carriers could enter a country without being detected
57 by the temperature screening at the airports. Lastly, Italian Brazilians are the largest group
58 of foreigners with full or partial Italian ancestry outside Italy, with São Paulo being the most
59 populous city with Italian ancestry in the world. In this paper, we investigated how the
60 growing epidemic in Italy affects the spread of the epidemic around the world by using
61 measurements of spatial proximity and travel volume between countries.

62 [Methods](#)

63 This study is based on the number of confirmed cases of COVID-19 in each country until
64 March 10th, 2020 obtained from World Health Organisation (WHO) situation reports [1] and
65 real-time online COVID-19 monitoring sites [11, 12]. The air travel data between countries
66 were obtained from the Official Airline Guide (OAG) database (www.oag.com). Preliminary
67 analysis includes Pearson's correlation analysis to examine the association between the
68 number of COVID-19 cases and travel influx (and arrival time) from the source country.
69 Spatial data reveal the degree of dependency among observations in geographical space
70 [13, 14]; consequently, spatial dependence measures such semi-variogram and Local
71 Indicator of Spatial Association (LISA) [15] were used to detect spatial patterns in COVID-19
72 data. For the spatial weights, we used Queen-style contiguity 1st order nearest neighbour
73 (i.e., two countries are neighbouring if they share common borders or a point). Additionally,
74 we developed a stochastic meta-population model of global spread using the 2018 IATA
75 flight data for travel volumes between each country. The country-specific compartmental
76 model is an adaptation of the classical SEIR (Susceptible-Exposed-Infectious-Recovered)
77 model based on reported dynamics of the novel coronavirus [16, 17] for global spread of
78 COVID-19 (see the appendix for model details and www.pandemic.org.au for deterministic
79 version) [18]. We used the cases on WHO situation report [1] to initialise the number of

80 people infected in Italy and assuming same for China to determine global risk from these
81 two epicenters under the same initial infection seed. Each country's risk of importation was
82 classified into quartiles: slight risk (<25%), moderate risk (25%-50%), high risk (50%-75%)
83 and extreme risk (>75%).

84 Results and Discussion

85 Up until the February 21st Italian outbreak, all reported European cases were
86 imported from China (Figure 1) [19]. The disease has now (as at March 10th) spread to 46
87 (excluding Vatican City and Liechtenstein) out of the 53 countries within the WHO European
88 region [1]. Italy has 9,172 confirmed cases and 463 deaths as of March 10th. The region
89 alone accounted for 46% of the total cases outside of China. There are currently three
90 COVID-19 epicenters –China, Italy, and Iran, which are mostly responsible for introducing
91 the virus globally (Figure 1).

92 We present in Figure 2, the distribution of time elapsed since 21st February when the
93 first imported case of COVID-19 from Italy was reported in each country against the
94 geographical distance from Italy. The outbreak in Italy is highly sporadic in the last three
95 weeks. While the shortest arrival time (after the Italy outbreak) was 5 days to Austria,
96 Switzerland, Brazil and Algeria, the virus spread to 44 countries within 17 days (median, 11
97 days). The global arrival times for three epicenters from the first reported cases in Wuhan,
98 China are shown in Figure S1.

99 The Pearson's correlation analysis ($r = -0.43$, $p = 0.004$) suggest negative association
100 between disease arrival time and number of cases for cases imported from Italy. That is,
101 countries with shorter introduction time have more number of cases. There was significant
102 association between the number of cases and daily travel influx from Italy ($r = 0.39$, $p = 0.011$).
103 The apparent spatial autocorrelation in the COVID cases within Europe (Moran's

104 Index=0.310, $p < 0.001$) is also significant (Figure S2). Notably, France and Italy are located
105 in the right upper quadrant, while Switzerland, Austria and Slovakia are on the left upper
106 quadrant, indicating a positive and negative spatial autocorrelation pattern respectively.
107 Additionally, semi-variograms indicated spatial autocorrelation of the disease incidence
108 exists up to a distance of 120 decimal degrees for Italy (Figure S3). Using bivariate LISA
109 analysis, we found evidence of spatial interaction between COVID-19 cases and arrival time
110 as well as travel influx (Moran's $I = 0.340$). This suggests that spatial variations within
111 European region were non-random, exhibiting effects of neighbouring interactions and
112 travel influx.

113 We simulated COVID-19 spread comparing two outbreak epicenters – Italy and
114 China. Figure 3 shows the relative risk of importation from the two epicenters. The results
115 show that whereas the Asia-Pacific is more at risk (high risk to extreme) from China
116 epicenter, Europe, South-America and Africa are more at risk from the Italian epicenter.
117 High population density and highly interconnected transportation networks connecting
118 tourism hubs in Northern Italy with major European cities have made it extremely difficult
119 to contain and reduce the number of infections. This aligns well with our prediction of
120 extreme risks of case importations from Italy to other European, while Latin American and
121 African countries have high risks (Figure 3). In the case of Italy, our modelling results could
122 provide an argument for the Italian government instituting a national quarantine and travel
123 restrictions on March 9th, 2020 to mitigate consequent spread of infection. Based on the
124 evidence from the Chinese policies of containment and quarantine that showed
125 considerable effectiveness [20], these interventions should slow the spread of the disease
126 within Europe and abroad.

127 Although most African countries have focused their COVID-19 surveillance efforts on
128 travelers from China, our analysis shows that this approach may need to be reconsidered as
129 the number of cases in Europe soar, making the risk of importation to Africa higher from
130 European countries than from China through stronger transportation connectivity and
131 migration flows (especially, with North African countries). As at now, Italy has put itself
132 under lockdown [21] and other countries such as Australia and the USA have restricted
133 flights from Italy [22, 23]. As new epicenters are emerging, countries are required to adapt
134 quickly and adjust their containment measures to reduce the spread of infection.

135 [Conclusion](#)

136 In summary, we estimated the risk of importation from two major epicenters to
137 other countries and showed that spatial proximity and mobility are important factors that
138 fuel disease importation. This analysis illustrates the potential development of the infection
139 spread and may be useful in countries' epidemic preparedness including large-scale
140 interventions such as travel restrictions and quarantine.

141

142 References


- 143 1. World Health Organization. *Coronavirus disease (COVID-2019) situation reports*. 2020 March
144 2, 2020]; Available from: [https://www.who.int/emergencies/diseases/novel-coronavirus-](https://www.who.int/emergencies/diseases/novel-coronavirus-2019/situation-reports)
145 [2019/situation-reports](https://www.who.int/emergencies/diseases/novel-coronavirus-2019/situation-reports).
- 146 2. Nextstrain. *Genomic epidemiology of novel coronavirus (hCoV-19)*. 2020 March 4, 2020];
147 Available from: <https://nextstrain.org/ncov>.
- 148 3. Pigato, M. and W. Tang, *China and Africa: Expanding economic ties in an evolving global*
149 *context. Investing in Africa Forum*. World Bank Working Paper, 2015. **95161**.
- 150 4. Shen, X. *Chinese Manufacturers Moving to Africa—Who? What? Where? Does Africa*
151 *Benefit?* in conference “China in the World Economy: Building a New Partnership with
152 *Africa.” World Bank, Washington, DC, May. 2014.*
- 153 5. Castelli, F. and G. Sulis, *Migration and infectious diseases*. *Clinical Microbiology and*
154 *Infection*, 2017. **23**(5): p. 283-289.
- 155 6. Khyatti, M., et al., *Infectious diseases in North Africa and north African immigrants to*
156 *Europe*. *The European Journal of Public Health*, 2014. **24**(suppl_1): p. 47-56.
- 157 7. Quilty, B.J. and S. Clifford, *Effectiveness of airport screening at detecting travellers infected*
158 *with novel coronavirus (2019-nCoV)*. *Eurosurveillance*, 2020. **25**(5).
- 159 8. Zou, L., et al., *SARS-CoV-2 viral load in upper respiratory specimens of infected patients*. *New*
160 *England Journal of Medicine*, 2020.
- 161 9. Rothe, C., et al., *Transmission of 2019-nCoV infection from an asymptomatic contact in*
162 *Germany*. *New England Journal of Medicine*, 2020.
- 163 10. Wu, Z. and J.M. McGoogan, *Characteristics of and important lessons from the coronavirus*
164 *disease 2019 (COVID-19) outbreak in China: summary of a report of 72 314 cases from the*
165 *Chinese Center for Disease Control and Prevention*. *Jama*, 2020.
- 166 11. Wu, T., X. Ge, and G. Yu, *An R package and a website with real-time data on the COVID-19*
167 *coronavirus outbreak*. medRxiv, 2020.
- 168 12. Worldometer. *COVID-19 CORONAVIRUS OUTBREAK*. 2020 [cited 2020 March 5, 2020];
169 Available from: <https://www.worldometers.info/coronavirus/>.
- 170 13. Liu, W., et al., *Spatial and temporal analysis of human infection with avian influenza A*
171 *(H7N9) virus in China, 2013*. *Eurosurveillance*, 2013. **18**(47): p. 20640.
- 172 14. Adegboye, O.A., D.H. Leung, and Y.G. Wang, *Analysis of spatial data with a nested*
173 *correlation structure*. *Journal of the Royal Statistical Society: Series C (Applied Statistics)*,
174 2018. **1**(67): p. 329-354.
- 175 15. Anselin, L., *A local indicator of multivariate spatial association: extending Geary's C*.
176 *Geographical Analysis*, 2019. **51**(2): p. 133-150.
- 177 16. Lin, Q., et al., *Modeling the spread of Middle East respiratory syndrome coronavirus in Saudi*
178 *Arabia*. *Statistical methods in medical research*, 2018. **27**(7): p. 1968-1978.
- 179 17. Funk, S. and R.M. Eggo, *Early dynamics of transmission and control of 2019-nCoV: a*
180 *mathematical modelling study*.
- 181 18. Bosco Ho, A.I.A., James Trauer, Emma McBryde *Global Pandemic Map*. 2019.
- 182 19. Stoecklin, S.B., et al., *First cases of coronavirus disease 2019 (COVID-19) in France:*
183 *surveillance, investigations and control measures, January 2020*. *Eurosurveillance*, 2020.
184 **25**(6): p. 2000094.
- 185 20. Chinazzi, M., et al., *The effect of travel restrictions on the spread of the 2019 novel*
186 *coronavirus (COVID-19) outbreak*. *Science*, 2020.
- 187 21. CNBC. *Italy expands its quarantine to the entire country as coronavirus cases and deaths*
188 *surge*. 2020 [cited 2020 11/03/2020]; Available from:
189 [https://www.cnbc.com/2020/03/09/italy-extends-its-quarantine-to-the-entire-country-pm-](https://www.cnbc.com/2020/03/09/italy-extends-its-quarantine-to-the-entire-country-pm-asks-residents-to-stay-at-home.html)
190 [asks-residents-to-stay-at-home.html](https://www.cnbc.com/2020/03/09/italy-extends-its-quarantine-to-the-entire-country-pm-asks-residents-to-stay-at-home.html).

191 22. Times, T.N.Y. *U.S. to Suspend Most Travel From Europe; N.B.A. Pauses After Player Gets*
192 *Virus*. 2020 [cited 2020 12/03/2020]; Available from:
193 <https://www.nytimes.com/2020/03/11/world/coronavirus-news.html>.
194 23. VOA. *Australia Extends Virus Travel Ban to Italy*. 2020 [cited 2020 12/03/2020]; Available
195 from: [https://www.voanews.com/science-health/coronavirus-outbreak/australia-extends-](https://www.voanews.com/science-health/coronavirus-outbreak/australia-extends-virus-travel-ban-italy)
196 [virus-travel-ban-italy](https://www.voanews.com/science-health/coronavirus-outbreak/australia-extends-virus-travel-ban-italy).
197 24. Riou, J., et al., *Adjusted age-specific case fatality ratio during the COVID-19 epidemic in*
198 *Hubei, China, January and February 2020*. medRxiv, 2020.
199 25. Team, R.C., *R: a language and environment for statistical computing, version 3.0. 2*. Vienna,
200 *Austria: R Foundation for Statistical Computing; 2013*. 2019.

201

202


203 **Figures legend**




204

205 Figure 1: Transmission routes of COVID-19 as at March 7th 2020. The lines represent
206 transmission routes from the source of COVID-19 into a country. Inset 1: European cases
207 originating from China (green lines) and Iran (orange lines). Inset 2: Cases originating from
208 Italy.

209




210




211

212 Figure 2: Distribution of arrival times of COVID-19 cases against geographic distance from
213 the source (Italy). The dots are proportional to the size of, (A) total cases, and (B) travel
214 influx from the source country.
215



216



217

218 Figure 3: Global risk of importation of COVID-19 from, (A) Italy (purple dot) and (B) China

219 (red dot) based on air travel influx from source country. The dots are proportional to the

220 cumulative relative risk that an infected individual will be arriving at a specific country for an

221 epicenter. Risk of importation: slight risk (<25%), moderate risk (25%-50%), high risk (50%-

222 75%) and extreme risk (>75%).

223

224 **Appendix 1**

225

226 **Table S1: State variable and parameter descriptions**


Variable	Description	
S	Susceptible class	
E_1	Latent stage not infectious	
E_2	Latent stage infectious	
I_1	First stage of symptomatic	
I_2	Second stage of symptomatic	
R	Recovered class	

Parameter	Description	Values (references)
$\beta_1, \beta_2, \beta_3$	Transmission rates for E_2, I_1, I_2 classes	0.3125, 0.5, 0.176 ([17])
σ_1	First stage incubation rate	0.3125 ([17])
σ_2	Second stage incubation rate	0.5 ([17])
γ_1	First stage of recovery	0.5 ([17])
γ_2	second stage of recovery	0.176 ([17])
c_h	COVID-19 case fatality	0.018 ([24])
μ	Death rate	Not used

227

228

229




230

231 Figure S1. Distribution of COVID-19 arrival times (days from first reported cases in China)


232 and distance from China. The size of the dot is proportional to the daily air travellers

233 between China and other countries as at March 10th, 2020.



234


235 Figure S2. Moran's Index for COVID-19 cases in Europe



236

237 Figure S3: Semi-variogram showing spatial autocorrelation for COVID-19 data up 0.000

238 decimal degrees (~1.2km) for Italy and ...for China.



239

240 Figure S4: Schematic diagram of the transmission dynamics

241

242

243

244 **Appendix 2: Mathematical Model details:**

245 The infectious classes are late latency E_2 and the two stages of symptomatic infectiousness
246 (I_1 and I_2). Patients either recover and moved to R class or die and are replaced to ensure
247 constant population. The two-country dynamical model is as shown below:

248
$$\frac{dS^1}{dt} = -\frac{(\beta_1^1 E_2^1 + \beta_2^1 I_1^1 + \beta_3^1 I_2^1)S^1}{N_1} + c_d^1 \gamma_2^1 I_2^1 - \frac{m_{12}S^1}{N_1} + \frac{m_{21}S^2}{N_2} \quad (1)$$

249
$$\frac{dE_1^1}{dt} = \frac{(\beta_1^1 E_2^1 + \beta_2^1 I_1^1 + \beta_3^1 I_2^1)S^1}{N_1} - \sigma_1 E_1^1 - \frac{m_{12}E_1^1}{N_1} + \frac{m_{21}E_1^2}{N_2}$$

250
$$\frac{dE_2^1}{dt} = \sigma_1 E_1^1 - \sigma_2 E_2^1 - \frac{m_{12}E_2^1}{N_1} + \frac{m_{21}E_2^2}{N_2}$$

251
$$\frac{dI_1^1}{dt} = \sigma_1 E_2^1 - \gamma_1^1 I_1^1 - \frac{m_{12}I_1^1}{N_1} + \frac{m_{21}I_1^2}{N_2}$$

252
$$\frac{dI_2^1}{dt} = \gamma_1^1 I_1^1 - \gamma_2^1 I_2^1$$

253
$$\frac{dR^1}{dt} = (1 - c_d^1) \gamma_2^1 I_2^1 - \frac{m_{12}R^1}{N_1} + \frac{m_{21}R^2}{N_2},$$

254 where superscript indicates country.

255 This model is extended to all countries and coded in R [25] using the infectious disease
256 node of the Australia Nectar Research cloud (www.nectar.org.au) as individual-based
257 model with a binomial distribution of the number of people experience an event at a
258 specific time step. The events are infection, migration, emigration, recovery and death
259 due to COVID-19. We neglected natural death, as this does not affect our result. We
260 further assumed that no countries except the epicenters are experiencing outbreak.
261 Hence the basic reproduction number at the epicentres is $R_0 = 2.68$. Each country
262 COVID-19 dynamics follows the schematic representation in Fig S4.

263

264