Estimating Spot Prevalence of COVID-19 from Daily Death Data in Italy

Ali Raheem[1*]

[1]South London and Maudsley NHS Foundation Trust

[*]To whom correspondence should be addressed; E-mail: ali.raheem(at)kcl.ac.uk.

COVID-19 like many infectious diseases has a variable severity. Due to limited testing capacity for new virus, the majority of infected showing mild or no symptoms it is likely current prevalence data is an underestimate. We present an estimate of the point prevalence of COVID-19 compared to the number of confirmed case in Italy based on the daily death rate and information about the incubation period, time from symptom onset to death and reported case fatality rate. Our model predicts that when the first 3 infected cases had been identified by Italian authories there were already nearly 30 cases in Italy, and by the 24th of February 2020 only 0.5% cases had been detected and confirmed by Italian authorities. While official statistics had 132 confirmed case we believe a more accurate estimate would be closer to 26000. With a case-doubling period of about 2.5 days.

Introduction

COVID-19 has now been declared a pandemic by the World Health Organisation. Caused by a betacoronvirus virus SARS-CoV2 which is related to the SARS and MERS virus.

This disease appears to have a mortality rate of approximated 1-15%. However there have been a wide variety in reported proportion of cases that are asymptomatic or only show mild non-specific symptoms. Making it difficult to estimate the prevalence of COVID-19 without widespread testing which has not yet been implemented in any country. Previously authors have used this to estimate the mortality of COVID-19 (1). Accurately estimating the prevalence of COVID-19 will allow organisations to make better informed decisions to control COVID-19.

Methods

We used a linear retrospective model to estimate past point prevalence using daily number of report deaths. This model required us to calculate a nominal time to death from infection. We based this value on data available from the World Health Organisation. The WHO report time from onset of symptoms to death of about 2 weeks (4). Additionally, the mean incubation period has been reported to be 6.4 days (2). We estimate that deaths on a given day should correlate with infections 3 weeks prior and use this with daily reported deaths to estimate the spot prevalence in the past. We obtained data on daily deaths and past cases from the European Centre for Disease Prevention and Control (3) cross referenced for accuracy with data from the World Health Organisation (4).

Data processing was carried out with R, the Juptyer notebook and Tidyverse software suites on a Debian 9.0 Stretch using the latest Jupyter/r-notebook docker image (jupyter/r-notebook:15a66513da30) (5) (6).

Using a case mortality of 7% was used based on the recent estimates.

$$Prevalence_{d-21} = \frac{Deaths_d}{Mortality}$$

Where d is time in days. Using a mortality of 7%

$$Prevalence_{d-21} = \frac{Deaths_d}{0.07}$$

The following data was generated using the reported number of deaths per day and new confirmed cases. From this a cumulative cases and cumulative deaths data was calculated and used to calculate the point prevalence according to the formula described above.

Using data up to the 16^{th} of March 2020 we can estimate the point prevalence 21 days into the past (the 24^{th} of February 2020). This value is subject to the inevitable jitter in deaths per day due to COVID-19, it therefore should be used to guide a trend line before interpretation.

Results

Figure 1 summaries the results graphically, the full results can be reviewed in the supplementary materials section in Table 1.

Italy initially confirmed 3 cases in Italy on the 31^{st} of January 2020, our model predicts in fact there were 28 COVID-19 cases in Italy. From the 31^{st} of January until the 22^{nd} of Febuary there was no detected transmission in Italy and the number confirmed cases remained at 3.

On the 24th of February when our prediction data ends there were 132 cases confirmed by Italian authorities but our model predicts there were near 26000 cases in reality.

Our model predicts that in this period there was undetected transmission resulting in a rise in cases from 28 to 18000. With a doubling period of about 2.5 days.

Discussion

The large disparity with estimated prevalence being much higher than confirmed cases indicates that either an increasing majority of cases are not detected. There seems to have been a period of several weeks where COVID-19 was transmitted in the Italian population undetected. Only a minority of cases appear to be confirmed at any point in time.

Conclusion

In this paper we present evidence that the currently confirmed cases of COVID-19 are a dramatic underestimate of the true point prevalence in Italy and a method to estimate point prevalence from daily deaths of COVID-19. Increasing the mortality would reduce the estimated prevalence but this alone could not make the estimates agree with the confirmed cases in order of magnitude. This methodology would be applicable to many other conditions and relies only on accurate estimate of deaths due to the condition which can easily be confirmed post-mortem and case mortality. Without incubation date or data on disease progression an accurate estimate can still be produced but will not provide temporal information but could be used to estimate the time from infection to death.

Limitations

This model used the spot daily reported deaths which may lag the true date of death due to delays in confirming and then reporting causes of death if COVID-19 was not diagnoses ante mortem. Our estimate of point prevalence varies proportionally to the error in deaths. Deaths due to infection not reported will cause an underestimate in prevalence. Estimating true mortality rates is difficult, and our estimate varies with

the reciprocal of the error in the mortality rate. Underestimating mortality will lead to increase in predicted spot prevalence. This is an evolving pandemic and due to the long incubation and time to death periods only a short period of point prevalence's can be estimated.

Competing Interests

The authors declare no competing interests.

References

1. D. Baud, et al., The Lancet Infectious Diseases (2020).

2. World Health Organisation, Coronavirus disease 2019 (COVID-19) Situation Report - 56 (2020). https://www.who.int/docs/default-source/coronaviruse/situation-reports/ 20200316-sitrep-56-covid-19.pdf?sfvrsn=9fda7db2_2[Online; accessed 17-March-2020].

3. J. A. Backer, D. Klinkenberg, J. Wallinga, Euro surveillance : bulletin Europeen sur les maladies transmissibles = European communicable disease bulletin 25, 2000062 (2020). 32046819[pmid].

4. European Centre for Disease Prevention and Control, Data on the geographic distribution of COVID-19 cases world-wide (2020). https://www.ecdc.europa.eu/en/publications-data/ download-todays-data-geographic-distribution-covid-19-cases-worldwide [Online; accessed 17-March-2020].

5. H. Wickham, tidyverse: Easily Install and Load the 'Tidyverse' (2017). R package version 1.2.1.

6. R Core Team, R: A Language and Environment for Statistical Computing, R Foundation for Statistical Computing, Vienna, Austria (2017).

Acknowledgements

I would like to thank the healthcare workers around the globe for their tireless efforts fighting COVID-19.

Supplementary materials

Table 1: Raw results data

Date	New Cases	New Deaths	Total Cases	Total Deaths	Est Total Cases
31/01/2020	3	0	3	0	-
01/02/2020	0	0	3	0	-
02/02/2020	0	0	3	0	28.57143
03/02/2020	0	0	3	0	28.57143
04/02/2020	0	0	3	0	85.71429
05/02/2020	0	0	3	0	157.14286
06/02/2020	0	0	3	0	171.42857
07/02/2020	0	0	3	0	242.85714
08/02/2020	0	0	3	0	300.00000
09/02/2020	0	0	3	0	414.28571
10/02/2020	0	0	3	0	500.00000
11/02/2020	0	0	3	0	742.85714
12/02/2020	0	0	3	0	1142.85714
13/02/2020	0	0	3	0	1528.57143
14/02/2020	0	0	3	0	2114.28571
15/02/2020	0	0	3	0	2814.28571
16/02/2020	0	0	3	0	3328.57143
17/02/2020	0	0	3	0	5228.57143
18/02/2020	0	0	3	0	6628.57143
19/02/2020	0	0	3	0	9014.28571

Table 1: Raw results data

Date	New Cases	New Deaths	Total Cases	Total Deaths	Est Total Cases
20/02/2020	0	0	3	0	11814.28571
21/02/2020	0	0	3	0	14514.28571
22/02/2020	14	0	17	0	18114.28571
23/02/2020	62	2	79	2	20585.71429
24/02/2020	53	0	132	2	25871.42857
25/02/2020	97	4	229	6	-
26/02/2020	93	5	322	11	-
27/02/2020	78	1	400	12	-
28/02/2020	250	5	650	17	-
29/02/2020	238	4	888	21	-
01/03/2020	240	8	1128	29	-
02/03/2020	561	6	1689	35	-
03/03/2020	146	17	1835	52	-
04/03/2020	667	28	2502	80	-
05/03/2020	587	27	3089	107	-
06/03/2020	769	41	3858	148	-
07/03/2020	778	49	4636	197	-
08/03/2020	1247	36	5883	233	-
09/03/2020	1492	133	7375	366	-
10/03/2020	1797	98	9172	464	-
11/03/2020	977	167	10149	631	-
12/03/2020	2313	196	12462	827	-
13/03/2020	2651	189	15113	1016	-
14/03/2020	2547	252	17660	1268	-
15/03/2020	90	173	17750	1441	-
16/03/2020	6230	370	23980	1811	-

Figure 1: Plotting Estimated Prevalence (circles), Confirmed Cases (filled triangles), Reported Deaths (triangles)