Insignificant effect of counter measure for coronavirus infectious disease -19 in Japan

Yoshiyuki Sugishita[1], Junko Kurita[2], Tamie Sugawara[1], Yasushi Ohkusa [1]

1) National Institute of Infectious Diseases, Tokyo, Japan

2) Center for Medical Sciences, Ibaraki Prefectural University of Health Sciences,

Ibaraki, Japan

Keywords: COVID-19, voluntary event cancellation, outbreak in Japan, SIR model, asymptomatic case

Corresponding author: Junko Kurita, kuritaj@ipu.ac.jp

Abstract

Background: To control COVID-19 outbreak in Japan, sports and entertainment events were canceled in Japan for two weeks from 26 February to 13 March. It has been designated as voluntary event cancellation (VEC).

Object: This study predicts the effectiveness of VEC enduring and after its implementation.

Method: We applied a simple susceptible–infected–recovery model to data of patients with symptoms in Japan during 14 January to VEC introduction and after VEC introduction to 8 March. We adjusted the reporting delay in the latest few days.

Results: Results suggest that the basic reproduction number, R_0, before VEC introduced as 2.29 with a 95% confidence interval (CI) was [2.19, 2.37] and the effective reproduction number, R_v, after VEC introduced as 1. 99; its 95% CI was [1.71,2.23].

Discussion and Conclusion: No significant effect of VEC was observed.

Introduction

The initial case of COVID-19 in Japan was that of a patient returning from Wuhan, China who showed symptoms on 3 January, 2020. Subsequently, as of 13 March, 2020, 714 cases were announced, including asymptomatic or those infected at abroad in countries such as China but excluding or those infected on a large cruise ship, the Diamond Princess [1].

Sports and entertainment events were canceled in Japan for two weeks from 26 February to 13 March according to a government advisory. At the same time, it was advised that small business and private meetings be cancelled voluntarily. The policy is designated as voluntary event cancellation (VEC). Moreover, since 3 March, almost all schools have been closed to control the spread until early April. Although schoolchildren are not of pestiferous age, the policy effects remain unknown. These policies must be evaluated as soon as possible. If the effective reproduction number, R_v under these measures is less than one, the outbreak can be contained. Alternatively, even given a low R_0 before these measures, if it was greater than one, it might prolong the outbreak. Nevertheless, one could expect to prevent some fatal cases over time by easing burdens on medical resources or developing a vaccine. The entire course of the outbreak might be altered if these measures were able to reduce infectiousness

considerably. Therefore, evaluation of these measures must be thorough when these

measures are commenced, continued, and ceased. The present study was conducted to

evaluate VEC before VEC is cancelled so as to contribute government's decision

making whether VEC will be continued or ceased.

Method

We applied a simple susceptible–infected–recovery (SIR) model [2] to the data

assuming an incubation period following its empirical distribution in the early stage of

the outbreak in Japan. Experiences of Japanese people living in Wuhan until the

outbreak provide information related to mild cases because complete laboratory

surveillance was administered for them. During January 29 – February 17, 2020, 829

Japanese people returned to Japan from Wuhan. All had received a test to detect

COVID-19; of them, 14 were found to be positive for COVID-19 [3]. Of those 14, 10

Japanese people had exhibited mild symptoms; the other 4 showed no symptom as of

February 25. Moreover, two Japanese residents of Wuhan exhibited severe symptoms:

one was confirmed as COVID-19. The other died, although no fatal case was confirmed

as COVID-19 by testing. In addition, two Japanese residents of Wuhan with mild

symptoms were refused re-entry to Japan even though they had not been confirmed as

infected. If one assumes that the Japanese fatal case in Wuhan and that the two rejected

re-entrants were infected with COVID-19, then 2 severe cases, 12 mild cases, and 4

asymptomatic cases were found to exist among these Japanese residents of Wuhan. We

therefore apply these proportions to the simulation.

Assuming that the power of infectivity among severe patients and mild patients

were equal among the asymptomatic cases, half of the symptomatic cases can be

assumed. This assumption about relative infectiousness among asymptomatic cases

compared with symptomatic cases was also assumed in simulation studies for influenza

[4–8]. We sought R_0 to fit the number of patients during 14 January – 28 February and

to minimize the sum of absolute values of discrepancies among the reported numbers

and the fitted values. Its 95% confidence interval (CI) was calculated using the 10000

iterations of bootstrapping for empirical distribution.

We used data of the community outbreak of patients with COVID-19 who showed

any symptom in Japan for 14 January – 13 March, 2020. We excluded some patients

who had returned from China, and who were presumed to be infected persons from the

Diamond Princess. They were presumed to be not community-acquired in Japan. During

this period, 544 cases with onset dates occurred.

Published information about COVID-19 patients with symptoms from the Ministry

of Labour, Health and Welfare (MLHW), Japan was usually adversely affected with

some delay caused by uncertainty during onset to visiting a doctor or in the timing of a

physician's suspicion of COVID-19. Therefore, published data of patients must be

adjust at least a few days. To adjust it, we applied the following regression. We denote

$Xt\text{-}k/t$ as the number of patients whose onset date was $t\text{-}k$ published on day t. The

dependent variables are the degree of reporting delay, $Xt\text{-}k\text{-}m/t \ / \ Xt\text{-}k\text{-}m/t\text{-}m$, where $k{>}m$

in several m and k. Here, m denotes the difference of the publishing dates between the

two published. Date t represents the publishing date of the latest publishing. The

explanatory variables were $1/k$, $1/m$, and $1/km$. The degree of reporting delay was

estimated as [estimated coefficient of constant term] + [estimated coefficient of $1/k$]$/k$,

when m was sufficiently large and time had passed. Therefore, this estimated degree of

reporting delay multiplied by the latest published data are expected to be predictions of

the number of patients whose onset date was $t\text{-}k$. We used this adjusted number of

patients in the latest few days including those after VEC was adopted. We used the

published data on 2,5,6, and 9-14 March,2020 by MLHW[1].

First, we estimated R_0 in Japan to fit the data of community outbreak before VEC

was introduced. Then, using the adjusted data of patients, we estimated R_v after VEC

was adopted. We discussed about R_v/R_0 through its distribution. We used 5% as

significant level.

Ethical consideration

All information used in the present study were published and thus there was no

ethical issue.

Results

During 14 January – 13 March in Japan, 640 community-acquired cases were

identified for whom the onset date was published. Figure 1 showed the empirical

distribution of incubation period among 62 cases whose exposed date and onset date

were published by MHLW. Its mode and median were six days and average was 6.74

days.

Figure 2 depicts the epidemic curves published at 3, 5, 6, 9 and 10 March. From

this information, we estimated the degree of reporting delay. Those results are presented

in Table 1. The table shows that $1/k$, $1/m$ and $1/km$ are all significant. When m is

sufficiently large, the effects of $1/m$ and $1/km$ converge to zero. Therefore, the estimated

degree of reporting delay consists of the term of $1/k$ and a constant term. Based on these

results, we predict the degrees of reporting delay as 19.3 for $k=1$, 9.64 for $k=2$, 6.42 for

k=3, and so on.

The value of R_0 before VEC introduced was estimated as 2.29 with a 95%CI was

[2.19, 2.37] and R_v, after VEC introduced as 2.03 and its 95% CI was [1.72, 2.32]. The

average and median of R_v/R_0 were 0.88 and its 95% CI was [0.79, 1.05].

Discussion

We applied a simple SIR model including asymptomatic cases that had not been

incorporated into the model to date. An earlier study [9] estimated R_0 for COVID-19 as

2.24–3.58 in Wuhan. Our R_0 obtained before VEC was similar.

Unfortunately, effectiveness of VEC was not observed even though it reduced

infectiousness by 13% in average. .

Conclusion

Results demonstrated that VEC can reduce infectiousness of COVID-19 by 13%,

but not significant. However, we have to note that the obtained results were just a

preliminary evaluation, and thus it is not a conclusive evaluation. VEC had been

extended until March 19 on March 10. We hope that the present study contribute

government's decision making for VEC. The present study is just the authors opinion,

but does not reflect any stance of our affiliation.

Conflict of interest

No author has any conflict of interest, financial or otherwise, to declare in relation to

this study.

Reference

1. Japan Ministry of Health, Labour and Welfare. Press Releases. h abroad such as

   China (in Japanese) [accessed on 12 March, 2020]

2. Japan Ministry of Health, Labour and Welfare. Press Releases.

   https://www.mhlw.go.jp/stf/houdou/houdou_list_202001.html. (in Japanese)

   [accessed on 12 March, 2020]

3. Ohkusa Y, Sugawara T, Taniguchi K, Okabe N. Real-time estimation and prediction

   for pandemic A/H1N1(2009) in Japan. J Infect Chemother. 2011;17:468-72.

4. Ferguson NM, Cummings DA, Cauchemez S, Fraser C, Riley S, Meeyai A, et al. :

   Strategies for containing an emerging influenza pandemic in Southeast Asia.

   Nature2005; 437:209-214.

5. Longini IM Jr, Nizam A, Xu S, Ungchusak K, Hanshaoworakul W, Cummings DA,

   Halloran ME: Containing Pandemic Influenza at the Source. Science2005;

   309:1083-1087.

6. Germann TC, Kadau L, Longini IM Jr, Macken CA: Mitigation strategies for

   pandemic influenza in the United States. Proc Natl Acad Sci USA 2006; 103:

   5935-5940.

7. Ferguson NM, Cummings DA, Fraser C, Cajka JC, Cooley PC, Burke DS:

Strategies for mitigating an influenza pandemic. Nature2006; 442: 448-452.

8. Ohkusa Y, Sugawara T: Simulation Model of Pandemic influenza in the Whole of Japan, Journal of Japanese Infectious Disease2009; 62:98-106.

9. Zhao S, Lin Q, Ran J, Musa SS, Yang G , Wang W, Lou Y, Gao D, Yang L, He D, Wang M. Preliminary Estimation of the Basic Reproduction Number of Novel Coronavirus (2019-nCoV) in China, From 2019 to 2020: A Data-Driven Analysis in the Early Phase of the Outbreak. Int J Infect Dis 2020[Online ahead of print]

10. Liu Y, Gayle AA, Wilder-Smith A, Rockly J. The reproductive number of COVID-19 is higher compared to SARS coronavirus. J Travel Med. 2020.DOI: 10.1093/jtm/taaa021

11. Lai C, Shih T, Ko W, Tang H, Hsueh P. Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) and Coronavirus disease-2019 (COVID-19): The Epidemic and the Challenges. Int J Antimicrob Agents. DOI:10.1016/j.ijantimicag.2020.105924

Table: Estimation results of the degree of reporting delay

	Estimated coefficient	*p*-value	95%CI	
			Lower bound	Upper bound
1/k	16.9	.000	14.7	19.0
1/m	1.01	.025	0.140	2.05
1/(km)	-21.8	.000	-26.7	-16.9
Constant	0.266	.236	-0.175	0.707

Note: The number of observations was 323. The coefficient of determination was 0.457.

k denotes the number of days until the last reported day. m denotes the difference between the publishing date in the past and the most recent publishing date. The dependent variable was the degree of reporting delay, which is defined as $X_{t-k-m|t}$ / $X_{t-k-m|t-m}$, where $X_{t-k|t}$ is the number of patients whose onset date was $t-k$ published on day t. We used the number of patients by onset day published on 2, 5, 6, 10, 11, 12, 13 and 14 March 2020 by Ministry of Labour, Health and Welfare, Japan.

Figure 1: Epirical distributionof incubation period published by Ministry of Labour,

Health and Welfare, Japan

(number of patients)

Notes: Bars indicates the number of patients by incubation period among 62 cases

whose exposed date and onset date were published by Ministry of Labour, Health and

Welfare, Japan.

Figure 2: Epidemic curves of COVID-19 in Japan published on 2, 6, 10, 12 and 14

March, 2020 by Ministry of Labour, Health and Welfare, Japan

(number of patients)

Note: The black bold, black thin, brioken gray, and gray bold lines respectively show

epidemic curves published on 14 March, 12 March, 10 March, 6 March, and 2 March by

Ministry of Labour, Health and Welfare, Japan.