

## Effects of voluntary event cancellation and school closure as countermeasures against COVID-19 outbreak in Japan

Yoshiyuki Sugishita<sup>1)</sup>, Junko Kurita<sup>2)</sup>, Tamie Sugawara<sup>1)</sup>, Yasushi Ohkusa<sup>2)</sup>

1) National Institute of Infectious Diseases, Tokyo, Japan

2) Department of Nursing Collage of Nursing, Tokiwa University, Ibaraki, Japan

Corresponding author: Yoshiyuki Sugishita, [yoshiyuk@gmail.com](mailto:yoshiyuk@gmail.com)

Keywords: COVID-19, voluntary event cancellation, outbreak in Japan, SIR model, asymptomatic case

## Abstract

**Background:** To control the COVID-19 outbreak in Japan, sports and entertainment events were canceled and schools were closed throughout Japan from February 26 through March 19. That policy has been designated as voluntary event cancellation and school closure (VECSC).

**Object:** This study assesses VECSC effectiveness based on predicted outcomes.

**Method:** A simple susceptible–infected–recovery model was applied to data of patients with symptoms in Japan during January 14 through March 25. The respective reproduction numbers were estimated before VECSC ( $R$ ), during VECSC ( $R_e$ ), and after VECSC ( $R_a$ ).

**Results:** Results suggest  $R$  before VECSC as 1.987 [1.908, 2.055],  $R_e$  during VECSC as 1.122 [0.980, 1.260], and  $R_a$  after VECSC as 3.086 [2.529, 3.739].

**Discussion and Conclusion:** Results demonstrated that VECSC can reduce COVID-19 infectiousness considerably, but the value of  $R$  rose to exceed 2.5 after VECSC.

1 Introduction

2 The initial case of COVID-19 in Japan was a patient returning from Wuhan, China who  
3 showed symptoms on January 3, 2020. Subsequently, as of April 6, 2020, the Ministry  
4 of Labour, Health and Welfare (MLHW) in Japan reported 3,906 cases in Japan,  
5 including asymptomatic cases but excluding those from a large cruise ship: the  
6 Diamond Princess [1].

7 Sports and entertainment events were canceled in Japan for two weeks from  
8 February 26 through March 11 in compliance with a government advisory. At that time,  
9 it was advised that small business and private meetings be cancelled voluntarily: the  
10 measure was not enacted as a law or enforced by authorities. Therefore, people were not  
11 arrested or cited even if they did not comply with this government advisory [2]; the  
12 effort depended entirely on voluntary compliance of individuals. Moreover, this was the  
13 first measure necessitating voluntary event cancellation. Those reasons complicate the  
14 *ex ante* prediction of the proportion of events that were cancelled and the extent to  
15 which contact among people was reduced. Moreover, as announced March 3, almost all  
16 schools were closed from the middle of March through spring vacation (early April) as  
17 a measure to control the spread of COVID-19. Even though young people can be  
18 infected and can transmit the virus to adults, school closure effects were questionable *ex*

19 *ante* because schoolchildren were not regarded as the most susceptible age class for  
20 COVID-19 [3–5]. Therefore, these policies must be evaluated *ex post*. The policies are  
21 known collectively as voluntary event cancellation and school closure (VECSC). After  
22 the VECSC period, VECSC ceased, but cancellation of large events was continued.

23 If the reproduction number under VECSC ( $R_e$ ) was less than one, then the outbreak  
24 can be expected to have been contained. Alternatively, even if  $R_e$  was markedly less  
25 than  $R_0$  but greater than one, one could have expected instead that the outbreak would  
26 be exacerbated.

27 This study was conducted to evaluate VECSC based on the epidemic curve because  
28 no evidence exists to reflect the number of cancelled events. The obtained result might  
29 be expected to contribute to a government decision to implement VECSC as a  
30 countermeasure in Japan.

31

## 32 Method

33 We applied a simple deterministic susceptible–infected–recovery(SIR) model [6] to  
34 the epidemic curve in Japan for its population of 120 million. We assume an incubation  
35 period following the empirical distribution from the early stage of the outbreak. A  
36 person on the  $i$ -th day of incubation will move to a symptomatic or an asymptomatic

37 state with probability of  $p(i)/\sum_{j=i}^L p(j)$ , where  $p(i)$  represents the proportion of the  $i$ -th day  
38 on the empirical distribution of the incubation period and  $L$  denotes the maximum  
39 length of the incubation period. Alternatively, a person moves to the  $i+1$  day in the  
40 incubation state with one minus  $p(i)/\sum_{j=i}^L p(j)$ .

41 Symptomatic and asymptomatic states were assumed to continue for seven days,  
42 with subsequent transition to a recovery state with probability of one [7]. This model  
43 ignores outcomes and necessary medical resources: states of death and hospitalization  
44 were not incorporated into the model.

45 Asymptomatic cases are not observable unless complete laboratory-based  
46 surveillance is performed. One exceptional study indicated them as 3/23 from a sample  
47 of elderly people [7]. We checked its robustness for the proportion of asymptomatic  
48 cases assuming a ratio of 4/19 from experiences of Japanese residents of Wuhan up to  
49 the outbreak [8].

50 Infectivity by severely infected patients and by mild patients was assumed to be  
51 equal. Moreover, we assumed that asymptomatic cases have the same power of  
52 infectivity as symptomatic cases [7]. That is merely an assumption. Therefore, we  
53 verified its robustness through sensitivity analysis of the infectiousness of asymptomatic  
54 cases, such as 50% assumed for simulation studies for influenza [9–13]. The distribution

55 of infectiousness in symptomatic and asymptomatic cases was assumed to be 30% on  
56 the onset day, 20% on the following day, and 10% for the subsequent five days [7].

57 Regarding its robustness, we also applied a uniform distribution for a week.

58 Because VECSC was conducted during February 27 – March 19, we divided the  
59 data period into three periods of those before VECSC, during VECSC, and after  
60 VECSC, with corresponding reproduction numbers represented as  $R_0$ ,  $R_e$ , and  $R_a$ . We  
61 modeled the change in the reproduction number given the prior day's reported number  
62 of persons susceptible, incubation, symptomatic, asymptomatic, and in a recovered  
63 state.

64 For this study,  $R_0$  was defined as the number of newly infected persons from one  
65 person infected over the infectious period when all persons were susceptible with no  
66 countermeasures. In addition,  $R_e$  and  $R_a$  were defined, respectively, as the numbers of  
67 newly infected persons from one person infected over the infectious period when all  
68 persons were susceptible under VECSC or during a time of countermeasures after the  
69 VECSC period. For estimation, the number of the newly infected persons was  
70 calculated using  $R_0$  ( $R_e$  or  $R_a$ )  $\times$  the proportion of susceptible persons among the  
71 population  $\times$  the number of persons with infectiousness weighted by their degree of  
72 infectiousness.

73           The values of  $R_0$ ,  $R_e$ , and  $R_a$  were sought to fit the data to minimize the sum of  
74 absolute values of discrepancies among the bootstrapped epidemic curve and the fitted  
75 values. The estimated distribution of three reproduction numbers was calculated using  
76 10,000 iterations of bootstrapping for empirical distribution of the data for symptomatic  
77 patients. In this sense, although this model is deterministic at each bootstrapping  
78 iteration, its range can be estimated through 10,000 iterations.

79           The bootstrapping procedure we used with fully replicated bootstrapping for a fixed  
80 number of initial cases. There were  $N$  patients in the data, with numbering of the  
81 patients from the initial case to the last case. Initially, no patient was on the  
82 bootstrapped epidemic curve. If a random variable drawn from a uniform distribution of  
83  $(0,1)$  was  $t$  included in the interval  $[i/(N-1), (i+1)/(N-1)]$ , then we added one to the onset  
84 date of  $i+1$ th patient to the bootstrapped epidemic curve. We replicated this procedure  
85  $N-1$  times. Thereby, we obtain the bootstrapped epidemic curve with  $N-1$  patients.  
86 Finally, we added the initial patient, for whom the onset date was January 14, to the  
87 bootstrapped epidemic curve. Consequently, we obtained a bootstrapped epidemic curve  
88 with  $N$  patients starting from January 14.

89           We estimated the curve sequentially as follows. First, we estimated  $R_0$  as the best fit  
90 to bootstrapped data for the pre-VECSC period. Then, based on the obtained  $R_0$  and the

91 course of the outbreak before the VECSC period, we estimated  $R_e$  as the best fit to  
92 bootstrapped data in the VECSC period. Finally, based on the obtained values of  $R_0$  and  
93  $R_e$ , we estimated  $R_a$  as the best fit to bootstrapped data for the post-VECSC period. In  
94 each step, reproduction numbers were grid searched in the interval of (0,10) by 0.001.

95 We conducted some sensitivity analyses for one way in addition to the base case as  
96 explained above: the infectiousness power of asymptomatic cases represented 50% of  
97 symptomatic cases; the proportion of asymptomatic cases was 4/19 as symptomatic  
98 cases; infectiousness and infectious patterns showed a uniform distribution for one week.  
99 We also estimated the parameters by minimizing the sum of the squared residuals or  
100 maximizing Poisson likelihood instead of the difference in absolute values and using the  
101 fitted incubation period as a log normal distribution instead of the empirical distribution.

102 Moreover, we examined the possibility of infection during two days before onset.  
103 Its probability can be represented as  $\sum_{i=2}^{L-1} p(i) / \sum_{j=i}^L p(j) x(i-1, t-1) + (1 - p(i) / \sum_{j=i}^L p(j))$   
104  $p(i+1) / \sum_{j=i+1}^L p(j) x(i-1, t-2) + x(L-1, t-1)$ , where  $x(i, t)$  represents the number of persons at  
105 the  $i$ -th incubation period on day  $t$ . They were assumed to have the same infectiousness  
106 in the first two days of symptomatic period. To meet the definition of the reproduction  
107 number, infectiousness in the incubation and (a) symptomatic period were adjusted such  
108 that the sum of infectiousness from two days before onset to one week following onset

109 would be one.

110 We also examined incorporation of the imported cases into the model. They were  
111 entered into the model as newly symptomatic cases with no history of incubation period  
112 when the imported cases were reported. The parameters in this model were estimated as  
113 minimizing the sum of the absolute difference among the number of patients including  
114 the imported cases and prediction.

115 We adopted 5% as the level for which we inferred significance. We used Matlab  
116 2014a to code the model as explained above.

117

118 Data source

119 The numbers of symptomatic patients during January 14 – March 25 were  
120 published by the MHLW [1] and were reported publicly from 46 prefectures and some  
121 cities  
122 ([https://www.pref.okinawa.lg.jp/site/hoken/chiikihoken/kekkaku/covid19\\_hasseijoukyo](https://www.pref.okinawa.lg.jp/site/hoken/chiikihoken/kekkaku/covid19_hasseijoukyo)  
123 [u.html](#);  
124 <https://www.pref.kagoshima.jp/ae06/kenko-fukushi/kenko-iryu/kansen/kansensho/coron>  
125 [avirus.html](#);  
126 [https://www.pref.miyazaki.lg.jp/kansensho-taisaku/covid-19/hassei\\_list.html](https://www.pref.miyazaki.lg.jp/kansensho-taisaku/covid-19/hassei_list.html);

- 127 <https://www.pref.oita.jp/site/covid19-oita/covid19-pcr.html>;
- 128 [https://www.pref.kumamoto.jp/kiji\\_32300.html](https://www.pref.kumamoto.jp/kiji_32300.html);
- 129 [https://www.pref.nagasaki.jp/bunrui/hukushi-hoken/kansensho/corona\\_nagasaki/corona\\_nagasaki\\_shousai/](https://www.pref.nagasaki.jp/bunrui/hukushi-hoken/kansensho/corona_nagasaki/corona_nagasaki_shousai/); <https://www.pref.saga.lg.jp/kiji00373220/index.html>;
- 130
- 131 <https://www.pref.fukuoka.lg.jp/contents/covid19-hassei.html>;
- 132 <https://www.city.fukuoka.lg.jp/hofuku/hokenyobo/health/kansen/cohs.html>;
- 133 <https://www.pref.kochi.lg.jp/soshiki/130401/2020022900049.html>;
- 134 <https://www.pref.ehime.jp/h25500/kansen/covid19.html#kansensha>;
- 135 [https://www.pref.kagawa.lg.jp/content/dir1/dir1\\_6/dir1\\_6\\_2/wt5q49200131182439.shtml#outbreak](https://www.pref.kagawa.lg.jp/content/dir1/dir1_6/dir1_6_2/wt5q49200131182439.shtml#outbreak);
- 136
- 137 <https://www.pref.yamaguchi.lg.jp/cms/a10000/korona2020/202004240002.html>;
- 138 <https://www.pref.hiroshima.lg.jp/soshiki/57/bukan-coronavirus.html>;
- 139 <https://www.pref.okayama.jp/page/645925.html#kennaijoukyou>;
- 140 [https://www.pref.shimane.lg.jp/bousai\\_info/bousai/kikikanri/shingata\\_taisaku/new\\_coronavirus\\_portal.html](https://www.pref.shimane.lg.jp/bousai_info/bousai/kikikanri/shingata_taisaku/new_coronavirus_portal.html); <https://www.pref.tottori.lg.jp/291425.htm>;
- 141
- 142 <https://www.pref.wakayama.lg.jp/prefg/041200/d00203387.html>;
- 143 <http://www.pref.nara.jp/module/1356.htm#moduleid1356>;
- 144 [https://web.pref.hyogo.lg.jp/kk03/corona\\_hasseijyokyo.html](https://web.pref.hyogo.lg.jp/kk03/corona_hasseijyokyo.html);

145 [http://www.pref.osaka.lg.jp/hodo/index.php?HST\\_TITLE1=%83R%83%8D%83i&SEA](http://www.pref.osaka.lg.jp/hodo/index.php?HST_TITLE1=%83R%83%8D%83i&SEARCH_NUM=10&searchFlg=%8C%9F%81@%8D%F5&site=fumin;)

146 [RCH\\_NUM=10&searchFlg=%8C%9F%81@%8D%F5&site=fumin;](http://www.pref.osaka.lg.jp/hodo/index.php?HST_TITLE1=%83R%83%8D%83i&SEARCH_NUM=10&searchFlg=%8C%9F%81@%8D%F5&site=fumin;)

147 <https://www.pref.kyoto.jp/kentai/news/novelcoronavirus.html#F;>

148 <https://www.pref.shiga.lg.jp/ippan/kenkouiryohukushi/yakuzi/310735.html;>

149 <https://www.pref.mie.lg.jp/YAKUMUS/HP/m0068000066.htm;>

150 <https://www.pref.aichi.jp/site/covid19-aichi/corona-kisya.html;>

151 <http://www.city.nagoya.jp/kenkofukushi/page/0000126920.html;>

152 <https://www.pref.shizuoka.jp/kinkyu/covid-19-tyuumokujouhou.html;>

153 [https://www.pref.gifu.lg.jp/kinkyu-juyo-joho/shingata\\_corona\\_kansendoko.html;](https://www.pref.gifu.lg.jp/kinkyu-juyo-joho/shingata_corona_kansendoko.html;)

154 [https://www.pref.nagano.lg.jp/hoken-shippei/kenko/kenko/kansensho/joho/corona-doko](https://www.pref.nagano.lg.jp/hoken-shippei/kenko/kenko/kansensho/joho/corona-doko.html;)

155 [.html;](https://www.pref.nagano.lg.jp/hoken-shippei/kenko/kenko/kansensho/joho/corona-doko.html;)

156 [https://www.pref.yamanashi.jp/koucho/coronavirus/info\\_coronavirus\\_prevention.html;](https://www.pref.yamanashi.jp/koucho/coronavirus/info_coronavirus_prevention.html;)

157 <https://www.pref.fukui.lg.jp/doc/kenkou/corona/jyoukyou.html;>

158 <https://www.pref.ishikawa.lg.jp/kansen/coronakennai.html;>

159 [http://www.pref.toyama.jp/cms\\_sec/1205/kj00021798.html;](http://www.pref.toyama.jp/cms_sec/1205/kj00021798.html;)

160 <https://www.pref.niigata.lg.jp/sec/kenko/covid19.html;>

161 [https://www.pref.kanagawa.jp/docs/ga4/bukanshi/occurrence\\_06.html;](https://www.pref.kanagawa.jp/docs/ga4/bukanshi/occurrence_06.html;)

162 <https://www.bousai.metro.tokyo.lg.jp/taisaku/saigai/1007261/index.html;>

163 <https://www.pref.chiba.lg.jp/shippei/press/2019/ncov-index.html>;

164 <https://www.pref.saitama.lg.jp/a0701/shingatacoronavirus.html>;

165 [https://www.pref.gunma.jp/07/z87g\\_00016.html](https://www.pref.gunma.jp/07/z87g_00016.html);

166 <http://www.pref.tochigi.lg.jp/e04/welfare/hoken-eisei/kansen/hp/coronakensahasseiyou>

167 <kyou.html>; <https://www.pref.ibaraki.jp/1saigai/2019-ncov/hassei.html>;

168 <https://www.pref.fukushima.lg.jp/sec/21045c/fukushima-hasseiyoukyou.html>;

169 <https://www.pref.yamagata.jp/ou/bosai/020072/kochibou/coronavirus/coronavirus.html>

170 #kensa;

171 <https://www.city.yamagata-yamagata.lg.jp/kakuka/kenkoiryo/kenkozoshin/sogo/kansen>

172 <syoud0409180023.html>; <https://www.pref.akita.lg.jp/pages/archive/47957>;

173 <https://www.pref.miyagi.jp/site/covid-19/02.html>;

174 <https://www.pref.iwate.jp/kurashikankyoku/iryoku/covid19/index.html>;

175 <https://www.pref.aomori.lg.jp/welfare/health/wuhan-novel-coronavirus2020.html>;

176 <http://www.pref.hokkaido.lg.jp/hf/kth/kak/hasseiyoukyou.htm#4/12>) as of April 6.

177 During this period, 1516 cases were recorded with onset dates. We excluded imported

178 cases and cases representing infected persons from the Diamond Princess because they

179 were presumed not to have been community-acquired in Japan.

180

181 Ethical considerations

182 All information used for this study has been published elsewhere [1]. There is  
183 therefore no ethical issue related to this study.

184

185 Results

186 Figure 1 depicts the empirical distribution of the incubation period among 91 cases  
187 for which the exposed date and onset date were published by MHLW. Its mode was six  
188 days. The average was 6.6 days. The maximum incubation period length was 17 days.

189 The value of  $R_0$  before introduction of VECSC was estimated as 2.534. Its range  
190 was [2.449, 2.598]. However,  $R_e$  during the VECSC period was estimated as 1.077  
191 [0.948, 1.228]. After VECSC,  $R_a$  was estimated as 4.455 [3.615, 5.255]. The null  
192 hypothesis that  $R_0$ ,  $R_e$ , and  $R_a$  were equal was therefore rejected.

193 Figure 2 depicts the observed epidemic curve and the predicted epidemic curve  
194 based on the estimated  $R_0$ ,  $R_e$ , and  $R_a$ . It showed goodness of fit that was pretty good.  
195 The small bump around February 26 reflects overshooting caused by the sharp decline  
196 of the reproduction number from  $R_0$  to  $R_e$ .

197 One-way sensitivity analysis revealed that if the infectiousness of asymptomatic  
198 cases was 50% of symptomatic cases instead of 100%, then  $R_0$  would be estimated as  
199 2.886 [2.801, 2.970],  $R_e$  would be 1.203 [1.059, 1.343], and  $R_a$  would be 4.985 [4.134,

200 5.893].

201 If the asymptomatic cases were 4/19 as symptomatic cases instead of 3/32, then  $R_0$   
202 would be estimated as 2.0492 [2.403, 2.557],  $R_e$  would be 1.091 [0.972, 1.237], and  $R_a$   
203 would be 4.428 [3.627, 5.049]. If the infectious pattern formed a uniform distribution  
204 for a week instead of the base case, then  $R_0$  would be estimated as 2534 [2.449, 2.598],  
205  $R_e$  would be 1.077 [0.948, 1.228], and  $R_a$  would be 4.455 [3.615, 5.255]. If we introduce  
206 infectiousness in the incubation period, then  $R_0$  would be estimated as 2.877 [2.765,  
207 2.980],  $R_e$  would be 1.278 [1.085, 1.478], and  $R_a$  would be 4.687 [3.621, 5.000].

208 When we estimate the parameters by minimizing the sum of the squared residuals  
209 instead of the absolute difference,  $R_0$  was estimated as 2.391 [2.505, 2.609],  $R_e$  was  
210 1.124 [0.935, 1.295], and  $R_a$  was 4.409 [3.168, 5.341]. In the case of using Poisson  
211 likelihood as evaluation, the parameters were estimated as 2.007 [1.957, 2.058], 1.141  
212 [1.030, 1.237], and 3.255 [2.697, 3.922]. However, when we estimate the parameters  
213 using a log normal distribution with mean of 1.66 and standard deviation of 0.558  
214 truncated as 17 days as incubation period instead of the empirical distribution,  $R_0$  was  
215 estimated as 2.2660 [2.179, 2.838],  $R_e$  was 1.116 [0.970, 1.277], and  $R_a$  was 3.574  
216 [2.897, 4.338].

217 However, we found 217 imported cases in this period, representing 14.3% of the

218 community acquired patients. When we incorporated the imported cases into the model,  
219  $R_0$  was estimated as 2.502 [2.412, 2.577],  $R_e$  was 1.113 [0.978, 1.298], and  $R_a$  was 4.311  
220 [3.603, 5.103]. Because the ranges of  $R_e$  and  $R_a$  were overlapped in all cases, these  
221 factors might not greatly affect both parameters, at least in the region with parameters  
222 and distribution that are considered reasonable. The estimated  $R_0$  was the most sensitive.  
223 When using Poisson likelihood for evaluation,  $R_0$  was significantly lower than in the  
224 base case. Moreover, infectiousness before onset raised  $R_0$  significantly. In other  
225 scenarios, the ranges of  $R_0$  were overlapped.

226

227

## 228 Discussion

229 We applied a simple deterministic SIR model including asymptomatic cases. A  
230 deterministic model might be simpler but more appropriate than a stochastic model for  
231 the present study including changing parameters and policy evaluation [14].

232 An earlier study [15–17] estimated  $R_0$  for COVID-19 as 2.24–3.58 in Wuhan. Our  
233  $R_0$  obtained for the period before VECSC was similar. However, an earlier study [18]  
234 found the average number of persons who were secondarily infected with COVID-19 in  
235 Japan as 0.6 before the VECSC period, with 80% of patients found to have infected no

236 one. If the average number of secondarily infected persons was less than one, then the  
237 outbreak might stall on its own. Given such an eventuality, no countermeasures would  
238 have been required. However, based on results of a study uploaded to the site of  
239 Japanese Public Health Association, health authorities in the public health sector or  
240 local government were urged to conduct contact tracing to detect clusters [19]. Because  
241 only a few patients infected other people, as found from this study, then the outbreak  
242 might be controlled if clusters could be suppressed.

243 Infectiousness two days before symptom onset has been described in the literature  
244 [7, 20]. We applied it as sensitivity analysis. It showed slightly higher reproduction  
245 numbers than the base model, although the estimated  $R_e$  and  $R_a$  were found to have a  
246 non-significant difference from the base case because a longer period of infectiousness  
247 implies fewer infected people on a day when the reproduction number does not change.  
248 However, the epidemic curve was given. Therefore, the estimated reproduction number  
249 probably increased by extending the period of infectiousness from seven days to nine  
250 days.

251 Moreover, when using Poisson likelihood for evaluation,  $R_0$  alone was found to be  
252 significantly lower than in the base case. Although the difference was not large, it might  
253 engender some difference attributable to the selection of the evaluation function.

254           Among children and younger adults, the proportion of asymptomatic infected  
255   people might be larger than among elderly people, but that point remains uncertain.  
256   Lower susceptibility of children [3–5] might imply a higher proportion of asymptomatic  
257   cases in children. In this sense, the finding of 3/23 among elderly people might be the  
258   lower bound of the proportion for the general population. Overall, sensitivity analysis  
259   led to similar estimation with the base case. Even though significant difference was  
260   found through comparison with the base case only in  $R_0$ , differences among them were  
261   not large, quantitatively speaking. Therefore, one might conclude that the obtained  
262   results were more or less robust.

263           The average incubation period applied in another study was 5.1 days [21], but it  
264   was 6.6 days for the present study. However, the latter number based on the epidemic  
265   curve considered in the present study, is certainly consistent with the data. It might  
266   reflect lower severity among patients in Japan than in other countries such as the US.  
267   The estimated value of the reproduction number probably would be smaller if we adopt  
268   a slightly shorter incubation period, which is a similar result to that found with  
269   sensitivity analysis for infectiousness before onset.

270           Many weeks passed since the end of the study period. Therefore, the reporting lag  
271   almost disappeared. If timely estimation, meaning estimation using data from a day

272 prior, was necessary, then we used an adjusted delay for reported data [8, 22, 23].

273 We used the minimized sum of the absolute values of discrepancies for the  
274 bootstrapped epidemic curve and the fitted values, instead of the minimized sum of  
275 squared residuals such as maximum likelihood estimation based on a normal  
276 distribution. In general, minimization of the sum of the absolute values is more robust  
277 than minimization of the sum of squares because the absolute value is less sensitive than  
278 squared values to the effects of outliers [24–26]. The daily epidemic curve sometimes  
279 exhibits spikes according to the day of the week and for other reasons. Especially, few  
280 patients per day were reported during the early stage of the outbreak. Therefore, spikes  
281 might be quite large. These spikes are probably outliers. They might overly affect the  
282 estimator. For this reason, we prefer minimization of the sum of the absolute values to  
283 minimization of the sum of squared values when analyzing daily data in the earlier stage  
284 of the outbreak.

285 Moreover, we chose non-parametric approaches using actual data in preference to  
286 parametric approaches assuming a particular distribution. Such an assumption might  
287 affect results through miss-specification. Therefore, we prefer to use the actual  
288 distribution of incubation periods or epidemic curves with no unnecessary or restrictive  
289 assumption. Sensitivity analysis using minimum least squares of errors or using Poisson

290 likelihood to maximize showed very similar results to those obtained with the base case.  
291 Although  $R_0$  and  $R_a$  were slightly higher,  $R_e$  were slightly lower. That result might  
292 indicate that some outlier in the pre-VECSC and post-VECSC period pulls the estimator  
293 in sensitivity analysis.

294 This study has some limitations. The first is that even though we evaluated VECSC,  
295 the respective effects of voluntary event cancellation and school closure cannot be  
296 discerned. To distinguish their respective effects, one would have to develop a model  
297 with several age classes. School closure mainly affects contact patterns among  
298 schoolchildren; voluntary event cancellation mainly affects patterns among adults.  
299 Therefore, studies of those respective age groups might elucidate the separate effects of  
300 these policies. That stands as a challenge for our future study.

301 The second point is underascertainment. Although the epidemic curves of  
302 COVID-19 in all countries are subject to underascertainment, it might be very difficult  
303 to evaluate the degree to which they are affected. It might bias the estimation result.

304 A third point is a lack of estimation of the outcomes such as dead or severe cases or  
305 necessary medical resources for the care of COVID-19 patients. We particularly  
306 examined how policies affect the reproduction number under countermeasures.  
307 Therefore, we ignored prediction of the entire course of the outbreak and its outcomes

308 such as the number of deaths. Nevertheless, outcomes are expected to be a primary  
309 concern for modelling. Moreover, the collapse of medical services can be expected to  
310 engender worse outcomes even if the reproduction number remains unchanged.  
311 Prediction of the effects of severe policies including lockdowns is anticipated as a  
312 challenge to be addressed in our future research.

313

#### 314 Conclusion

315 Results have demonstrated that VECSC can reduce the infectiousness of  
316 COVID-19 considerably: approximately to one. However, the figure is probably greater  
317 than one. Outbreaks might continue for a long time. Therefore, lockdown policies are  
318 expected to be as effective as VECSC if they are executed carefully. After VECSC, the  
319 reproduction number escalated again beyond that before VECSC. Similar phenomena  
320 might be observed after a lockdown. It is our earnest hope that results of the present  
321 study can contribute to governmental policy-making related to lockdown measures or  
322 other countermeasures to combat the spread and destruction of COVID-19.

323 The present study was based on the authors' opinions. Neither the results nor  
324 implications reflect any stance or policy of professionally affiliated bodies.

325

326 Acknowledgments

327 We acknowledge the great efforts of all staff at public health centers, medical  
328 institutions, and other facilities who are fighting the spread and destruction associated  
329 with COVID-19.

330

331 Reference

- 332 1. Japan Ministry of Health, Labour and Welfare. Press Releases.  
333 [https://www.mhlw.go.jp/stf/newpage\\_10723.html](https://www.mhlw.go.jp/stf/newpage_10723.html) (in Japanese) [accessed on  
334 April 10, 2020]
- 335 2. Japan Times on February 29  
336 [https://www.japantimes.co.jp/news/2020/02/29/national/politics-diplomacy/japan-s](https://www.japantimes.co.jp/news/2020/02/29/national/politics-diplomacy/japan-s-hinzo-abe-coronavirus-2/#.Xr4T9meP604)  
337 [hinzo-abe-coronavirus-2/#.Xr4T9meP604](https://www.japantimes.co.jp/news/2020/02/29/national/politics-diplomacy/japan-s-hinzo-abe-coronavirus-2/#.Xr4T9meP604) [accessed on May 15, 2020]
- 338 3. Lee PI, Hu YL, Chen PY, Huan YC, Hsueh PR, Lee PI, et al. Are children less  
339 susceptible to COVID-19? *J Microbiol Immunol Infect.* 2020 Feb  
340 25:S1684-1182;30039-6.
- 341 4. Kelvin A, Halperin S. COVID-19 in children: the link in the transmission chain.  
342 *The Lancet Infectious Diseases* ([https://doi.org/10.1016/S1473-3099\(20\)30236-X](https://doi.org/10.1016/S1473-3099(20)30236-X))
- 343 5. Qiu H, Wu J, Liang H, Yunling L, Song Q, Chen D. Clinical and epidemiological  
344 features of 36 children with coronavirus disease 2019 (COVID-19) in Zhejiang,  
345 China: an observational cohort study. *Lancet Infect Dis* 2020; published online  
346 March 25. [https://doi.org/10.1016/S1473-3099\(20\)30198-5](https://doi.org/10.1016/S1473-3099(20)30198-5).
- 347 6. Ohkusa Y, Sugawara T, Taniguchi K, Okabe N. Real-time estimation and  
348 prediction for pandemic A/H1N1(2009) in Japan. *J Infect Chemother.*  
349 2011;17:468-72.
- 350 7. Kimball A, Hatfield KM, Arons M, James A, Taylor J, Spicer K, Bardossy AC,  
351 Oakley LP, Tanwar S, Chisty Z, Bell JM, Methner M, Harney J, Jacobs JR, Carlson  
352 CM, McLaughlin HP, Stone N, Clark S, Brostrom-Smith C, Page LC, Kay M,

- 353 Lewis J, Russell D, Hiatt B, Gant J, Duchin JS, Clark TA, Honein MA, Reddy SC,  
354 Jernigan JA; Public Health ? Seattle & King County; CDC COVID-19  
355 Investigation Team. Asymptomatic and Presymptomatic SARS-CoV-2 Infections in  
356 Residents of a Long-Term Care Skilled Nursing Facility - King County,  
357 Washington, March 2020. *Morb Mortal Wkly Rep.* 2020;69:377-81. doi:  
358 10.15585/mmwr.mm6913e1.
- 359 8. Kurita J, Sugawara T, Ohkusa Y. Forecast of the COVID-19 outbreak and effects of  
360 self-restraint in going out in Tokyo, Japan.  
361 <https://medrxiv.org/cgi/content/short/2020.04.02.20051490v3>
- 362 9. Ferguson NM, Cummings DA, Cauchemez S, Fraser C, Riley S, Meeyai A, et al. :  
363 Strategies for containing an emerging influenza pandemic in Southeast Asia.  
364 *Nature*2005; 437:209-214.
- 365 10. Longini IM Jr, Nizam A, Xu S, Ungchusak K, Hanshaoworakul W, Cummings DA,  
366 Halloran ME: Containing Pandemic Influenza at the Source. *Science*2005;  
367 309:1083-1087.
- 368 11. Germann TC, Kadau L, Longini IM Jr, Macken CA: Mitigation strategies for  
369 pandemic influenza in the United States. *Proc Natl Acad Sci USA* 2006; 103:  
370 5935-5940.
- 371 12. Ferguson NM, Cummings DA, Fraser C, Cajka JC, Cooley PC, Burke DS:  
372 Strategies for mitigating an influenza pandemic. *Nature*2006; 442: 448-452.
- 373 13. Ohkusa Y, Sugawara T: Simulation Model of Pandemic influenza in the Whole of  
374 Japan, *Journal of Japanese Infectious Disease*2009; 62:98-106.

- 375 14. Britton T. Stochastic model review. <https://arxiv.org/pdf/0910.4443> [accessed on  
376 May 16]
- 377 15. Zhao S, Lin Q, Ran J, Musa SS, Yang G, Wang W, Lou Y, Gao D, Yang L, He D,  
378 Wang M. Preliminary Estimation of the Basic Reproduction Number of Novel  
379 Coronavirus (2019-nCoV) in China, From 2019 to 2020: A Data-Driven Analysis in  
380 the Early Phase of the Outbreak. *Int J Infect Dis* 2020[Online ahead of print]
- 381 16. Liu Y, Gayle AA, Wilder-Smith A, Rockliff J. The reproductive number of  
382 COVID-19 is higher compared to SARS coronavirus. *J Travel Med.* 2020.DOI:  
383 10.1093/jtm/taaa021
- 384 17. Lai C, Shih T, Ko W, Tang H, Hsueh P. Severe Acute Respiratory Syndrome  
385 Coronavirus 2 (SARS-CoV-2) and Coronavirus disease-2019 (COVID-19): The  
386 Epidemic and the Challenges. *Int J Antimicrob Agents.*  
387 DOI:10.1016/j.ijantimicag.2020.105924
- 388 18. Nishiura H, Oshitani H, Kobayashi T, Saito T, Sunagawa T, Wakita T, MHLW  
389 COVID-19 Response Team, Suzuki M. Closed environments facilitate secondary  
390 transmission of coronavirus disease 2019 (COVID-19). doi:  
391 <https://doi.org/10.1101/2020.02.28.20029272>
- 392 19. Oshitani H. Concept of the Counter Measure for COVID-19  
393 [https://www.jsph.jp/covid/files/gainen\\_0402.pdf](https://www.jsph.jp/covid/files/gainen_0402.pdf) (in Japanese) [accessed on My  
394 16, 2020]
- 395 20. He X, Lau EHY, Wu P, Deng X, Wang J, Hao X, Lau YC, Wong JY, Guan Y, Tan X,  
396 Mo X, Yanqing Chen Y, Liao B, Chen W, Hu F, Zhang Q, Zhong M, Yanrong Wu Y,

- 397 Zhao L, Zhang F, Cowling BJ, Li F, Leung GM. Temporal Dynamics in Viral  
398 Shedding and Transmissibility of COVID-19. *Nat Med* 2020;26:672-5.
- 399 21. Lauer SA, Grantz KH, Bi Q, Jones FK, Zheng Q, Meredith HR, Azman AS,  
400 Reich NG, Lessler J. The Incubation Period of Coronavirus Disease 2019  
401 (COVID-19) From Publicly Reported Confirmed Cases: Estimation and  
402 Application. *Annals of Internal Medicine* 2020. <https://doi.org/10.7326/M20-0504>
- 403 22. Hohle M, an der Heiden M. Bayesian nowcasting during the STEC O104: H4  
404 outbreak in Germany, 2011. *Biometrics*. 2014;70(4):993.1002.
- 405 23. Glockner S, Krause G, Hohle M. Now-casting the COVID-19 epidemic: The use  
406 case of Japan, March 2020. medRxiv preprint doi:  
407 <https://doi.org/10.1101/2020.03.18.20037473>.
- 408 24. Narula SC. Wellington JF. The Minimum Sum of Absolute Errors Regression: A  
409 State of the Art Survey. *International Statistical Review* 1982. 50: 317-26.  
410 doi:10.2307/1402501. JSTOR 1402501.
- 411 25. Phillips RF. Least absolute deviations estimation via the EM algorithm. *Statistics*  
412 and Computing 2002. 12: 281-5. doi:10.1023/A:1020759012226.
- 413 26. Siemsen E, Bollen KA. Least Absolute Deviation Estimation in Structural Equation  
414 Modeling. *Sociological Methods & Research* 2007. 36: 227-65.  
415 doi:10.1177/0049124107301946

416

417

418

419

420

421 Figure legend

422 Figure 1: Empirical distribution of incubation period published by Ministry of Labour,  
423 Health and Welfare, Japan

424 Notes: Bars shows distribution of incubation period among 91 cases whose exposed  
425 date and onset date were published by Ministry of Labour, Health and Welfare, Japan.  
426 Because the patients whose incubation was longer than 14 days were just 4%, they were  
427 included as bars on 14.

428


429 Figure 2: Observed epidemic curve of COVID-19 patients and predicted epidemic curve  
430 from the model based on the estimated reproduction number

431 Note: Bar indicates the observed epidemic curve and bold line indicates the predicted  
432 line based on the estimated reproduction numbers. Thin lines indicate range of the best  
433 fit line at each bootstrapping iteration. Between two vertical lines indicates voluntary  
434 event cancellation and school closure period.

435


436

1 Figure 1: Empirical distribution of incubation period published by Ministry of Labour,  
2 Health and Welfare, Japan  
3 (%)



6 Notes: Bars shows distribution of incubation period among 91 cases whose exposed  
7 date and onset date were published by Ministry of Labour, Health and Welfare, Japan.

1 Figure 2: Observed epidemic curve of COVID-19 patients and predicted epidemic curve  
2 from the model based on the estimated reproduction number  
3 (patients)



6 Note: Bar indicates the observed epidemic curve and bold line indicates the predicted  
7 line based on the estimated reproduction numbers. Thin lines indicate range of the best  
8 fit line at each bootstrapping iteration. Between two vertical lines indicates voluntary  
9 event cancellation and school closure period.

10  
11