Climate Affects Global Patterns Of Covid-19 Early Outbreak Dynamics

Gentile Francesco Ficetola[1,2,*] & Diego Rubolini[1*]

1 Dipartimento di Scienze e Politiche Ambientali, Università degli Studi di Milano, via Celoria 26,

I- 20133 Milano, Italy

2 Université Grenoble Alpes, CNRS, Université Savoie Mont Blanc, LECA, Laboratoire d'Ecologie

Alpine, F-38000 Grenoble

*Contributed equally to this work; order was decided with a coin toss

Correspondence: francesco.ficetola@gmail.com, diego.rubolini@unimi.it

Phone numbers
Gentile Francesco Ficetola: 0039 338 1844255
Diego Rubolini: 0039 02 503 14718

Abstract

Environmental factors, including seasonal climatic variability, can strongly impact on spatio-

temporal patterns of infectious disease outbreaks. We assessed the effects of temperature and

humidity on the global patterns of COVID-19 early outbreak dynamics during January-March 2020.

Climatic variables were the best drivers of global variation of confirmed COVID-19 cases growth

rates. Growth rates peaked in temperate regions of the Northern Hemisphere with mean temperature

of ~5°C and humidity of ~0.6-1.0 kPa during the outbreak month, while they decreased in warmer

and colder regions. The strong relationship between local climate and COVID-19 growth rates

suggests the possibility of seasonal variation in the spatial pattern of outbreaks, with temperate

regions of the Southern Hemisphere becoming at particular risk of severe outbreaks during the next

months.

A recently discovered coronavirus, SARS-CoV-2, is the aethiological agent of a pandemic disease,

Covid-19, causing severe pneumonia outbreaks at the global scale[1]. Covid-19 cases are now

reported in more than 155 countries and regions worldwide[2]. Three months after the discovery of

SARS-CoV-2, the global pattern and the early dynamics of Covid-19 outbreaks seem highly

variable. Some countries have been experiencing limited growth and spread of Covid-19 cases,

while others are suffering widespread community transmission and nearly exponential growth of

infections[2]. Given the impact of environmental conditions on the transmission of many pathogens,

we tested the hypothesis that the severity of Covid-19 outbreaks across the globe is affected by

spatial variation of key environmental factors, such as temperature, air humidity and pollution[3-7].

We then evaluated if this could help to illustrate global variation in the risk of severe Covid-19

outbreaks in the coming months.

Relying on a publicly available global dataset[2], we computed the daily growth rates r of

confirmed Covid-19 cases (Covid-19 growth rate hereafter) for 121 countries/regions (see the

Methods section and Table S6 in the Supplementary Appendix). We limited our measure of

epidemics growth rate to the first 5 days after reaching a minimum threshold of confirmed cases

(25, 50 or 100), as the mean incubation period of Covid-19 is ca. 5 days[8] and, immediately after the

first confirmed cases, many countries put in place unprecedented containment measures to mitigate

pathogen spread and community transmission[9]. Variation at these early epidemic growth rates

should best reflect the impact of local environmental conditions on disease spread. We restricted

analyses to data reported before March 19, as during that week many regions of the world adopted

stringent containment measures even in absence of large numbers of reported cases. For instance,

on March 17, 37 US states closed schools to prevent disease spread, including several states with

less than 25 confirmed Covid-19 cases[10]. We also considered additional factors that could affect

SARS-CoV-2 transmission dynamics, such as human population density, government per-capita

health expenditure, and average air pollution levels (fine particulate matter; see Methods).

Covid-19 growth rates showed high variability at the global scale (Fig. 1A-C). The observed

daily growth rate after reaching 50 cases (r_{50}) was on average 0.18 [95% CI 0.16-0.19], and ranged

from 0.01 (Kuwait) to 0.44 (Denmark). The highest growth rates were observed in temperate

regions of the Northern Hemisphere (Fig. 1C). Growth rates calculated using different minimum

thresholds of confirmed cases (25 or 100) were strongly positively correlated (see Methods),

indicating robustness of our results to the choice of thresholds.

Climate variables were the most important predictors of Covid-19 growth rate (Table S1).

The best-fitting linear mixed model suggested that r_{50} is non-linearly related to spatial variation in

mean temperature of the outbreak month (Fig. 1A, Tables S2-S3). Growth rates peaked in regions

with mean temperature of ~5°C during the outbreak month, and decreased both in warmer and

colder climates (Fig. A, Table S3). The comparison of models with different combinations of

predictors confirmed temperature as the variable with the highest relative importance in explaining

variation of r_{50} (Table S1), and temperature was the only parameter included in the best-fitting

model (Tables S2-S3). Temperature and humidity of the outbreak month showed a strong, positive

relationship across regions (Fig. S1), thus they could not be included as predictors in the same

model. When we repeated the analyses including humidity instead of temperature, r_{50} varied

significantly and non-linearly with humidity, peaking at ~0.6-1.0 kPa (Fig. 1B, Tables S4-S5). The

best model also showed slightly larger growth rates in countries with greater health expenditure

(Table S5), possibly because of more efficient early reporting and/or faster diagnosis of Covid-19

cases.

Results were highly consistent if we calculated growth rates after minimum thresholds of 25

or 100 cases (r_{25} and r_{100}, respectively) instead of 50 (Tables S3 and S5). Human population density

and air pollution showed very limited relative importance values (always < 0.50; Table S1).

We then displayed potential seasonal changes in Covid-19 growth rates by projecting our

best model of r_{50} in relation to temperature under the average temperature conditions of the current

(March) and next (June and September) months (Fig. 1C-E). The predicted global distribution of

Covid-19 growth rates based on March temperatures showed favorable conditions for disease

spread in most temperate regions of the Northern Hemisphere, and matched well with the observed

spatial distribution of Covid-19 growth rates during the January-March global outbreak (Fig. 1C).

The expected seasonal rise in temperatures during the next months could results in less suitable

conditions for Covid-19 spread in these areas. Conversely, seasonally decreasing temperatures

could accelerate disease spread in large areas of the Southern Hemisphere, including south

America, south Africa, eastern Australasia and New Zealand (Fig. 1D-E).

The management of Covid-19 outbreaks is undoubtedly one of the biggest challenges

governments will face in the coming months. Our spatially-explicit analysis suggests that, at least in

some parts of the world, ongoing containment efforts could benefit from the interplay between

pathogen spread and local climate. We do not claim that climate is the single major driver of Covid-

19 spread. The huge variation of Covid-19 growth rates among regions with similar climate indeed

suggests that diverse and complex social and demographic factors, as well as stochasticity, may

strongly contribute to determine the severity of Covid-19 outbreaks. Yet, climate can contribute to

explain the variability in global patterns of Covid-19 growth rates. In the coming months, we may

thus expect that large areas of the Southern Hemisphere will show environmental conditions

promoting severe Covid-19 outbreaks.

 Materials and methods

 Covid-19 dataset

 We downloaded the time series of confirmed Covid-19 cases from the Johns Hopkins University

 Center For Systems Science and Engineering (JHU-CSSE) GitHub repository

 (https://github.com/CSSEGISandData/Covid-19/: file 'time_series_19-covid-Confirmed.csv') [11].

 This datafile is updated once a day (at 23:59 UTC) and reports, for each day since January 22, 2020,

 all confirmed Covid-19 cases at the country level or at the level of significant geographical units

 belonging to the same country, which we defined here as 'regions' (e.g. US states or China

 provinces), whenever separate Covid-19 cases data for these regions are available. Initially, US data

 were reported by county but, as of March 9, they were reported at the state level. We therefore

 merged all US county data before March 9 to state level, and used state-level time series for

 subsequent calculations. With the exception of US data, in all other cases we maintained the

 original country/region information adopted by the JHU-CSSE. The datafile considered for the

 analyses was downloaded on March 19, 2020, and included confirmed Covid-19 cases until March

 18, 2020. From this dataset, we selected data for all countries / regions for which local outbreaks

 were detected. We define a local outbreak event when at least 50 positive cases were detected in a

 given country / region, and calculated the growth rate of confirmed Covid-19 cases between day 1

 and day 5, when day 1 was the day at which the 50 cases threshold was reached. We calculate the

 daily growth rate r of confirmed Covid-19 cases for each country/region, assuming an exponential

 growth as: $r = [\ln(n \text{ cases}_{day 5}) - \ln(n \text{ cases}_{day 1})] / 5$. We checked the robustness of our estimates of

 growth rate by calculating daily growth rate after the first 25, 50 or 100 cases (r_{25}, r_{50} and r_{100},

 respectively). Growth rates estimated at different thresholds were strongly positively correlated

 (Pearson's correlation coefficients, r_{25} vs. r_{50}: r = 0.74; r_{50} vs. r_{100}: r = 0.81).

 The dataset does not report information on containment measures, and these may be highly

 heterogeneous among countries/regions. We decided to calculate growth rate on the basis of the

first five days, in order to obtain an estimate of the non-intervened spread of the disease (i.e. before

stringent containment measures are undertaken). Five days provides a reasonable trade-off between

having to unreliable estimates of growth rates (if calculated on the basis of a smaller number of

days, e.g. 3), and obtaining growth rates influenced by the enforcement of heavy containment

measures (such as immediate isolation of confirmed cases). Five days is the median estimated time

spanning before the onset of symptoms [8], implying that infected patients might spread the virus for

5 days undetected in absence of preventive control measures. The mean estimated growth rate of

confirmed Covid-19 cases showed a tendency to decrease from r_{25} to r_{100} (mean and 95% c.i.: $r_{25} =$

0.21 [0.19-0.22, n = 121], $r_{50} = 0.18$ [0.16-0.19, n = 90], $r_{100} = 0.16$ [0.14-0.18, n = 69]), possibly

because of the progressive effect of containment measures that were adopted in different countries

at different times and different minimum thresholds after the onset of the local outbreak. We

excluded from analyses countries/regions with less than 100000 inhabitants (in our dataset, San

Marino only). As of March 19, 2020, the JHU-CSSE dataset provided information for a total of 121

countries/regions for the calculation of r_{25}, 90 for r_{50}, and 69 for r_{100}. The final list of

countries/regions included in the analyses, together with estimated confirmed Covid-19 growth

rates at different thresholds, is reported in Table S6.

Environmental and socio-economic variables

We considered two climatic variables that are known to affect the spread of viruses: mean air

temperature and vapor pressure, which is a measure of absolute humidity. Previous studies showed

that, for coronaviruses and influenza viruses, survival is generally higher at low temperature and

low values of absolute humidity [5,6,12-14]. For each country/region, we thus calculated the mean

monthly values for temperature (°C) and vapor pressure (kPa) for January, February and March on

the basis of the WorldClim 2.1 raster layers at 10 arc-minutes resolution [15]. We relied on

WorldClim climatic data because homogeneous data on conditions for the period January-March

2020 are not yet available at a global scale (see e.g.

https://www.ecmwf.int/en/forecasts/datasets/reanalysis-datasets/era5), and spatial variation among

areas of the world is generally much stronger than inter-annual variation for the same region [16]. As

additional predictors, we considered mean human population density [17] (population density

hereafter, expressed in inhabitants/km^2) and per-capita government health expenditure (health

expenditure hereafter) (indicator 'Domestic General Government Health Expenditure (GGHE-D)

per Capita in US$; average of 2015-2017 values downloaded from the World Health Organization

database at https://apps.who.int/nha/database). Health expenditure was available at country-level

only: hence, regions within countries were assigned the same health expenditure value. Finally, it

has been proposed that air pollution, and especially fine atmospheric particulate, could enhance the

persistence and transmission of coronaviruses [3,18]. We therefore extracted values of annual

concentration (µg/m^3) of ground-level fine particulate matter (PM2.5) for 2016 from the NASA

Socioeconomic Data and Applications Center [19], and calculated the mean abundance of PM2.5 for

each country/region. We performed all spatial analyses using the raster package in R [20].

Statistical analyses

We used linear mixed models (LMMs) to relate the global variation of r_{50}, r_{25} and r_{100} to the five

environmental predictors (temperature and humidity of outbreak month; population density; health

expenditure and PM2.5). To associate climate variables to the estimated r-values for each

country/region, we first extracted the mean month of the 5 days over which we computed the r-

values (rounded to the nearest integer) (outbreak month). We then assigned to the r-values of each

country/region the mean temperature and humidity of the month during which the outbreak

occurred. Country was included as a random factor to take into account potential non-independence

of growth rates from regions belonging to the same country. Non-linear relationships between

climatic factors and ecological variables are frequent [21], and in exploratory plots we detected a clear

non-linear relationship between r-values and climate. Therefore, for climatic variables, we included

in models both linear and quadratic terms. Humidity, population density, health expenditure and

PM2.5 were $\log_{10}$-transformed to reduce skewness and improve normality of model residuals.

We adopted a model selection approach to identify the variables most likely to affect the

global variation of Covid-19 growth rate [22]. We built models representing the different

combinations of independent variables, and ranked them on the basis of Akaike's Information

Criterion (AIC). AIC trades-off explanatory power vs. number of predictors; parsimonious models

explaining more variation have the lowest AIC values and are considered to be the "best models" [22].

For each candidate model, we calculated the Akaike weight ω_i, representing the probability of the model given the data

[23]. We then calculated the relative variable importance of each variable (RVI) as the sum of ω_i of the models where

each variable is included. RVI can be interpreted the probability that a variable should be included in the best model

[22,24]. Model selection analyses and the calculation of RVI can be heavily affected by collinearity among variables. In our

dataset, temperature and humidity showed a very strong positive correlation (Fig. S1 and Table S7); furthermore,

population density was strongly positively associated with PM2.5 (Figure S1 and Table S7). Therefore, temperature and

humidity, or population density and PM2.5, could not be considered together in the same models [24,25]. All other

predictors showed weak correlations and should not cause collinearity issues [25] (Table S7). We therefore repeated the

model selection for different combinations of uncorrelated variables. First, we considered temperature, health

expenditure and population density as independent variables. Then we repeated the analysis using humidity instead of

temperature, and we calculated the RVI of variables separately for these two model selection analyses. Finally, to assess

the role of PM2.5, we repeated these two model selections analyses using PM2.5 instead of health expenditure. The

RVI values for all tested models are reported in Table S1. Due to low RVI of PM2.5 in all models (Table S1), we

subsequently report detailed results of models including population density instead of PM2.5 (Tables S2-S5). To test the

robustness of our conclusion to subjective thresholds for the minimum number of cases, all analyses were repeated

considering the three estimates of Covid-19 growth rate as dependent variables (r_{25}, r_{50} and r_{100}).

LMMs were fitted using the lmer function of the lme4 R package [26], while tests statistics were calculated using

the lmerTest package [27]. To confirm that spatial autocorrelation did not bias the outcome of our

analyses, we calculated the spatial autocorrelation (Moran's I) of the residuals of best-fitting models

using the EcoGenetics package in R [28] at lags of 1000 km up to a maximum distance of 5000 km.

Model residuals did not show significant spatial autocorrelation at any lag (in all cases, |Moran's I|

< 0.10 and $P > 0.11$), suggesting that spatial autocorrelation was not a major issue in our analyses

[29].

References

1.   World Health Organization. Coronavirus disease (COVID-2019) situation reports:

https://www.who.int/emergencies/diseases/novel-coronavirus-2019/situation-reports/ [accessed 1

March 2020]; 2020.

2.   Dong E, Du H, Gardner L. An interactive web-based dashboard to track COVID-19 in real

time. The Lancet Infectious Diseases 2020:DOI:https://doi.org/10.1016/S473-3099(20)30120-1.

Data available at https://github.com/CSSEGISandData/COVID-19.

3.   Setti L, Passarini F, de Gennaro G, et al. Relazione circa l'effetto dell'inquinamento da

particolato atmosferico e la diffusione di virus nella popolazione.

http://www.simaonlus.it/wpsima/wp-content/uploads/2020/03/COVID19_Position-

Paper_Relazione-circa-l%E2%80%99effetto-dell%E2%80%99inquinamento-da-particolato-

atmosferico-e-la-diffusione-di-virus-nella-popolazione.pdf: Società Italiana di Medicina

Ambientale; 2020.

4.   Sajadi MM, Habibzadeh P, Vintzileos A, Shokouhi S, Miralles-Wilhelm F, Amoroso A.

Temperature and Latitude Analysis to Predict Potential Spread and Seasonality for COVID-19.

2020:Available at SSRN: https://ssrn.com/abstract=3550308.

5.   Kampf G, Todt D, Pfaender S, Steinmann E. Persistence of coronaviruses on inanimate

surfaces and their inactivation with biocidal agents. Journal of Hospital Infection 2020;104:246-51.

6.   Tamerius JD, Shaman J, Alonso WJ, et al. Environmental Predictors of Seasonal Influenza

Epidemics across Temperate and Tropical Climates. PLoS Pathog 2013;9:12.

7.   Wang J, Tang K, Feng K, Lv W. High Temperature and High Humidity Reduce the

Transmission of COVID-19 (March 9, 2020). Available at SSRN

2020:http://dx.doi.org/10.2139/ssrn.3551767

8.   Li Q, Guan X, Wu P, et al. Early Transmission Dynamics in Wuhan, China, of Novel

Coronavirus–Infected Pneumonia. New England Journal of Medicine 2020.

9.   Hellewell J, Abbott S, Gimma A, et al. Feasibility of controlling COVID-19 outbreaks by

isolation of cases and contacts. The Lancet Global Health 2020;8:e488-e96.

10.   Maxouris C. These states have some of the most drastic restrictions to combat the spread of

coronavirus. CNN 2020:https://edition.cnn.com/2020/03/17/us/states-measures-coronavirus-

spread/index.html.

11.   Dong E, Du H, Gardner L. An interactive web-based dashboard to track COVID-19 in real

time. The Lancet Infectious Diseases 2020.

12.   Shaman J, Kohn M. Absolute humidity modulates influenza survival, transmission, and

seasonality. Proceedings of the National Academy of Sciences 2009;106:3243.

13.   Lowen AC, Steel J. Roles of Humidity and Temperature in Shaping Influenza Seasonality.

Journal of Virology 2014;88:7692-5.

14.   Lowen AC, Mubareka S, Steel J, Palese P. Influenza virus transmission is dependent on

relative humidity and temperature. PLoS Pathog 2007;3:1470-6.

15.   Fick SE, Hijmans RJ. WorldClim 2: new 1-km spatial resolution climate surfaces for global

land areas. Int J Climatol 2017;37:4302-15.

16.   Harris I, Jones PD, Osborn TJ, Lister DH. Updated high-resolution grids of monthly

climatic observations – the CRU TS3.10 Dataset. Int J Climatol 2014;34:623-42.

17.   Center for International Earth Science Information Network CCU. Gridded Population of

the World, Version 4 (GPWv4): Population Density, Revision 11. Palisades, NY: NASA

Socioeconomic Data and Applications Center (SEDAC); 2018.

18.   van Doremalen N, Bushmaker T, Morris D, et al. Aerosol and surface stability of HCoV-19

(SARS-CoV-2) compared to SARS-CoV-1. medRxiv 2020;2020.03.09.20033217:doi:

https://doi.org/10.1101/2020.03.09.20033217.

19.   van Donkelaar A, Martin A, Brauer M, et al. Global Annual PM2.5 Grids from MODIS,

MISR and SeaWiFS Aerosol Optical Depth (AOD) with GWR, 1998-2016. Palisades NY: NASA

Socioeconomic Data and Applications Center (SEDAC). https://doi.org/10.7927/H4ZK5DQS;

2018.

20.   Hijmans RJ. raster: Geographic data analysis and modeling. R package version 3.0-7:

http://CRAN.R-project.org/package=raster; 2019.

21.   Legendre P, Legendre L. Numerical Ecology. Amsterdam: Elsevier; 2012.

22.   Burnham KP, Anderson DR. Model selection and multimodel inference: a practical

information-theoretic approach. New York: Springer Verlag; 2002.

23.   Lukacs PM, Thompson WL, Kendall WL, et al. Concerns regarding a call for pluralism of

information theory and hypothesis testing. J Appl Ecol 2007;44:456-60.

24.   Giam XL, Olden JD. Quantifying variable importance in a multimodel inference framework.

Methods Ecol Evol 2016;7:388-97.

25.   Dormann CF, Elith J, Bacher S, et al. Collinearity: a review of methods to deal with it and a

simulation study evaluating their performance. Ecography 2013;36:27-46.

26.   Bates D, Maechler M, Bolker BM, Walker S. Fitting Linear Mixed-Effects Models Using

lme4. J Stat Softw 2015;67:1-48.

27.   Kuznetsova A, Brockhoff B, Christensen HB. lmerTest Package: Tests in Linear Mixed

Effects Models. J Stat Softw 2017;82:1-26.

28.   Roser LG, Ferreyra LI, Saidman BO, Vilardi JC. EcoGenetics: An R package for the

management and exploratory analysis of spatial data in landscape genetics. Mol Ecol Resour

2017;17:e241-e50.

29.   Beale CM, Lennon JJ, Yearsley JM, Brewer MJ, Elston DA. Regression analysis of spatial

data. Ecol Lett 2010;13:246-64.

Figure 1. Variation of Covid-19 growth rates in relation to climate, and spatial predictions for
different months.
270
Variation of confirmed Covid-19 cases growth rates for the first 5 days after reaching a minimum
threshold of 50 cases (r_{50}) during the January-March 2020 pandemic outbreak (n = 90
countries/regions, see list in Table S6) in relation to the mean temperature (**Panel A**) and to the
mean absolute humidity of the outbreak month (**Panel B**). The lines are obtained from the best-
fitting linear mixed models (LMMs) of r_{50} in relation to temperature or humidity, respectively (see
Tables S3 and S5). The quadratic terms of both temperature and humidity were highly significant
(temperature: $F_{1,87} = 14.4$, $P < 0.001$; humidity: $F_{1,84} = 7.82$, $P = 0.006$; full details in Tables S3 and
S5). Shaded areas are 95% confidence band. **Panel C** shows the global patterns of r_{50}, with the size
of dots is proportional to the observed r_{50} value. The background shows the spatial prediction of
growth rates according to mean March temperatures[15]. Predictions are based on the best-fitting
LMM of r_{50} in relation to mean temperature of the outbreak month (Table S3). **Panels D and E**
show the spatial prediction of growth rates according to mean June and September temperatures[15],
highlighting that optimal conditions for disease spread appear in temperate regions of the Southern
Hemisphere.
285