Modeling COVID-19 epidemics in an Excel spreadsheet: Democratizing

the access to first-hand accurate predictions of epidemic outbreaks

Mario Moisés Alvarez[1,2,*], Everardo González-González[1,2], and Grissel Trujillo-de

Santiago[1,3]

[1] Centro de Biotecnología-FEMSA, Tecnologico de Monterrey, Monterrey 64849, NL, México

[2] Departamento de Bioingeniería, Escuela de Ingeniería y Ciencias, Tecnologico de Monterrey,

Monterrey 64849, NL, México

[3] Departamento de Ingeniería Mecatrónica y Eléctrica, Escuela de Ingeniería y Ciencias,

Tecnologico de Monterrey, Monterrey 64849, NL, México

(*) corresponding author: *mario.alvarez@tec.mx*

Abstract

COVID-19, the first pandemic of this decade and the second in less than 15 years, has

harshly taught us that viral diseases do not recognize boundaries; however, they truly do

discriminate between aggressive and mediocre containment responses.

We present a simple epidemiological model that is amenable to implementation in Excel

spreadsheets and sufficiently accurate to reproduce observed data on the evolution of the

COVID-19 pandemics in different regions (i.e., Italy, Spain, and New York City (NYC)).

We also show that the model can be adapted to closely follow the evolution of COVID-19

in any large city by simply adjusting two parameters related to (a) population density and

(b) aggressiveness of the response from a society/government to epidemics. Moreover, we

show that this simple epidemiological simulator can be used to assess the efficacy of the

response of a government/society to an outbreak.

The simplicity and accuracy of this model will greatly contribute to democratizing the

availability of knowledge in societies regarding the extent of an epidemic event and the

efficacy of a governmental response.

Keywords: *COVID-19, coronavirus, SARS-CoV2, mathematical modeling, epidemic,*

pandemic, Excel

32   Preprint medRxiv

34 **Introduction**

A SARS-CoV2 (COVID-19) pandemic was declared by the World Health Organization in

March 2020. More than 100,000 positive cases of COVID-19 infection had been declared

worldwide at that point, mainly in China, Italy, Iran, Spain, and other European countries.

By the end of March 2020, the official cumulative number of infected worldwide ascended

to more than 700,000, with a toll of death higher than 32,000 and a strong presence in Las

Americas, mainly in the USA[1]. COVID-19, the first pandemic of this decade and the

second in less than 15 years, has harshly taught us that viral diseases do not recognize

boundaries; however, they truly do discriminate between aggressive and mediocre

containment responses.

Indeed, three months have passed since the emergence of COVID-19, and we have been

able to observe exemplary responses from some Asian countries (i.e., China[2], South

Korea[3], Singapore[4], and Japan), some highly aggressive responses in Europe and America

(i.e., Germany and USA), and several delayed or not so effective responses from other

regions (i.e., Italy and Spain)[5]. At this point, some territories in Latin America are just

experiencing the "lag phase" of the COVID-19 pandemic at home and do not appear having

yet implemented proper containment measures as rapidly as needed.

The gap between developed and developing countries may explain some of the differences

in the scale of the responses that we are observing. Countries that are better equipped than

others in terms of high-end scientific development, diagnostics technology, and health care

infrastructure may respond more efficaciously to a pandemic scenario. However, other

tools, such as mathematical modeling, are much more widely available and may be of

extraordinary value when managing epidemic events such as the COVID-19 pandemics. To

date, many papers have reported the use of mathematical models and simulators to evaluate

the progression of COVID-19 in local or more global settings[63,7–9]. Predictions on the

possible evolution of COVID-19 based on mathematical modeling could therefore represent

important tools for designing and/or evaluating countermeasures[8,10–12].

However, mathematical modeling may (and probably should) become a much more

available tool in the case of public health emergencies—one ideally widely available to

practically any citizen in any of our societies. One decade ago, during the influenza

pandemics, mathematical modeling of epidemic events was the realm of privileged

epidemiologists who had (a) a fast computer, (b) programing experience, and (c) and access

to epidemiological data. Today, those three ingredients are now reduced to a convectional

laptop, very basic differential equation-solving skills, and access to a website with reliable

online statistical information on the epidemics.

The main purpose of this contribution is to demonstrate that a simple mathematical model,

amenable to implementation in an Excel spreadsheet, can accurately predict the evolution

of an epidemic event at a local level (i.e., in any major urban area). This may be extremely

valuable for government officials who must predict, with high fidelity, the progression of

an epidemic event to better design their action strategies. Moreover, the democratization of

the modeling of complex epidemic events will empower citizens, enabling them to forecast,

decide, and evaluate. For instance, using this simple model, virtually any citizen could

assess, in real time, the efficacy of the actions of her/his society in the face of an outbreak.

Rationale of the model formulation

Here, we construct a very simple epidemiological model for the propagation of COVID-19

in urban areas. The model is based on a set of differential equations. The first equation of

the set (equation 1) states that the rate of accumulation of infected habitants in an urban

area (assumed to be a closed system) is given by the sum of the number of new infections

(positive contribution), the number of recovered patients (negative contribution), and the

number of deaths (negative contribution). A second differential equation states that the rate

of accumulation of the infected but asymptotic population is proportional to the population

of infected and symptomatic subjects (equation 2). Two additional equations relate the

number of deaths and recovered patients with the number of newly infected ones (equation

3 and 4). Finally, the rate of depletion of the pool of the population susceptible to infection

is given by the sum of recovered patients, asymptomatic infected, and deaths (equation 5).

Recent experimental evidence suggests that rhesus macaques that recovered from SARS-

CoV-2 infection could not be reinfected[13]. However, at this point, the acquisition of full

immunity to reinfection has not been proved in humans, although it is well documented for

other coronavirus infections, such as SARS, and MERS[14,15]. The analysis of sera of one

COVID-19 patient showed a peak production of specific IgGs against SARS-COV-2 by

two weeks after the onset of symptoms [16]. Based on immunological information on SARS

and MERS epidemiology and the limited evidence on the nature of the host immune

response to SARS-COV-2, we assume here that recovered patients become immune to

reinfection.

$dX_s/dt = R_{Infected-s} - R_{Recovered} - R_{Death}$ Equation (1)

$dX_{as}/dt = R_{Infected-as} = (2.5/1.0) * R_{Infected-s}$ Equation (2)

$dD/dt = R_{Death} = 0.023 * R_{Infected-s}$ Equation (3)

$dR/dt = R_{Recovered} = 0.977 * R_{Infected-s}$ Equation (4)

$dP_s/dt = - R_{Infected-as} - R_{Recovered} - R_{Death}$ Equation (5)

This system is equivalent to:

$dX_s/dt = R_{Infected-s} - R_{Recovered} - R_{Death}$ Equation (1)

where:

$R_{Infected-as} = (2.5/1.0) * R_{Infected-s}$ Equation (2´)

$R_{Death} = 0.023 * R_{Infected-s}$ Equation (3´)

$R_{Recovered} = 0.977 * R_{Infected-s}$ Equation (4´)

$P_{s\_n} = P_{s\_n-1} - (X_{as} + R + D)$ Equation (5´)

In this system, all equations depend on $R_{Infected-s}$. Here, we propose a simple formulation for

the evaluation of $R_{Infected-s}$ at the onset of a local epidemic event.

$R_{Infected-s} = dI_s/dt = \mu_o I_s$ Equation (6)

where μ_o is the specific rate of infection of a population in a large and vastly uninfected

urban area. We further propose that μ_o may be calculated from actual epidemiological data

corresponding to the first exponential stage of COVID-19 local epidemics. We determined

the appropriate ranges of values for μ_o by analyzing publicly available data from different

websites that continuously monitor the progression of confirmed cases of COVID-19 for

different nations (Table 1).

Table 1. Websites displaying COVID-19 data in practically real time.

• Our World in data:
https://ourworldindata.org/coronavirus
• El País
https://elpais.com/sociedad/2020/03/16/actualidad/1584360628_538486.html
• Coronavirus COVID-19 Global Cases by the Center for Systems Science and Engineering
(CSSE) at Johns Hopkins University (JHU).
https://gisanddata.maps.arcgis.com/apps/opsdashboard/index.html#/bda7594740fd40299423467b48e9ecf6
• Wikipedia, The Free Encyclopedia
https://en.wikipedia.org/wiki/2020_coronavirus_pandemic_in_Iran

This model correctly describes the evolution of the number of newly infected during the

initial stage of the epidemic episode. For later times, the rate of new infections is corrected

by a term that depends on the demographic density (Dd) of the region. Therefore:

$R_{Infected-s} = dI_s/dt = \mu_o \, I_s \, (Dd/Dd_{ref})$ Equation (7)

In equation (7), $Dd = P_s/A$, where A is the surface area of the region subject to analysis. In

this formulation, Dd is the total number of inhabitants of the region who are susceptible to

infection, while Dd_{ref} is a value of demographic density in a densely populated urban area

that the model uses as a reference. In this work, the demographic density of the city of

Madrid is used as Dd_{ref}. Furthermore, since Dd is a function that considers only the

population susceptible to infection, a counter is needed to continuously update the number

of recovered patients, asymptomatic patients, and deaths. Therefore, at each time step

during the numerical integration, the susceptible population is updated by subtracting the

number of number of recovered patients, asymptomatic patients, and deaths. Note that in

our Excel spreadsheet, we use Dd/Dd_{ref} = density_Factor (Supplemental Excel File 1).

Defining an expression for $R_{Infected-s}$ enables stepwise numerical integration, for example by

the Euler method. We have implemented this solution in a spreadsheet. To that aim,

differential equations (1) and (7) should be converted into their corresponding equations of

differences:

$\Delta X_s = \{R_{Infected-s} - R_{Recovered} - R_{Death}\}\, \Delta t$ Equation (8)

$\Delta I_s = \{\mu_o\, I_s\, (Dd/Dd_{ref})\}\, \Delta t$ Equation (9)

For all the simulation results presented here, we set Δt=1h= 1/24 day. We have solved this

differential set, step by step, updating the values of $R_{Infected-s}$, $R_{Recovered}$, R_{Death}, and P_s,

according to equations (2´) to (5´). The ratio (Dd/Dd_{ref}) is also recalculated at each time

step using the updated value of P_s from equation (5´).

Selection of relevant epidemiological parameters for COVID-19

The number of asymptomatic inhabitants was calculated under the assumption that only

~30.0% of the infected population develops symptomatology (2.5 asymptomatic subjects

per 1.0 symptomatic subject). This assumption should be regarded as speculative, since

very limited information specific for the ration between symptomatic and asymptomatic

COVID-19 patients is available at this point.[17,18] The percentage of asymptomatic

infections during pandemic Influenza A/H1N1/2009, based on epidemiology studies

founded in serological analysis in a vast range of geographical settings, has been estimated

has been between 65 and 85%[19]. These values are also consistent with the high number of

asymptomatic infected subjects estimated for other pandemic events. For instance, in the

context of pandemic influenza A/H1N1/2009, up to 20–40% of the population in urban

areas (i.e., Monterrey, México, and Pittsburgh, USA) [20,21] exhibited specific antibodies

regardless of experiencing symptoms, while the fraction of confirmed symptomatic

infections was lower than less than 10%.

In addition, the average time of sickness was set at 14 days in our simulations, within the

range reported from 14 to 32 days[22], with a median time to recovery of 21 days[23].

Therefore, the number of patients recovered (R) is calculated as a fraction of 0.977 of those

infected 14 days previously. Similarly, asymptomatic patients are only removed from the

pool of susceptible after full recovery. Note that, in the current version of our model,

asymptomatic patients are not considered part of the population capable of transmitting

COVID-19, despite recently reported evidence that suggests that asymptomatic subjects (or

minimally symptomatic patients) may exhibit similar viral loads[24] to those of symptomatic

patients and may be active transmitters of the disease[2,25]. The number of deceased patients

was calculated as 0.023 of those infected 14 days before. This mortality percentage (case

fatality rate) lies within the range reported in recent literature for COVID-19[9,26–28]. The

time lapse of 14 days between the onset of disease and death was statistically estimated by

Linton et al. in a recent report [29].

The straightforward implementation of the model in Excel (Supplemental Excel File 1),

using the set of parameters described before, allows the calculation of all populations (I_s, X_s,

X_{as}, D, R, and P_s) every hour. Note that this model enables the description of the

progressive exhaustion of the epidemic, as expected by the progressive depletion of the

susceptible population. Next, we discuss criteria for selection of the values of μ_o based on

the initial behavior of the COVID-19 Pandemic at different urban areas around the globe.

Estimation of specific epidemic rate values

Figure 1A shows the progression on the number of COVID-19 positive cases in different

regions, namely Spain (mainly Madrid), Iran (mainly Tehran), and New York City (NYC).

We have selected these three data sets to illustrate that the evolution of the epidemic has a

local flavor that mainly depends on the number of initial infected persons, the demographic

density, and the set of containment measures taken by government officials and society.

Figure 1B shows the natural log of the cumulative number of infections over time for the

same set of countries. This simple plotting strategy is highly useful for analyzing the local

rate of progression of the pandemic. If the local epidemic progression is consistent with a

simple first order exponential model where $dI/dt = \mu*I$, then the integral form of this

equation renders the linear equation: $\ln I/I_o = \mu*t$. During the exponential phase, a straight

line should be observed, and the slope of that line denotes the specific rate (μ) of the

epidemic spreading. Note that COVID-19 has exhibited a wide range of spreading rates in

different countries (from ~0.3 to ~0.9 day^{-1}). Note also that μ is related to the doubling time

(t_d), often reported in population and epidemiological studies, by the equation $t_d = \ln 2 / \mu$.

Therefore, ranges of doubling times between 0.75 and 2.45 days are observed just among

these three regional cases.

Figure 1. Epidemiological data related to the onset of a COVID-19 pandemic in different regions. (A) Cumulative number of positive cases of COVID-19 infection in Spain (yellow circles), Iran (green squares), and NYC (blue triangles) during the first days after the outbreak. (B) Natural logarithm of the cumulative number of positive cases of COVID-19 infection in Spain (yellow circles), Iran (green squares), and NYC (blue triangles). (C) Cumulative number of positive cases of COVID-19 infection in Italy (blue squares) and South Korea (red circles). (D) Natural logarithm of the cumulative number of positive cases of COVID-19 infection in Italy (blue squares) and South Korea (red circles). Two clearly distinctive exponential stages are observed in the case of South Korean progression.

Different exponential stages, perfectly distinguishable by their exhibition of different slopes

(Table 2), may be observed within the same time series. For instance, the outbreak in NYC

(Figure 1B; blue symbols) was first described by an extremely high slope (μ_o = 0.926 day$^-$

^{1}). However, after a series of measures adopted in NYC by the federal, state, and local

governments, the specific growth rate of the epidemics fell to μ = 0.308 day^{-1}.

The last point is extremely important, since two drastically different slopes can be observed

before and after a package of adequate measures within the same territory. In addition, two

localities that experienced similar initial specific epidemic rates may exhibit dramatically

different evolutions as a function of the initial response of government and society (Figure

1C,D). For instance, while the COVID-19 epidemics in Italy and South Korea exhibited

practically equal μ_o values, the Italian outbreak has maintained the same growth rate

throughout 20 days, while South Korea has set an example by effectively and rapidly

lowering the specific epidemic rate to nearly 0 in just two weeks.

Table 2. Specific infection rates (μ_o) and associated doubling times (t_d) for COVID-19
infection in different geographic regions.
246

Territory	Temporality	μ	t_d
Spain (Madrid)	initial	0.358	1.937
Italy	initial	0.326	2.128
Italy	after stringent measures	0.119	5.849
Iran	initial	0.491	1.411
Iran (Tehran)	initial	0.506	1.370
Germany	initial	0.280	2.474
NYC	initial	0.591	1.173
NYC	after measures	0.293	2.362
South Korea	initial	0.293	2.362
South Korea	after stringent measures; massive testing	0.000	ND[*]
France	initial	0.379	1.828
France	after measures	0.161	4.311
Mexico	initial	0.209	3.324

247   (*) Not determinable
248

Validation and predictions

We have run different scenarios to validate the predictive capabilities of our epidemic

model for COVID-19. Overall, the model is capable of closely reproducing the progression

of reported cases for urban areas of more than 5×10^6 inhabitants (i.e., Iran, the city of

Tehran in Iran, Spain, and NYC). We found that, adapting the model to a particular locality

is straightforward and only requires (a) the calculation of the population and the surface

area of the urban area, and (b) the selection of a t_d value (time to doubling the name of

infections). Note that our model is formulated in terms of values of the specific epidemic

growth rate (μ_o for the onset of the epidemic and μ for later times). However, expressing

the specific epidemic rate in terms of doubling time ($t_d = \text{Ln } 2/\mu$) is more practical and

simpler to communicate and understand (Table 2).

The selection of μ_o (t_d) can be easily done by fitting the prediction to the initial set of

reported cases of infection. In our experience, four to five reliable data points are needed

for a good fit. For instance, Figure 2 shows the predicted trend of the pandemic in NYC

during the first 28 days of March, 2020. In addition, we set ($Dd/Dd_{ref} = 1.90$), since the

population density in NYC is 1.90-fold higher than that in Madrid. A value of $t_d = 2.25$ was

also set for the first week of this simulation. Later, at day 7 (March 7), we reset the value of

t_d to 3.75 to reflect the modification of the slope of the local epidemic event in NYC (Figure

1d), due to the implemented measures of containment. Based on this exercise, we foresee

that this simple modeling tool can be used to evaluate the efficacy of containment

strategies. In other words, the value of μ_o required in the simulation to adapt the predicted

data to the actual trend of the local epidemic provides an indicator of the local rate of

spreading of the pandemic.

Figure 2. Progression of the COVID-19 Pandemic in NYC. (A) Initial evolution of the number

of positive cases of COVID-19 in NYC. Actual data points, as officially reported, are shown using

red circles. Simulation predictions are described by the blue dotted line. (B) Model prediction of

the total number of symptomatic patients through the months of March and April. (C) Model

prediction of new cases of COVID-19 during the period from March 1 to May 20, 2020 if no

further containment actions are adopted.

Therefore, the differences between μ_o before and after interventions provide a real-time

quantitative measure of the effectiveness of that set of measures. This can be extremely

useful when assessing the efficacy of control of epidemics. For example, for NYC, this

simple model states that the set of containment measures adopted during the first week of

March in NYC diminished the specific rate of the epidemic by increasing the doubling time

of infections from a value of 2.25 to 3.75 days.

The ability to make close predictions of the progression of cases in a particular region has

profound and enabling implications. For example, in March 15[th], our simulations predicted

that, in absence of more aggressive containment measures (yellow trend in Figure 2A), the

peak of infections in NYC will be reached by April 10, 2010, after reaching the

unprecedented value of 11,000 new cases per day, and a cumulative number of 1×10^6

citizens infected. However, we observed a deviation from this prediction by the third week

of March that we attribute to the stringent measures of social distancing established in NYC

earlier that week. Accordingly, we multiplied the value of (Dd/Dd_{ref}) in our simulations by

a factor of 0.50 to properly fit the new trend on actual cases (blue trend in Figure 2A). Note

that his suggest that the measures of social distancing imposed in NYC were equivalent to

decrease the effective demographic density to 50%. At the end of March, after this

adjustment, our model forecasts a peak of infections of nearly 5,000 new cases per day (less

than half than the prediction before social distancing), and a cumulative number of 1×10^6

citizens infected.

Effect of social distancing

Social distancing has been regarded as the one of the most effective buffering measures for local COVID-19 epidemics[30,31]. Next, we evaluate the effect of different degrees of social distancing on the shape of the epidemic curve for NYC, one of the most densely urban areas worldwide. This evaluation is straightforward, since the formulation of our model explicitly considers the demographic density of the region as the most important modifier of the rate of progression of the epidemics.

In the Excel implementation of the model, we multiply the demographic ration (Dd/Dd_{ref}) by 0.75 to calculate the impact of distancing measures that would diminish social contact by 25%. Similarly, we multiply (Dd/Dd_{ref}) by 0.50 to simulate the effect of a scenario of social distancing that would diminish close social interaction by 50%. Figure 4 shows the effect of three different degrees/levels of social distancing on the cumulative number of infections (Figure 3A) and on the number of new cases of infection per day (Figure 3B).

Social distancing has a clear buffering effect on the epidemics, delaying the occurrence of the peak of infections and distributing the number of cases across a longer time span. This is remarkably important as it provides time for proper attention to patients with severe symptomatology[5].

For instance, our results suggest that, for an urban area such as NYC, imposing measures that guarantee a social distance equivalent to a decrease in demographic density of 50% will delay the peak of maximum number of infections by 25 days and will decrease its intensity from 23000 to 9000 new cases of infection per day. In turn, this implies a lower demand for hospital beds per day during the epidemics and may mark the difference between a manageable crisis and a public health catastrophe[5,30].

Figure 3. Prediction of the effect of social distancing on the progression of the COVID-19 pandemics in New York City (NYC). (A) Model prediction of the total number of symptomatic patients from March 1 to May 31, 2020 for different scenarios of social distancing: social distancing as in March 20, 2010 (blue line, current prediction); social distancing effective on March 20, whereby the effective demographic density in NYC is reduced by 25% (light green line); social distancing effective on March 20 whereby the effective demographic density in NYC is reduced by 50% (red line); and social distancing effective on March 10, whereby the effective demographic density in NYC is reduced by 25% (dark green line). (B) Model prediction for the number of new infections per day for each of the scenarios of social distancing described.

339 Interestingly, the effect of anticipating measures of social distancing has a moderate effect

340 on retarding the infection curve but not on decreasing the cumulative number of infections.

341 This moderate gain of time provides additional leeway for planning interventions or

342 allocating resources, with time being gold during pandemic events.

Prediction in real time

We are currently following the onset of the COVID-19 pandemic in Monterrey, the second

most industrialized city in México and the third most populated. Monterrey has a similar

demographic density to that of Madrid (Dd/Dd_{ref}=0.95). In addition, we set $t_d = 2.5$, based

on proper fitting to the first set of official values of COVID-19 infected announced for

Monterrey by the local authorities from March 15 to March 19, 2020. Remarkably, the

simulation results have accurately predicted the nine subsequent actual values, as officially

reported from March 19 to March 28 (Figure 4 A). Monitoring actual data, while

comparing with model predictions, enables real time assessment of the effectiveness of the

containment measures. In turn, this empowers officials, scientists, health care providers,

and citizens. Moreover, friendly and widely available mathematical modeling enables

rational planning. For instance, according to the pandemic scenario predicted for

Monterrey, in the absence of further containment measures and stricter social distancing,

the total number of symptomatic infected will surpass 650,000 persons (Figure 4B), and the

number of new infections per day (Figure 4C) will exhibit a peak of 2000 by the end of

April. The simulation may be used to forecast the demand of beds during the month of

April 2020, which is estimated to exhibit a peak of nearly 50000.

Figure 4. Progression of the COVID-19 pandemic in the metropolitan area of Monterrey, Nuevo León, Mexico. (A) Initial evolution of the number of positive cases of COVID-19 in the metropolitan area of Monterrey. Actual data points, as officially reported, are shown using red circles. Simulation predictions are described by the blue dotted line. (B) Model prediction of the total number of symptomatic patients from March 1 to May 20, 2020. (C) Model prediction of new cases of COVID-19 during the period from March 1 to May 20, 2020 if no further containment actions are adopted. (D) Estimation of the number of beds needed during the month of April 2020 in Monterrey, based on the number of patients that will require hospitalization according to the model predictions.

This estimate considers that only 10.0% of the symptomatic patients will require

hospitalization, which may be optimistic. Reports based in 44000 COVID-19 cases in

China indicate that the percentage of patients with severe symptoms may be 14%, with a

5% of critical cases[32]. In prospective, the total number of beds in the Mexican public health

sector is estimated in 20,000 (for the whole country).

Concluding remarks

We used a set of differential equations, recent epidemiological data regarding the evolution

of COVID-19 infection in a reduced set of regions (i.e., Spain, Iran, and NYC), and basic

information on the characteristics of COVID-19 infection (i.e., time from infection to

recovery, case mortality rate) to accurately recreate the onset of the COVID-19 in two

urban areas with different demographic characteristics (i.e., NYC and Monterrey, México).

We showed that the model can be adapted to closely follow the evolution of COVID-19 in

densely populated urban areas by simply adjusting two parameters related to (a) population

density and (b) aggressiveness of the response from a society/government to epidemics.

Scenarios such as those currently unfolding in Iran, Italy, or Spain emphasize the

importance of planning ahead during epidemic events. The availability of a simple model

may be highly enabling for local governments, physicians, civil organizations, and citizens

as they struggle in their endeavor to accurately forecast the progression of an epidemic and

formulate a plan of action. As previously stated, the use of simple/user-friendly models to

evaluate in (practically) real time the effectiveness of containment strategies or programs

may be a powerful tool for analyzing and facing epidemic events[6,12]. This contribution

shows the prediction potential of an extremely simple simulation tool that can be used by

practically any citizen with basic training in Excel.

Acknowledgments

MMA, EGG, and GTdS acknowledge the funding received from CONACyT (Consejo

Nacional de Ciencia y Tecnología, México).

Author contributions

404   MMA, EGG, and GTdS collected and analyzed epidemiology data. MMA formulated the

405   model and run the simulations. MMA and GTdS wrote the manuscript. All authors

406   reviewed and approved the manuscript.

Competing interest

The authors declare no competing interests.

References:

1.   Holshue, M. L. *et al.* First Case of 2019 Novel Coronavirus in the United States. *N.*

Engl. J. Med. (2020). doi:10.1056/nejmoa2001191

2.   MacIntyre, C. R. Global spread of COVID-19 and pandemic potential. *Glob.*

Biosecurity **1**, (2020).

3.   Choi, S. C. & Ki, M. Estimating the reproductive number and the outbreak size of

Novel Coronavirus disease (COVID-19) using mathematical model in Republic of

Korea. *Epidemiol. Health* e2020011 (2020). doi:10.4178/epih.e2020011

4.   Wong, J. E. L., Leo, Y. S. & Tan, C. C. COVID-19 in Singapore-Current

Experience: Critical Global Issues That Require Attention and Action. *JAMA* (2020).

doi:10.1001/jama.2020.2467

5.   Remuzzi, A. & Remuzzi, G. COVID-19 and Italy: what next? *Lancet* (2020).

doi:10.1016/s0140-6736(20)30627-9

6.   Roosa, K. *et al.* Real-time forecasts of the COVID-19 epidemic in China from

February 5th to February 24th, 2020. *Infect. Dis. Model.* **5**, 256–263 (2020).

7.   Peng, L., Yang, W., Zhang, D., Zhuge, C. & Hong, L. Epidemic analysis of COVID-

19 in China by dynamical modeling. (2020).

8.   Kucharski, A. J. *et al.* Early dynamics of transmission and control of COVID-19: a

mathematical modelling study. *Lancet Infect. Dis.* (2020). doi:10.1016/S1473-

3099(20)30144-4

9.   Jung, S. *et al.* Real-Time Estimation of the Risk of Death from Novel Coronavirus

(COVID-19) Infection: Inference Using Exported Cases. *J. Clin. Med.* **9**, 523 (2020).

10.   Hellewell, J. *et al.* Feasibility of controlling COVID-19 outbreaks by isolation of

cases and contacts. *Lancet Glob. Heal.* **8**, e488–e496 (2020).

11.   Gostic, K., Gomez, A. C. R., Mummah, R. O., Kucharski, A. J. & Lloyd-Smith, J. O.

Estimated effectiveness of symptom and risk screening to prevent the spread of

COVID-19. *Elife* **9**, (2020).

12.   Cauchemez, S., Hoze, N., Cousien, A., Nikolay, B. & ten bosch, Q. How Modelling

Can Enhance the Analysis of Imperfect Epidemic Data. *Trends in Parasitology* **35**,

369–379 (2019).

13.   Bao, L. *et al.* Reinfection could not occur in SARS-CoV-2 infected rhesus

macaques. *bioRxiv* 2020.03.13.990226 (2020). doi:10.1101/2020.03.13.990226

14.   Prompetchara, E., Ketloy, C. & Palaga, T. Allergy and Immunology Immune

444   responses in COVID-19 and potential vaccines: Lessons learned from SARS and

445   MERS epidemic. doi:10.12932/AP-200220-0772

446   15.   Liu, W. *et al.* Two-Year Prospective Study of the Humoral Immune Response of

447   Patients with Severe Acute Respiratory Syndrome. *J. Infect. Dis.* **193**, 792–795

448   (2006).

449   16.   Zhou, P. *et al.* A pneumonia outbreak associated with a new coronavirus of probable

450   bat origin. *Nature* **579**, 270–273 (2020).

451   17.   Mizumoto, K., Kagaya, K., Zarebski, A. & Chowell, G. Estimating the

452   asymptomatic proportion of coronavirus disease 2019 (COVID-19) cases on board

453   the Diamond Princess cruise ship, Yokohama, Japan, 2020. *Eurosurveillance* **25**,

454   2000180 (2020).

455   18.   Nishiura, H. *et al.* Estimation of the asymptomatic ratio of novel coronavirus

456   infections (COVID-19). *medRxiv* 2020.02.03.20020248 (2020).

457   doi:10.1101/2020.02.03.20020248

458   19.   Leung, N. H. L., Xu, C., Ip, D. K. M. & Cowling, B. J. The fraction of influenza

459   virus infections that are asymptomatic: a systematic review and meta-analysis.

460   doi:10.1097/EDE.0000000000000340

461   20.   Elizondo-Montemayor, L. *et al.* Seroprevalence of antibodies to influenza

462   A/H1N1/2009 among transmission risk groups after the second wave in Mexico, by

463   a virus-free ELISA method. *Int. J. Infect. Dis.* **15**, e781–e786 (2011).

464   21.   Zimmer, S. M. *et al.* Seroprevalence Following the Second Wave of Pandemic 2009

465   H1N1 Influenza in Pittsburgh, PA, USA. doi:10.1371/journal.pone.0011601

22.   Lan, L. *et al.* Positive RT-PCR Test Results in Patients Recovered From COVID-19.

JAMA (2020). doi:10.1001/jama.2020.2783

23.   Bi, Q. *et al.* Epidemiology and Transmission of COVID-19 in Shenzhen China:

Analysis of 391 cases and 1,286 of their close contacts. *medRxiv*

2020.03.03.20028423 (2020). doi:10.1101/2020.03.03.20028423

24.   Zou, L. *et al.* SARS-CoV-2 Viral Load in Upper Respiratory Specimens of Infected

Patients. *N. Engl. J. Med.* **382**, 1177–1179 (2020).

25.   Bai, Y. *et al.* Presumed Asymptomatic Carrier Transmission of COVID-19. *JAMA*

(2020). doi:10.1001/jama.2020.2565

26.   Lai, C. C., Shih, T. P., Ko, W. C., Tang, H. J. & Hsueh, P. R. Severe acute

respiratory syndrome coronavirus 2 (SARS-CoV-2) and coronavirus disease-2019

(COVID-19): The epidemic and the challenges. *International Journal of*

Antimicrobial Agents **55**, 105924 (2020).

27.   Xu, Z. *et al.* Pathological findings of COVID-19 associated with acute respiratory

distress syndrome. *Lancet Respir. Med.* **0**, (2020).

28.   Porcheddu, R., Serra, C., Kelvin, D., Kelvin, N. & Rubino, S. Similarity in Case

Fatality Rates (CFR) of COVID-19/SARS-COV-2 in Italy and China. *J. Infect. Dev.*

Ctries. **14**, 125–128 (2020).

29.   Linton, N. M. *et al.* Epidemiological characteristics of novel coronavirus infection:

A statistical analysis of publicly available case data.

doi:10.1101/2020.01.26.20018754

30.   Anderson, R. M., Heesterbeek, H., Klinkenberg, D. & Hollingsworth, T. D. How

will country-based mitigation measures influence the course of the COVID-19

epidemic? *The Lancet* **395**, 931–934 (2020).

31.   Isolation, quarantine, social distancing and community containment: pivotal role for

old-style public health measures in the novel coronavirus (2019-nCoV) outbreak |

Journal of Travel Medicine | Oxford Academic. Available at:

https://academic.oup.com/jtm/article/27/2/taaa020/5735321. (Accessed: 24th March

2020)

32.   Wu, Z. & McGoogan, J. M. Characteristics of and Important Lessons from the

Coronavirus Disease 2019 (COVID-19) Outbreak in China: Summary of a Report of

72314 Cases from the Chinese Center for Disease Control and Prevention. *JAMA - J.*

Am. Med. Assoc. (2020). doi:10.1001/jama.2020.2648