One size does not fit all. Genomics differentiates among binge-eating disorder,

bulimia nervosa, and anorexia nervosa

Christopher Hübel[1,2,3,4]*, Mohamed Abdulkadir[5,6]*, Moritz Herle[7], Ruth J.F. Loos[8],

Gerome Breen[1,2], Cynthia M. Bulik[4,9,10,11] & Nadia Micali[5,6]

[1]Social, Genetic & Developmental Psychiatry Centre, Institute of Psychiatry, Psychology

& Neuroscience, King's College London, UK

[2]UK National Institute for Health Research (NIHR) Biomedical Research Centre for

Mental Health, South London and Maudsley Hospital, London, UK

[3]National Centre for Register-based Research, Aarhus Business and Social Sciences,

Aarhus University, Aarhus, Denmark

[4]Department of Medical Epidemiology and Biostatistics, Karolinska Institutet,

Stockholm, Sweden

[5]Department of Pediatrics Gynaecology and Obstetrics, Faculty of Medicine, University

of Geneva, Geneva, Switzerland

[6]Department of Psychiatry, Faculty of Medicine, University of Geneva, Geneva,

Switzerland

[7]Department of Biostatistics & Health Informatics, Institute of Psychiatry, Psychology &

Neuroscience, King's College London, UK

[8]Icahn School of Medicine at Mount Sinai, New York, New York, USA

[9]Department of Psychiatry, University of North Carolina at Chapel Hill, Chapel Hill, NC,

USA

[10]Department of Nutrition, University of North Carolina at Chapel Hill, Chapel Hill, NC,

USA

[11]Great Ormond Street Institute of Child Health, University College London, London, UK

*contributed equally to this work

Correspondence to:

Dr. Micali, Department of Psychiatry, Faculty of Medicine, University of Geneva, Geneva,

Switzerland, e-mail: nadia.micali@hcuge.ch

Keywords: Avon Longitudinal Study of Parents and Children (ALSPAC); binge eating

disorder; polygenic scores; UK Biobank

Abstract (162/250, unstructured)

Objective. Genome-wide association studies have identified multiple genomic regions

associated with anorexia nervosa. Relatively few or no genome-wide studies of other

eating disorders, such as bulimia nervosa and binge-eating disorder, have been

performed, despite their substantial heritability. Exploratively, we aimed to identify

traits that are genetically associated with binge-type eating disorders.

Method. We calculated genome-wide polygenic scores for 269 trait and disease

outcomes using PRSice v2.2 and their association with anorexia nervosa, bulimia

nervosa, and binge-eating disorder in up to 640 cases and 17,050 controls from the UK

Biobank. Significant associations were tested for replication in the Avon Longitudinal

Study of Parents and Children (up to 217 cases and 3018 controls).

Results. Individuals with binge-type eating disorders had higher polygenic scores than

controls for other psychiatric disorders, including depression, schizophrenia, and

attention deficit hyperactivity disorder, and higher polygenic scores for body mass

index.

Discussion. Our findings replicate some of the known comorbidities of eating disorders

on a genomic level and motivate a deeper investigation of shared and unique genomic

factors across the three primary eating disorders.

Introduction

Eating disorders are complex and heritable psychiatric conditions (Yilmaz et al., 2015). The most studied is anorexia nervosa, which is characterised by dangerously low body weight and extreme fear of gaining weight, while engaging in behaviours that induce negative energy balance, such as fasting or, in some cases, excessive exercise (Treasure et al., 2015). In contrast, individuals with bulimia nervosa and binge-eating disorder experience episodes of excessive overeating (i.e., binge eating) accompanied by a sense of loss of control, in which they consume unusually large amounts of food in a short period of time. Individuals with bulimia nervosa also engage in recurrent compensatory behaviors (e.g., self-induced vomiting, laxative, or diuretic abuse) to counteract the effects of binge eating (American Psychiatric Association, 2013).

Twin, family, and adoption studies over the past 30 years have shown that eating disorders are heritable (Yilmaz et al., 2015). The largest genome-wide association study (GWAS) to date identified eight genomic regions associated with anorexia nervosa and also found genetic correlations with obsessive-compulsive disorder, schizophrenia, anxiety, and major depressive disorder. This implies that anorexia nervosa shares genetic risk variants with these phenotypes. However, the study also found that anorexia nervosa has significant negative genetic correlations with anthropometric traits, including BMI, fat mass, and fat-free mass. This means that anorexia nervosa risk variants are enriched for variants associated with lower BMI, fat mass, and fat-free mass. (Duncan et al., 2017; Hübel et al., 2019; Watson et al., 2019). The GWAS additionally highlighted a shared genetic basis between anorexia nervosa and high-density lipoprotein cholesterol, fasting insulin, as well as insulin sensitivity, suggesting that anorexia nervosa may have a metabolic component. It is unclear, however, if binge-type eating disorders may also have a metabolic component.

Although twin and family studies demonstrate a positive genetic correlation

(~0.46) between anorexia nervosa and bulimia nervosa, indicating considerable shared

genetics (Bulik et al., 2010; Yao, Larsson, et al., 2019), GWASs of bulimia nervosa or

binge-eating disorder with sufficient size have yet to be conducted to allow well

powered investigations of their molecular genetic correlations with each other. On a

phenotypic level, binge-eating disorder co-occurs with anxiety, depression, as well as

attention deficit hyperactivity disorder, substance abuse, and personality disorders. In

addition, individuals with binge-eating disorder commonly are overweight or obese

(Hilbert, 2019). We have previously suggested that increased genetic liability for

obesity was associated with increased risk of binge eating in adolescence (Abdulkadir et

al., 2020). In contrast, it is not well understood the extent to which binge-type eating

disorders share genetics with other psychiatric, anthropometric, and metabolic traits.

To explore differences in the genetics of eating disorders and generate new

hypotheses regarding their genetic overlap with other traits, we conducted polygenic

score analyses to identify traits that are genetically associated with either of the three

primary eating disorders: anorexia nervosa, bulimia nervosa, or binge-eating disorder,

in a subsample of the UK Biobank (Sudlow et al., 2015) with attempted replication in

the Avon Longitudinal Study of Parents and Children (ALSPAC) cohort.

Methods

Discovery sample of the UK Biobank

The UK Biobank (ukbiobank.ac.uk) is a unique epidemiological resource to

improve prevention, diagnosis, and treatment of psychiatric and somatic illnesses. The

UK Biobank recruited participants from the general population between 2006–2010. All

participants were between 40 to 69 years old, were registered with a general

practitioner through the United Kingdom's National Health Service, and lived within

traveling distance of one of the assessment centers. The UK Biobank is approved by the

North West Multi-centre Research Ethics Committee. Genomewide array data for this

study were available for 488,363 individuals. All participants gave written consent. We

identified non-European participants by 4-means clustering on the first two principal

components derived from the genotype data, and excluded related individuals (KING

relatedness metric >0.088, equivalent to a relatedness value of 0.25; N = 7,765).

We identified 1,488 participants who either self-reported any of the three

primary eating disorders in the mental health questionnaire (Davis et al., 2020) or had

an International Classification of Diseases, version 10 (ICD-10) (World Health

Organization, 1992) hospital diagnosis of F50.0 or F50.1 for anorexia nervosa, or F50.2

or F50.3 for bulimia nervosa. This resulted in 768 (4.7%) participants with anorexia

nervosa, 423 (2.7%) with bulimia nervosa, and 561 (3.5%) with binge-eating disorder.

If participants self-reported more than one diagnosis, they were assigned to all groups.

We randomly sampled one set of controls in the ratio of 1 to 10, resulting in 15,500

controls by using the following exclusion criteria: Controls must have answered the

mental health questionnaire, were not diagnosed with a psychiatric disorder (i.e., self-

report or ICD-10 diagnosis), or taking any psychotropic medication. The final analysis

included 17,050 (92% female) European participants (**Supplementary Table 1**)

representing 3.4% of the genotyped UK Biobank participants (n = 502,682). This study

was completed under the UK Biobank approved study application 27546.

Sample from the Avon Longitudinal Study of Parents and Children (ALSPAC)

The ALSPAC is a population-based sample of pregnant women and their children based

in the former county of Avon, UK (Boyd et al., 2013; Fraser et al., 2013; Golding et al.,

2001). Women expected to deliver from April 1, 1991, until December 31, 1992, were

invited to participate. Children from 14,541 pregnancies were enrolled and 13,988 alive

at 1 year. Additionally, 913 children were enrolled at age 7 years. All women gave

written informed consent. Children at age 14 (wave 14, n = 10,581), 16 (wave 16, n =

9,702), and 18 years (wave 18, n = 9,505) that had not withdrawn consent were

followed up. Parents answered questionnaires on 7,025 adolescents at wave 14 and on

5,656 at wave 16. With 6,140 (58%) responding at wave 14, 5,069 (52%) at wave 16,

and 3,228 (34%) at wave 18 which was used to augment the validity of the probable

eating disorder diagnoses (Micali et al., 2015). The study website contains details of all

the data that is available through a fully searchable data dictionary and variable search

tool (www.bristol.ac.uk/alspac/researchers/our-data/). To reduce potential

confounding through genetic relatedness in our analyses, we removed randomly one

individual of each pair that is closely related (φ > 0.2) using PLINK v1.90 (Chang et al.,

2015), excluding 75 individuals that were duplicates, monozygotic twins, first-degree

relatives (i.e., parent-offspring and full siblings), or second-degree relatives (i.e., half-

siblings, uncles, aunts, grandparents, and double cousins, **Supplementary Table 2**).

We derived probable eating disorder diagnoses as previously reported

(**Supplementary Table 3**) (Micali et al., 2015), using a combination of adolescent self-

report as well as information on the adolescents provided by their parents, a gold

standard for childhood psychiatric disorders.

Ethical approval for the ALSPAC participants of this study was obtained from the

ALSPAC Ethics and Law Committee and the Local Research Ethics Committees:

www.bristol.ac.uk/alspac/researchers/research-ethics/. Consent for biological samples

has been collected in accordance with the Human Tissue Act (2004) and informed

consent for the use of data collected via questionnaires and clinics was obtained from

participants following the recommendations of the ALSPAC Ethics and Law Committee

at the time.

Polygenic risk scoring on eating disorder phenotypes in UK Biobank and

replication in ALSPAC

We used PRSice (Choi & O'Reilly, 2019), version 2.2.3. We clumped the single

nucleotide polymorphisms (SNPs) that were present both in the summary statistics of

the trait and in the genotype data of the UK Biobank (i.e., overlapping SNPs) to obtain

genetically independent SNPs. We retained the SNP with the smallest p value in each

250 kilobase window of all those in linkage disequilibrium ($r^2 > 0.1$). We calculated 269

polygenic scores at their optimal p value threshold in each individual by scoring the

number of effect alleles (weighted by the allele effect size) across the set of remaining

SNPs (for a full list, see **Supplementary Table 4**). We calculated the polygenic score

using the high-resolution scoring (i.e., incrementally across a large number of p value

thresholds) method to identify the p value threshold at which the polygenic score is

optimally associated with the outcome and explains the most variance (i.e., resulting in

the highest adjusted R^2 for continuous outcomes and Nagelkerke's R^2 on the liability

scale for binary outcomes). We evaluated the associations between polygenic score and

eating disorder diagnosis using logistic regressions, adjusted for sex and the first six

principal components that were calculated on the European subsample. To adjust for

overfitting, we permuted case-control status at every p value threshold 10,000 times

and, hence, calculated empirical p values. We converted the observed R^2 to the liability

scale assuming following population prevalences of 3% for anorexia nervosa, 1.8% for

bulimia nervosa, and 3% for binge-eating disorder (Micali et al., 2017; Smink et al.,

2014).

To correct for multiple testing (i.e., 269 polygenic score regression models), we

calculated Q values using the false discovery rate approach (Benjamini & Hochberg,

1995; Benjamini & Yekutieli, 2001). We did not stratify analyses by sex because of the

low number of male eating disorder cases, but included sex as a covariate. We used the

UK Biobank sample as our discovery cohort and the SNPs that were included in the

strongest associated polygenic score were used to derive polygenic scores in the

ALSPAC sample, in which we repeated the analysis also adjusting for sex, and the first

six ancestry-informative principal components.

Results

Descriptive statistics of the analyses samples included from UK Biobank and ALSPAC

are listed in Supplementary Tables 1 and 2.

Polygenic scoring on eating disorder phenotypes in UK Biobank and ALSPAC

After correcting for multiple testing using the false discovery rate adjustment, 18

polygenic scores were significantly associated with the three primary eating disorders

in the subsample of the UK Biobank (**Figure 1**).

196   **Psychiatric disorders and neurological disease.** Five polygenic scores

reflecting polygenic liability for psychiatric disorders and one neurological disease

showed significant associations with eating disorders (**Supplementary Table 4 & 5**).

The schizophrenia polygenic score was positively associated with anorexia nervosa

(odds ratio [OR] per standard deviation in polygenic score = 1.18, 95% confidence

interval [CI]: 1.09, 1.27; Q = 0.03) and binge-eating disorder (OR = 1.23, 95% CI: 1.13,

1.35; Q = 0.01). The major depressive disorder polygenic score was positively

associated with anorexia nervosa (OR = 1.20, 95% CI: 1.11, 1.29; Q = 0.01) and binge-

eating disorder (OR = 1.25, 95% CI: 1.14, 1.36; Q = 0.01). Additionally, the attention

deficit hyperactivity disorder polygenic score was positively associated with binge-

eating disorder (OR = 1.28, 95% CI: 1.17, 1.39; Q = 0.01). The bipolar disorder polygenic

score (OR = 1.17, 95% CI: 1.08, 1.26; Q = 0.04), the anorexia nervosa polygenic score

(OR = 1.32, 95% CI: 1.22, 1.42; Q = 0.01), and the migraine polygenic score (OR = 1.17,

95% CI: 1.09, 1.26; Q = 0.04) were positively associated with anorexia nervosa.

210   **Somatic traits.** Several polygenic scores indexing polygenic load for

anthropometric traits, including waist circumference (OR = 1.43, 95% CI: 1.31, 1.56; Q =

0.01), hip circumference (OR = 1.39, 95% CI: 1.27, 1.51; Q = 0.01), overweight (OR =

1.35, 95% CI: 1.24, 1.47; Q = 0.01), obesity class 1 (OR = 1.38, 95% CI: 1.27; 1.51; Q =

0.01), class 2 (OR = 1.33, 95% CI: 1.22; 1.45; Q = 0.01) and class 3 (OR = 1.21, 95% CI:

1.11, 1.32; Q = 0.04), BMI (OR = 1.44, 95% CI: 1.32; 1.56; Q = 0.01) and extreme BMI (OR

= 1.32, 95% CI: 1.22, 1.44; Q = 0.01) were positively associated with binge-eating

disorder. However, it is important to note that these polygenic scores were correlated

with each other with r = .10 to r = .67 (**Supplementary Figure 1**). Additionally, the

childhood obesity polygenic score (OR = 1.22, 95% CI: 1.12, 1.33; Q = 0.04) was

positively associated with binge-eating disorder, while the age at menarche polygenic

score (OR = 0.79, 95% CI: 0.72, 0.86; Q = 0.01) was negatively associated with binge-

eating disorder. Neither the childhood obesity nor the age at menarche polygenic score

were associated with anorexia nervosa.

The overweight polygenic score (OR = 1.26, 95% CI: 1.14, 1.38; Q = 0.01) and the

waist circumference polygenic score (OR = 1.25, 95% CI: 1.13, 1.38; Q = 0.02) were also

positively associated with bulimia nervosa.

Sex and age differences. The adult obesity polygenic score was more strongly

associated with anorexia nervosa than the childhood obesity polygenic score. The

difference in the log odds ratios was statistically significant (difference = -0.15, p =

0.004). We did not detect age differences in the associations of the obesity polygenic

scores with bulimia nervosa or binge-eating disorder (**Supplementary Table 6**). We

found no evidence for sex differences in the associations of the body mass index

polygenic score with anorexia nervosa, bulimia nervosa, or binge-eating disorder.

Sensitivity analyses. We compared logistic regression results when assigning

participants to all possible eating disorder diagnoses with assigning participants

exclusively to one eating disorder diagnosis in the UK Biobank sample. The effect sizes

of both analyses correlated with r = 0.98, indicating that the analyses were not sensitive

to participant overlap (**Supplementary Figure 2**).

Independent replication. In ALSPAC (**Supplementary Table 7**), the

overweight polygenic score (OR = 1.38, 95% CI: 1.13, 1.69; $p_{empirical}$ = 0.05) and the

obesity class 1 polygenic score (OR = 1.23, 95% CI: 1.00; 1.50; $p_{empirical}$ = 0.001) were

positively associated with binge-eating disorder. Correlations between these polygenic

scores are presented in **Supplementary Figure 3**.

Discussion

Here, we present the first molecular genetic evidence that the underlying biology

differs between binge-type eating disorders and anorexia nervosa and these differences

can be captured at the genomic level. We find that polygenic scores for psychiatric

disorders and anthropometric traits are associated with binge-type eating disorders

(see **Figure 1**). Our UK Biobank analyses propose strong positive associations between

bulimia nervosa/binge-eating disorder and anthropometric polygenic scores (e.g.,

overweight and waist circumference), suggesting that binge eating shares genomic

variants with overweight and obesity. In contrast, for anorexia nervosa, the direction of

these associations was reversed. Binge-eating disorder also showed an association with

the childhood obesity polygenic score while anorexia nervosa did not, further

highlighting the genomic differences between binge-type eating disorders and anorexia

nervosa. The genetic relationship between binge eating and BMI had been studied

previously (Abdulkadir et al., 2020; Bulik et al., 2003) and is replicated by our analyses.

It is notable that previous work suggests that genomic variants associated with BMI are

predominantly expressed in brain tissue (Finucane et al., 2018). Our results also suggest

that these BMI-associated genomic variants are relevant for eating disorders, but may

act in opposite directions in binge-type eating disorders and anorexia nervosa.

Using the UK Biobank data, we also found that psychiatric polygenic scores (e.g.,

schizophrenia and major depressive disorder) were positively associated with both

anorexia nervosa and binge-eating disorder (**Figure 1, Panel A**). The ADHD polygenic

score was only significantly associated with binge-eating disorder, while the anorexia

polygenic score was not associated with bulimia nervosa or binge-eating disorder.

Our findings are in agreement with previous studies that showed genetic

correlations between binge eating and depression (Munn-Chernoff et al., 2015) as well

as genetic correlations between anorexia nervosa and depression (Wade et al., 2000;

Watson et al., 2019), migraine (Mustelin et al., 2014), schizophrenia (Duncan et al.,

2017; Watson et al., 2019), bipolar disorder (Stahl et al., 2019) and anthropometric

traits (Duncan et al., 2017; Watson et al., 2019). In line with our findings, previous

GWASs showed no genomic association between anorexia nervosa and ADHD, whereas

an ADHD polygenic score was associated with binge eating in a Swedish twin sample

(Capusan et al., 2017; Yao, Kuja-Halkola, et al., 2019). Moreover, a large evidence-base

highlights phenotypic overlap between ADHD and binge eating (Cortese et al., 2016;

Leventakou et al., 2016; Sonneville et al., 2015).

Large GWASs have recently suggested the involvement of a metabolic component

in anorexia nervosa through genetic correlations (Duncan et al., 2017; Watson et al.,

2019); however, at the current sample size, we did not observe associations between

metabolic polygenic scores and the three primary eating disorders in the UK Biobank

most probably due to limited statistical power (**Figure 1, Panel B**).

The association between polygenic liability for higher body mass and eating

disorders is further complicated as it shows age dependence (Hübel et al., 2019) and

confounding through age at menarche. In both the UK Biobank and the ALSPAC cohort,

we found that age at menarche and BMI polygenic scores were negatively correlated

with each other. However, the age at menarche polygenic score was only negatively

associated with binge-eating disorder but not with anorexia nervosa (**Figure 1**, **Panel**

C). Phenotypic associations between early menarche and bulimia nervosa have

previously been reported (Fairburn et al., 1999; Reichborn-Kjennerud et al., 2004), but

do not always replicate (Algars et al., 2014). On a genetic level, binge eating correlated

negatively with age at menarche in a twin study (Baker et al., 2012) and anorexia

nervosa was not genetically correlated with age at menarche in our polygenic score

analysis nor in previous GWASs (Duncan et al., 2017; Watson et al., 2019). This suggests

that the association between age at menarche and eating disorders may be driven by

symptoms related to binge eating rather than restriction.

Furthermore, ADHD and BMI show a positive genetic correlation (Demontis et

al., 2019; Hübel et al., 2019) and in our analysis the ADHD polygenic score was

associated with binge-eating disorder. Clinically, the ADHD medication

lisdexamfetamine is used to treat binge-eating disorder (Hudson et al., 2017). This

combined evidence suggests that the shared biology between ADHD and BMI may

contribute to binge eating. These novel findings warrant further research; investigating

how genomic variants associated with ADHD, commonly characterized by inattention

and impulsivity, are also associated with binge-eating disorder and contribute to the

high comorbidity between ADHD and obesity (Cortese et al., 2016; Leventakou et al.,

2016; Sonneville et al., 2015).

Our findings must be interpreted in the light of the following limitations: the

number of individuals with eating disorders in both samples was relatively low, some of

the eating disorder diagnoses and symptoms were self-reported, and the samples only

included white British participants. Furthermore, we were unable to differentiate

between binge-eating/purging and restricting anorexia nervosa as subtype information

was not available. However, symptoms and derived diagnoses in the ALSPAC cohort

were obtained both from parents and adolescents, representing the gold standard in

child and adolescent psychiatry, strengthening the validity of the diagnoses. Further,

replication studies are essential to lend confidence to the results reported here.

323   Our findings show for the first time that similarities exist in the genomic

324 psychiatric liability for binge-type eating disorders and anorexia nervosa. However, we

325 find clear differences between binge-type eating disorders and anorexia nervosa in the

326 underlying biology of body mass regulation at the genomic level. These findings open

327 important avenues for translational research relevant to eating disorder phenotypes

328 and overlap between eating disorders. Specifically, it remains unknown which biological

329 mechanisms are shared between ADHD and binge eating.

Acknowledgements

We are deeply grateful to all the families who took part in this study, the midwives for

their help in recruiting them, and the whole ALSPAC team, which includes interviewers,

computer and laboratory technicians, clerical workers, research scientists, volunteers,

managers, receptionists and nurses. This study was completed as part of approved UK

Biobank study applications 27546 to Prof Breen.

Funding

This study represents independent research part funded by the UK National Institute

for Health Research (NIHR) Biomedical Research Centre at South London and Maudsley

NHS Foundation Trust and King's College London. The views expressed are those of the

author(s) and not necessarily those of the UK NHS, the NIHR or the Department of

Health. High performance computing facilities were funded with capital equipment

grants from the GSTT Charity (TR130505) and Maudsley Charity (980). This work was

supported by the UK Medical Research Council and the Medical Research Foundation

(ref: MR/R004803/1). The UK Medical Research Council and Wellcome (Grant ref:

102215/2/13/2 and 217065/Z/19/Z) and the University of Bristol provide core

support for ALSPAC. A comprehensive list of grants funding is available on the ALSPAC

website (http://www.bristol.ac.uk/alspac/external/documents/grant-

acknowledgements.pdf); This research was specifically funded by the NIHR

(CS/01/2008/014), the NIH (MH087786-01). GWAS data was generated by Sample

Logistics and Genotyping Facilities at Wellcome Sanger Institute and LabCorp

(Laboratory Corporation of America) using support from 23andMe. NM and CB

acknowledge funding from the National Institute of Mental Health (R21 MH115397). CB

acknowledges funding from the Swedish Research Council (VR Dnr: 538-2013-8864),

the National Institute of Mental Health (R21MH115397; R01 MH109528;

R01MH120170; R01MH119084; R01MH120170), and Lundbeckfonden. MH is

supported by fellowship from the Medical Research Council UK (MR/T027843/1). The

content is solely the responsibility of the authors and does not necessarily represent the

official views of the National Institutes of Health. The funders were not involved in the

design or conduct of the study; collection, management, analysis, or interpretation of

the data; or preparation, review, or approval of the manuscript.

Availability of data and materials

366   This study is based on data from the UK Biobank. Researchers can apply for the data via

367   the following website: https://www.ukbiobank.ac.uk/. This study is based on data

368   from the ALSPAC study (http://www.bristol.ac.uk/alspac/). Interested researchers can

369   apply for data access with the University of Bristol, UK. Analysis scripts can be

370   requested from the authors.

Declaration of interest

Dr. Breen has received grant funding from and served as a consultant to Eli Lilly, has

received honoraria from Illumina and has served on advisory boards for Otsuka. Dr.

Bulik is a grant recipient from and has served on advisory boards for Shire and is a

consultant for Idorsia. She receives royalties from Pearson. She is a grant recipient from

Lundbeckfonden. All other authors have indicated they have no conflicts of interest to

disclose.

Author Contribution

References

Abdulkadir, M., Herle, M., De Stavola, B. L., Hübel, C., Santos Ferreira, D. L., Loos, R. J. F., Bryant-

Waugh, R., Bulik, C. M., & Micali, N. (2020). Polygenic Score for Body Mass Index Is

Associated with Disordered Eating in a General Population Cohort. *Journal of Clinical*

Medicine Research, *9*(4). https://doi.org/10.3390/jcm9041187

Algars, M., Huang, L., Von Holle, A. F., Peat, C. M., Thornton, L. M., Lichtenstein, P., & Bulik, C. M.

(2014). Binge eating and menstrual dysfunction. *Journal of Psychosomatic Research*, *76*(1),

19–22.

American Psychiatric Association. (2013). *Diagnostic and Statistical Manual of Mental Disorders*

(5th ed.). American Psychiatric Association.

Baker, J. H., Thornton, L. M., Bulik, C. M., Kendler, K. S., & Lichtenstein, P. (2012). Shared genetic

effects between age at menarche and disordered eating. *The Journal of Adolescent Health:*

Official Publication of the Society for Adolescent Medicine, *51*(5), 491–496.

Benjamini, Y., & Hochberg, Y. (1995). Controlling the False Discovery Rate: A Practical and

Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society. Series B,*

Statistical Methodology, *57*(1), 289–300.

Benjamini, Y., & Yekutieli, D. (2001). The control of the false discovery rate in multiple testing

under dependency. *Annals of Statistics*, *29*(4), 1165–1188.

Boyd, A., Golding, J., Macleod, J., Lawlor, D. A., Fraser, A., Henderson, J., Molloy, L., Ness, A., Ring,

S., & Davey Smith, G. (2013). Cohort Profile: the "children of the 90s"--the index offspring of

the Avon Longitudinal Study of Parents and Children. *International Journal of Epidemiology*,

42(1), 111–127.

Bulik, C. M., Sullivan, P. F., & Kendler, K. S. (2003). Genetic and environmental contributions to

obesity and binge eating. *The International Journal of Eating Disorders*, *33*(3), 293–298.

Bulik, C. M., Thornton, L. M., Root, T. L., Pisetsky, E. M., Lichtenstein, P., & Pedersen, N. L. (2010).

Understanding the relation between anorexia nervosa and bulimia nervosa in a Swedish

national twin sample. *Biological Psychiatry*, *67*(1), 71–77.

Capusan, A. J., Yao, S., Kuja-Halkola, R., Bulik, C. M., Thornton, L. M., Bendtsen, P., Marteinsdottir,

I., Thorsell, A., & Larsson, H. (2017). Genetic and environmental aspects in the association

between attention-deficit hyperactivity disorder symptoms and binge-eating behavior in

adults: a twin study. *Psychological Medicine*, *47*(16), 2866–2878.

Chang, C. C., Chow, C. C., Tellier, L. C., Vattikuti, S., Purcell, S. M., & Lee, J. J. (2015). Second-

generation PLINK: rising to the challenge of larger and richer datasets. *GigaScience*, *4*, 7.

Choi, S. W., & O'Reilly, P. F. (2019). PRSice-2: Polygenic Risk Score software for biobank-scale

data. *GigaScience*, *8*(7). https://doi.org/10.1093/gigascience/giz082

Cortese, S., Moreira-Maia, C. R., St Fleur, D., Morcillo-Peñalver, C., Rohde, L. A., & Faraone, S. V.

(2016). Association Between ADHD and Obesity: A Systematic Review and Meta-Analysis.

The American Journal of Psychiatry, *173*(1), 34–43.

Davis, K. A. S., Coleman, J. R. I., Adams, M., Allen, N., Breen, G., Cullen, B., Dickens, C., Fox, E.,

Graham, N., Holliday, J., Howard, L. M., John, A., Lee, W., McCabe, R., McIntosh, A., Pearsall, R.,

Smith, D. J., Sudlow, C., Ward, J., … Hotopf, M. (2020). Mental health in UK Biobank -

development, implementation and results from an online questionnaire completed by 157

366 participants: a reanalysis. *BJPsych Open*, *6*(2), e18.

Demontis, D., Walters, R. K., Martin, J., Mattheisen, M., Als, T. D., Agerbo, E., Baldursson, G.,

Belliveau, R., Bybjerg-Grauholm, J., Bækvad-Hansen, M., Cerrato, F., Chambert, K.,

Churchhouse, C., Dumont, A., Eriksson, N., Gandal, M., Goldstein, J. I., Grasby, K. L., Grove, J.,

… Neale, B. M. (2019). Discovery of the first genome-wide significant risk loci for attention

deficit/hyperactivity disorder. *Nature Genetics*, *51*(1), 63–75.

Duncan, L., Yilmaz, Z., Gaspar, H., Walters, R., Goldstein, J., Anttila, V., Bulik-Sullivan, B., Ripke, S.,

Eating Disorders Working Group of the Psychiatric Genomics Consortium, Thornton, L.,

Hinney, A., Daly, M., Sullivan, P. F., Zeggini, E., Breen, G., & Bulik, C. M. (2017). Significant

Locus and Metabolic Genetic Correlations Revealed in Genome-Wide Association Study of

Anorexia Nervosa. *The American Journal of Psychiatry*, *174*(9), 850–858.

Fairburn, C. G., Cooper, Z., Doll, H. A., & Welch, S. L. (1999). Risk factors for anorexia nervosa:

three integrated case-control comparisons. *Archives of General Psychiatry*, *56*(5), 468–476.

Finucane, H. K., Reshef, Y. A., Anttila, V., Slowikowski, K., Gusev, A., Byrnes, A., Gazal, S., Loh, P.-R.,

Lareau, C., Shoresh, N., Genovese, G., Saunders, A., Macosko, E., Pollack, S., Brainstorm

Consortium, Perry, J. R. B., Buenrostro, J. D., Bernstein, B. E., Raychaudhuri, S., ... Price, A. L.

(2018). Heritability enrichment of specifically expressed genes identifies disease-relevant

tissues and cell types. *Nature Genetics*, *50*(4), 621–629.

Fraser, A., Macdonald-Wallis, C., Tilling, K., Boyd, A., Golding, J., Davey Smith, G., Henderson, J.,

Macleod, J., Molloy, L., Ness, A., Ring, S., Nelson, S. M., & Lawlor, D. A. (2013). Cohort Profile:

the Avon Longitudinal Study of Parents and Children: ALSPAC mothers cohort.

International Journal of Epidemiology, *42*(1), 97–110.

Golding, J., Pembrey, M., Jones, R., & ALSPAC Study Team. (2001). ALSPAC--the Avon

Longitudinal Study of Parents and Children. I. Study methodology. *Paediatric and Perinatal*

Epidemiology, *15*(1), 74–87.

Hübel, C., Gaspar, H. A., Coleman, J. R. I., Hanscombe, K. B., Purves, K., Prokopenko, I., Graff, M.,

Ngwa, J. S., Workalemahu, T., ADHD Working Group of the Psychiatric Genomics

Consortium, Meta-Analyses of Glucose and Insulin-related traits consortium (MAGIC),

Autism Working Group of the Psychiatric Genomics Consortium, Bipolar Disorder Working

Group of the Psychiatric Genomics Consortium, Eating Disorders Working Group of the

Psychiatric Genomics Consortium, Major Depressive Disorder Working Group of the

Psychiatric Genomics Consortium, OCD & Tourette Syndrome Working Group of the

Psychiatric Genomics Consortium, PTSD Working Group of the Psychiatric Genomics

Consortium, Schizophrenia Working Group of the Psychiatric Genomics Consortium, Sex

Differences Cross Disorder Working Group of the Psychiatric Genomics Consortium, ...

Breen. (2019). Genetic correlations of psychiatric traits with body composition and

glycemic traits are sex- and age-dependent. *Nature Communications*, *10*(1), 5765.

Leventakou, V., Micali, N., Georgiou, V., Sarri, K., Koutra, K., Koinaki, S., Vassilaki, M., Kogevinas,

M., & Chatzi, L. (2016). Is there an association between eating behaviour and attention-

deficit/hyperactivity disorder symptoms in preschool children? *Journal of Child Psychology*

and Psychiatry, and Allied Disciplines, *57*(6), 676–684.

Micali, N., Martini, M. G., Thomas, J. J., Eddy, K. T., Kothari, R., Russell, E., Bulik, C. M., & Treasure,

J. (2017). Lifetime and 12-month prevalence of eating disorders amongst women in mid-

life: a population-based study of diagnoses and risk factors. *BMC Medicine*, *15*(1), 12.

Micali, N., Solmi, F., Horton, N. J., Crosby, R. D., Eddy, K. T., Calzo, J. P., Sonneville, K. R., Swanson,

S. A., & Field, A. E. (2015). Adolescent Eating Disorders Predict Psychiatric, High-Risk

Behaviors and Weight Outcomes in Young Adulthood. *Journal of the American Academy of*

Child and Adolescent Psychiatry, *54*(8), 652–659.e1.

Munn-Chernoff, M. A., Grant, J. D., Agrawal, A., Koren, R., Glowinski, A. L., Bucholz, K. K., Madden,

P. A. F., Heath, A. C., & Duncan, A. E. (2015). Are there common familial influences for major

depressive disorder and an overeating-binge eating dimension in both European American

and African American female twins? *The International Journal of Eating Disorders*, *48*(4),

375–382.

Mustelin, L., Raevuori, A., Kaprio, J., & Keski-Rahkonen, A. (2014). Association between eating

disorders and migraine may be explained by major depression. *The International Journal of*

Eating Disorders, *47*(8), 884–887.

Reichborn-Kjennerud, T., Bulik, C. M., Sullivan, P. F., Tambs, K., & Harris, J. R. (2004). Psychiatric

and medical symptoms in binge eating in the absence of compensatory behaviors. *Obesity*

Research, *12*(9), 1445–1454.

Smink, F. R. E., Hoeken, D., Oldehinkel, A. J., & Hoek, H. W. (2014). Prevalence and severity of

DSM-5 eating disorders in a community cohort of adolescents. *The International Journal of*

Eating Disorders, *47*(6), 610–619.

Sonneville, K. R., Calzo, J. P., Horton, N. J., Field, A. E., Crosby, R. D., Solmi, F., & Micali, N. (2015).

Childhood hyperactivity/inattention and eating disturbances predict binge eating in

adolescence. *Psychological Medicine*, *45*(12), 2511–2520.

Stahl, E. A., Breen, G., Forstner, A. J., McQuillin, A., Ripke, S., Trubetskoy, V., Mattheisen, M., Wang,

Y., Coleman, J. R. I., Gaspar, H. A., de Leeuw, C. A., Steinberg, S., Pavlides, J. M. W.,

Trzaskowski, M., Byrne, E. M., Pers, T. H., Holmans, P. A., Richards, A. L., Abbott, L., … Bipolar

Disorder Working Group of the Psychiatric Genomics Consortium. (2019). Genome-wide

association study identifies 30 loci associated with bipolar disorder. *Nature Genetics*, *51*(5),

793–803.

Sudlow, C., Gallacher, J., Allen, N., Beral, V., Burton, P., Danesh, J., Downey, P., Elliott, P., Green, J.,

Landray, M., Liu, B., Matthews, P., Ong, G., Pell, J., Silman, A., Young, A., Sprosen, T., Peakman,

T., & Collins, R. (2015). UK biobank: an open access resource for identifying the causes of a

wide range of complex diseases of middle and old age. *PLoS Medicine*, *12*(3), e1001779.

Treasure, J., Zipfel, S., Micali, N., Wade, T., Stice, E., Claudino, A., Schmidt, U., Frank, G. K., Bulik, C.

M., & Wentz, E. (2015). Anorexia nervosa. *Nature Reviews. Disease Primers*, *1*, 15074.

Wade, T. D., Bulik, C. M., Neale, M., & Kendler, K. S. (2000). Anorexia nervosa and major

depression: shared genetic and environmental risk factors. *The American Journal of*

Psychiatry, *157*(3), 469–471.

Watson, H. J., Yilmaz, Z., Thornton, L. M., Hübel, C., Coleman, J. R. I., Gaspar, H. A., Bryois, J.,

Hinney, A., Leppä, V. M., Mattheisen, M., Medland, S. E., Ripke, S., Yao, S., Giusti-Rodríguez, P.,

Anorexia Nervosa Genetics Initiative, Hanscombe, K. B., Purves, K. L., Eating Disorders

Working Group of the Psychiatric Genomics Consortium, Adan, R. A. H., … Bulik, C. M.

(2019). Genome-wide association study identifies eight risk loci and implicates metabo-

psychiatric origins for anorexia nervosa. *Nature Genetics*, *51*(8), 1207–1214.

World Health Organization. (1992). *ICD-10: International Statistical Classification of Diseases and*

Related Health Problems: 10th revision. World Health Organization.

Yao, S., Kuja-Halkola, R., Martin, J., Lu, Y., Lichtenstein, P., Norring, C., Birgegård, A., Yilmaz, Z.,

Hübel, C., Watson, H., Baker, J., Almqvist, C., Eating Disorders Working Group of the

Psychiatric Genomics Consortium, Thornton, L. M., Magnusson, P. K., Bulik, C. M., & Larsson,

H. (2019). Associations Between Attention-Deficit/Hyperactivity Disorder and Various

Eating Disorders: A Swedish Nationwide Population Study Using Multiple Genetically

Informative Approaches. *Biological Psychiatry*, *86*(8), 577–586.

Yao, S., Larsson, H., Norring, C., Birgegård, A., Lichtenstein, P., D☐Onofrio, B. M., Almqvist, C.,

Thornton, L. M., Bulik, C. M., & Kuja-Halkola, R. (2019). Genetic and environmental

contributions to diagnostic fluctuation in anorexia nervosa and bulimia nervosa.

Psychological Medicine, 1–8.

Yilmaz, Z., Hardaway, J. A., & Bulik, C. M. (2015). Genetics and Epigenetics of Eating Disorders.

Advances in Genomics and Genetics, *5*, 131–150.

Figure 1. Polygenic scores associated with eating disorders in the UK Biobank.

Panel A *shows psychiatric and behavioral,* *Panel B* *metabolic, and* *Panel C*

anthropometric polygenic scores and their association estimates with self-reported or

hospital-diagnosed eating disorders in the UK Biobank sample (n = 17,050). Filled dots are

statistically significant after adjustment for multiple testing through the false discovery

approach. Dots represent odds ratios (ORs) and error bars index 95% confidence intervals

obtained via logistic regression and 10,000 permutations to obtain empirical p values.