

1 **One size does not fit all. Genomics differentiates among binge-eating disorder,**  
2 **bulimia nervosa, and anorexia nervosa**

3

4 Christopher Hübel<sup>1,2,3,4\*</sup>, Mohamed Abdulkadir<sup>5,6\*</sup>, Moritz Herle<sup>7</sup>, Ruth J.F. Loos<sup>8</sup>,  
5 Gerome Breen<sup>1,2</sup>, Cynthia M. Bulik<sup>4,9,10,11</sup> & Nadia Micali<sup>5,6</sup>

6

7 <sup>1</sup>Social, Genetic & Developmental Psychiatry Centre, Institute of Psychiatry, Psychology  
8 & Neuroscience, King's College London, UK

9 <sup>2</sup>UK National Institute for Health Research (NIHR) Biomedical Research Centre for  
10 Mental Health, South London and Maudsley Hospital, London, UK

11 <sup>3</sup>National Centre for Register-based Research, Aarhus Business and Social Sciences,  
12 Aarhus University, Aarhus, Denmark

13 <sup>4</sup>Department of Medical Epidemiology and Biostatistics, Karolinska Institutet,  
14 Stockholm, Sweden

15 <sup>5</sup>Department of Pediatrics Gynaecology and Obstetrics, Faculty of Medicine, University  
16 of Geneva, Geneva, Switzerland

17 <sup>6</sup>Department of Psychiatry, Faculty of Medicine, University of Geneva, Geneva,  
18 Switzerland

19 <sup>7</sup>Department of Biostatistics & Health Informatics, Institute of Psychiatry, Psychology &  
20 Neuroscience, King's College London, UK

21 <sup>8</sup>Icahn School of Medicine at Mount Sinai, New York, New York, USA

22 <sup>9</sup>Department of Psychiatry, University of North Carolina at Chapel Hill, Chapel Hill, NC,  
23 USA

24 <sup>10</sup>Department of Nutrition, University of North Carolina at Chapel Hill, Chapel Hill, NC,  
25 USA

26 <sup>11</sup>Great Ormond Street Institute of Child Health, University College London, London, UK

27

28 \*contributed equally to this work

29

30 Correspondence to:

31 Dr. Micali, Department of Psychiatry, Faculty of Medicine, University of Geneva, Geneva,

32 Switzerland, e-mail: [nadia.micali@hcuge.ch](mailto:nadia.micali@hcuge.ch)

33

34 Keywords: Avon Longitudinal Study of Parents and Children (ALSPAC); binge eating

35 disorder; polygenic scores; UK Biobank

36 **Abstract (162/250, unstructured)**

37 **Objective.** Genome-wide association studies have identified multiple genomic regions  
38 associated with anorexia nervosa. Relatively few or no genome-wide studies of other  
39 eating disorders, such as bulimia nervosa and binge-eating disorder, have been  
40 performed, despite their substantial heritability. Exploratively, we aimed to identify  
41 traits that are genetically associated with binge-type eating disorders.

42 **Method.** We calculated genome-wide polygenic scores for 269 trait and disease  
43 outcomes using PRSice v2.2 and their association with anorexia nervosa, bulimia  
44 nervosa, and binge-eating disorder in up to 640 cases and 17,050 controls from the UK  
45 Biobank. Significant associations were tested for replication in the Avon Longitudinal  
46 Study of Parents and Children (up to 217 cases and 3018 controls).

47 **Results.** Individuals with binge-type eating disorders had higher polygenic scores than  
48 controls for other psychiatric disorders, including depression, schizophrenia, and  
49 attention deficit hyperactivity disorder, and higher polygenic scores for body mass  
50 index.

51 **Discussion.** Our findings replicate some of the known comorbidities of eating disorders  
52 on a genomic level and motivate a deeper investigation of shared and unique genomic  
53 factors across the three primary eating disorders.

54

## 55 **Introduction**

56 Eating disorders are complex and heritable psychiatric conditions (Yilmaz et al.,  
57 2015). The most studied is anorexia nervosa, which is characterised by dangerously low  
58 body weight and extreme fear of gaining weight, while engaging in behaviours that  
59 induce negative energy balance, such as fasting or, in some cases, excessive exercise  
60 (Treasure et al., 2015). In contrast, individuals with bulimia nervosa and binge-eating  
61 disorder experience episodes of excessive overeating (i.e., binge eating) accompanied by  
62 a sense of loss of control, in which they consume unusually large amounts of food in a  
63 short period of time. Individuals with bulimia nervosa also engage in recurrent  
64 compensatory behaviors (e.g., self-induced vomiting, laxative, or diuretic abuse) to  
65 counteract the effects of binge eating (American Psychiatric Association, 2013).

66 Twin, family, and adoption studies over the past 30 years have shown that eating  
67 disorders are heritable (Yilmaz et al., 2015). The largest genome-wide association study  
68 (GWAS) to date identified eight genomic regions associated with anorexia nervosa and  
69 also found genetic correlations with obsessive-compulsive disorder, schizophrenia,  
70 anxiety, and major depressive disorder. This implies that anorexia nervosa shares  
71 genetic risk variants with these phenotypes. However, the study also found that  
72 anorexia nervosa has significant negative genetic correlations with anthropometric  
73 traits, including BMI, fat mass, and fat-free mass. This means that anorexia nervosa risk  
74 variants are enriched for variants associated with lower BMI, fat mass, and fat-free  
75 mass. (Duncan et al., 2017; Hübel et al., 2019; Watson et al., 2019). The GWAS  
76 additionally highlighted a shared genetic basis between anorexia nervosa and high-  
77 density lipoprotein cholesterol, fasting insulin, as well as insulin sensitivity, suggesting  
78 that anorexia nervosa may have a metabolic component. It is unclear, however, if binge-  
79 type eating disorders may also have a metabolic component.

80           Although twin and family studies demonstrate a positive genetic correlation  
81   (~0.46) between anorexia nervosa and bulimia nervosa, indicating considerable shared  
82   genetics (Bulik et al., 2010; Yao, Larsson, et al., 2019), GWASs of bulimia nervosa or  
83   binge-eating disorder with sufficient size have yet to be conducted to allow well  
84   powered investigations of their molecular genetic correlations with each other. On a  
85   phenotypic level, binge-eating disorder co-occurs with anxiety, depression, as well as  
86   attention deficit hyperactivity disorder, substance abuse, and personality disorders. In  
87   addition, individuals with binge-eating disorder commonly are overweight or obese  
88   (Hilbert, 2019). We have previously suggested that increased genetic liability for  
89   obesity was associated with increased risk of binge eating in adolescence (Abdulkadir et  
90   al., 2020). In contrast, it is not well understood the extent to which binge-type eating  
91   disorders share genetics with other psychiatric, anthropometric, and metabolic traits.

92           To explore differences in the genetics of eating disorders and generate new  
93   hypotheses regarding their genetic overlap with other traits, we conducted polygenic  
94   score analyses to identify traits that are genetically associated with either of the three  
95   primary eating disorders: anorexia nervosa, bulimia nervosa, or binge-eating disorder,  
96   in a subsample of the UK Biobank (Sudlow et al., 2015) with attempted replication in  
97   the Avon Longitudinal Study of Parents and Children (ALSPAC) cohort.

98

## 99   **Methods**

### 100   **Discovery sample of the UK Biobank**

101           The UK Biobank ([ukbiobank.ac.uk](http://ukbiobank.ac.uk)) is a unique epidemiological resource to  
102   improve prevention, diagnosis, and treatment of psychiatric and somatic illnesses. The  
103   UK Biobank recruited participants from the general population between 2006–2010. All  
104   participants were between 40 to 69 years old, were registered with a general

105 practitioner through the United Kingdom's National Health Service, and lived within  
106 traveling distance of one of the assessment centers. The UK Biobank is approved by the  
107 North West Multi-centre Research Ethics Committee. Genomewide array data for this  
108 study were available for 488,363 individuals. All participants gave written consent. We  
109 identified non-European participants by 4-means clustering on the first two principal  
110 components derived from the genotype data, and excluded related individuals (KING  
111 relatedness metric  $>0.088$ , equivalent to a relatedness value of 0.25;  $N = 7,765$ ).

112 We identified 1,488 participants who either self-reported any of the three  
113 primary eating disorders in the mental health questionnaire (Davis et al., 2020) or had  
114 an International Classification of Diseases, version 10 (ICD-10) (World Health  
115 Organization, 1992) hospital diagnosis of F50.0 or F50.1 for anorexia nervosa, or F50.2  
116 or F50.3 for bulimia nervosa. This resulted in 768 (4.7%) participants with anorexia  
117 nervosa, 423 (2.7%) with bulimia nervosa, and 561 (3.5%) with binge-eating disorder.  
118 If participants self-reported more than one diagnosis, they were assigned to all groups.  
119 We randomly sampled one set of controls in the ratio of 1 to 10, resulting in 15,500  
120 controls by using the following exclusion criteria: Controls must have answered the  
121 mental health questionnaire, were not diagnosed with a psychiatric disorder (i.e., self-  
122 report or ICD-10 diagnosis), or taking any psychotropic medication. The final analysis  
123 included 17,050 (92% female) European participants (**Supplementary Table 1**)  
124 representing 3.4% of the genotyped UK Biobank participants ( $n = 502,682$ ). This study  
125 was completed under the UK Biobank approved study application 27546.

126

### 127 **Sample from the Avon Longitudinal Study of Parents and Children (ALSPAC)**

128 The ALSPAC is a population-based sample of pregnant women and their children based  
129 in the former county of Avon, UK (Boyd et al., 2013; Fraser et al., 2013; Golding et al.,

130 2001). Women expected to deliver from April 1, 1991, until December 31, 1992, were  
131 invited to participate. Children from 14,541 pregnancies were enrolled and 13,988 alive  
132 at 1 year. Additionally, 913 children were enrolled at age 7 years. All women gave  
133 written informed consent. Children at age 14 (wave 14,  $n = 10,581$ ), 16 (wave 16,  $n =$   
134  $9,702$ ), and 18 years (wave 18,  $n = 9,505$ ) that had not withdrawn consent were  
135 followed up. Parents answered questionnaires on 7,025 adolescents at wave 14 and on  
136 5,656 at wave 16. With 6,140 (58%) responding at wave 14, 5,069 (52%) at wave 16,  
137 and 3,228 (34%) at wave 18 which was used to augment the validity of the probable  
138 eating disorder diagnoses (Micali et al., 2015). The study website contains details of all  
139 the data that is available through a fully searchable data dictionary and variable search  
140 tool ([www.bristol.ac.uk/alspac/researchers/our-data/](http://www.bristol.ac.uk/alspac/researchers/our-data/)). To reduce potential  
141 confounding through genetic relatedness in our analyses, we removed randomly one  
142 individual of each pair that is closely related ( $\varphi > 0.2$ ) using PLINK v1.90 (Chang et al.,  
143 2015), excluding 75 individuals that were duplicates, monozygotic twins, first-degree  
144 relatives (i.e., parent-offspring and full siblings), or second-degree relatives (i.e., half-  
145 siblings, uncles, aunts, grandparents, and double cousins, **Supplementary Table 2**).

146 We derived probable eating disorder diagnoses as previously reported  
147 (**Supplementary Table 3**) (Micali et al., 2015), using a combination of adolescent self-  
148 report as well as information on the adolescents provided by their parents, a gold  
149 standard for childhood psychiatric disorders.

150 Ethical approval for the ALSPAC participants of this study was obtained from the  
151 ALSPAC Ethics and Law Committee and the Local Research Ethics Committees:  
152 [www.bristol.ac.uk/alspac/researchers/research-ethics/](http://www.bristol.ac.uk/alspac/researchers/research-ethics/). Consent for biological samples  
153 has been collected in accordance with the Human Tissue Act (2004) and informed  
154 consent for the use of data collected via questionnaires and clinics was obtained from

155 participants following the recommendations of the ALSPAC Ethics and Law Committee  
156 at the time.

157

## 158 **Polygenic risk scoring on eating disorder phenotypes in UK Biobank and** 159 **replication in ALSPAC**

160 We used PRSice (Choi & O'Reilly, 2019), version 2.2.3. We clumped the single  
161 nucleotide polymorphisms (SNPs) that were present both in the summary statistics of  
162 the trait and in the genotype data of the UK Biobank (i.e., overlapping SNPs) to obtain  
163 genetically independent SNPs. We retained the SNP with the smallest  $p$  value in each  
164 250 kilobase window of all those in linkage disequilibrium ( $r^2 > 0.1$ ). We calculated 269  
165 polygenic scores at their optimal  $p$  value threshold in each individual by scoring the  
166 number of effect alleles (weighted by the allele effect size) across the set of remaining  
167 SNPs (for a full list, see **Supplementary Table 4**). We calculated the polygenic score  
168 using the high-resolution scoring (i.e., incrementally across a large number of  $p$  value  
169 thresholds) method to identify the  $p$  value threshold at which the polygenic score is  
170 optimally associated with the outcome and explains the most variance (i.e., resulting in  
171 the highest adjusted  $R^2$  for continuous outcomes and Nagelkerke's  $R^2$  on the liability  
172 scale for binary outcomes). We evaluated the associations between polygenic score and  
173 eating disorder diagnosis using logistic regressions, adjusted for sex and the first six  
174 principal components that were calculated on the European subsample. To adjust for  
175 overfitting, we permuted case-control status at every  $p$  value threshold 10,000 times  
176 and, hence, calculated empirical  $p$  values. We converted the observed  $R^2$  to the liability  
177 scale assuming following population prevalences of 3% for anorexia nervosa, 1.8% for  
178 bulimia nervosa, and 3% for binge-eating disorder (Micali et al., 2017; Smink et al.,  
179 2014).

180           To correct for multiple testing (i.e., 269 polygenic score regression models), we  
181   calculated  $Q$  values using the false discovery rate approach (Benjamini & Hochberg,  
182   1995; Benjamini & Yekutieli, 2001). We did not stratify analyses by sex because of the  
183   low number of male eating disorder cases, but included sex as a covariate. We used the  
184   UK Biobank sample as our discovery cohort and the SNPs that were included in the  
185   strongest associated polygenic score were used to derive polygenic scores in the  
186   ALSPAC sample, in which we repeated the analysis also adjusting for sex, and the first  
187   six ancestry-informative principal components.  
188

## 189 **Results**

190 Descriptive statistics of the analyses samples included from UK Biobank and ALSPAC  
191 are listed in Supplementary Tables 1 and 2.

### 192 **Polygenic scoring on eating disorder phenotypes in UK Biobank and ALSPAC**

193 After correcting for multiple testing using the false discovery rate adjustment, 18  
194 polygenic scores were significantly associated with the three primary eating disorders  
195 in the subsample of the UK Biobank (**Figure 1**).

196 **Psychiatric disorders and neurological disease.** Five polygenic scores  
197 reflecting polygenic liability for psychiatric disorders and one neurological disease  
198 showed significant associations with eating disorders (**Supplementary Table 4 & 5**).  
199 The schizophrenia polygenic score was positively associated with anorexia nervosa  
200 (odds ratio [OR] per standard deviation in polygenic score = 1.18, 95% confidence  
201 interval [CI]: 1.09, 1.27;  $Q = 0.03$ ) and binge-eating disorder (OR = 1.23, 95% CI: 1.13,  
202 1.35;  $Q = 0.01$ ). The major depressive disorder polygenic score was positively  
203 associated with anorexia nervosa (OR = 1.20, 95% CI: 1.11, 1.29;  $Q = 0.01$ ) and binge-  
204 eating disorder (OR = 1.25, 95% CI: 1.14, 1.36;  $Q = 0.01$ ). Additionally, the attention  
205 deficit hyperactivity disorder polygenic score was positively associated with binge-  
206 eating disorder (OR = 1.28, 95% CI: 1.17, 1.39;  $Q = 0.01$ ). The bipolar disorder polygenic  
207 score (OR = 1.17, 95% CI: 1.08, 1.26;  $Q = 0.04$ ), the anorexia nervosa polygenic score  
208 (OR = 1.32, 95% CI: 1.22, 1.42;  $Q = 0.01$ ), and the migraine polygenic score (OR = 1.17,  
209 95% CI: 1.09, 1.26;  $Q = 0.04$ ) were positively associated with anorexia nervosa.

210 **Somatic traits.** Several polygenic scores indexing polygenic load for  
211 anthropometric traits, including waist circumference (OR = 1.43, 95% CI: 1.31, 1.56;  $Q =$   
212 0.01), hip circumference (OR = 1.39, 95% CI: 1.27, 1.51;  $Q = 0.01$ ), overweight (OR =  
213 1.35, 95% CI: 1.24, 1.47;  $Q = 0.01$ ), obesity class 1 (OR = 1.38, 95% CI: 1.27; 1.51;  $Q =$

214 0.01), class 2 (OR = 1.33, 95% CI: 1.22; 1.45;  $Q = 0.01$ ) and class 3 (OR = 1.21, 95% CI:  
215 1.11, 1.32;  $Q = 0.04$ ), BMI (OR = 1.44, 95% CI: 1.32; 1.56;  $Q = 0.01$ ) and extreme BMI (OR  
216 = 1.32, 95% CI: 1.22, 1.44;  $Q = 0.01$ ) were positively associated with binge-eating  
217 disorder. However, it is important to note that these polygenic scores were correlated  
218 with each other with  $r = .10$  to  $r = .67$  (**Supplementary Figure 1**). Additionally, the  
219 childhood obesity polygenic score (OR = 1.22, 95% CI: 1.12, 1.33;  $Q = 0.04$ ) was  
220 positively associated with binge-eating disorder, while the age at menarche polygenic  
221 score (OR = 0.79, 95% CI: 0.72, 0.86;  $Q = 0.01$ ) was negatively associated with binge-  
222 eating disorder. Neither the childhood obesity nor the age at menarche polygenic score  
223 were associated with anorexia nervosa.

224 The overweight polygenic score (OR = 1.26, 95% CI: 1.14, 1.38;  $Q = 0.01$ ) and the  
225 waist circumference polygenic score (OR = 1.25, 95% CI: 1.13, 1.38;  $Q = 0.02$ ) were also  
226 positively associated with bulimia nervosa.

227 **Sex and age differences.** The adult obesity polygenic score was more strongly  
228 associated with anorexia nervosa than the childhood obesity polygenic score. The  
229 difference in the log odds ratios was statistically significant (difference = -0.15,  $p =$   
230 0.004). We did not detect age differences in the associations of the obesity polygenic  
231 scores with bulimia nervosa or binge-eating disorder (**Supplementary Table 6**). We  
232 found no evidence for sex differences in the associations of the body mass index  
233 polygenic score with anorexia nervosa, bulimia nervosa, or binge-eating disorder.

234 **Sensitivity analyses.** We compared logistic regression results when assigning  
235 participants to all possible eating disorder diagnoses with assigning participants  
236 exclusively to one eating disorder diagnosis in the UK Biobank sample. The effect sizes  
237 of both analyses correlated with  $r = 0.98$ , indicating that the analyses were not sensitive  
238 to participant overlap (**Supplementary Figure 2**).

239           **Independent replication.** In ALSPAC (**Supplementary Table 7**), the  
240   overweight polygenic score (OR = 1.38, 95% CI: 1.13, 1.69;  $p_{\text{empirical}} = 0.05$ ) and the  
241   obesity class 1 polygenic score (OR = 1.23, 95% CI: 1.00; 1.50;  $p_{\text{empirical}} = 0.001$ ) were  
242   positively associated with binge-eating disorder. Correlations between these polygenic  
243   scores are presented in **Supplementary Figure 3**.

244

245

246

247

## 248 **Discussion**

249 Here, we present the first molecular genetic evidence that the underlying biology  
250 differs between binge-type eating disorders and anorexia nervosa and these differences  
251 can be captured at the genomic level. We find that polygenic scores for psychiatric  
252 disorders and anthropometric traits are associated with binge-type eating disorders  
253 (see **Figure 1**). Our UK Biobank analyses propose strong positive associations between  
254 bulimia nervosa/binge-eating disorder and anthropometric polygenic scores (e.g.,  
255 overweight and waist circumference), suggesting that binge eating shares genomic  
256 variants with overweight and obesity. In contrast, for anorexia nervosa, the direction of  
257 these associations was reversed. Binge-eating disorder also showed an association with  
258 the childhood obesity polygenic score while anorexia nervosa did not, further  
259 highlighting the genomic differences between binge-type eating disorders and anorexia  
260 nervosa. The genetic relationship between binge eating and BMI had been studied  
261 previously (Abdulkadir et al., 2020; Bulik et al., 2003) and is replicated by our analyses.  
262 It is notable that previous work suggests that genomic variants associated with BMI are  
263 predominantly expressed in brain tissue (Finucane et al., 2018). Our results also suggest  
264 that these BMI-associated genomic variants are relevant for eating disorders, but may  
265 act in opposite directions in binge-type eating disorders and anorexia nervosa.

266 Using the UK Biobank data, we also found that psychiatric polygenic scores (e.g.,  
267 schizophrenia and major depressive disorder) were positively associated with both  
268 anorexia nervosa and binge-eating disorder (**Figure 1, Panel A**). The ADHD polygenic  
269 score was only significantly associated with binge-eating disorder, while the anorexia  
270 polygenic score was not associated with bulimia nervosa or binge-eating disorder.

271 Our findings are in agreement with previous studies that showed genetic  
272 correlations between binge eating and depression (Munn-Chernoff et al., 2015) as well

273 as genetic correlations between anorexia nervosa and depression (Wade et al., 2000;  
274 Watson et al., 2019), migraine (Mustelin et al., 2014), schizophrenia (Duncan et al.,  
275 2017; Watson et al., 2019), bipolar disorder (Stahl et al., 2019) and anthropometric  
276 traits (Duncan et al., 2017; Watson et al., 2019). In line with our findings, previous  
277 GWASs showed no genomic association between anorexia nervosa and ADHD, whereas  
278 an ADHD polygenic score was associated with binge eating in a Swedish twin sample  
279 (Capusan et al., 2017; Yao, Kuja-Halkola, et al., 2019). Moreover, a large evidence-base  
280 highlights phenotypic overlap between ADHD and binge eating (Cortese et al., 2016;  
281 Leventakou et al., 2016; Sonnevile et al., 2015).

282

283       Large GWASs have recently suggested the involvement of a metabolic component  
284 in anorexia nervosa through genetic correlations (Duncan et al., 2017; Watson et al.,  
285 2019); however, at the current sample size, we did not observe associations between  
286 metabolic polygenic scores and the three primary eating disorders in the UK Biobank  
287 most probably due to limited statistical power (**Figure 1, Panel B**).

288       The association between polygenic liability for higher body mass and eating  
289 disorders is further complicated as it shows age dependence (Hübel et al., 2019) and  
290 confounding through age at menarche. In both the UK Biobank and the ALSPAC cohort,  
291 we found that age at menarche and BMI polygenic scores were negatively correlated  
292 with each other. However, the age at menarche polygenic score was only negatively  
293 associated with binge-eating disorder but not with anorexia nervosa (**Figure 1, Panel**  
294 **C**). Phenotypic associations between early menarche and bulimia nervosa have  
295 previously been reported (Fairburn et al., 1999; Reichborn-Kjennerud et al., 2004), but  
296 do not always replicate (Algars et al., 2014). On a genetic level, binge eating correlated  
297 negatively with age at menarche in a twin study (Baker et al., 2012) and anorexia

298 nervosa was not genetically correlated with age at menarche in our polygenic score  
299 analysis nor in previous GWASs (Duncan et al., 2017; Watson et al., 2019). This suggests  
300 that the association between age at menarche and eating disorders may be driven by  
301 symptoms related to binge eating rather than restriction.

302

303         Furthermore, ADHD and BMI show a positive genetic correlation (Demontis et  
304 al., 2019; Hübel et al., 2019) and in our analysis the ADHD polygenic score was  
305 associated with binge-eating disorder. Clinically, the ADHD medication  
306 lisdexamfetamine is used to treat binge-eating disorder (Hudson et al., 2017). This  
307 combined evidence suggests that the shared biology between ADHD and BMI may  
308 contribute to binge eating. These novel findings warrant further research; investigating  
309 how genomic variants associated with ADHD, commonly characterized by inattention  
310 and impulsivity, are also associated with binge-eating disorder and contribute to the  
311 high comorbidity between ADHD and obesity (Cortese et al., 2016; Leventakou et al.,  
312 2016; Sonnevile et al., 2015).

313

314         Our findings must be interpreted in the light of the following limitations: the  
315 number of individuals with eating disorders in both samples was relatively low, some of  
316 the eating disorder diagnoses and symptoms were self-reported, and the samples only  
317 included white British participants. Furthermore, we were unable to differentiate  
318 between binge-eating/purging and restricting anorexia nervosa as subtype information  
319 was not available. However, symptoms and derived diagnoses in the ALSPAC cohort  
320 were obtained both from parents and adolescents, representing the gold standard in  
321 child and adolescent psychiatry, strengthening the validity of the diagnoses. Further,  
322 replication studies are essential to lend confidence to the results reported here.

323           Our findings show for the first time that similarities exist in the genomic  
324 psychiatric liability for binge-type eating disorders and anorexia nervosa. However, we  
325 find clear differences between binge-type eating disorders and anorexia nervosa in the  
326 underlying biology of body mass regulation at the genomic level. These findings open  
327 important avenues for translational research relevant to eating disorder phenotypes  
328 and overlap between eating disorders. Specifically, it remains unknown which biological  
329 mechanisms are shared between ADHD and binge eating.

330

331

332 **Acknowledgements**

333 We are deeply grateful to all the families who took part in this study, the midwives for  
334 their help in recruiting them, and the whole ALSPAC team, which includes interviewers,  
335 computer and laboratory technicians, clerical workers, research scientists, volunteers,  
336 managers, receptionists and nurses. This study was completed as part of approved UK  
337 Biobank study applications 27546 to Prof Breen.

338

339 **Funding**

340 This study represents independent research part funded by the UK National Institute  
341 for Health Research (NIHR) Biomedical Research Centre at South London and Maudsley  
342 NHS Foundation Trust and King's College London. The views expressed are those of the  
343 author(s) and not necessarily those of the UK NHS, the NIHR or the Department of  
344 Health. High performance computing facilities were funded with capital equipment  
345 grants from the GSTT Charity (TR130505) and Maudsley Charity (980). This work was  
346 supported by the UK Medical Research Council and the Medical Research Foundation  
347 (ref: MR/R004803/1). The UK Medical Research Council and Wellcome (Grant ref:  
348 102215/2/13/2 and 217065/Z/19/Z) and the University of Bristol provide core  
349 support for ALSPAC. A comprehensive list of grants funding is available on the ALSPAC  
350 website ([http://www.bristol.ac.uk/alspac/external/documents/grant-](http://www.bristol.ac.uk/alspac/external/documents/grant-acknowledgements.pdf)  
351 [acknowledgements.pdf](http://www.bristol.ac.uk/alspac/external/documents/grant-acknowledgements.pdf)); This research was specifically funded by the NIHR  
352 (CS/01/2008/014), the NIH (MH087786-01). GWAS data was generated by Sample  
353 Logistics and Genotyping Facilities at Wellcome Sanger Institute and LabCorp  
354 (Laboratory Corporation of America) using support from 23andMe. NM and CB  
355 acknowledge funding from the National Institute of Mental Health (R21 MH115397). CB  
356 acknowledges funding from the Swedish Research Council (VR Dnr: 538-2013-8864),

357 the National Institute of Mental Health (R21MH115397; R01 MH109528;  
358 R01MH120170; R01MH119084; R01MH120170), and Lundbeckfonden. MH is  
359 supported by fellowship from the Medical Research Council UK (MR/T027843/1). The  
360 content is solely the responsibility of the authors and does not necessarily represent the  
361 official views of the National Institutes of Health. The funders were not involved in the  
362 design or conduct of the study; collection, management, analysis, or interpretation of  
363 the data; or preparation, review, or approval of the manuscript.

364

### 365 **Availability of data and materials**

366 This study is based on data from the UK Biobank. Researchers can apply for the data via  
367 the following website: <https://www.ukbiobank.ac.uk/>. This study is based on data  
368 from the ALSPAC study (<http://www.bristol.ac.uk/alspac/>). Interested researchers can  
369 apply for data access with the University of Bristol, UK. Analysis scripts can be  
370 requested from the authors.

371

### 372 **Declaration of interest**

373 Dr. Breen has received grant funding from and served as a consultant to Eli Lilly, has  
374 received honoraria from Illumina and has served on advisory boards for Otsuka. Dr.  
375 Bulik is a grant recipient from and has served on advisory boards for Shire and is a  
376 consultant for Idorsia. She receives royalties from Pearson. She is a grant recipient from  
377 Lundbeckfonden. All other authors have indicated they have no conflicts of interest to  
378 disclose.

379

### 380 **Author Contribution**

381 CH, MA, MH analysed the data. CH, MA, MH drafted the manuscript. CMB, NM, RFL, and  
382 GB supervised the work. All authors substantially contributed to the conception and  
383 interpretation of the work, revised the manuscript for important intellectual content  
384 and approved the final version. All authors agree to be accountable for all aspects of this  
385 work.  
386  
387

## 388   **References**

- 389   Abdulkadir, M., Herle, M., De Stavola, B. L., Hübel, C., Santos Ferreira, D. L., Loos, R. J. F., Bryant-  
390       Waugh, R., Bulik, C. M., & Micali, N. (2020). Polygenic Score for Body Mass Index Is  
391       Associated with Disordered Eating in a General Population Cohort. *Journal of Clinical*  
392       *Medicine Research*, 9(4). <https://doi.org/10.3390/jcm9041187>
- 393   Algars, M., Huang, L., Von Holle, A. F., Peat, C. M., Thornton, L. M., Lichtenstein, P., & Bulik, C. M.  
394       (2014). Binge eating and menstrual dysfunction. *Journal of Psychosomatic Research*, 76(1),  
395       19–22.
- 396   American Psychiatric Association. (2013). *Diagnostic and Statistical Manual of Mental Disorders*  
397       (5th ed.). American Psychiatric Association.
- 398   Baker, J. H., Thornton, L. M., Bulik, C. M., Kendler, K. S., & Lichtenstein, P. (2012). Shared genetic  
399       effects between age at menarche and disordered eating. *The Journal of Adolescent Health: Official Publication of the Society for Adolescent Medicine*, 51(5), 491–496.
- 400   Benjamini, Y., & Hochberg, Y. (1995). Controlling the False Discovery Rate: A Practical and  
401       Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society. Series B, Statistical Methodology*, 57(1), 289–300.
- 402   Benjamini, Y., & Yekutieli, D. (2001). The control of the false discovery rate in multiple testing  
403       under dependency. *Annals of Statistics*, 29(4), 1165–1188.
- 404   Boyd, A., Golding, J., Macleod, J., Lawlor, D. A., Fraser, A., Henderson, J., Molloy, L., Ness, A., Ring,  
405       S., & Davey Smith, G. (2013). Cohort Profile: the “children of the 90s”--the index offspring of  
406       the Avon Longitudinal Study of Parents and Children. *International Journal of Epidemiology*,  
407       42(1), 111–127.
- 408   Bulik, C. M., Sullivan, P. F., & Kendler, K. S. (2003). Genetic and environmental contributions to  
409       obesity and binge eating. *The International Journal of Eating Disorders*, 33(3), 293–298.
- 410   Bulik, C. M., Thornton, L. M., Root, T. L., Pisetsky, E. M., Lichtenstein, P., & Pedersen, N. L. (2010).  
411       Understanding the relation between anorexia nervosa and bulimia nervosa in a Swedish  
412       national twin sample. *Biological Psychiatry*, 67(1), 71–77.
- 413   Bulik, C. M., Thornton, L. M., Root, T. L., Pisetsky, E. M., Lichtenstein, P., & Pedersen, N. L. (2010).  
414       Understanding the relation between anorexia nervosa and bulimia nervosa in a Swedish  
415       national twin sample. *Biological Psychiatry*, 67(1), 71–77.

- 415 Capusan, A. J., Yao, S., Kuja-Halkola, R., Bulik, C. M., Thornton, L. M., Bendtsen, P., Marteinsdottir,  
416 I., Thorsell, A., & Larsson, H. (2017). Genetic and environmental aspects in the association  
417 between attention-deficit hyperactivity disorder symptoms and binge-eating behavior in  
418 adults: a twin study. *Psychological Medicine*, *47*(16), 2866–2878.
- 419 Chang, C. C., Chow, C. C., Tellier, L. C., Vattikuti, S., Purcell, S. M., & Lee, J. J. (2015). Second-  
420 generation PLINK: rising to the challenge of larger and richer datasets. *GigaScience*, *4*, 7.
- 421 Choi, S. W., & O'Reilly, P. F. (2019). PRSice-2: Polygenic Risk Score software for biobank-scale  
422 data. *GigaScience*, *8*(7). <https://doi.org/10.1093/gigascience/giz082>
- 423 Cortese, S., Moreira-Maia, C. R., St Fleur, D., Morcillo-Peñalver, C., Rohde, L. A., & Faraone, S. V.  
424 (2016). Association Between ADHD and Obesity: A Systematic Review and Meta-Analysis.  
425 *The American Journal of Psychiatry*, *173*(1), 34–43.
- 426 Davis, K. A. S., Coleman, J. R. I., Adams, M., Allen, N., Breen, G., Cullen, B., Dickens, C., Fox, E.,  
427 Graham, N., Holliday, J., Howard, L. M., John, A., Lee, W., McCabe, R., McIntosh, A., Pearsall, R.,  
428 Smith, D. J., Sudlow, C., Ward, J., ... Hotopf, M. (2020). Mental health in UK Biobank -  
429 development, implementation and results from an online questionnaire completed by 157  
430 366 participants: a reanalysis. *BJPsych Open*, *6*(2), e18.
- 431 Demontis, D., Walters, R. K., Martin, J., Mattheisen, M., Als, T. D., Agerbo, E., Baldursson, G.,  
432 Belliveau, R., Bybjerg-Grauholm, J., Bækvad-Hansen, M., Cerrato, F., Chambert, K.,  
433 Churchhouse, C., Dumont, A., Eriksson, N., Gandal, M., Goldstein, J. I., Grasby, K. L., Grove, J.,  
434 ... Neale, B. M. (2019). Discovery of the first genome-wide significant risk loci for attention  
435 deficit/hyperactivity disorder. *Nature Genetics*, *51*(1), 63–75.
- 436 Duncan, L., Yilmaz, Z., Gaspar, H., Walters, R., Goldstein, J., Anttila, V., Bulik-Sullivan, B., Ripke, S.,  
437 Eating Disorders Working Group of the Psychiatric Genomics Consortium, Thornton, L.,  
438 Hinney, A., Daly, M., Sullivan, P. F., Zeggini, E., Breen, G., & Bulik, C. M. (2017). Significant  
439 Locus and Metabolic Genetic Correlations Revealed in Genome-Wide Association Study of  
440 Anorexia Nervosa. *The American Journal of Psychiatry*, *174*(9), 850–858.
- 441 Fairburn, C. G., Cooper, Z., Doll, H. A., & Welch, S. L. (1999). Risk factors for anorexia nervosa:

442 three integrated case-control comparisons. *Archives of General Psychiatry*, 56(5), 468–476.

443 Finucane, H. K., Reshef, Y. A., Anttila, V., Slowikowski, K., Gusev, A., Byrnes, A., Gazal, S., Loh, P.-R.,  
444 Lareau, C., Shores, N., Genovese, G., Saunders, A., Macosko, E., Pollack, S., Brainstorm  
445 Consortium, Perry, J. R. B., Buenrostro, J. D., Bernstein, B. E., Raychaudhuri, S., ... Price, A. L.  
446 (2018). Heritability enrichment of specifically expressed genes identifies disease-relevant  
447 tissues and cell types. *Nature Genetics*, 50(4), 621–629.

448 Fraser, A., Macdonald-Wallis, C., Tilling, K., Boyd, A., Golding, J., Davey Smith, G., Henderson, J.,  
449 Macleod, J., Molloy, L., Ness, A., Ring, S., Nelson, S. M., & Lawlor, D. A. (2013). Cohort Profile:  
450 the Avon Longitudinal Study of Parents and Children: ALSPAC mothers cohort.  
451 *International Journal of Epidemiology*, 42(1), 97–110.

452 Golding, J., Pembrey, M., Jones, R., & ALSPAC Study Team. (2001). ALSPAC--the Avon  
453 Longitudinal Study of Parents and Children. I. Study methodology. *Paediatric and Perinatal  
454 Epidemiology*, 15(1), 74–87.

455 Hübel, C., Gaspar, H. A., Coleman, J. R. I., Hanscombe, K. B., Purves, K., Prokopenko, I., Graff, M.,  
456 Ngwa, J. S., Workalemahu, T., ADHD Working Group of the Psychiatric Genomics  
457 Consortium, Meta-Analyses of Glucose and Insulin-related traits consortium (MAGIC),  
458 Autism Working Group of the Psychiatric Genomics Consortium, Bipolar Disorder Working  
459 Group of the Psychiatric Genomics Consortium, Eating Disorders Working Group of the  
460 Psychiatric Genomics Consortium, Major Depressive Disorder Working Group of the  
461 Psychiatric Genomics Consortium, OCD & Tourette Syndrome Working Group of the  
462 Psychiatric Genomics Consortium, PTSD Working Group of the Psychiatric Genomics  
463 Consortium, Schizophrenia Working Group of the Psychiatric Genomics Consortium, Sex  
464 Differences Cross Disorder Working Group of the Psychiatric Genomics Consortium, ...  
465 Breen. (2019). Genetic correlations of psychiatric traits with body composition and  
466 glycaemic traits are sex- and age-dependent. *Nature Communications*, 10(1), 5765.

467 Leventakou, V., Micali, N., Georgiou, V., Sarri, K., Koutra, K., Koinaki, S., Vassilaki, M., Kogevinas,  
468 M., & Chatzi, L. (2016). Is there an association between eating behaviour and attention-

- 469 deficit/hyperactivity disorder symptoms in preschool children? *Journal of Child Psychology*  
470 *and Psychiatry, and Allied Disciplines*, 57(6), 676–684.
- 471 Micali, N., Martini, M. G., Thomas, J. J., Eddy, K. T., Kothari, R., Russell, E., Bulik, C. M., & Treasure,  
472 J. (2017). Lifetime and 12-month prevalence of eating disorders amongst women in mid-  
473 life: a population-based study of diagnoses and risk factors. *BMC Medicine*, 15(1), 12.
- 474 Micali, N., Solmi, F., Horton, N. J., Crosby, R. D., Eddy, K. T., Calzo, J. P., Sonnevile, K. R., Swanson,  
475 S. A., & Field, A. E. (2015). Adolescent Eating Disorders Predict Psychiatric, High-Risk  
476 Behaviors and Weight Outcomes in Young Adulthood. *Journal of the American Academy of*  
477 *Child and Adolescent Psychiatry*, 54(8), 652–659.e1.
- 478 Munn-Chernoff, M. A., Grant, J. D., Agrawal, A., Koren, R., Glowinski, A. L., Bucholz, K. K., Madden,  
479 P. A. F., Heath, A. C., & Duncan, A. E. (2015). Are there common familial influences for major  
480 depressive disorder and an overeating-binge eating dimension in both European American  
481 and African American female twins? *The International Journal of Eating Disorders*, 48(4),  
482 375–382.
- 483 Mustelin, L., Raevuori, A., Kaprio, J., & Keski-Rahkonen, A. (2014). Association between eating  
484 disorders and migraine may be explained by major depression. *The International Journal of*  
485 *Eating Disorders*, 47(8), 884–887.
- 486 Reichborn-Kjennerud, T., Bulik, C. M., Sullivan, P. F., Tambs, K., & Harris, J. R. (2004). Psychiatric  
487 and medical symptoms in binge eating in the absence of compensatory behaviors. *Obesity*  
488 *Research*, 12(9), 1445–1454.
- 489 Smink, F. R. E., Hoeken, D., Oldehinkel, A. J., & Hoek, H. W. (2014). Prevalence and severity of  
490 DSM-5 eating disorders in a community cohort of adolescents. *The International Journal of*  
491 *Eating Disorders*, 47(6), 610–619.
- 492 Sonnevile, K. R., Calzo, J. P., Horton, N. J., Field, A. E., Crosby, R. D., Solmi, F., & Micali, N. (2015).  
493 Childhood hyperactivity/inattention and eating disturbances predict binge eating in  
494 adolescence. *Psychological Medicine*, 45(12), 2511–2520.
- 495 Stahl, E. A., Breen, G., Forstner, A. J., McQuillin, A., Ripke, S., Trubetskoy, V., Mattheisen, M., Wang,

496 Y., Coleman, J. R. I., Gaspar, H. A., de Leeuw, C. A., Steinberg, S., Pavlides, J. M. W.,  
497 Trzaskowski, M., Byrne, E. M., Pers, T. H., Holmans, P. A., Richards, A. L., Abbott, L., ... Bipolar  
498 Disorder Working Group of the Psychiatric Genomics Consortium. (2019). Genome-wide  
499 association study identifies 30 loci associated with bipolar disorder. *Nature Genetics*, *51*(5),  
500 793–803.

501 Sudlow, C., Gallacher, J., Allen, N., Beral, V., Burton, P., Danesh, J., Downey, P., Elliott, P., Green, J.,  
502 Landray, M., Liu, B., Matthews, P., Ong, G., Pell, J., Silman, A., Young, A., Sprosen, T., Peakman,  
503 T., & Collins, R. (2015). UK biobank: an open access resource for identifying the causes of a  
504 wide range of complex diseases of middle and old age. *PLoS Medicine*, *12*(3), e1001779.

505 Treasure, J., Zipfel, S., Micali, N., Wade, T., Stice, E., Claudino, A., Schmidt, U., Frank, G. K., Bulik, C.  
506 M., & Wentz, E. (2015). Anorexia nervosa. *Nature Reviews. Disease Primers*, *1*, 15074.


507 Wade, T. D., Bulik, C. M., Neale, M., & Kendler, K. S. (2000). Anorexia nervosa and major  
508 depression: shared genetic and environmental risk factors. *The American Journal of*  
509 *Psychiatry*, *157*(3), 469–471.

510 Watson, H. J., Yilmaz, Z., Thornton, L. M., Hübel, C., Coleman, J. R. I., Gaspar, H. A., Bryois, J.,  
511 Hinney, A., Leppä, V. M., Mattheisen, M., Medland, S. E., Ripke, S., Yao, S., Giusti-Rodríguez, P.,  
512 Anorexia Nervosa Genetics Initiative, Hanscombe, K. B., Purves, K. L., Eating Disorders  
513 Working Group of the Psychiatric Genomics Consortium, Adan, R. A. H., ... Bulik, C. M.  
514 (2019). Genome-wide association study identifies eight risk loci and implicates metabo-  
515 psychiatric origins for anorexia nervosa. *Nature Genetics*, *51*(8), 1207–1214.

516 World Health Organization. (1992). *ICD-10: International Statistical Classification of Diseases and*  
517 *Related Health Problems: 10th revision*. World Health Organization.

518 Yao, S., Kuja-Halkola, R., Martin, J., Lu, Y., Lichtenstein, P., Noring, C., Birgegård, A., Yilmaz, Z.,  
519 Hübel, C., Watson, H., Baker, J., Almqvist, C., Eating Disorders Working Group of the  
520 Psychiatric Genomics Consortium, Thornton, L. M., Magnusson, P. K., Bulik, C. M., & Larsson,  
521 H. (2019). Associations Between Attention-Deficit/Hyperactivity Disorder and Various  
522 Eating Disorders: A Swedish Nationwide Population Study Using Multiple Genetically

- 523 Informative Approaches. *Biological Psychiatry*, 86(8), 577–586.
- 524 Yao, S., Larsson, H., Norring, C., Birgegård, A., Lichtenstein, P., D'Onofrio, B. M., Almqvist, C.,  
525 Thornton, L. M., Bulik, C. M., & Kuja-Halkola, R. (2019). Genetic and environmental  
526 contributions to diagnostic fluctuation in anorexia nervosa and bulimia nervosa.  
527 *Psychological Medicine*, 1–8.
- 528 Yilmaz, Z., Hardaway, J. A., & Bulik, C. M. (2015). Genetics and Epigenetics of Eating Disorders.  
529 *Advances in Genomics and Genetics*, 5, 131–150.
- 530



**Figure 1. Polygenic scores associated with eating disorders in the UK Biobank.**

**Panel A** shows psychiatric and behavioral, **Panel B** metabolic, and **Panel C** anthropometric polygenic scores and their association estimates with self-reported or hospital-diagnosed eating disorders in the UK Biobank sample ( $n = 17,050$ ). Filled dots are statistically significant after adjustment for multiple testing through the false discovery approach. Dots represent odds ratios (ORs) and error bars index 95% confidence intervals obtained via logistic regression and 10,000 permutations to obtain empirical  $p$  values.