Correlation between universal BCG vaccination policy and reduced mortality for COVID-
19

Aaron Miller, Mac Josh Reandelar, Kimberly Fasciglione, Violeta Roumenova, Yan Li, and
Gonzalo H. Otazu*

Department of Biomedical Sciences, NYIT College of Osteopathic Medicine, New York Institute
of Technology, Old Westbury, New York, USA

*Correspondence to: gotazual@nyit.edu

Abstract

COVID-19 has spread to most countries in the world. Puzzlingly, the impact of the disease varies
in different countries. This variation is attributed to differences in cultural norms, mitigation efforts,
and health infrastructure. Here, we propose that national differences in COVID-19 impact could
be partially explained by different national policies with respect to Bacillus Calmette-Guérin (BCG)
vaccination. BCG vaccination has been reported to offer broad protection from other respiratory
infections besides tuberculosis. We compared BCG vaccination policies with the morbidity and
mortality for COVID-19 for middle-high and high-income countries. We found that countries
without universal policies of BCG vaccination (Italy, the Netherlands, USA) have been more
severely affected compared to countries with universal and long-standing BCG policies. The
difference cannot be accounted for by differences in disease onset, adoption of early social
distancing policies, state of health services, nor income level. Reduced mortality suggests BCG
vaccination could be a potential new tool in the fight against COVID-19.

Introduction

The COVID-19 pandemic originated in China and has quickly spread across the world, affecting
nearly every country. However, there are some striking differences in how COVID-19 is behaving
in different countries. For instance, in Italy there has been strong curtailing of social interactions
but COVID-19 mortality is high (29315 deaths as of May 5th, 2020). In contrast, Japan had some
of the earlier cases, but the mortality is low (556 deaths as of May 5th, 20202) despite not having
adopted some of the more restrictive social isolation measurements ("Japan Was Expecting a
Coronavirus Explosion. Where Is It? | The Japan Times" n.d.). These puzzling differences have
been adjudicated to different cultural norms as well as differences in medical care standards. In
addition, it has been proposed that the reduced mortality in children might be due to recent
vaccination(Cao et al. 2020). Here we propose that the country-by-country difference in COVID-
19 mortality can be partially explained by national policies on Bacillus Calmette-Guérin (BCG)
vaccination.

BCG is a live attenuated strain derived from an isolate of *Mycobacterium bovis* used widely across
the world as a vaccine for Tuberculosis (TB), with many nations, including Japan and China,
having a universal BCG vaccination policy in newborns("WHO | Tuberculosis" n.d.). Other
countries such as Spain, France, and Switzerland, have discontinued their universal vaccine
policies due to comparatively low risk for developing *M. bovis* infections as well as the proven
variable effectiveness in preventing adult TB. Countries such as the United States, Italy, and the
Netherlands, have yet to adopt universal vaccine policies for similar reasons.

Several vaccines including the BCG vaccination have been shown to produce positive
"heterologous" or non-specific immune effects leading to improved response against other non-

mycobacterial pathogens. For instance, in severe combined immunodeficiency (SCID) mice,
which lack functional B- and T-cells, BCG vaccination protected against a secondary non-
mycobacterial challenge, demonstrating the ability of innate immune cells to mount a non- specific
'memory-like' response(Kleinnijenhuis et al. 2014). This phenomenon was named 'trained
immunity' and is proposed to be caused by metabolic and epigenetic changes leading to the
promotion of genetic regions encoding for pro-inflammatory cytokines(Netea et al. 2016). BCG
vaccination significantly increases the secretion of pro-inflammatory cytokines, specifically IL-1B,
which has been shown to play a vital role in antiviral immunity(Kleinnijenhuis et al. 2014).
Additionally, a study in Guinea-Bissau found that children vaccinated with BCG were observed to
have a 50% reduction in overall mortality, which was attributed to the vaccine's effect on reducing
respiratory infections and sepsis (Kristensen, Aaby, and Jensen 2000) i.

Reports on the duration of BCG vaccination protection against TB vary between 10 years(Sterne,
Rodrigues, and Guedes 1998) and 60 years (Aronson et al. 2004). BCG protection for conditions
other than TB has also shown long-lasting protection: a 60-year follow up of a clinical trial
suggested that individuals who received the BCG vaccination during early childhood subsequently
demonstrated a 2.5 fold reduction of lung cancer development (Usher et al. 2019). Whether the
mechanism responsible for the BCG vaccination's long-standing effectiveness against
mycobacterial pathogens carries over as a means of protection against the novel COVID-19
remains unknown, but the potential opens up an interesting area for exploration.

Given our current understanding of the BCG vaccine's nonspecific immunotherapeutic
mechanisms and by analyzing current epidemiological data, this investigation aims to identify a
possible correlation between the existence of universal BCG vaccine policies and mortality
associated to COVID-19 infections all over the world.

Methods

We collected the BCG vaccination policies across countries from the BCG World Atlas(Zwerling
et al. 2011), available from http://www.bcgatlas.org/. We complemented the database with
respect to the dates of initiation of BCG vaccination. The additional references are in the adjunct
table. Data on COVID-19 cases and death per country were obtained from
https://google.org/crisisresponse/covid19-map on May 5th, 2020. We included in the analysis
only countries with more than 1 million inhabitants. The mortality rate might be influenced by
multiple factors including a country's economics and the age distribution of the population. To
account for economical differences, we classified countries according to their GNI per capita in
2018   using   the   World   Bank   data
(https://datahelpdesk.worldbank.org/knowledgebase/articles/906519-world-bank-country-and-
lending-groups).   Countries were divided into three categories: low income (L) with an annual
income of 1,025 dollars or less, lower-middle income with an income between 1,026 and 3,995
dollars, and middle-high and high-income countries, which included countries with annual
incomes over 3,996 dollars.

COVID-19 mortality increases steeply with age(Zhou et al. 2020). Universal vaccination policies
are more common in developing countries which tend to have younger populations. A younger
population would show spurious reduced mortality for COVID-19.   To control for this variable, we
did not use the total number of inhabitants per country to calculate the mortality rate, but we
considered the susceptible population as the inhabitants over 65 years of age per country(Shet
et al. n.d.). Medians between populations were compared using the Wilcoxon rank sum test.

Multivariate linear analyses were done using the function *fitlm* in Matlab (R2017b). The Matlab
script is available as supplementary material.

Results

Initially, we compared countries that never had in place a universal BCG vaccination policy (Italy,
USA, Lebanon, Nederland, and Belgium), with countries that have a current universal BCG
vaccination policy. Low-income countries report a median of 12.6 deaths per million people over
65 years of age. In fact, 23% (4/17) of these countries reported zero deaths attributed to COVID-
19, consistent with the hypothesis for a protective role of BCG vaccination. This effect, however,
could be attributed to significantly lower testing rates in low income countries (Spearnman
correlation between median income and tests per million inhabitants, rho=0.8083,p<1e-6, n=101
countries). Therefore, we will only consider middle-high and high-income countries for our
analysis, which should compare countries with similar levels of medical care standards, sanitation,
and testing levels for COVID-19. Middle high and high income countries with universal policies have a
high BCG coverage, with a median of 97% with the 10 percentile being 87% and the 90 percentile being
99%.

Middle-high and high-income countries that have a current universal BCG policy (55 countries)
had a median of 84.5 deaths per million people over 65 years of age (see **Figure 1**). In contrast,
middle-high and high-income countries that never had a universal BCG policy (5 countries) had
a higher mortality rate, with a median of 1583.2 deaths per million people over 65 years of age.
This difference between the median number of deaths was significant (p=0.008, Wilcoxon rank
sum test).

An alternative explanation for the reduced number of deaths in countries with universal BCG
vaccination policies could be that COVID-19 arrived late in these countries and that they are in
the earlier stages of the epidemic. To evaluate for the significance of a possible confounding effect
of epidemic stage, we plotted the number of deaths per million inhabitants over 65 years of age
versus the number of days since the first reported case of COVID-19 per country (see **Figure 2**).
As expected, there is a general increase in the number of reported deaths (as of 05/05/20) as
more days pass since a country's first reported case. Additionally, middle-high and high-income
countries *with* universal BCG vaccination policies that saw larger case volumes relatively early
have a median of 105.1 deaths per million people over 65 years of age. This is significantly lower
(p= 0.0059, Wilcoxon rank sum test) than the median of 1583.2 for countries in the same
economic category that never implemented a universal policy and were affected at comparable
times or even later (with at least 68 days passing since the first reported case). Therefore, the
increased mortality for countries that never implemented a universal BCG vaccination policy
cannot be explained by an earlier epidemic onset.

122
Although we have stratified the countries and compared only those with middle-high and high-
income levels that should be relatively homogeneous, differences in the strength of the health
system, poverty rates, and quality of public health policy in response to COVID-19 could still be
residual confounders that explain the observed increased death rates in countries that never
implemented a universal BCG vaccination policy. In order to address these confounders directly,
we used a multivariate linear regression analysis (see Table 1) on the middle-high and high-
income countries (55 with BCG policy+6 that never had a policy). We included the days since the
first reported case as covariates to account for differences in the stage of the epidemic in a country.
We included two variables that accounted for the quality of the public health policy with respect
to COVID-19. One was the number of tests performed per million people in a country which would
correlate with the strength of a country's response. The other was the speed of a country's
reaction to the pandemic which we quantified as the number of days between the first detected

case and schools closing. We used the median life expectancy in a country as a surrogate for the
strength of the health system. Although the countries that we considered were middle-high and
high-income countries, we also included the median income per country. Although median income
might fail to capture the degree of resources assigned to effective COVID-19 mitigation measures,
it would serve as an indicator of possible measures that could be taken.   Finally, we included the
BCG vaccination policy as a binary variable.
141

	Estimate	Standard Error	T statistic	P value
Intercept	9.4735	3.6302	2.6096	0.016
Number tests per million people	7.2609e-06	6.4418e-06	1.1272	0.27182
Log10(days since first COVID-19 case)	-4.4502	1.8847	-2.3612	0.027487
Log10(days from first case to school closing + 2)	0.39123	0.41023	0.95369	0.3506
Median life expectancy (years)	0.023614	0.029294	0.80611	0.42881
Median income (US dollars)	-6.7105e-06	1.1589e-05	-0.57904	0.56845
BCG policy (1 current, 0 never)	-1.4656	0.35403	-4.1396	0.000429

142
143 Table 1. Multivariate analysis of log10(deaths per million inhabitants over 65 years of age).
144 Number of observations: 29, Error degrees of freedom: 22
145 Root Mean Squared Error: 0.494
146 R-squared: 0.599, Adjusted R-Squared 0.49
147 F-statistic vs. constant model: 5.48, p-value = 0.00134
148

Consistent with our previous analysis, we found that the only significant covariate (p=0.000429)
was the presence of universal BCG vaccination policy in reducing the mortality of COVID-19.
None of the other factors reached significance (p>0.01). However, there still might be other factors
(genetic or environmental) that might be producing the correlation between BCG vaccination
policy and reduced deaths due to COVID-19.

The number of deaths per million people over the age of 65 years did not correlate with the
number of tests performed (p=0.27182), indicating that the number of deaths was a robust
indicator that did not depend on a country's testing response. Nevertheless, we have used data
from COVID-19 attributed deaths as reported by each country. There still might be systematic
differences across countries on whether a death is attributed to COVID-19. For instance, if
countries with universal BCG vaccination would tend to assign COVID-19 deaths to other causes
(atypical pneumonia), that could create a spurious correlation between BCG vaccination and
reduced mortality. In order to address this concern, we used a smaller dataset of European
countries (https://www.euromomo.eu/) that calculated excess deaths in 2020. We extracted the
data from weeks 12 to 17 of the year 2020 (see **Figure 3**) when the epidemic was peaking in

Europe. Countries with universal BCG vaccination policy had median excess deaths per week of
0.42 (n=5), whereas countries without universal BCG vaccination policies had significantly higher
median excess deaths of 16.02 per week (n=3, p=0.035, Wilcoxon rank sum test). Albeit a small
sample, this independent measurement supports the idea that the reported correlation between
a country's BCG vaccination and reduced mortality due to COVID-19 is not an artifact of national
differences in COVID-19 death assignation.

A possible mechanism for a putative protective effect of BCG in a population might include direct
protection in the elderly population, which is the most affected group(Zhou et al. 2020), as well as
herd immunity acting over the whole population, both young and old. Using data from March 21st,
2020(Miller et al. 2020), we reported a negative correlation between the year start of the
vaccination policy and mortality both in middle high and high income countries with current
policies, as well as countries that had stopped their policies. This was consistent with BCG
vaccination producing a long-lasting protection to the elderly population from BCG vaccination
received in childhood. However, using the current data from May 5th, there is no significant
correlation between start year of the vaccination and mortality per million people over 65 years of
age (see **Fig. 4**). We do not know if the early reported correlation was spuriously produced by
the evolving nature of the pandemic, or if earlier response to the disease are less affected by a
country's pandemic mitigation responses and depend more on intrinsic immunity in the elderly
population to COVID-19. This would be consistent with a recent pre-print report (Berg et al. 2020)
suggesting that BCG vaccination policy slows down the spread of COVID-19, making it easier to
observe differences in the protective effect of BCG earlier in the epidemic.

A second possibility for a putative protective effect of BCG for COVID-19 could be a reduction in
the spread of the disease, acting across the population. This would be reflected as a reduction in
the number of COVID-19 reported cases. Consistent with this hypothesis, the countries with low-
income levels that have a universal policy (16 countries) reported a smaller number of cases of
COVID-19 per million inhabitants with a median of 8.85. However, the issue of underreporting
might be more critical for estimating the number of cases and we have excluded the low-income
countries from further analysis. Middle-high and high-income countries that have a current
universal BCG vaccination policy (55 countries) had a median of 313.6 cases per million
inhabitants (see **Figure 5**). Consistent with the role of BCG in slowing spread of COVID-19;
middle-high and high-income countries that never had a universal BCG policy (5 countries) had
about 10 times the number of cases, with a median of 3522.0 cases per million inhabitants. This
difference in the medians between countries was significant (p=0.0283, Wilcoxon rank sum test),
suggesting that broad BCG vaccination along with other measures could help slow the spread of
COVID-19.   However, multivariate analysis of the number of cases (see **Table 2**) revealed that
although the BCG vaccination policy had a negative correlation with the log$_{10}$ number of COVID-
19 reported cases, the factor did not reach significance (p=0.12). The most significant factor
(p=0.00599) in determining the number of cases per million inhabitants was the number of tests
performed per million inhabitants. Conclusions based on the number of deaths per million
inhabitants over 65 years of age are more robust than conclusions based on the number of cases
per million inhabitants as they do not depend on the number of tests performed.

	Estimate	Standard Error	T statistic	P value
Intercept	5.9208	2.7057	2.1883	0.039559
Number tests per million people	1.46e-05	4.8013e-06	3.0408	0.005999
Log10(days since first case)	-2.8594	1.4047	-2.0356	0.054015
Log10(days from first case To school closing + 2)	0.3817	0.30576	1.2484	0.22502
Median life expectancy (years)	0.02121	0.021834	0.97144	0.34189
Median income(US dollars)	1.1029e-05	8.6378e-06	1.2769	0.21497
BCG policy (1 current, 0 never)	-0.42549	0.26387	-1.6125	0.12111

Table 2. Multivariate analysis of log10(number of cases per million inhabitants). Number of
observations: 29, Error degrees of freedom: 22

Root Mean Squared Error: 0.368

R-squared: 0.678, Adjusted R-Squared 0.591

F-statistic vs. constant model: 7.73, p-value = 0.000148

Discussion

We have shown that differences in mortality produced by COVID-19 across countries are
correlated with a country's BCG vaccination policy. We centered our analysis in middle-high and
high-income countries because they constitute a relatively homogenous group of countries and
under the assumption that reported COVID-19 statistics from these countries would be more
reliable than statistics from less wealthy countries. We performed multivariate analysis on those
countries that controlled for the age distribution of the population, income per capita, stage of the
epidemic of a country, and quality of the medical care. We also included as covariates factors
related to a country's response to COVID-19. These other factors did not explain the lower
mortality in countries with mandatory BCG vaccination. The correlation was confirmed using an
independent measure, excess deaths, from a smaller dataset from Europe.

Although these confounders did not explain the observed trend, other unaddressed factors could
be the causal link between BCG vaccination and reduced mortality. This includes complex factors
such as the genetic makeup of the population, as well as environmental factors. Therefore, our
correlational study should not be the basis for changes in clinical practice nor public health policies.
Those changes, if warranted, should be based on the results of well-controlled clinical trials that
could demonstrate a causal relationship between BCG vaccination and COVID-19 outcomes. At
this   time,   there   are   currently   14   clinical   trials   registered   at   the   NIH

(https://clinicaltrials.gov/ct2/results?cond=covid&term=bcg&cntry=&state=&city=&dist=)   exploring
this possibility.

We did not find a correlation between early implementation of a universal vaccination policy and
reduced mortality in middle-high and-high income countries. We did not include in our analysis
middle-low or low-income countries because our database might not capture the exact BCG
vaccination policies of former European colonies in Asia and Africa during colonial times. For
instance, BCG Atlas lists for Vietnam a start date of 1985 for universal BCG vaccination, but there
are reports of widespread use of BCG vaccination during the French colonial period (Monnais
2006). Interestingly, Vietnam with a population of 95 million has reported zero fatalities due to
COVID-19.

Although COVID-19 deaths are concentrated in the elderly population(Zhou et al. 2020), recent
BCG vaccination in children might play a role in reducing mortality by decreasing the transmission
of the disease to the vulnerable population. Asymptomatic children in Germany(Jones et al., n.d.),
where BCG is not applied to children since 1998, had viral loads that were as high as adults,
which suggests that unvaccinated children may be as infectious as adults. A reduction of the viral
load in children might act as a plausible mechanism by which BCG vaccination in children could
reduce infections and mortality in a country.

Looking at an individual's BCG vaccination history offers the possibility to perform better
epidemiological studies than country-wide comparisons to determine vaccination schedules and
strains that might optimize protection against COVID-19 as well as possible mechanisms of BCG
protection. For instance, a recent study determined that BCG vaccination in childhood did not
reduce COVID-19 infection in young adults, a population with reduced mortality due to COVID-19
(Hamiel, Kozer, and Youngster 2020). Patients with bladder cancer that have been treated with
multiple doses of BCG (Alexandroff et al. 1999) constitute a population that is worth studying for
possible protective effects against COVID-19.

If BCG were protective for COVID-19, why did COVID-19 spread in China despite having a
universal BCG policy since the 1950s? During the Cultural Revolution (1966-1976), tuberculosis
prevention and treatment agencies were disbanded and weakened("Development and
Expectation of Tuberculosis Service System in China" n.d.). We speculate that this could have
created a pool of potential hosts that would be affected by and spread COVID-19. Currently,
however, the situation in China seems to be improving.

Our data suggest that BCG vaccination is correlated with reduced mortality associated with
COVID-19. However, there is still not proof that BCG inoculation at old age would boost defenses
in elderly humans against COVID-19, but it seems to do so in Guinea pigs against *M. tuberculosis*
(Komine-Aizawa et al. 2010).

BCG vaccination has been shown to produce broad protection against viral infections and
sepsis(Moorlag et al. 2019), raising the possibility that the protective effect of BCG may not be
directly related to actions on COVID-19 but associated co-occurring infections or sepsis. However,
we also found that BCG vaccination correlated with a reduction in the number of COVID-19
reported infections in a country suggesting that BCG might confer some protection specifically
against COVID-19. The broad use of the BCG vaccine across a population could reduce the
number of carriers, and combined with other measures could act to slow down or stop the spread
of COVID-19. However, an alternative explanation would be that COVID-19 in a person with BCG

vaccination would have a milder presentation, reducing the possibility that such a case would be
even detected in the first place.

Different countries use different BCG vaccination schedules(Zwerling et al. 2011), as well as
different strains of the bacteria (Horwitz et al. 2009). We have not divided the data depending on
the strain used to determine whether different strains are better at stopping the spread of infection,
as well as reducing mortality in the elderly population, as suggested by Jun Sato("If I Were North
American/West European/Australian, I Would Take BCG Vaccination Now against the Novel
Coronavirus Pandemic." n.d.). As each country might have used the same strain for the whole
population, the difference in strains for different purposes should be gathered in randomized
control trials with different subjects from the same population tested with different strains.

Randomized controlled trials using BCG vaccination are required to determine how fast an
immune response develops that protects against COVID-19. BCG is generally innocuous with the
main side effect of inflammation at the site of injection. However, BCG is contraindicated in
immunocompromised people as well as pregnant women("Fact Sheets | Infection Control &
Prevention | Fact Sheet - BCG Vaccine | TB | CDC" n.d.), so care should be taken when applying
this possible intervention for COVID-19.

Acknowledgments

The authors would like to thank Raddy Ramos, Randy Stout, Isaac Kurtzer, Martin Gerdes, Kurt
Amsler, Brian Harper, Madhukar Pai, Emily MacLean, Emma Risson and students, postdocs and
faculty at the Immunology Institute of the Icahn School of Medicine, Mount Sinai for comments on
the manuscript. We would like to thank Kashif Zafar for highlighting the widespread use of BCG
vaccination by colonial powers in Asia and Africa between 1920s to 1950s, a critical piece of
information.

References

Alexandroff, Anton B., Andrew M. Jackson, Michael A. O'Donnell, and Keith James. 1999. "BCG
Immunotherapy of Bladder Cancer: 20 Years On." *Lancet*. Elsevier Limited.
https://doi.org/10.1016/S0140-6736(98)07422-4.

Aronson, Naomi E., Mathuram Santosham, George W. Comstock, Robin S. Howard, Lawrence
H. Moulton, Everett R. Rhoades, and Lee H. Harrison. 2004. "Long-Term Efficacy of BCG
Vaccine in American Indians and Alaska Natives: A 60-Year Follow-up Study." *Journal of
the American Medical Association*. JAMA. https://doi.org/10.1001/jama.291.17.2086.

Berg, Martha K, Qinggang Yu, Cristina E Salvador, Irene Melani, and Shinobu Kitayama. 2020.
"Mandated Bacillus Calmette-Guérin (BCG) Vaccination Predicts Flattened Curves for the
Spread of COVID-19." https://doi.org/10.1101/2020.04.05.20054163.

Cao, Qing, Yi Ching Chen, Chyi Liang Chen, and Cheng Hsun Chiu. 2020. "SARS-CoV-2
Infection in Children: Transmission Dynamics and Clinical Characteristics." *Journal of the
Formosan Medical Association*. Elsevier B.V. https://doi.org/10.1016/j.jfma.2020.02.009.

"Development and Expectation of Tuberculosis Service System in China." n.d. Accessed March
24, 2020. http://www.zgflzz.cn/EN/Y2012/V34/I9/557.

"Fact Sheets | Infection Control & Prevention | Fact Sheet - BCG Vaccine | TB | CDC." n.d.

Accessed March 23, 2020.
https://www.cdc.gov/tb/publications/factsheets/prevention/bcg.htm.

Hamiel, Uri, Eran Kozer, and Ilan Youngster. 2020. "SARS-CoV-2 Rates in BCG-Vaccinated and Unvaccinated Young Adults." *JAMA*, May. https://doi.org/10.1001/jama.2020.8189.

Horwitz, Marcus A., Günter Harth, Barbara Jane Dillon, and Saša Masleša-Galić. 2009. "Commonly Administered BCG Strains Including an Evolutionarily Early Strain and Evolutionarily Late Strains of Disparate Genealogy Induce Comparable Protective Immunity against Tuberculosis." *Vaccine* 27 (3): 441–45. https://doi.org/10.1016/j.vaccine.2008.10.058.

"If I Were North American/West European/Australian, I Would Take BCG Vaccination Now against the Novel Coronavirus Pandemic." n.d. Accessed May 31, 2020. https://www.jsatonotes.com/2020/03/if-i-were-north-americaneuropeanaustral.html.

"Japan Was Expecting a Coronavirus Explosion. Where Is It? | The Japan Times." n.d. Accessed March 23, 2020. https://www.japantimes.co.jp/news/2020/03/20/national/coronavirus-explosion-expected-japan/#.XnllWahKjIU.

Jones, Terry C, Barbara Mühlemann, Talitha Veith, Marta Zuchowski, Jörg Hofmann, Angela Stein, Anke Edelmann, Victor Max Corman, Christian Drosten, and Professor Christian Drosten. n.d. "An Analysis of SARS-CoV-2 Viral Load by Patient Age."

Kleinnijenhuis, Johanneke, Jessica Quintin, Frank Preijers, Christine Stabell Benn, Leo A.B. Joosten, Cor Jacobs, Joke Van Loenhout, et al. 2014. "Long-Lasting Effects of Bcg Vaccination on Both Heterologous Th1/Th17 Responses and Innate Trained Immunity." *Journal of Innate Immunity* 6 (2): 152–58. https://doi.org/10.1159/000355628.

Komine-Aizawa, Shihoko, Toshio Yamazaki, Tsuyoshi Yamazaki, Shin Ichiro Hattori, Yuji Miyamoto, Naoki Yamamoto, Shinji Haga, et al. 2010. "Influence of Advanced Age on Mycobacterium Bovis BCG Vaccination in Guinea Pigs Aerogenically Infected with Mycobacterium Tuberculosis." *Clinical and Vaccine Immunology* 17 (10): 1500–1506. https://doi.org/10.1128/CVI.00190-10.

Kristensen, I., P. Aaby, and H. Jensen. 2000. "Routine Vaccinations and Child Survival: Follow up Study in Guinea-Bissau, West Africa." *British Medical Journal* 321 (7274): 1435–39. https://doi.org/10.1136/bmj.321.7274.1435.

Miller, Aaron, Mac Josh Reandelar, Kimberly Fasciglione, Violeta Roumenova, Yan Li, and Gonzalo H Otazu. 2020. "Correlation between Universal BCG Vaccination Policy and Reduced Morbidity and Mortality for COVID-19: An Epidemiological Study." *MedRxiv*, March, 2020.03.24.20042937. https://doi.org/10.1101/2020.03.24.20042937.

Monnais, Laurence. 2006. "PREVENTIVE MEDICINE AND 'MISSION CIVILISATRICE' Uses of the BCG Vaccine in French Colonial Vietnam between the Two World Wars." Vol. 2.

Moorlag, S. J.C.F.M., R. J.W. Arts, R. van Crevel, and M. G. Netea. 2019. "Non-Specific Effects of BCG Vaccine on Viral Infections." *Clinical Microbiology and Infection*. Elsevier B.V. https://doi.org/10.1016/j.cmi.2019.04.020.

Netea, M. G., L. A. B. Joosten, E. Latz, K. H. G. Mills, G. Natoli, H. G. Stunnenberg, L. A. J. ONeill, and R. J. Xavier. 2016. "Trained Immunity: A Program of Innate Immune Memory in Health and Disease." *Science* 352 (6284): aaf1098–aaf1098.

https://doi.org/10.1126/science.aaf1098.

Shet, Anita, Debashree Ray, Neelika Malavige, Mathuram Santosham, and Naor Bar-Zeev. n.d.
"Differential COVID-19-Attributable Mortality and BCG Vaccine Use in Countries."
Accessed June 5, 2020. https://doi.org/10.1101/2020.04.01.20049478.

Sterne, J. A.C., L. C. Rodrigues, and I. N. Guedes. 1998. "Does the Efficacy of BCG Decline
with Time since Vaccination?" *International Journal of Tuberculosis and Lung Disease* 2
(3): 200–207.

Usher, Nicholas T., Suyoung Chang, Robin S. Howard, Adriana Martinez, Lee H. Harrison,
Mathuram Santosham, and Naomi E. Aronson. 2019. "Association of BCG Vaccination in
Childhood With Subsequent Cancer Diagnoses: A 60-Year Follow-up of a Clinical Trial."
JAMA Network Open 2 (9): e1912014.
https://doi.org/10.1001/jamanetworkopen.2019.12014.

"WHO | Tuberculosis." n.d. Accessed March 23, 2020.
https://www.who.int/immunization/diseases/tuberculosis/en/.

Zhou, Fei, Ting Yu, Ronghui Du, Guohui Fan, Ying Liu, Zhibo Liu, Jie Xiang, et al. 2020.
"Articles Clinical Course and Risk Factors for Mortality of Adult Inpatients with COVID-19 in
Wuhan , China : A Retrospective Cohort Study." *The Lancet* 6736 (20): 1–9.
https://doi.org/10.1016/S0140-6736(20)30566-3.

Zwerling, Alice, Marcel A. Behr, Aman Verma, Timothy F. Brewer, Dick Menzies, and Madhukar
Pai. 2011. "The BCG World Atlas: A Database of Global BCG Vaccination Policies and
Practices." *PLoS Medicine* 8 (3). https://doi.org/10.1371/journal.pmed.1001012.

Figure 1: Higher death rates were presented in countries that never implemented a universal BCG vaccination policy. Bars indicate the median value of a given group of countries.

Figure 2: Date of onset of the epidemic does not explain higher number of deaths in countries
395 that did not establish a universal BCG policy.

400 **Figure 3**: Higher excess deaths were presented in European countries that never implemented
401 a universal BCG vaccination policy. Bars indicate the median value of a given group of countries.

Figure 4: Earlier date of the start of vaccination does not correlate with reduced mortality . Left panel correspond to upper middle income and high income countries with current universal BCG vaccination policy. The right panel corresponds to countries that do not have a current universal vaccination policy.

Figure 5: Higher number of COVID-19 cases were presented in countries that never implemented a universal BCG vaccination policy. Bars indicate the median value of a given group of countries.

438