Clinical features and outcomes of 197 adult discharged patients with COIVD-19 in Yichang, Hubei

Fating Zhou[1,2*], Xiaogang Yu[1,2], Xiaowei Tong, Rong Zhang[1,2]

1.The First College of Clinical Medical Science, China Three Gorges University, Yichang, Hubei, 443003, China

2. Emergency Department, Yichang Central People's Hospital, Yichang, Hubei, 443003, China

3. Yichang third People's Hospital, Yichang, Hubei, 443003, China

Correspondence to: Rong Zhang, *E*-mail: 13972046539@ 163.com

Abstract Purpose To investigate the epidemiology an clinical feactures of discharged adult patients with COIVD-19 in Yicang. **Method** The retrospective study recruited 197 cases of COIVD-19 discharged from Yichang Central People's Hospital and Yichang Third People's Hospital from Jan 17 to Feb 26, 2020. All cases were confirmed by real-time RT-PCR or chest computer tomography (CT). The survivors were followed up untill March 4,2020. Clinical data, including demographic charateristic, presentation, underlying diseas, exposure history, laboratory examination, radiology and prognosis were enrolled and analysed by SPSS 19.0 software. **Results** There were 197 adult discharged patients with COIVD-19 in this study. Statistical analysis indicated that the average age was 55.94 years, and femal patients were 99(50.3%).Those patients mainly resided in urban with exposure history in 2 weeks, while 7 medical staffs were infected. Fever (77.6%%), cough (43.6%) and weakness (14.7%) were the common symptoms. Leucocyetes were mainly normal or decreased in 185 patients (92.9%), both lymphocytes and eosinophils were below normal range, the ratios were 56.9% and 50.3%, repectively. On the contrary, lactate dehydrogenases raised in 65 patients. C-reactive protein (72.4%) elevated in the most of patients. The sensitivity of RT-PCR was 63.5%. Chest CT indicated that bilateral patchy shadows (69.0%) were the most common imaging manifestations.169 (85.8%) patients recovered and transferred to a designated hospital for obeservation, and the others (14.2%) turned worst and died of acute respirtory failure. **Conclusion** COIVD-19 infection with highly contagious have become a life-threaten public healthy problem, the sensitivity of RT-PCR was limited. Chest CT scan was recommended for the suspected patients. Furthermore, lymphocytopenia and eosnophils declining without leukocytosis may be considered as a useful evidence for the diagnosis.

Keywords: Coronvirus infection, Acute respirtory distress syndrome, pneumonia, COVID-19

Introduction

Since several cases of 2019 novel coronavirus (2019-nCoV) pneumonia was reported in Wuhan [1].This disease was identified by the Chinase Center for Disease Control and Prevention(CDC) On Janurary 3, 2020. And then was named as COVID-19 by WHO [2]. At the end of Juanuary, this disease spreaded rapidly to other cities of adjacent to Wuhan in Hubei province, including Huanggang, Xiaogan, Jingzhou and Yichang. Yichang, the 2[nd] lagerst city with Three Gorges Dam in Hubei province, was about 300 kilometers far away from Wuhan, and many residents who worked or studied at Wuhan, returned to hometown for Spring Festival. By March 4, 2020, official reported there was 931 confirmed cases with COVID-19 in Yichang.

Similar to the server acute respiratory syndrome (SARS) and Middle East respiratory (MERS),

2019-nCoV can also cause serve acute respiratory syndrome [3]. Numerous patients can be cured effectively at the early stage, but some patients with COVID-19 may develop pulmonary oedema, ARDS, or multiple organ failure, and even died abruptly [4]. Currently, researches about epidemilogy and clinical fearures of COVID-19 focused on the hospitalized patients [5], but the discharged patients was still scarely investigate. Therefore, this study was aimed to clarify the epidemeiology and clinical features of discharged patients with COVID-19.

Methods

This study was approved by the Ethics Committee of Yichang Central people's hospital. All adult discharged patients with COVID-19 from Yichang Central People's Hospital and The third People's Hospital from Jan 17 to Feb 26, 2020. Those patients with epedimiology history were confirmed by chest CT scans or RT-PCR assays. Throat-swab or broncholveolr lvge fluid sample were collected from all the suspected patients at admission, and RT-PCR assays were perfomed at clinical laboratory. According to the fifih edition expert conference by Chinese healhty commission, except being indetified by RT-PCR, those suspected patients presenting with lung imaging fearure also regard as clinicially diagonsis cases in Hubei province. ARDS was identified in accordance with the Berlin definition [6], and acute kidney injury was verified according to Improve Global Outcome definition [7]. If the serum levels of cardiac biomarkers ascended to the 99th percentile upper reference limit or new abnormalities were displayed in echocardiography or electrocardiography (ECG), the patient was regarded as acute cardiac injury. The discharged criteria included that symptom disappeared and the lesions eliminated evidently in the chest CT. Furthermore, RT-PCR was negative 2 consecutive beyond 24 hours. After discharged from hosptial, the patients went to the designated hospital for observeration. All patients except deaths were monitored up to March 4, 2020.

Statistical Analysis

Clinical datas, including date, ages, occupation,location, presentation, underlying diseas, laboratory test, RT-PCR ，chest CT, treatment and outcome were collected from electronic medical record(EMR) . If data were missing, we collect data by direct communication with attending doctor. Then the data were inputted and analyzed by SPSS 19.0 software. Descriptive statistics were presented as median with interquartile range or frequency with percentage.

Results

Epidemiology of COVID-19 patients inYichang

A total of 197 adult discharged patients with COVID-19 were recruited in Yichang Central People's Hospital（101, 51.3%）and Yichang third people's hospital（86，48.7%）.The first cases was a 65-years old woman from Wuhan on Januray 15, 2020. The quantity of admitted patients increased sharply on the 15 days later, ascended to the peak on the Februray 2, and decreased obviously on February 7 (See the Figure1).

The average age of dischaged patients was 55.94±18.83 (IQR, 40.25-70.75) years, and percentage of 49.7 (98) was femal. Those cases were mainly occurred in the urban (178, 90.4%), espiciallly Xiling Distric, because numerous patients were employees of China Gezhouba Group Corporation (CCGC) returning from Wuhan. Except for seven medical staffs, most of them were employees (58, 29.4%), retirees (55, 27.4%), and agriculture workers (29, 14.7%). A total of 187 patients were clustered and

had clear explosure histories. Among of them, 75(38.7%) patients residesed in the community of COVID-19 patients, 71(36.8%) patients had ever contacted with relaties or friends from Wuhan, 41(20.8%) patients had a trip to Wuhan.

Clincal feactures of adult discharged patients with COVID-19

Fever (153, 77.6%) was the most common presentations, following cough (89, 45.2%) and weakness (29, 14.7%), respectively. Other less common symptoms were sore throat (11, 5.6%), shortness of breath (10, 5.1%), and muscle ache (9, 4.6%, Table 2). However, 3 patients without any complaints were detected by chest CT scan. Among all those patients, the average days from exposure to sympton onset was 6.14±9.27 (IQR, 2-7) days. The longest incubation period was 55 days. 48 patients were accompanyed with underlaying illnes, including cardiovascular (24.4%) and cerebrovascular diseases (4.6%), endocrine system diseases (9.1%), chronic renal diseas (1.5%), maglinant tumor(1.5%), and digestive system diseases (1.0%), nervous system diseases. Considering vital sign and CT manifestations, 56 patient were admited to intensive cure unite （ICU）, and 141 patients were transferred to general isolation wards.

Laboratory examinations of adult discharged patients with COVID-19

The laboratory exmamination was underwent after admission, the blood routine results revealed that monocytes and platelet kept at normal range in numerous patients, leucocytes were at the normal range in 117 (59.4%) patients, and below the normal range in 66 (33.5%) patients. While lymphocytes decreased in many patients. Intersting, eosinophils displayed a simmilar change tendcy as lymphocytes, it decreased obviously over half of them. 44 (22.3%) patients had different degrees of liver function abnormality, with elevating of aspartate aminotransferase (AST) or alanine aminotransferase (ALT). Total bilirubin (TBIL) was above the normal range in 24(12.2%) patients, and direct bilirubin (DBIL) was above in 10 (5.1%) patient. Most of those patients had a abnormal myocardial zymogram, which indicated that Creatine kinase was above in 10 (5.1%) patients, Creative kinase MB in 19 (9.6%) patients, and Lactate dehydrogenase in 65(33.0%) patients. For the coagulation tests, Prothrombin time (PT) and activated partial thromboplastin time (APTT) were at the normal range in the majority of patients, while fib was above the normal range in the 78 (39.6%) patients, D-dimmer was above the normal range in the 99 (47.2%) patients. 65 (33.0%) patients had different degrees of renal function damage, with elevating blood urea nitrogen or creatin. Blood gas analysis was examined in 69 cases, and the result demonstrated that PaO_2 was decreased in the 28(40.5%) patients. Regarding the infection index, procalcitonin mainly kept at the normal range. On the contrary, c-reative protein (CRP), erythrocyte sedimentation rate (ESR) and interleukin-6 (IL-6) increased in most of patients.

Although RT-PCR was regarded the most accurate examinations for COVID-19, the sensitivity was limited. RT-PCR postive rate was 63.5%, the first test was 46.7%, and the second test was 11.2%.Comparing with PCR , almost all patients had been discover lesions in the chest CT scans, 32 (15.3%)patients displayed unilateral pneumonia, 136 (69.0%)patients displayed bilateral pneumonia, and only 25(12.7%) patients displayed multiple monttling and ground-glass opcity(Table 4,Figure 2).

Treatment and outcome of adult discharged patients with COVID-19

To avoid to hypoxemia and acute respiratory distress, 147(76.6%) patients received oxygen by nasal cathete, 34(15.3%) patients received oxygen by mask, and 16 (8.1%) patients were

treated with invasive vetilation in the ICU. Common complication among the 197 discharged patients included acute liver injury, septic shock, acute myocardiall injury, acute renal injury and ARDF. Most patients were treated with antiviral drugs (Oseltamivir), 151(76.6%) patients received antibiotics(moxifloxacin and cefperazone-sulbactam), and several patients were treated with glucocorticoids (75, 38.1%) and immunoglobulin(52, 26.9%). Additionally, 81(41.1%) patients were treated with Chinese tradditional medicine. By the end of Febururay 26, 28 (14.2%) patients had died, and 169(85.2%) patient was discharged and transferred to designated hosptial for observation. The average of hospital stay was 15.08 days (IQR, 4-24, Table 4).

Disscussion

Coronavirus was a large RNA virus, including SARS-CoV, MERS-CoV, HCoV-OC43, HCoV-229E, HCoV-NL63 and HCoV-HKU1. SARS-CoV and MERS-CoV may induce sever repiratory syndrome, but the remains only caused mild upper repiratoy infectious. 2019-nCoV was a new subtype, with high incidence and rapid infection, it has spreaded widely in China and abroad, involving Italy, German, Korean, Iran, and Japan [8-9]. At present, there was no vaccine and antiviral drugs against this disease. Therefore, Extensive studies of clinical features on COVID-19 have not only a better understanding of coronavirus transimission, but have also been key to detect quickly and accurately.

COVID-19 diagnosis was based on exposures, symptoms, laboratory tests, chest CT and RT-PCR. In the early pahase of the epidemic, almost all patients worked at or lived around the Huanan Seafood Whole Market in Wuhan [10]. With COVID-19 spreaded rapidly in Hubei province, Wuhan-related exposures was regarded as a key criterion. Previous retrospective research demonstrated that 86% cases had a Wuhan-related exposures history [11]. Similarly, our study revealed most of infected patients lived around Xiling strict, because of many residents working at China Gezhouba Group Corporation, returning from Wuhan. The common symptoms of COVID-19 patients were fever, fatigue and cough, some patients may present with abdominal symptoms [12]. While a few patients without any symptoms were detected accidently by chest CT [11]. Thus, it is difficult to suspect the disease without auxiliary examinations.

The significant of laboratory examintions was blood routine, the results suggested leucocyetes were mainly normal or decreased, and lymphocytes reduced evidently in many COVID-19 patients [13]. A recent research indicated that lymphocytes,especially $CD3^+$ lymphocytes and $CD45^+$ lymphocytes, decreased significantly in compasrion with non-COVID patients, while there was no significantly difference in quantities and proportions of neutrophils and monocytes[14].Results of this analysis showed lymphocytes and eostinophils decreased obviously without leukocytosis in most patients, and Neutrophils,monocytes and platelet mianly kept at normal range,it was consistent with previous reports. In term of chest CT, the typical image features were periphearl, subpleural ground glass opacities in the lower lobes. However, a study of CT mianifestions demonstrated that it was approximately 14% COVID-19 patients with atypical CT features [15]. Real time RT-PCR was considered a standard assesment for COVID-19，but the sensitivity was lower than chest CT scans, RT-PCR was 71%, chest CT scans reached 98%[16].Thus, it was essential to perform chest CT for detecting the suspected patients.

As described before, there was currently no clinical approved antiviral drugs and vanccine

for coronavirus infections except for supportive treatment [17]. Immunoglobulin may enhance the ability of anti-infection in severely ill patient, and steroids are considered in those patients with ARDS. Oxygen inhalation was important to avoid to ARDS for the COVID-19 paitents with hypoxemia. In this study, 76.6% patients receivd oxygen by nasal cathete, and 90% received antiviral therapy, and 38.1% received glucocorticoids. CDC reported that the case-fatality rate (CFR) of COVID-19 the overall case--fatality rate (CFR) of COVID-19 was 2.3% in confirmed cases, and CFR among critical cases may reach 49.0% [11]. Nevertheless, most of patients were still treated in the hospital. Anoter retrospective study suggest the CFR was about 11% [13]. Statistical analysis of this study found that the CFR was 14.2%, and most of them were over 70 year-old patients accompanying with chronic illness.In the past one month, numerous effective measures, including blocking cities and roads, closing supermaket, monitoring temperature, building a new hosptial and seting up medical assistance team, have been carried out to controlling infection source, preventting transmission and reducing death.Until now, there was no new confirmed patients for 13 days,and hospitalized patients decreased sharply.

In conclusion, this retrospective study, based on clinical data, revealed that COIVD-19 mainly infected in older people with comorbidities, and may cause acute severe respiratory syndrome.Chest CT was key to screen those suspected patients, and lymphocytopenia and eosnophils declining without leukocytosis may be regarded as an novel evidence in the dignosis of COVID-19.

Conflicts of interest

The authors have declared that no conflict of interest exists.

Funding

No funding is provided with this study

Acknowledgment

None.

Reference

1. Li X, Wang W, Zhao X, et al. Transmission dynamics and evolutionary history of 2019-nCoV[J]. Journal of Medical Virology, 2020:1-9. https://doi.org/10.1002/jmv.25701.

2. World Health Organization. Clinical management of severe acute respiratory infection when novel coronavirus (nCoV) infection is suspected: interim guidance. Published January 28, 2020. Accessed January 31, 2020. https://www. who.int/publications-detail/clinical-management-of-severe-acute-respiratory-infection-when-novel-coronavirus-(ncov)-infection-is-suspected.

3. Graham R L, Donaldson E F, Baric R S. A decade after SARS: strategies for controlling emerging coronaviruses[J]. Nature Reviews Microbiology, 2013, 11(12): 836-848. https://doi.org/oi:10.1038/nrmicro3143.

4. Xu Z, Shi L, Wang Y, et al. Pathological findings of COVID-19 associated with acute respiratory distress syndrome[J]. The Lancet Respiratory Medicine, 2020. https://doi.org/10.1016/S2213-2600(20)30076-X.

5. Wang D, Hu B, Hu C, et al. Clinical characteristics of 138 hospitalized patients with 2019 novel coronavirus–infected pneumonia in Wuhan, China[J]. Jama, 2020. https://doi.org/10.1001/jama.2020.1585.

6. Force A D T, Ranieri V M, Rubenfeld G D, et al. Acute respiratory distress syndrome[J]. Jama,

2012, 307(23): 2526-2533. https://doi.org/10.1001/jama.2012.5669.

7. Stevens P E, Levin A. Evaluation and management of chronic kidney disease: synopsis of the kidney disease: improving global outcomes 2012 clinical practice guideline[J]. Annals of internal medicine, 2013, 158(11): 825-830. https://doi.org/10.7326/0003-4819-158-11-201306040-00007.

8. Rothe C, Schunk M, Sothmann P, et al. Transmission of 2019-nCoV infection from an asymptomatic contact in Germany[J]. New England Journal of Medicine, 2020. https://doi.org/110.1056/NEJMc2001468.

9. Ki M. Epidemiologic characteristics of early cases with 2019-nCoV disease in Republic of Korea[J]. Epidemiology and health, 2020: e2020007. https://doi.org/10.4178/epih.e2020007.

10. Huang C, Wang Y, Li X, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China[J]. The Lancet, 2020, 395(10223): 497-506. https://doi.org/10.1016/S0140-6736(20)30183-5.

11. Wu Z, McGoogan J M. Characteristics of and important lessons from the coronavirus disease 2019 (COVID-19) outbreak in China: summary of a report of 72 314 cases from the Chinese Center for Disease Control and Prevention[J]. Jama, 2020. https://doi.org/110.1001/jama.2020.2648.

12. Zhang H, Kang Z, Gong H, et al. The digestive system is a potential route of 2019-nCov infection: a bioinformatics analysis based on single-cell transcriptomes[J]. BioRxiv, 2020. https://doi.org/110.1101/2020.01.30. 927806.

13. Chen N, Zhou M, Dong X, et al. Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study[J]. The Lancet, 2020, 395(10223): 507-513. https://doi.org/10.1016/S0140-6736(20)30211-7

14. Zheng Y, Huang Z, Ying G, et al. Comparative study of the lymphocyte change between COVID-19 and non-COVID-19 pneumonia cases suggesting uncontrolled inflammation might not be the main reason of tissue injury[J]. medRxiv, 2020. https://doi.org/10.1101/2020.02.19.20024885.

15. Pan Y, Guan H, Zhou S, et al. Initial CT findings and temporal changes in patients with the novel coronavirus pneumonia (2019-nCoV): a study of 63 patients in Wuhan, China[J]. European radiology, 2020: 1-4. https://doi.org/10.1007/s00330-020-06731-x.

16. Fang Y, Zhang H, Xie J, et al. Sensitivity of chest CT for COVID-19: comparison to RT-PCR[J]. Radiology, 2020: 200432.http://doi.org/10.1148/radiol.2020200432.

17. Zumla A, Chan J F W, Azhar E I, et al. Coronaviruses—drug discovery and therapeutic options[J]. Nature reviews Drug discovery, 2016, 15(5): 327-347. https://doi.org/110.1038/nrd.2015.37.

Fig 1. The distribution of admission date and crucial event in Yichang.

	Patients(n=197)
Age,years	
Mean($\overline{X}\pm$SD)	55.94±18.83
Range	18-91
≤30	21(10.7%)
30-60	96(48.7%)
＞60	80(40.6%)
Sex	
Female	98（49.7%）
Male	99（50.3%）
Month	
January	66(33.5%)
February	131(66.5)
Occupation	
Agricultural worker	29(14.7%)
Retired	55(27.9%)
Employee	58(29.4%)
Self-employed	22(11.2%)
Medical staff	7(3.6%)
Others	26(13.2%)
Location	
Rural	19(9.6%)
Urban	178(90.4%)
Exposure history (<2 weeks)	
A trip to Wuhan	41（20.8%）
Contacting with relatives or friends from Wuhan	71（36.8%）
2019-nCov patients in their community	75（38.7%）

Table1. Demographics and baseline characteristics of 197 discharged patients with COVID-19.

.

	Patients(n=197)
Signs and symptoms at adimission	
Fever	153(77.66%)
Cough	89(45.2%)
Weakness	29(14.7%)
Sore throat	11(5.6%)
Shortness of breath	10(5.1%)
Muscle ache	9(4.6%)
Days from exposure to sympom onset	
Means	6.14±9.27
Range	1-55
Chronic medical illnes	
Cardiovascular diseases	48(24.4%)
Diabetes	18(9.1%)
Cerebrovascular diseases	9(4.6%)
Chronic renal diseases	3(1.5%)
Malignant tumor	3(1.5%)
Digestive system diseases	2(1.0%)
Respiratory system diseases	2(1.0%)
Admission to intensive care unit	56(29.4%)

Table2. Clinical characteristics of 197 discharged patients' with COVID-19.

	Average	Normal Range	Increased	Decreased
Blood routine				
Leucocytes	5.79±3.13	4-10 ($\times 10^9$ / L)	14(7.1%)	66(33.5%)
Neutrophils	4.18±3.06	1.8-6.3 ($\times 10^9$ / L)	33(16.8%)	23(11.7%)
Eosinophils	0.09±0.51	0.02-0.52 ($\times 10^9$/ L)	1(5.1%)	99(50.3%)
Lymphocytes	1.13±0.74	1.1-3.2 ($\times 10^9$ /L)	7（3.6%）	112（56.9%）
Monocytes	0.41±0.22	0.1-0.6 ($\times 10^9$/L)	4(2.0%）	31(15.7%)
Platelet	165.35±67.97	100-400 ($\times 10^9$/L)	1(0.5%)	30(15.2%)
Blood chemistry				
Alanine aminotransferase	38.40±44.42	9-50 (U/ L)	36(18.3%)	7(35.5%)
Aspartate aminotransferase	38.84±44.42	15-40 (U/ L)	44(22.3%)	11(5.6%)
Total bilirubin	16.31±13.86	5.1-28 (μmol/L)	24(12.2%)	2(1.0%)
Direct bilirubin	6.15±12.35	0-10.0 (μmol /L)	10(5.1%)	–
Blood urea nitrogen	7.69±9.86	3.1-8.0 (μmol /L)	47(23.9%)	52(26.4%)
Serum creatinine	106.80±150.53	57-97 (μmol /L)	35(17.8%)	53(26.9%)
Myocardial enzymes				
Creatine kinase	133.67±223.48	50-310 (IU/L)	10(5.1%)	39(19.8%)
Creative kinase MB	21.11±57.00	0-25 (IU/ L)	19(9.6%)	–
Lactate dehydrogenase	266.20±153.70	120-250 (IU/L)	65(33.0%)	8(4.1%)
Hydroxybutyrate dehydrogenase	189.96±116.88	95-250 (IU /L)	30(15.2%)	14(7.1%)
Coagulation function				
Prothrombin time	11.18±1.72	10.0-14.5(s)	6(3.0%)	14(7.1%)
Activated partial thromboplastin time	32.66±7.15	20.0-45.0(s)	10(5.1%)	2(1.0%)
Fibrinogen	4.13±2.64	2.00-4.00(g /L)	78(39.6%)	–
D-dimer	2309.36±8217.50	0-500(ng/ mL)	93(47.2%)	–
Blood gas analysis				
pH	7.40±0.10	7.35-7.45	15/69(21.7%)	11/69(15.9%)
PaO$_2$	81.77±37.73	80-100 (mmHg)	4/69(5.8%)	28/69(40.5%
PaCO$_2$	37.73±9.53	35-45 (mmHg)	8/69(11.6%)	25/69(36.2)
Infection related biomarkers				
Procalcitonin	0.46±2.17	0-0.05 (ng/ mL)	60/176(34.1%)	–
C-reactive protein	54.97±58.76	0-10 (mm /h)	80/123(72.4%)	–
Erythrocyte sedimentation rate	32.00±27.50	0-10 (mm/h)	59/82(71.2%)	–
Interleukin-6	112.69±254.68	0-7 (pg/mL)	23/35(65.7%)	–

Table3. Results of laboratory examinations in the 197 adult discharged patients with COVID-19.

Total positive of RT-PCR	63.5%
The first test	46.7%
The Second test	11.2%
The third test	4.1%
Chest CT scan or X-ray	
Abnormities	193(98.0%)
Unilateral pneumonia	32(16.24%)
Bilateral pneumonia	136(69.0%)
Multiple mottling and ground-glass opacity	25 (12.7%)
Treatment	
Oxygen therapy	147(76.6%)
No-invasive	34(15.3%)
invasive	16(8.1%)
Antibiotic	151(76.6%)
Antiviral	179(90.9%)
Glucocorticoids	75(38.1%)
Intravenous immunoglobulin	52(26.9%)
Chinese Traditional medicine	81(41.1%)
Complication	
ARDS	26(13.2%)
Acute renal injury	16(8.1%)
Acute myocardial injury	14(7.1%)
Septic shock	12(6.1%)
Acute liver injury	8(4.1%)
Outcome	
Death	28(14.2%)
Survival	169(85.8%)
Hospital stays (days)	
Means	15.08±17.47
0-7 (days)	79(40.1%)
7-14 (days)	31(15.7%)
>14 (days)	87(44.2%)

Table 4. PCR, chest CT scans and treatment of discharged patients.

a. Chest CT images on 9 days after symptom onset

b. Chest CT images on 19 days after symptom onset

c. Chest CT images on 26 days after symptom onset

Fig2. Chest CT images of a 26-year-old patient with COVID-19. a Chest CT images obtained on February 2, 2020, showed ground glass opacity in both lungs on day 5 after symptom onset. b Images taken on February 12,2020, indicated the lesions increased obviously. C Images taken on February 19, 2020.revealed bilateral ground glass opacity absorbed dramatically, and the patient discharged from hospital and return to the local hospital.

a. Chest CT images on 5 days after symptom onset

b. Chest CT images on 14 days after symptom onset

Fig3. Chest CT images of a 62-year-old patient with COVID-19. a Chest CT images obtained on February 2, 2020, showed ground glass opacity in both lungs on day 5 after symptom onset. b Images taken on February 11,2020, revealed the lesions increased rapidly, and the patient died on February 16,2020.