A single holiday was the turning point of the COVID-19 policy of Israel

Klausner, Ziv PhD[1][†]; Fattal, Eyal PhD[1*][†]; Hirsch, Eitan PhD[2];
Shapira, Shmuel C., MD, MPH[3]

[1] Department of Applied Mathematics, Israel Institute for Biological
Research, Ness-Ziona, Israel
[2] Environmental Sciences Division, Israel Institute for Biological
Research, Ness-Ziona, Israel
[3] Director general, Israel Institute for Biological Research, Ness-Ziona,
Israel

* Correspondence to: Applied Math department, Israel Institute for Biological Research, P.O. Box 19,
Ness-Ziona, 7410001, ISRAEL. Tel.: +972 8 9381794; fax: +972 8 9381432; e-mail: eyalf@iibr.gov.il

† These two authors contributed equally to the article

Abstract

Background:

The impact of COVID-19 has been profound, and the public health challenge seem to be the most serious regarding respiratory viruses since the 1918 H1N1 influenza pandemic. In the absence of effective vaccine or biomedical treatment, the basic rules of public health measures have not changed, namely public distancing.

Methods:

We analyzed epidemiological investigation reports during the first month of the outbreak in Israel. In addition, we present a deterministic compartment model and simulations of several scenarios emphasizing quarantine and isolation policies given their efficiency.

Results:

We identify an abrupt change from controlled epidemic regime to an exponential growth ($R_0 = 2.19$) in light of the actual policy-makers decisions and public behavior in Israel. Our analysis show that before the abrupt change, the new cases trend was due to returning citizens infected abroad. The abrupt change followed a holiday in which social distancing was clearly inefficient and many public gatherings were held. We further discuss three different modeled scenarios of quarantine efficiency: high-, medium-, and low-efficiency.

Conclusions:

Israel early lessons show that there is no allowance to compromise with the directive of social distancing. Even before the onset of the pandemic in Israel, fine-tuned but determined early decisions were taken by policy makers to monitor flight arrivals from Covid-19 affected regions and to limit public gatherings. Our analysis show that one particular holiday has shifted the occurrence curve from controlled

67 regime to exponential growth. Therefore, even a short lapse in public responsiveness

68 can have a dramatic effect.

1. Introduction

Since its emergence the impact of COVID-19 has been profound, and the

public health challenge seem to be the most serious seen in a respiratory virus since

the 1918 H1N1 influenza pandemic (Soper 1919). In this study we present the results

of epidemiological data and modelling of one month since the onset of the outbreak in

Isreal, addressing public events occurring during this period and the sensitivity to a

number of public health measures focusing on social distancing (quarantine and

isolation).

The epidemiological data studied, consisting of 381 laboratory confirmed

COVID-19 cases, has been obtained from the epidemiological investigation reports

that were released by the Israeli Ministry of Health (Israeli Ministry of Health, 2020).

This allowed us to identify and separate the incoming from abroad (which policy

required to enter quarantine on arrival) to the local cases, thus analyzing the net

contribution of local infections.

In addition, we present an extended deterministic SEIR (Susceptible, Exposed,

Infectious, and Recovered) model to simulate disease outbreak scenarios. In

particular, the model includes quarantine of asymptomatic suspected population

(exposed) and isolation of symptomatic and infectious patients. The model takes into

account the efficiency of the quarantine and isolation measures. We discuss three

different quarantine efficiency scenarios: high-efficiency, medium-efficiency, and

low-efficiency. The resulting analysis from the epidemiological cases data are

discussed in light of public events and compared to model simulations. We analyze

and discuss an abrupt change from controlled epidemic regime to an exponential

growth regime in light of policy makers' decisions and public behavior.

2. Methods

The dynamics of spread of epidemics as well as the quarantine-isolation policy

of Israel was modeled using the SEQIJR model (e.g. Gumel, et al., 2004). This is a

deterministic compartmental model which allows the implicit inclusion of biological

epidemiological phases (including incubation period) as well as governmental

interventions such as quarantine and their actual efficiency of implementation. A

successful a posteriori implementation of this model to the transmission dynamics and

control of the SARS epidemics in Toronto, Hong Kong, Singapore and Beijing is

given in Gumel, et al., 2004. The model consists of a system of 7 dynamical equations

and 15 parameters. For details of the model and its parameters see the online

supplementary information.

The data analyzed in this study was obtained from the epidemiological

investigation reports that were released by the Israeli Ministry of Health (Israeli

Ministry of Health, 2020). From the total of 883 PCR laboratory confirmed COVID-

19 cases we analyzed the 384 cases that were investigated epidemiologically,

allowing us to separate the incoming from abroad (quarantined on arrival by policy)

to local cases. This allowed us to separate the imported cases (travelers arriving from

abroad) from the locally infected cases. The data spans over the first month of the

COVID-19 outbreak in Israel, beginning in February 21[st] 2020 and going until March

20[th]. We further note that during the examined period the number of PCR tests

performed rose daily, reaching 1869 at the end of the period. However, the proportion

of positive tests remained approximately the same, as 7.9% (Israeli Ministry of

Health, 2020).

The incidence curve was modelled as a fit to an exponential growth function

(de Silva, et al., 2009; Zhao, et al., 2020). Several serial interval distributions that

were estimated for COVID-19 were examined (Nishiura, et al., 2020; Ganyani, et al.,

2020; Zhao, et al., 2020).

3. Results

The dynamics of epidemic spread in Israel in terms of daily reported

confirmed cases is shown in fig. 1. There are two distinct regimes in the curve. The

first ended in March 12, 2020 in which the local to travel associated infection ratio

was around 1:3 (up to 1:2.5), i.e., for every three imported cases, there was one local

infection. The second period, began a day later. The first distinct sign for the change

was evident on March 14[th] were the ratio rose to 1:2, then continued to rise on March

17[th] to 1:1.5 and finally, after a month into the epidemic in Israel, on March 21[st] there

were more local infections in Israel than imported cases (the ratio was above 1:1).

March 12[th] is two days after the Jewish holiday of Purim, which this year was

celebrated on March 10[th]. Purim customs include wearing masks and costumes and

holding public celebrations, parades as well as religious gatherings. Because Purim is

also a school holiday, many celebration are held a day or two earlier.

FIG. 1. Accumulated reported confirmed COVID-19 cases in Israel (blue bars) and the ratio of local infections to travel associated infections (orange line)

The effective reproduction number, R_0, was estimated for the daily new cases data of the first month of the COVID-19 epidemic. The lowest estimated R_0 was 2.08 (95%CI 1.93-2.6) for the Gamma distributed serial interval with mean 4.4 and SD 3 days (Zhao, et al., 2020). The highest estimate was 2.37 (95%CI 2.16-2.61) for the Gamma distributed serial interval with mean 5.2 and SD 2.8 days (Ganyani, et al., 2020). The mean R_0 overall the serial interval distributions examined, was 2.19.

The SEQIJR model solutions are characterized by 3 locally stable equilibrium points in parameters space. These refer to the following regimes or types of dynamics of epidemic spread: controlled (decaying), flattened, uncontrolled (baseline SEIR model; further details can be found in the supplementary material). In this study we simulated three scenarios, the first corresponding to the first equilibrium and two corresponding to the second:

155   1. A controlled high-efficiency quarantining (decaying green curve in

156   Fig. 2). This regime is characterized by an early entry of asymptomatic

157   suspected population to home quarantine. Moreover we assume the

158   infectiousness in home quarantine is one sixth compared to free

159   asymptomatic. The efficiency of isolation is 70%.

160   2. A flattened for medium-efficiency quarantining (purple curve in Fig.

161   2). This regime is characterized by a late entry of asymptomatic

162   suspected population to home quarantine. Moreover we assume the

163   infectiousness in home quarantine is one third compared to free

164   asymptomatic. The efficiency of isolation is 70%.

165   3. A flattened for low-efficiency quarantining (light-blue curve in Fig. 2).

166   This regime is characterized by a late entry of asymptomatic suspected

167   population to home quarantine. Moreover we assume the

168   infectiousness in home quarantine is similar to free asymptomatic. The

169   efficiency of isolation is 30%.

As discussed above, the dynamics of epidemic spread in Israel until March 8, 2020

corresponds to the controlled regime characterizes with R_0 of the order of half (green

curve in Fig. 2). On the other hand, after March 8, 2020 the regime corresponds to the

flattened regime with low-efficiency quarantining and $R_0 = 2.19$.

Fɪɢ. 2. Total number of symptomatic cases in Israel from SEQIJR model

simulations for 3 scenarios: green – high-efficiency quarantining, purple – medium-

efficiency quarantining, light blue – low-efficiency quarantining.

4. Discussion

Israel has begun early its mitigation policy against the COVID-19 epidemic.

From the beginning of February, it was decided: to close all border passages (via land,

sea and air) to people that are not Israeli citizens or resident that have visited China

recently; to stop all direct flights from Israel to China and to require Israeli citizens

returning from China or that were in touch with a confirmed COVID-19 patients to a

14 days home quarantine. A short while later, the quarantine requirement was

expanded to other Asian countries. The first COVID-19 patients in Israel were two

passengers that returned from the "Diamond Princess", in 21 and 23 of February.

They entered directly to hospital isolation. A week later, a passenger from Italy was

diagnosed as a COVID-19 patient, and Italy was added to the list of countries that

require 14 days quarantine.

The quarantine-isolation policy succeeded in keeping the rate of daily new

cases small, up until March 9[th]. Then, 4 days later, a sudden change of regime has

occurred, which was manifested by the distinct change of the epidemic curve of Israel

towards exponential growth. March 15[th] marks the first time that the daily new cases

of locally infected were higher than the new travel associated cases.

The timing of this abrupt change is not of coincidence. Regarding the cases

arriving from abroad, a requirement for home quarantine affecting all travelers

arriving began on Match 9[th]. Moreover, during the period between March 9[th] to 11[th] a

Jewish holiday, Purim, was celebrated. This holiday is characterized by big parades

organized by local municipalities, as well as religious gatherings and privately

organized parties. Although authorities cancelled the public parades, many privately

organized and religious crowding had occurred. Regrettably, these drove Israel from a

controlled, mitigated regime to an exponential growth, as described in the results

section. Therefore, despite its intense efforts, Israel's effective R_0 for the period

ending in March 20[th] stands on around 2.19, slightly smaller than the R_0 of 2.6-3.2

estimated for the republic of Korea and Italy, for the period ending in March 5[th].

Such abrupt transition based on social behavior emphasizes the fragility of mitigation

policies.

We therefore emphasis the importance of early fine-tuned but intense

directives for social distancing and isolation measures. This study clearly

demonstrates the lesson learned from the Israeli policy, that even a short lapse in

public responsiveness can have a dramatic effect on public health during pandemic

outbreak.

References

1. Israel Ministry of Health. Information on confirmed patients and COVID-19

press releases. https://govextra.gov.il/ministry-of-health/corona/corona-

virus/spokesman-messages-corona/. Accessed Mar 22, 2020.

2. De Silva UC, Warachit J, Waicharoen S, Chittaganpitch M. 2009. A

preliminary analysis of the epimemiology of Influenza A(H1N1)v virus

infection in Thailand from early outbreak data. *Eurosurveillance*: **14**(31).

3. Nishiura H, Linton NM, Akhmetzhanov AR. 2020. Serial interval of novel

coronavirus (COVID-19) infections. *International Journal of Infectious*

Diseases DOI: 10.1016/j.ijid.2020.02.060

4. Zhao S, Gao D, Zhuang Z, Chong MKC, Cai Y, Ran J, Cao P, Wang K, Lou

Y, Wang W, Yang L, He D. 2020. Estimating the serial interval of the novel

coronavirus disease (COVID-19): A statistical analysis using the public data in

Hong Kong from January 16 to February 15, 2020. preprint is under

consideration at Infectious Diseases of Poverty. DOI: 10.21203/rs.3.rs-

18805/v1

5. Ganyani T, Kremer C, Dongxuan C, Torneri A, Faes C, Wallinga J, Hens N.

Estimating the generation interval for COVID-19 based on symptom onset

data. medRxiv doi:10.1101/2020.03.05.20031815

6. Gumel AB, Ruan S, Day T, Watmough J, Brauer F, van den Driessche P,

Gabrielson D, Bowman C, Alexander ME, Ardal S, Wu J, Sahai BM.

Modelling strategies for controlling SARS outbreaks. Proc Biol Sci. 2004;

271(1554):2223-32. doi:10.1098/rspb.2004.2800

7. Soper GA. 1919. The lessons of the Pandemic. *Science* **49**(1274): 501-506.

DOI: 10.1126/science.49.1274.501

8. Zhuang Z, Zhao S, Lin Q, Cao P, Lou Y, Yang L, Yang S, He D, Xiao L.

2020. Preliminary estimating the reproduction number of the coronavirus

disease (COVID-19) outbreak in Republic of Korea and Italy by 5 March

2020. medRxiv preprint DOI: https://doi.org/10.1101/2020.03.02.20030312

246