Triage assessment of cardiorespiratory risk status based on measurement of the anaerobic threshold, and estimation by activity limitation in patients with pulmonary arteriovenous malformations and hereditary haemorrhagic telangiectasia

[1,2]Saranya Thurairatnam, [3]Filip Gawecki, [4]Timothy Strangeways, [5]Joseph Perks BSc, [6]Vatshalan Santhirapala MB ChB*, [7]Jonathan Myers, [8]Hannah C Tighe BSc; [9]Luke SGE Howard DPhil FRCP, and [10]Claire L. Shovlin PhD FRCP.

[1] School of Medicine, Imperial College London UK, and [2] Hull/York School of Medicine, UK ‡; [3] School of Medicine, Imperial College London UK†; [4] School of Medicine, Imperial College London UK†; [5]Respiratory Medicine, Imperial College Healthcare NHS Trust, London UK; [6]Respiratory Medicine, Imperial College Healthcare NHS Trust, London UK; [7]Respiratory Medicine, Imperial College Healthcare NHS Trust, London UK‡; [6]Imaging, Imperial College Healthcare NHS Trust, London UK; [6]School of Medicine, Imperial College London UK*; [7] Division of Cardiology, Veterans Affairs Palo Alto Health Care System (VAPAHCS) and Stanford University, Palo Alto, CA, [8]Respiratory Medicine, Imperial College Healthcare NHS Trust, London UK; [9]National Pulmonary Hypertension Service,, Imperial College Healthcare NHS Trust, London UK; [10]NHLI Cardiovascular Sciences Imperial College London, UK; VASCERN HHT European Reference Centre and Respiratory Medicine, Imperial College Healthcare NHS Trust, London UK

Corresponding author:
Claire L. Shovlin PhD FRCP, Professor of Practice (Clinical and Molecular Medicine), NHLI Cardiovascular Sciences, Imperial Centre for Translational and Experimental Medicine, Imperial College London, Hammersmith Campus, Du Cane Road, London W12 0NN, UK. c.shovlin@imperial.ac.uk

Short running title   Rare disease triage- pulmonary AVMs and HHT

Abstract- 300 words, Text 2729 words

Key Messages

What is already known:

- Alongside age, pre-existing medical conditions are perceived negatively during triage assessments, particularly if rare, and/or theoretically expected to influence cardiorespiratory risk;

- Anaesthetists use cardiopulmonary exercise testing to categorise patients to higher and lower risk independently to diagnostic labels, but this is not feasible in acute settings;

- Pulmonary arteriovenous malformations are an exemplar of a condition where, due to expected or measured abnormalities (hypoxaemia- low PaO_2 SpO_2), poor physiological capacity might be predicted.

What this study adds

- Neither age nor usual SpO_2 predicted lower/higher risk categories by anaerobic threshold, but haemoglobin-dependent indices of oxygen delivery to the tissues were associated with higher risk, offering opportunities for improvement by attention to anaemia and aerobic conditioning;

- Baseline exercise tolerance may override age and diagnostic labels in triage settings: the 13-point VSAQ Veterans Specific Activity Questionnaire (VSAQ) is suggested as a rapid screening tool for cardiorespiratory risk assessment.

Key words:

anaesthesia, assessment, general, rare disease, respiratory, triage

Abstract

BACKGROUND: Rapid triaging, as in the current COVID-19 pandemic, focuses on age and pre-existing medical conditions. In contrast, preoperative assessments use cardiopulmonary exercise testing (CPET) to categorise patients to higher and lower risk independent of diagnostic labels. Since CPET is not feasible in population-based settings, our aims included evaluation of a triage/screening tool for cardiorespiratory risk.

METHODS: CPET-derived anaerobic thresholds were evaluated retrospectively in 26 patients with pulmonary arteriovenous malformations (AVMs) who represent a challenging group to risk-categorise. Pulmonary AVM-induced hypoxaemia secondary to intrapulmonary right-to-left shunts, anaemia from underlying hereditary haemorrhagic telangiectasia and metabolic equivalents derived from the 13-point Veterans Specific Activity Questionnaire (VSAQ) were evaluated as part of routine clinical care. Pre-planned analyses evaluated associations and modelling of the anaerobic threshold and patient-specific variables.

RESULTS: In the 26 patients (aged 21-77, median 57 years), anaerobic threshold ranged from 7.6-24.5 (median 12.35) $ml.min^{-1}kg^{-1}$ and placed more than half of the patients (15, 57.7%) in the >11 $ml.min^{-1}kg^{-1}$ category suggested as "lower-risk" for intra-abdominal surgeries. Neither age nor baseline SpO_2 predicted anaerobic threshold, or lower/higher risk categories, either alone or in multivariate analyses, despite baseline oxygen saturation (SpO_2) ranging from 79 to 99 (median 92)%, haemoglobin from 108 to 183 (median 156)$g.L^{-1}$. However, lower haemoglobin, and particularly, arterial oxygen content and oxygen pulse were associated with increased cardiorespiratory risk: Modelling a haemoglobin increase of 25$g.L^{-1}$ placed a further 7/26 (26.9%) patients in a lower risk category. For patients completing the VSAQ, derived metabolic equivalents were strongly associated with anaerobic threshold enabling risk evaluations through a simple questionnaire.

CONCLUSIONS: Baseline exercise tolerance may override age and diagnostic labels in triage settings. These data support approaches to risk reduction by aerobic conditioning and attention to anaemia. The VSAQ is suggested as a rapid screening tool for cardiorespiratory risk assessment to implement during triage/screening.

Introduction

Difficult triage decisions need to be made in many clinical settings involving large numbers of critically ill patients, as during the current COVID-19 pandemic. Such decisions are based on factors such as age and pre-existing medical conditions, in addition to acute observations and measurements. With the exception of certain common disease states, there is little evidence regarding associations with specific infections or complication risks. Diagnostic labels are generally linked under a single, negative umbrella of "pre-existing medical conditions". There is particular concern that as for health insurance, lack of familiarity with rare diseases may lead to an inappropriately negative weighting, with no time to redress in an acute triage setting.

In pre-operative assessments, anaesthetists increasingly use cardiopulmonary exercise testing (CPET) to identify patients who may be unable to appropriately respond to increased cardiorespiratory demands of surgery due to reduced cardiorespiratory reserve.[1,2] The CPET-derived measure of anaerobic threshold (AT) of <11 ml.min^{-1} kg^{-1} has been identified in multiple studies and systematic reviews to be associated with adverse outcomes, mortality, and longer lengths of stay in a variety of surgeries, including intra-abdominal and intra-thoracic procedures.[2-4] The AT represents the point where ATP generation cannot be met by mitochondrial metabolism. It is considered a good measure as it reflects oxygen delivery and patient conditioning, and is not dependent on the patient's motivation during exercise.[2]

The rapid assessment tool selected for evaluation was the Veteran's Specific Activity Questionnaire (VSAQ [5]). This is a simple 13-point scale of activities of increasing difficulty whereby the user indicates which activity normally causes them to stop when performed for a period of time.[5] The activities correspond to metabolic equivalents (METs), and numerous studies show a good correlation with AT derived from cardiopulmonary exercise testing [6-8], mortality [9] and postoperative complications [10].

We focussed on one particular rare disease that provides an instructive example of potentially over-

called cardiorespiratory risk. Pulmonary arteriovenous malformations (AVMs) are abnormal vascular

connections between pulmonary arteries and veins, resulting in an anatomic right-to-left shunt.[11]

Patients with pulmonary AVMs can demonstrate pronounced physiological abnormalities, including

significant hypoxaemia, [12-18] increased minute ventilation ($\dot{V}E$) for given increases in CO_2

production ($\dot{V}E/\dot{V}CO_2$ slope) [16,17], high cardiac output states [19], and often iron deficiency and

anaemia due to inadequate replacement of haemorrhagic iron losses from underlying hereditary

haemorrhagic telangiectasia (HHT) [20,21] There is no published guidance on management of

individuals with pulmonary AVMs or HHT undergoing anaesthesia, and each year, our service receives

requests regarding suitability for surgery and insurance, and/or reports that surgery or insurance has

been withheld because of the perceived risks of pulmonary AVMs/HHT.

Our goal was to evaluate commonly used assessment criteria and examine the potential role for a rapid

assessment tool that could distinguish lower risk individuals in an emergency setting, based on usual

cardiorespiratory status. The detailed study aims were to explore which variables may be associated

with cardiorespiratory risk defined by the anaerobic threshold in order to inform triage and develop

approaches to help guide pre-exposure [23] or pre-operative [1-4,10] management. Having recently

applied the VSAQ to observational studies in patients with pulmonary AVMs and HHT, [17,18] we

hypothesised that this could prove to be a useful risk categorising tool for triage purposes across wider

patient groups.

Methods

General patient evaluations

With ethical approvals (Hammersmith and Queen Charlotte's and Chelsea Research Ethics Committee

(LREC 2000/5792), patient indices derived as part of the clinical assessment process in a pulmonary

AVM service at a single centre were examined as described elsewhere [14,15,17,18]. These included

arterial oxygen content (CaO_2) derived from SpO_2 values measured in the erect posture breathing room

air using the established formula:

$$CaO_2 = \frac{1.34 \times Haemoglobin \times SpO_2}{100}$$

Cardiopulmonary exercise tests

Previously reported cardiopulmonary exercise tests in patients with pulmonary AVMs where there had

been striking variability in anaerobic thresholds [16,17] were reanalysed with a focus on triage/pre-

operative assessments methods. Ethical approval had been granted by the National Research Ethics

Service (NRES) Committee South West London REC3: 11/H0803 and NRES London Riverside

Committee:15/L0/0590. Written informed consent had been obtained from all participants. Full

methodological details are provided in [16,17].

Veteran's Specific Activity Questionnaire (VSAQ)

The VSAQ was administered as part of routine clinical care to patients for independent completion,

using a modified version as presented in *Figure 1*. Patient-reported activity limitations in the 13-point

scale were converted to metabolic equivalents (METs) in which 1 MET equals the consumption of 3.5

ml O_2 per kilogram of body weight. Metabolic equivalents (METs) were calculated from the VSAQ as

in the original protocol,[5] and subsequent validations [6-10], by the formula:

$$Predicted\ METs = 4.74 + (0.97 \times VSAQ\ score) - (0.06 \times Age)$$

Data Analysis

Statistical analyses were performed in Microsoft Excel and Stata IC versions 14 and 15 (Statacorp,

Texas). Two-way analyses used Mann Whitney U test and three-way analyses Kruskal Wallis tests.

Prior to data analyses, patients were pre-categorised based on the published anaerobic threshold

delimiter of 11 ml.min^{-1} kg^{-1}.[24] Since the risk categories may change in the future as more evidence

becomes available, patients were also pre categorised above and below the median AT value.[24].

Additionally, to further support the robustness of data analysis, regression analyses were performed

using A and log-transformed AT as the outcome variables (log-transformed AT had a more normal

distribution, data not shown).

Patient and Public Involvement statement

Patients were involved in earlier testing of the VSAQ [17,18] and aspects of design of the CPET

protocols. Focussing of our data towards the triaging of patients was an outcome of inputs from British

patients contacting us in March 2020, focussing on the question "Am I at High Risk?

Results

CPET Participant Demographics

The 26 patients with pulmonary AVMs comprised 16 male, 10 females, and were aged 21-77 (median

57) years. SpO_2 ranged from 79 to 99 (median 92)%, haemoglobin from 108 to 183 (median 156)g.L^{-1},

and body mass index (BMI) from 20 to 35.7 (median 26.1) kg.m^{-2}. Comorbidities were present in 11

patients: three had known asthma or chronic obstructive pulmonary disease (COPD), one had sleep

apnoea, and two had type 2 diabetes mellitus. In addition, three had suffered a previous stroke, transient

ischaemic attack, or venous thromboemboli, one was in atrial fibrillation, one had well controlled

hypertension, one was hypercholesterolaemic, one was significantly depressed, and two had benign

prostatic hypertrophy.

CPET Demographics identify a low risk group

As presented in *Table 1*, based on the established anaerobic threshold delimiter of 11 ml.min^{-1} kg^{-1},

more than half of the cohort with pulmonary AVMs (15/26, 57.7%) were categorised as pre-operative

"low risk", comprising 13 males and 2 females. The low risk group achieved a median 97% of their

predicted maximum work rate compared to the high risk group median of 68% predicted (Table 1).

Similarly, the median peak $\dot{V}O_2$ in the low risk group was 160% of the median in the high risk group.

The CPET-evaluated total oxygen consumption at peak exercise ($\dot{V}O_2$), has also been used for high risk

anaesthetic categorisation, noting that for reliable peak $\dot{V}O_2$ measurements, patients need to meet their

point of maximal exercise. In the current study, all low risk patients identified by anaerobic threshold

were also in a low risk category if defined by peak $VO_2 < 20$ ml.min^{-1}.kg^{-1} [2] (data not shown).

Age and usual SpO$_2$ not associated with cardiorespiratory risk or anaerobic threshold

There was no difference in age between the low and high risk groups categorised by an anaerobic

threshold delimiter of 11 ml.min^{-1} kg^{-1} (low risk mean 52.2 [95% confidence interval CI 43.8, 60.6]

years, versus high risk mean 53.9 [95% CI 42.9, 64.9] years). *Table 1* and *Figure 2A* display the

median values, the interquartile ranges (IQR), and 2 standard deviations. There was also no difference

in age between the low and high risk groups categorised by upper/lower 50th percentiles (lower risk

mean 52.6 [95% CI 43.8, 61.5] years, versus higher risk mean 53.2 [95% CI 43.2, 63.3] years (*Table*

2)). In keeping with this, there was no detectable association between age and the absolute or log-

transformed anaerobic threshold values (*p*-values >0.62, data not shown).

Despite some very low resting SpO$_2$ measurements, there was also no difference in pulse oximetry-

measured oxygen saturation (SpO$_2$) between the low and high risk groups: Categorised by an anaerobic

threshold delimiter of 11 ml.min^{-1} kg^{-1} , the means [95% CI] were 92 [90, 94]% for the low risk group,

versus 91 [86, 95]% for the high risk group (*Table 1*, *Figure 2B*). Categorised by the upper/lower 50th

percentile groups, the respective means [95% CI] were 92.5 [90.3, 94.6]% for lower risk, versus 90.5

[87.1, 94.0]% for higher risk (*Table 2*). In crude and age-adjusted regression there was no detectable

association between SpO$_2$ and either category, or the absolute anaerobic threshold in ml.min^{-1} kg^{-1} or

the log transformed values (*p*-values > 0.48, data not shown).

We concluded that assuming the study had sufficient power, neither age, nor more surprisingly SpO$_2$

(in the setting of a compensated baseline state), in themselves, would be markers of a higher risk state

in the cohort.

Low haemoglobin, arterial oxygen content and oxygen pulse indicative of higher risk status

A different picture emerged when examining markers of oxygen delivery to the tissues, thus confirming

that the study did have sufficient power to be discerning:

First, the higher risk groups had a trend towards lower haemoglobin, although there was some overlap

in confidence intervals. Categorised by an anaerobic threshold delimiter of 11 ml.min^{-1} kg^{-1}, the mean

[95% CI] values were 159 [14.9, 17.0] g.L^{-1} for the low risk group compared to 144 [13.1, 15.6] g.L^{-1}

for the high risk group (*Table 1, Figure 3*). By the upper/lower 50th percentiles, the respective means

[95% CI] were 158 [146, 171] g.L^{-1} for the lower risk group compared to 147 [135, 159] g.L^{-1} for the

higher risk group (*Table 2*).

More strikingly, the higher risk groups had significantly lower arterial oxygen content (CaO$_2$)

representing the oxygen content per unit volume of arterial blood, and calculated based on the oxygen

carriage of 1.34mls per gram of fully saturated haemoglobin: Categorised by an anaerobic threshold

delimiter of 11 ml.min^{-1}.kg^{-1} **,** the mean [95% CI] values were 19.6[18.4, 21.0] ml.dL^{-1} for the low risk

group, but 17.4 [15.8, 19.0] ml.dL^{-1} for the high risk group (*Table 1, Figure 3*). Categorised by the

upper/lower 50th percentiles, the means [95% CI] were 19.6 [18.1, 21.0] ml.L^{-1} for the lower risk group

but only 17.8 [16.4, 19.3] ml.L^{-1} for the higher risk group (*Table 2*).

Similarly, the higher risk group had a significantly lower mean oxygen pulse, representing the amount

of oxygen extracted/ delivered per heart beat: Categorised by an anaerobic threshold delimiter of 11

ml.min^{-1} kg^{-1} , the mean [95% CI] values were 14.4 [11.0, 17.3] ml.beat^{-1} for the low risk group, but 9.7

[6.7, 12.6] ml.beat^{-1} for the high risk group (*Table 1*). Categorised by the upper/lower 50th percentiles,

the mean [95% CI] values were 14.6[12.3, 16.9] ml.L^{-1} for the lower risk group but only 9.7 [7.2, 12.1]

ml.L^{-1} for the higher risk group (*Table 2*).

Anaemia is very common, and readily correctable in clinical practice. We modelled whether increasing

haemoglobin alone might allow patients to move from a high to lower risk category. In multivariate

regression analysis, haemoglobin explained 57.3% of the variance in log-transformed AT (adjusted R^2

0.57). The regression coefficient of 0.053 (95% confidence interval 0.005, 0.100, p=0.031) implied that

for each 1 g.dL^{-1} (10 g./L^{-1}) rise in haemoglobin, the AT would rise by 0.76 ml.min^{-1}.kg^{-1}, and a

haemoglobin rise of 2.5 g.dL^{-1} would increase AT by 1.9 ml.min^{-1}.kg^{-1}, moving 7 (63.6%) of patients

from high to low risk.   In the existing dataset, it was not possible to preselect all patients who would

benefit using single resting demographics (data not shown).

Other measurements that were associated with the higher risk status were higher serum bicarbonate and

higher minute ventilation ($\dot{V}E$) for given increases in CO_2 production ($\dot{V}E$/ $\dot{V}CO_2$ slope, Supplementary

Figure 1). These are not currently amenable to therapeutic correction.

VSAQ Score association with anaerobic threshold and risk categorisation

Having demonstrated that at least half of the patients with the rare pulmonary vascular abnormality

would have their good exercise capacity and lower risk status readily identified were it feasible to

perform CPET, we were conscious that in standard clinical practice, it is impractical to perform CPET

on every patient. Usual activity could however be analysed by the VSAQ. As in data published for other

general population cohorts,[5-9] for pulmonary AVM patients also completing the VSAQ, there was a

good association between the previous CPET-derived AT and METs derived from the VSAQ

(Supplementary Figure 3).

We used the derived relationship between and AT and VSAQ to model the expected cut off by age on

the VSAQ that might indicate an individual in a lower risk category based on the established AT of 11

$ml.min^{-1}.kg^{-1}$. As noted in *Figure 4*, this differs by patient age such that a VSAQ of 8 would be

suggestive of lower risk irrespective of age, whereas older individuals in the "best" New York Heart

Association (NYHA) category [25] could still fall into the higher risk category defined by AT

$<11ml.min^{-1}.kg^{-1}$. In younger individuals, lower risk would be assigned even in the setting of more

limited exercise capacity, (VSAQ 4-7, NYHA II). In other words, the VSAQ provided sufficient

granularity to indicate where age-related physiology would be 'offset' in particularly active older adults,

and where there may be more concern for a much less active younger individual.

Discussion

We demonstrate that high proportions of patients with a label that might be expected to mean "pre-

existing cardio-respiratory condition" do not fall into classical high risk categories when more carefully

evaluated. Furthermore, age and usual resting SpO_2 were not associated with the anaerobic threshold

and therefore cardiorespiratory risk status in continuous, or categorical analyses. However, lower

haemoglobin, and haemoglobin-dependent indices of arterial oxygen content and delivery were

important predictors of lower anaerobic threshold and a higher risk state: our *post hoc* calculations

suggested increasing haemoglobin could have moved many of the high risk group into a lower risk

category. We also demonstrate that a simple patient-based metric, the VSAQ, could allocate patients to

lower and higher cardiorespiratory risk categories as based on the anaerobic threshold of <11 $ml.min^{-1}$

kg^{-1}.

The study numbers are small but notably demonstrated non-overlapping confidence intervals for key

variables of oxygen tissue delivery. The current findings build on substantial previously published data

and analyses on pulmonary AVM [16,17] and general population [5-10] cohorts. Furthermore, if the

American Society of Anaesthetists (ASA) Physical Status Classification System [1] was employed,

most pulmonary AVM patients would not fall into a high risk category because in the absence of other

diseases, individuals with pulmonary AVMs rarely complain of respiratory symptoms.[14,17,22]

Compensatory mechanisms are so effective that in one study, work rate and oxygen consumption on

maximal CPET did not improve following embolisation treatments that obliterated the pulmonary

AVM(s) and improved SpO_2.[16]   However, the issue is how best to capture this good exercise

tolerance. We have previously reported the ease of use with the VSAQ in a pre-assessment clinic.[18]

We have now adjusted to send the VSAQ to patients by email so they can report back the lowest number

at which they needed to stop at a subsequent teleconsultation, thus conveying complex physiological

information in seconds (*Onabanjo et al, manuscript in preparation*). While measurements in acute

settings do not reflect the patient's baseline, usual activity could be captured by the VSAQ either before

or at the time of triage assessment.

Potential mechanisms for the association between lower anaerobic threshold and less successful surgical

outcomes have been put forward, including the suggestion that regular exercise stimulates ischaemic

preconditioning and lessens surgical demand by enabling the body to adjust to ischaemia and better

utilise oxygen. Additionally, endurance exercise has been found to increase mitochondrial mass, which

can therefore delay the start of anaerobic respiration by enhancing the utilisation of oxygen by

mitochondria.[6] "Prehabilitation" or pre-operative exercise therapy has been found to improve post-

operative outcomes in other disease groups and has been proposed to help prepare for COVID-19

infection.[23] Herein we also show that addressing anaemia is likely to be an additional strategy to

reduce cardiorespiratory risk status.

In summary, high proportions of patients with a label that might be expected to mean "pre-existing

cardiorespiratory condition" do not fall into classical high risk categories when more carefully

evaluated. Given the need for appropriate allocation of ward/critical care resources, whether for

surgery, or in infective setting, we suggest the VSAQ offers a cost-effective tool that can be easily

integrated into triages or anaesthetic pre-assessments to assist with rapid evaluation of cardiorespiratory

risk.

DECLARATIONS

ETHICS APPROVAL AND CONSENT TO PARTICIPATE:

- Ethical approvals in place:

(1) CPET Studies: National Research Ethics Service (NRES) Committee South West London REC3:

11/H0803 and NRES London Riverside Committee:15/L0/0590, written informed consent for all

(2) Case Notes review: Hammersmith and Queen Charlotte's and Chelsea Research Ethics Committee

(LREC 2000/5792, approved without requiring consent.

CONSENT FOR PUBLICATION:

- Not applicable

AVAILABILITY OF DATA AND MATERIAL:

- The datasets analysed during the current study are available from the corresponding authors on reasonable request

COMPETING INTERESTS:

- The authors have no competing interest to declare

FUNDING :

-Funding was received from the European Respiratory Society (2012 Rare Disease Achievement Award to

CLS); National Institute of Health Research London (NW) Comprehensive Local Research Network; HHT

patient donations; Imperial College BSc project funds (to CLS for ST, FG, TS, and VS), and the National

Institute of Health Research Biomedical Research Centre Scheme (Imperial BRC). The views expressed are those of the authors and not necessarily those of funders, the NHS, the NIHR, or the Department of Health and Social Care.

AUTHORS' CONTRIBUTIONS:

-Conception and design: ST, FG, TS, CLS. Analysis and interpretation: ST, FG, TS, JP, BM, JEJ, ST, VS, JM, HCT, LH and CLS. Drafting the manuscript for important intellectual content: ST, FG, CLS. *In detail:* ST devised the CPET analyses focussing on anaesthetic risk, performed literature studies, added to the pulmonary AVM CPET database, performed the data analysis, generated Figures 2, 3 and Supplementary Figures 1 and 2, and wrote the first draft of the manuscript. FG introduced the VSAQ clinic assessments, performed literature studies, generated the observational database, drafted manuscript sections and generated Supplementary Figure 3. TS performed literature studies, assisted in obtaining ethical approvals to recruit patients with airflow obstruction, and added to the pulmonary AVM CPET database. JP co-supervised ST, and FG, and performed and interpreted CPET data measurements. VS performed literature studies, contributed to generation of the observational pulmonary AVM database, assisted in obtaining initial ethical approvals to recruit patients without airflow obstruction, and initiated the pulmonary AVM CPET database. JM generated the VSAQ, advised on physiological concepts, and contributed to data interpretation. HT co-supervised ST, and FG, and performed and interpreted CPET data measurements. LH co-supervised TS and VS, and performed CPET data measurements. CLS supervised all students, devised the initial CPET study, reviewed all patients, performed literature searches, analysed and interpreted data, performed additional presented data analyses, generated other Figures, and wrote the final manuscript. All authors contributed to and approved the final version of this manuscript.

ACKNOWLEDGEMENTS:

THURAIRATNAM S, GAWECKI F, STRANGEWAYS T, PERKS J, SANTHIRAPALA V, MYERS J, TIGHE HC; HOWARD LSGE, SHOVLIN CL.
TRIAGE ASSESSMENT OF CARDIORESPIRATORY RISK STATUS BASED ON MEASUREMENT OF THE ANAEROBIC THRESHOLD, AND
ESTIMATION BY PATIENT-REPORTED ACTIVITY LIMITATION. *UPLOADED 16.04.2020, VERSION 2 MEDRXIV/2020/045203.*

-The authors thank the patients for their willing cooperation in these studies.

REFERENCES

National Institute for Health and Care Excellence Routine preoperative tests for elective surgery Available from: https://wwwniceorguk/guidance/ng45

Agnew N. Preoperative cardiopulmonary exercise testing. Continuing Education in Anaesthesia Critical Care and Pain. 2010; 10(2): 33-37. Available from: doi: 10.1093/bjaceaccp/mkq001.

Ridgway ZA, Howell SJ Cardiopulmonary exercise testing: a review of methods and applications in surgical patients European Journal of Anaesthesiology 2010;27(10): 858–865 Available from: doi:101097/EJA0b013e32833c5b05

Moran J, Wilson F, Guinan E, Mccormick P, Hussey J, Moriarty J Role of cardiopulmonary exercise testing as a risk-assessment method in patients undergoing intra-abdominal surgery: a systematic review British Journal of Anaesthesia 2016;116(2):177-191 Available from: doi:101093/bja/aev454

Myers J, Do D, Herbert W, Ribisl P, Froelicher VF. A nomogram to predict exercise capacity from a specific activity questionnaire and clinical data. Am J Cardiol. 1994;73(8): 591-596.

Myers J, Bader D, Madhavan R, Froelicher V. Validation of a specific activity questionnaire to estimate exercise tolerance in patients referred for exercise testing. Am Heart J 2001;142(6): 1041-1046.

Athanasopoulos LV, Dritsas A, Doll HA, Cokkinos DV.The role of NT-proBNP in explaining the variance in anaerobic threshold and VE/VCO(2) slope. J Cardiopulm Rehabil Prev. 2011 Sep-Oct;31(5):316-21.

Jette M, Sidney K, Blümchen G. Metabolic equivalents (METS) in exercise testing, exercise prescription, and evaluation of functional capacity. Clin Cardiol. 1990;13(8): 555-565.

McAuley P, Myers J, Abella J, Froelicher V. Evaluation of a specific activity questionnaire to predict mortality in men referred for exercise testing. Am Heart J. 2006;151(4): 890. e1-890. e7.

Snowden CP, Prentis JM, Anderson HL, Roberts DR, Randles D, Renton M, Manas DM. Submaximal cardiopulmonary exercise testing predicts complications and hospital length of stay in patients undergoing major elective surgery. Ann Surg. 2010 Mar;251(3):535-41.

Shovlin CL, Condliffe R, Donaldson JW, Kiely DG, Wort SJ; British Thoracic Society British Thoracic Society Clinical Statement on Pulmonary Arteriovenous Malformations Thorax 2017 Dec;72(12):1154-1163

Shovlin CL. Pulmonary arteriovenous malformations. Am J Respir Crit Care Med. 2014 Dec 1;190(11):1217-28.

Whyte MK, Hughes JM, Jackson JE, Peters AM, Hempleman SC, Moore DP, et al. Cardiopulmonary response to exercise in patients with intrapulmonary vascular shunts. J Appl Physiol 1993;75:321–328

Santhirapala V, Williams LC, Tighe HC, Jackson JE, Shovlin CL. Arterial oxygen content is precisely maintained by graded erythrocytotic responses in settings of high/normal serum iron levels, and predicts exercise capacity. An observational study of hypoxaemic patients with pulmonary arteriovenous malformations. PLoS ONE 2014 Mar 17;9(3):e90777.

Rizvi A, Babawale L, Tighe HC, Macedo P, Hughes JMB, Jackson JE, Shovlin CL. Hemoglobin is a vital determinant of arterial oxygen content in hypoxemic patients with pulmonary arteriovenous malformations. Ann Am Thorac Soc. 2017 Jun;14(6):903-91.

Howard LSGE, Santhirapala V, Murphy K, Mukherjee B, Busbridge M, Tighe HC, et al. Cardiopulmonary exercise testing demonstrates maintenance of exercise capacity in patients with hypoxemia and pulmonary arteriovenous malformations. Chest. 2014;146(3): 709–718. Available from: doi:10.1378/chest.13-2988.

Gawecki F, Strangeways T, Amin A, Perks J, Wolfenden H, Thurainatnam S, Rizvi A, Jackson JE, Santhirapala V, Myers J, Brown J, Howard LSGE, Tighe HC, Shovlin CL. Exercise capacity reflects airflow limitation rather than hypoxaemia in patients with pulmonary arteriovenous malformations. QJM. 2019 Jan 17. doi: 10.1093/qjmed/hcz023.

Gawecki F, Myers J., Shovlin CL. Veterans Specific Activity Questionnaire (VSAQ): a new and efficient method of assessing exercise capacity in patients with pulmonary arteriovenous malformations BMJ Open Respir Res. 2019 Mar 1;6(1):

e000351http://bmjopenrespres.bmj.com/cgi/content/abstract/6/1/e000351

Vorselaars VM, Velthuis S, Mager JJ, Snijder RJ, Bos WJ, Vos JA, et al. Direct haemodynamic effects of pulmonary arteriovenous malformation embolisation. Neth Heart J 2014;22:328–333.

Shovlin CL, Buscarini E, Kjeldsen AD, Mager HJ, Sabba C, Droege F, Geisthoff U, Ugolini S, Dupuis-Girod S. European Reference Network For Rare Vascular Diseases (VASCERN) Outcome Measures For Hereditary Haemorrhagic Telangiectasia (HHT). Orphanet J Rare Dis. 2018 Aug 15;13(1):136. doi: 10.1186/s13023-018-0850-2.

Finnamore H, Le Couteur J, Hickson M, Busbridge M, Whelan K, Shovlin CL. Hemorrhage-adjusted iron requirements, hematinics and hepcidin define hereditary hemorrhagic telangiectasia as a model of hemorrhagic iron deficiency. PLoS One. 2013 Oct 16;8(10):e76516.

Santhirapala V, Chamali B, McKernan H, Tighe HC, Williams LC, Springett JT, Bellenberg HR, Whitaker AJ, Shovlin CL. Orthodeoxia and postural orthostatic tachycardia in patients with pulmonary arteriovenous malformations: a prospective 8-year series. Thorax. 2014 Nov;69(11):1046-7.

Shovlin CL, Moorthy K, Lees C. Covid-19: Home based exercise activities could help during self isolation BMJ 2020: https://blogs.bmj.com/bmj/2020/03/16/covid-19-home-based-exercise-activities-could-help-during-self-isolation/

Thurainatnam S. Pre-Operative Insights from Cardiopulmonary Exercise Testing in Patients with Pulmonary Arteriovenous Malformations. BSc Project, Imperial College London (2017).

Criteria Committee, New York Heart Association , Inc. Diseases of the Heart and Blood Vessels. Nomenclature and Criteria for diagnosis, 6th edition Boston, Little, Brown and Co. 1964, p 114.

FIGURES

Figure 1: The UK-modified Veterans Specific Activity Questionnaire (VSAQ)

It is important for us to understand whether you have normal, limited or even high levels of activity. On this list, first please circle <u>any activity</u> that when performed for a period of time, would typically <u>make you to want to stop.</u> Then record in the box, the smallest number that you circled an activity for:	1	Eating, getting dressed or working at a desk
	2	Taking a shower, shopping, cooking, walking down eight steps
	3	Walking slowly on a flat surface for 100-200 meters A moderate amount of work around the house such as vacuuming, sweeping the floors or carrying groceries
	4	Light gardening work Painting or light carpentry
	5	Walking briskly e.g. four miles per hour Social dancing, washing the car
	6	Playing nine holes of golf carrying your own clubs. Heavy carpentry. Pushing a lawn mower
	7	Performing heavy outdoors work e.g. digging Walking uphill, tennis singles, carrying a 4-5 year old child
	8	Moving heavy furniture, jogging slowly on the flat, carrying a toddler up stairs
	9	Cycling at a moderate pace, sawing wood
	10	Brisk swimming, cycling uphill, walking briskly up hill, jog six miles per hour
	11	Cross country skiing, carrying a heavy load up two flights of stairs, cycling briskly
	12	Running briskly, continuously
	13	Any competitive activity, including those which involve intermittent sprinting Running or rowing competitively, cycling races

(adapted from [18] with authors' permission)

Figure 2: Age and oxygen saturation in low and high risk groups.

*Box plots comparing values between the low risk and high risk anaerobic threshold (AT) groups for 26 pulmonary AVM patients, where high risk status was defined by AT lower than 11ml.min⁻¹kg⁻¹: **A)** Age (ys), **B)** Resting SpO₂ (%).Boxes indicate the median and interquartile range (IQR), and error bars represent 2 standard deviations, with dots at the extremes representing outliers. P-values were calculated by Mann Whitney U test.*

Figure 3: Haemoglobin and arterial oxygen content in low and high risk groups

*Box plots comparing values between the low risk and high risk anaerobic threshold (AT) groups for 26 pulmonary AVM patients, where high risk status was defined by AT lower than 11ml.min⁻¹kg⁻¹: **A)** Haemoglobin (g.dL⁻¹); **B)** CaO₂ (ml.dL⁻¹). Boxes indicate the median and interquartile range (IQR), and error bars represent 2 standard deviations, with dots at the extremes representing outliers. P-values were calculated by Mann Whitney U test.*

Figure 4: Age-VSAQ method suggesting lower and higher risk AT categories.

The lowest 10 VSAQ scores and associated exercise limitations are plotted against age. Horizontal bands indicate the respective New York Heart Association (NYHA[25]) categories of I (no symptoms on ordinary physical activity), II (limited on ordinary activity), III (limited at 20-100m), and IV (limited at rest). To indicate lower and higher risk categories, the regression line is plotted for an anaerobic threshold of 11 ml.min^{-1}.kg^{-1}. To provide an indication of confidence limits, and the direction and scale of variation if this threshold scale were to be adjusted, regression lines for anaerobic thresholds of 10 and 12 ml.min^{-1}.kg^{-1} are also plotted

Triage assessment of cardiorespiratory risk status based on measurement of the anaerobic threshold, and estimation by activity limitation in patients with pulmonary arteriovenous malformations and hereditary haemorrhagic telangiectasia

Saranya Thurairatnam, Filip Gawecki, Timothy Strangeways, Joseph Perks Vatshalan Santhirapala, Jonathan Myers Hannah C Tighe; Luke SGE Howard, Claire L. Shovlin.

Supplementary Data

Supplementary Methods

METS VSAQ AT Formulae

Since

$$Predicted\ METs = 4.74 + (0.97 \times VSAQ\ score) - (0.06 \times Age)$$

And

$$(0.8368381\ METs) + 5.1992178 = AT$$

Then

$$\frac{AT - 5.1992178}{0.8368381} = 4.74 + (0.97 \times VSAQ\ score) - (0.06 \times Age)$$

And for the AT cut off of 11ml.min^{-1}kg^{-1}

$$\frac{11 - 5.1992178}{0.8368381} = 4.74 + (0.97 \times VSAQ\ score) - (0.06 \times Age)$$

$$6.931785491 = 4.74 + (0.97 \times VSAQ\ score) - (0.06 \times Age)$$

$$2.191785491 = (0.97 \times VSAQ\ score) - (0.06 \times Age)$$

$$2.191785491 - (0.97 \times VSAQ\ score) = -(0.06 \times Age)$$

$$(0.97 \times VSAQ\ score) - 2.191785491 = (0.06 \times Age)$$

So by Excel, Tested ages where the relevant VSAQ score would yield a figure <2.19:

VSAQ	VSAQx.97	(VSAQx.97)-2.191785491	(VSAQx.97)-2.191785491/0.06
1	0.97	-1.221785491	-20.3631
2	1.94	-0.251785491	-4.19642
3	2.91	0.718214509	11.97024
4	3.88	1.688214509	28.13691
5	4.85	2.658214509	44.30358
6	5.82	3.628214509	60.47024
7	6.79	4.598214509	76.63691
8	7.76	5.568214509	92.80358
9	8.73	6.538214509	108.9702
10	9.7	7.508214509	125.1369
11	10.67	8.478214509	141.3036
12	11.64	9.448214509	157.4702
13	12.61	10.41821451	173.6369

Supplementary Figure 1

Comparison of CPET-derived variables related to ventilation in low and high risk groups delimited by anaerobic threshold (AT) of 11ml.min^{-1}kg^{-1}: A) serum bicarbonate (mmol/L) available in N=17, **B)** V_E/V_{CO_2} slope. Boxes indicate the median and interquartile range (IQR), and error bars represent 2 standard deviations, with dots at the extremes representing outliers. P-values were calculated by Mann Whitney *U* test. Note that the lower risk group displayed lower bicarbonate indicative of more exuberant ventilation, yet had less steep V_E/V_{CO2} slope consistent with greater ventilatory efficiency.

Supplementary Figure 2

Comparison of CPET-derived variables related to ventilation in low and high risk groups delimited by above and below median anaerobic threshold (AT): A) Serum bicarbonate (mmol/L) available in N=17, **B)** V_E/V_{CO_2} slope. Additional details as for Supplementary Figure 1 above.

Supplementary Figure 3

Relationship between METs and Anaerobic Threshold. Anaerobic threshold plotted against predicted METs from VSAQ score. The regression coefficient was 0.84 (95% confidence intervals 0.12, 1.56), p= 0.032.