1 **Herd immunity vs suppressed equilibrium in COVID-19 pandemic: different**
2 **goals require different models for tracking**
3
4

5 **Authors**:
6 Norden E. Huang[#], Fangli Qiao[#], Wang Qian[+] and Ka-Kit Tung[*]
7
Affiliations:
[#] *Data Analysis Laboratory, FIO, Qingdao* 266061*, China*
[+] *Shanghai Jiao Tong University, Shanghai, 200240, China*
[*] *Department of Applied Mathematics, University of Washington, Seattle, WA* 98195
[#] Co-first authors: These authors contribute equally to this work.
FQ: qiaofl@fio.org.cn. NEH: norden@ncu.edu.tw
*Corresponding author: ktung@uw.edu

KEYWORDS
Covid-19; Epidemiology; herd immunity; suppressed equilibrium; Data-driven
approach; prediction of turning point; Active Infected Cases; peak total
hospitalizations; duration of hospital stay; end of epidemic; local-in-time metric.
20
21

22   **ABSTRACT**
23
New COVID-19 epicenters have sprung up in Europe and US as the epidemic in
China wanes. Many mechanistic models' past predictions for China were
widely off the mark (1, 2), and still vary widely for the new epicenters, due to
uncertain disease characteristics. The epidemic ended in Wuhan, and later in
South Korea, with less than 1% of their population infected, much less than
that required to achieve "herd immunity". Now as most countries pursue the
goal of "suppressed equilibrium", the traditional concept of "herd immunity"
in epidemiology needs to be re-examined. Traditional model predictions of
large potential impacts serve their purpose in prompting policy decisions on
contact suppression and lockdown to combat the spread, and are useful for
evaluating various scenarios. After imposition of these measures it is
important to turn to statistical models that incorporate real-time information
that reflects ongoing policy implementation and degrees of compliance to
more realistically track and project the epidemic's course. Here we apply such
a tool, supported by theory and validated by past data as accurate, to US and
Europe. Most countries started with a Reproduction Number of 4 and declined
to around 1 at a rate highly dependent on contact-reduction measures.
41
42
43

1. Introduction.

Almost one hundred years ago, a classic paper published in Proceedings of Royal Society by Kermack and McKendrick (3), entitled "A contribution of mathematical theory of epidemic", started the tradition of mathematical modelling on the spread of infectious disease among a susceptible population. Current thinking in epidemiology is still deeply rooted in concepts introduced in that paper, some of which are still relevant, while others need to be modified. The mechanistic model they introduced is called the SIR model, for Susceptible-Infected-Recovered/Removed: A susceptible population is infected by the introduction of a few infected individuals. The infected population eventually recovered, and in the process acquiring immunity to the original infectious disease, or dead (removed). An epidemic ends when susceptibles are exhausted as most of the population is immunized this way, achieving "herd immunity". There have been many variants to this basic model. One common modification is to add an extra population of E for exposed individuals who are not yet infectious, in the so-called SEIR model. Other---agent-based---- models take advantage of the modern computing power to further subdivide the population into many subgroups and even simulate movements of individuals. But basic concepts are similar to the SIR model. These mechanistic models are of critical importance before the outbreak starts, for the models could be used to explore various scenarios for policy decisions on social distancing and lockdown. Once the outbreak starts, a different kind of models are needed----so far not well developed----which are capable of using real-time data to provide the needed parameters for epidemic management, predict various turning points and peak medical resource needs, and monitor the effect of compliance of the quarantine. In this paper, we introduce such a statistical model, supported by theory and validated by observed data.

There are actually two possible end states of an epidemic: one is through "herd immunity" mentioned above when we discuss the mechanistic models, and the other is an unstable state achieved by suppressing contacts among individuals, called "suppressed equilibrium" here, which is pursued by most countries in this pandemic, though many policy makers may not be aware of the distinction between the two. This second state is (parametrically) unstable because if the social distancing measures are relaxed and the businesses reopened, the disease could initiate a second wave, as most of the population has not acquired immunity. Even if the epidemic ends in one country, there could still be subsequent waves of infection by imports from abroad unless there is strict quarantine of cross-border travelers. There is also a third route: to suppress just enough so that the maximum cases of hospitalizations are kept below the maximum capacity of the medical system in each district, delaying long enough either for the population to eventually achieve herd immunity or for a vaccine to be developed for the disease. More people would be infected and more would die in this approach than the suppressed equilibrium, but less than that in the herd immunity scenario, where the hospitals may become

89 overwhelmed. Which approach to take is the difficult decision confronting policy
90 makers at the beginning of an epidemic.
91
It was reported (2) that UK first contemplated not suppressing the epidemic
through lockdowns, fearing that doing so would only lead to a larger second
outbreak because most of the population would not have gained immunity. So the
plan was to let the epidemic run its course while protecting the elderly. But when
shown a model prediction (4) that such a "do-nothing scenario" would lead to
500,000 deaths and 81% of the population infected, policy makers changed course
and imposed strict counter measures. This is an important and proper role for a
model, to prompt policy actions to combat the spread of the disease. Once the
outbreak started, the accuracy of the mechanistic model predictions cannot be
verified as the forecast forever changed the course of the epidemic in UK. Health
officials in Sweden didn't believe in models and decided to pursue "herd immunity"
starting 12 March.
104
The number of people that will have to be infected before achieving herd immunity
depends on how contagious the disease is. The 16 March report of Ferguson et al.(4)
assumed an infection rate, expressed in terms of Basis Reproductive Number R_0 of
2.4. For US it predicted that 81% of the population would have to be infected in this
do-nothing scenario, or about 250 million, resulting in 2.2 million dead. Later
updates in the 30 March report of Flaxman et al. (5) suggested that R_0 should be
above 4 for European countries studied. Estimating this number will be one of the
tasks in the present work so that one can evaluate what it entails for the herd-
immunity approach. We will later show, directly from data, that this estimate of R_0
~4 also holds for the US, and in fact approximately so for every country we
examined. So over 90% of the US population would need to be infected before herd
immunity could be achieved. COVID-19 turns out to be much more contagious than
originally thought. See also (6).
118
Since a "suppressed equilibrium" is achieved in a very different manner than that for
the "herd immunity", our estimate of the end date of the epidemic as a consequence
of contact suppression is not based on the number of susceptibles, *S,* approaching a
small critical value (i.e. when most of the population is infected, hence acquiring
immunity), but the daily new cases *N(t)* approaching zero and remaining so for two
incubation periods. For prediction purpose, the date when the *N(t)* is near zero is
estimated by 3 standard deviations from its peak. These two quantities, the peak
and the standard deviation, can be extracted from the data as the epidemic is
developing. Our estimate of the end of the epidemic is earlier, usually significantly
so, because it does not depend on a high percentage of the population having been
infected to achieve herd immunity.
130
For South Korea, the epidemic in that country ended with just 0.02% of its
population infected. Wuhan, the epidemic ended with less than 0.5% of its
population infected, both less than 1% of that required to achieve herd immunity as
predicted by most mechanistic models. Even if these numbers are multiplied by a

factor of 10 (as some in the media suggested that the Wuhan data may be an
underestimate), it is still one order of magnitude less. Furthermore, it is generally
accepted that South Korea's data are good. Indeed, the situation in these countries
represents early examples of the "suppressed equilibrium". Because of its much
lower number of deaths, such an end state is a goal that most countries have
decided to pursue, despite the enormous toll on the economy due to the much
reduced business activity for the two to three months that it would take to achieve it.
142
The above-quoted numbers for Wuhan and South Korea are for the confirmed cases,
and do not include the asymptomatic infected. Recently (in early April) about 3,330
individuals in Santa Clara County in California were tested for antibodies to the
COVID-19 virus in their blood (7). 50 were tested positive (meaning that they were
exposed in the past to the virus), yielding a crude prevalence rate of 1.5%. When
weighed by demographics and extrapolated statistically to the whole county's
population, it was calculated that 2.8% of the population could have been infected,
with a 95% confidence range of 1.3-4.7%. These numbers, less than 5%, are also
much less than what is required to achieve herd immunity, in a county where the
epidemic is waning at the time.
153
In pursuing the "suppressed equilibrium" and monitoring the progress towards it, a
second type of statistical models is required. This type of models should reflect the
contact-reduction measures already in place and the degree of compliance by the
population in each region. Unlike China and South Korea, many European countries
imposed these measures in stages, and so the contact rate among the population
was reduced in a time-dependent way. In US, even before the epidemic is over, some
states are starting to reopen businesses. Only a real-time, data driven model can
reflect these changes. Such a model should be based on sound epidemiological
principles. Prediction based on a purely statistical model without an epidemiological
foundation, even though using real-time and past data, is akin to watching the daily
stock market fluctuations and performing "technical analysis" to ask when the peak
is for an investor to time a sell order. A prediction could be made but the
uncertainty would be so large that it is just likely for the prediction to be right as for
it to be wrong.
168
Each type of models has its strengths and weaknesses. For the mechanistic model,
such as SIR and SEIR or agent-based versions, a key parameter, the infection rate, is
not known for an emerging disease such as SARS-CoV-2, and this has been a source
of difficulty with predictions using such models. For the second type of models,
especially the purely statistical models without epidemiological basis, it is not
known which quantity of the epidemic is predictable. For example, there have been
many empirical models based on the assumption that the progression of daily cases
follow a Gaussian "epidemic curve" in time, starting with the early model of William
Farr in 1840: "Law of Epidemics" in his second annual report to the Registrar
General of England and Wales (8). Lacking the epidemiological mechanism that
Kermack and McKendrick (3) later proposed, the "law" simply reflected Farr's
conviction that the observed deceleration of the rate of increase of infected would

not lead to an impending catastrophe but to a crest and then accelerated decline.
The latest is that of the Institute of Health Metrics and Evaluation (IHME)(9). It
turns out that fitting three parameters that define a Gaussian to a short time series
and then using that Gaussian to predict the peaks of the epidemic is an ill-posed
problem(10). The uncertainty for next-day prediction is near 100%, and should
increase further for predictions a few days out (11). In this work we pay special
attention to which epidemiological property is predictable, and we quantify the
uncertainty of our prediction.
189

The search for the correct "geometry of epidemic curves" has a long history in
statistical modelling. Farr's law is purely descriptive. Farr did not realize that his
epidemic curve is Gaussian but nevertheless his descriptive law of second ratios
could be used for prediction, though not very accurate. For example (see (12)), if
$x_1, x_2, x_3, x_4, x_5, x_6 ....$ are the successive weekly incidence (i.e. new cases) or mortality,
his law says that the ratio of successive ratios of these numbers is a constant:

$$\frac{x_4/x_3}{x_2/x_1} = \frac{x_5/x_4}{x_3/x_2} = ... = K,$$

which is less than 1. That is, there is a constant deceleration of the rate of growth of
the cases. After measuring this constant from the early weeks' data, future
incidence values can be predicted. It was John Brownlee in 1907 (13) who realized
that the above formula, when logarithm is taken----turning the ratio of ratios to
difference of differences----is a finite difference form of the second order time-
derivative of $\log x$ being a negative constant (12):

$$\frac{d^2}{dt^2}\log x = \log K < 0.$$

Integrating twice and then taking exponentiation lead to $x(t)$ being a Gaussian form.
Brownlee thought this normal form for the epidemic curve is a fundamental law in
epidemiology, but his proposed explanation for the declining growth of the
incidence of an epidemic as due to decreasing "infectivity" was not well-received by
epidemiologists at the time.
209

Brownlee (13) provided examples of several epidemics showing that there was fore-
aft symmetry in their epidemic curves. For COVID-19, we find that the epidemic
curve for Wuhan, China follows a Gaussian, with near fore-aft symmetry, but that for
US has a rapid rise but slow decline, definitely not Gaussian. While it may be
possible that without human intervention, a solitary outbreak may follow a
Gaussian curve, in the modern era of contact suppression, the epidemic curve is
shaped by such interventions. We shall explain Wuhan's shape as due to the fact
that the contact suppression measures were consistently imposed throughout the
course of the outbreak, while in the case of US, its states and the populace, were
relaxing earlier measures on the aft side of the curve, when the new cases declined,
creating a fore-aft asymmetry. Therefore, one should take into account that in the
modern era, as countries pursue a "suppressed equilibrium" at great economic cost,
there is a tendency in countries with decentralized state governments to relax the

countermeasures to various degrees once the disease crested, giving rise to
subsequent waves of infection. [A second reason is that the US data is an aggregate
of data for different epicenters, with staggered recovery. A third reason for the
asymmetry may be artificial: In many countries there is usually an increase in the
testing of the population as the production of the test kits and facilities to process
them ramp up after the initial shortage. Therefore it is to be expected that there
would be more cases found later in the course of the epidemic than in the beginning.]
230
Brownlee rejected the idea of herd immunity, that the epidemic's decline was due to
"an exhaustion of susceptibles, enshrined 20 years later in the SIR model of
Kermack and McKendrick (3). His alternative, "infectivity" idea was based on the
thinking that the decline was due to "the loss of infecting power on the part of the
organism"(13), and that this biological property of the pathogen ("organism")
should follow some fundamental law. This biological property of the virus has not
been observed in the current COVID-19 pandemic, and does not appear to be a
factor. However, Brownlee's idea can be resurrected by modifying the definition of
"infectivity" to include social factors, since how many people one infected individual
can infect, as measured by the Effective Reproduction Number, R_t, depends on the
product of the number of persons contacted during the infectious period and the
probability of the contacted person contracting the disease. After implementation of
contact-reduction measures, we can actually see from the data (in section 2) the
decline of this measure of "infectivity". Furthermore the decline is steeper in
countries that have the more stringent contact-reduction policies and
implementation. Although both the mechanisms of loss of susceptibles and
decrease in "infectivity" are likely at play, with the extremely small percentage of
the population infected in the current pandemic, the second mechanism appears to
be the dominant one as countries strive to achieve the "suppressed equilibrium".
Given this situation, model predictions of the decline of the epidemic based the
number of susceptibles decreasing, as in SIR and SEIR models, may be missing the
main cause for the observed progression of the disease in the current pandemic.
253
Brownlee's idea, with modification expressed above, can be cast in a mathematical
form as:
$x_{t+1} = R_t x_t$,
where x_t is the incidence (new cases) at time t , and x_{t+1} is the incidence one
infectious period later. An infection period is the duration an infected person
remains infectious. R_t is defined earlier as the number of people one infected
individual would infect during the period when he is infectious. If R_t is a constant,
$R_t = R_0$, the solution to the above finite difference equation is:
$x_t = x_0 (R_0)^t$;
the solution is an unimpeded exponential growth (since $(R_0)^t = \exp\{t \log R_0\}$) for the
relevant case of $R_0 > 1$. Brownlee(14) commented that such an epidemic form is
contrary to the facts: "The assumption that the infectivity of an organism is constant,

leads to epidemic forms which have no accordance with the actual facts." With R_t as
a decreasing function of time, which we find is actually the case in section 2, the
above solution becomes Gaussian-like. Specifically, if R_t decreases by a factor $q<1$
after each period (12, 14), due to a "loss of infecting power", i.e. $R_t = R_0 q^{(t-1)}$, then the
solution is Gaussian:

$$x_t = x_0 (R_0)^t q^{t(t-1)/2} \ ,$$

(since $x_t = x_0 \exp\{t \log R_0 + \frac{1}{2} t(t-1)\log q\}$, noting $\log q < 0$).

The exponential growth from $(R_0)^t$ is eventually overtaken by the more rapid

decrease of q^{t^2}. Note that in this argument, no mention is made of the decrease of
the number of susceptibles; this is not needed when the decrease is so small
compared to the population as a whole.
277
This paper is organized as follows: We first introduce the relationship between the
net infection rate and the Reproduction Numbers. The epidemiological basis for our
model is discussed. Then, we will present a suite of prediction tools for epidemic
management. We will also provide a summary of the COVID-19 inferred
epidemiological characteristics for various countries in Asia, Europe and US. Finally,
a discussion and conclusion will be given. The theoretical support is given in the
appendix.
285
2. **The net infection rate and the Reproduction Numbers**
Before we discuss our prediction model, we first discuss diagnosing a key parameter
used as input in most mechanistic epidemiological models, the infection rate, or
equivalently expressed as the *Basic Reproduction Number* or the time-dependent
Effective Reproduction Number. Using real-time data, we diagnose it for different
countries in the world, which actually reflects the underlying influence factors.
292
We define in general the *net infection rate* $\alpha(t)$ as the time-varying exponential
growth rate of the active infected cases (15):

$$\alpha(t) = \frac{d}{dt}\log I(t) = \frac{\frac{d}{dt}I(t)}{I(t)} \ .$$

The active infected number, $I(t)$ is given by the equation that describes its rate of
increase as the daily new infected cases $N(t)$ minus the daily recovered/removed
case, $R(t)$:

$$\frac{d}{dt}I = N(t) - R(t).$$

The dead ("removed") is included in $R(t)$ in our calculations. The peak number of
active infected cases is a key parameter in the planning for hospital resources. This
turning point, denoted by t_p, can be located in a local-in-time manner by when R
starting to exceed N, without first accumulate the data in time to find $I(t)$. Maximum

demand for hospital resources occurs at its peak, and not at the peak of $N(t)$,
although the latter is a more commonly reported quantity. *R(t)* is not a factor in the
initial rise of the outbreak, nor is it needed to explain why the rise slows and
eventually crests. However in the waning phase, when the new cases are smaller,
the rising recovered cases need to be taken into account to determine t_p.
309
In traditional mechanistic models, such as the SIR model(3) , there is also a time-
dependent net infection rate, which at *t=0* , when the population is completely
susceptible, is related to the *Basic Reproduction Number R_0*. See ref (16) for a
discussion of the complexities associated with this key parameter. We will not be
using the SIR model but it is useful to relate our general definition to what is
traditionally used. The equation for *I* in the SIR model is:
316

$$dI/dt = aSI - bI = bI(aS/b - 1),$$

where $aS(t)$ is the infection rate and b is the recovery/removal rate.

Therefore $\quad \alpha(t) = \dfrac{dI/dt}{I} = b(\dfrac{aS(t)}{b} - 1) = b(R_t - 1).$

So for the SIR model $R_t = \dfrac{aS(t)}{b}$. So as the population of the susceptible $S(t)$

decreases---"exhaustion of the susceptibles" as hosts for the disease----the Effective
Reproduction Number decreases. More relevant for our discussion is the possibility
that the infection rate *a* could change as a result of contact reduction measures in
place.
325
326 Initially when the whole population is not yet infected, the Basic Reproduction

327 Number is $R_0 = \dfrac{aS(0)}{b} = R_t(0) = \alpha(0)b + 1$.

328
The above-defined time-dependent net infection rate generalizes this concept to be
independent of the SIR or other models: If in the course of an epidemic, $\alpha(t)$ is
positive, the number of infectives will grow exponentially, reaching a peak number
of infectives when $\alpha(t) = 0$ at $t = t_p$, which is a critical turning point mentioned
above. Then the total number of active infectives will decrease exponentially. In
terms of $R_t = \alpha(t)/b + 1$, if this number is greater (less) than 1 the total number of
active infectives will grow (decrease) at time t . We will here use $\alpha(t)$ directly. R_t
however is the more watched quantity by the mainstream modelers (16). It can be
calculated from the net infection rate, but will require a parameter *b*, the recovery
rate, which may be different for different regions. Furthermore, many countries do
not keep adequate records on those who recovered, and so there is an uncertainty in
estimating *b.* In Figure 2, R_t is obtained by estimating this parameter as
$b \approx 1/\sigma_R \approx 1/\sigma_N$, where σ_R is the standard deviation for the distribution of the
daily recovered and σ_N is that for the daily newly infected numbers. These

parameters are calculated for each country (see Table 1). R_0 is obtained from R_t in
the initial period, before there is significant recovered population.
345
346 For an emerging disease such as SARS-CoV-2, there is not enough statistics for
347 estimating the true infection rate (e.g. the parameter a in the SIR model). Models
348 usually assume a value of R_0 or perform scenario calculations for a range of values of
349 R_0.
350
Figure 1 shows the net infection rate for several countries. Since the official data
that we use include only the confirmed cases ("cases" for short), and these tend to
have more serious symptoms that require hospitalization, the reported I cases are
commonly referred to as *total hospitalizations*. The peak of total hospitalizations is
closely watched by hospital administrators and policy makers. $\alpha(t)$ is commonly
referred to as *rate of hospitalization*. Its inverse gives the e-folding time in days for
the cases in an outbreak. A value, of say α between 0.3 and 0.4, where most
countries cluster in the initial period, implies an e-folding time of about 3 days
(doubling time of 2 days). The much higher values α for many regions in the
beginning of our data record may not be due to indigenous disease infection. See
later discussion.
362
363

364
Figure 1. The time-dependent net infection rate (in units of 1/day) as a function of
time starting on the date (listed in the inset) when the newly confirmed case
number first exceeds 100 for each region. To obtain the actual calendar date, add
the dates on the horizontal axis to the starting date indicated in the inserted legend.
The number of confirmed cases on the starting date is listed at the top. Three day
averaging on the raw data has been used.

The time for different countries is aligned in Figure 1 to begin the time series when
each region reached 100 new cases. This way, the progression of the epidemic in
each country can be compared. Figure 1 reveals the effects of different policy
measures each country adopted. First, South Korea and China exHubei have similar
net infection rates (until past their respective turning point); both are much lower
than other countries. In the case of South Korea, the government identified early
that its epicenter of the epidemic was at church gatherings in the city of Daegu and
North Gyeongsang province, where 90% of the initial cases were found. Specifically,
a confirmed COVID-19 patient was reported to have attended the Shincheonji
Church of Jesus services twice on February 9th and 16th. Then aggressive contact
tracing was pursued. After the turning point, South Korea soon experienced some
second wave episodes, which were successfully contained. These two regions'
rigorously implemented contact reduction and aggressive pursuit of 'test-trace-treat'
measures led to them being the extreme examples of the "suppressed equilibrium".
386

Germany and Italy have similar exponential growth rates of the net infected case
numbers, both slightly higher than Wuhan. More surprisingly, US has the highest
exponential net infection rate, 1.5 times that of Germany and Italy and twice that of
Wuhan. This can be attributed to the fact that US so far does not have a nation-wide
lockdown, and Europe has had partial lockdowns in phases. Germany took a week
longer than Wuhan to reach its turning point, while US will take weeks longer than
Germany. China outside Hubei reached its turning point early, in fact 9 days earlier
than the epicenter, Wuhan. This fact is significant, for it is qualitatively different
than many mechanistic model predictions, which had the epicenter achieving its
turning point 1-2 weeks earlier than China outside Hubei (17), probably based on
the herd immunity concept.
398

The net infection rates for China outside Hubei and South Korea are more
monotonic than other regions shown. This is due to the fact that there was not a
piece-meal imposition of social-distancing measures, unlike other western countries.
The strict measures were imposed and enforced throughout the course. [For
Wuhan, the large spike on day 23(12 February) was due to a change in the
diagnostic criteria from a positive nucleic acid test to chest scans, booking in one
day more than 13,000 cases.]
406

The case of Sweden needs a special explanation. The epidemic in Sweden initially
grows with an e-folding time of around 3 days, in line with other countries. Then on
12 March the government announced that because of limited resources, it no longer
would test for the COVID-19 infection, except for those with serious symptoms
already in the hospitals who furthermore are also in the high-risk group. As a result
the new cases took a nose-dive on that day, leading to an artificially low net
infection rate of 0.1, implying a 10 day e-folding time. The denominator in the
calculation for $\alpha(t)$ is $I(t)$, which is an accumulated quantity, and includes those
who tested positive prior to 12 March under more liberal criteria, and so this

situation leads to a flat, low level of $\alpha(t)$ just above 0. It would eventually cross 0
with a large enough death numbers. Sweden's policy decision to pursue "herd
immunity" (while protecting the elderly) has been touted as a viable and perhaps
preferable approach to those of other countries in their pursuit of "suppressed
equilibrium". It only encouraged those over 70 to stay home and banned visits to the
nursing home and gatherings with over 50 people, while business, stores,
restaurants and kindergarten through grade nine are open. The success or failure of
this approach cannot be evaluated by the incomplete data. Intriguingly, based on
the recorded death number, it shows that Sweden's toll is 5 and 11 times that of its
neighbors Denmark and Norway, respectively. The population in each of these
neighbors is half that of Sweden.
427

428
429
430
Figure 2. Effective Reproduction Number for each country or region. The horizontal
axis denotes days since day 0 (the corresponding calendar date is given in the inset),
which is the starting date for our calculation. This date is determined by the
threshold that the accumulated number of infectives first exceeds 100. The actual
number for each region on that day is listed at the top.   The thick dashed curve is
the average of the curves for USA and European countries, including Russia.
437
Figure 2 converts Figure 1 to show R_t for each country. It shows that R_t clusters
around 4 for all countries in the initial period. Because of the problems for the data
in the initial period, the curves cannot be extended back in time to deduce R_0. But
based on R_t a few days later, R_0 for COVID-19 should be around 4, similar to that for
SARS. It was originally thought that COVID-19's R_0 was between 2.0 to 2.5 (18),
seemingly much less contagious than SARS at 4 (19). It is also much more

contagious than the 2009 swine flu pandemic, caused by the H1N1 virus, whose R_0
was estimated(20) to be 1.4 to 1.6.
446
447 In deducing the Reproduction Numbers we should not count the large spike for
448 China outside Hubei on day 4. That increase was not due to indigenous
449 transmission, for most of the initial cases were imported from Hubei. This should
450 not be used to infer the Reproduction Number. Similarly for South Korea in the first
451 few days shown.
452
As is in Figure 1, the decrease of R_t from 4 to 1 for different countries reflects
different level of contact-reduction measures adopted and enforced. With China
outside Hubei and South Korea sloping more steeply than Europe than US. The
behavior of these numbers for the European countries are rather similar to the
model results of Flaxman et al.(5) of Imperial College (their Figure 2), who imposed
these measures in their model on the dates they were actually imposed. With their
emphasis on more accurately predicting the mortality, when $R_0 \sim 4$ is used in their
SEIR model the modeled death numbers are close to the reported deaths, but the
number of infected in the model is an order of magnitude higher (Figure 2 in (5)).
This is reasonable and consistent with our earlier statement that the data of infected
cases do not include the asymptomatics or those with mild symptoms who were not
tested, but reflect those with more serious symptoms that required hospitalization,
and therefore more prone to die.
466
UK's record for the recovered is almost nonexistent, and what is available shows
that the recovered number is only a few percent of the deaths, which does not
appear to be reasonable. Without the recovered in the data, UK's R_t hovers above 1.
There appears to be a similar situation in some other countries, such as Italy.
Therefore, the behavior of R_t in the later stages of the epidemic (in the
neighborhood of the turning point) is probably not correctly depicted by the data
shown for these countries. Nevertheless, in the initial period, when the number of
recovered is small, the data shown can be used to estimate R_t and R_0.
475
SEIR model was also used to deduce the Reproduction Number by Institute of
Disease Modeling (21) in an effort to monitor the effect of social distancing
measures adopted near Seattle (King County, Washington). They found that R_t was
reduced from 2.7 to 1.4. Since it was not below 1, the Institute's report advised
continuing the measures in place. Since the report rate p, that is, the ratio of the
number of reported cases vs. the true infected number, was unknown, the authors
assumed a range of values and obtained, $R_0 \sim 2.7 \pm 0.9$. One can see from this
application of the Reproduction Numbers how important it is to monitor in real-
time the progress of policy measures to determine whether it is time to relax the
measures in place. And also how difficult it is to infer these numbers.
486
The report rate p is the ratio of the confirmed cases to the true infected numbers. If
it were known, the case numbers can be divided by p to yield the true infected

numbers. Because it is largely unknown at the present time, it creates the
aforementioned uncertainty in estimating Reproduction Numbers using models.
However, since our net infection rate $\alpha(t)$ is a ratio of the derivative of *I* and *I*,
dividing each by *p* does not change the ratio. We therefore recommend using the
net infection rate for real-time monitoring instead of the Reproduction Number.
The assumption is that the report rate has not changed in time, at least not during
the two-week period prior to *t* that the accumulation takes place for the
denominator (see Appendix). This same consideration is also partly behind our later
using the ratio of new cases to recovered cases for prediction.
498
499 The problem with the data for Sweden is that in addition to the change in the report
500 rate, the record for the recovered is not available.
501
This discussion also highlights the difficulty of using mechanistic models for real-
time monitoring. In addition to not knowing the report rate *p* to compare model
output with the reported case numbers, a key parameter needed in the mechanistic
models, the infection rate *a*, is largely unknown for the emerging disease. There is a
large population of asymptomatic, untested and unreported infectives. This infected
population is nevertheless infectious and produces some of the infected cases
reported. It would usually lead to an overestimate of the infection rate *a* when only
the reported cases are used in the estimation. Our tools do not need to use an
infection rate for prediction.
511
3. Epidemiological basis
After establishing the relationship between the net infection rate and the
Reproduction Numbers, we briefly summarize the main ingredients to establish the
epidemiological basis of our model. Details of our model are given in the Appendix.
The model is used here only to infer general properties of an outbreak, and to
discover which properties can be predicted.
518
Based on epidemiological theory, the infected population is governed by the Von
Foerster equation in an age-structure population model (see (22)), where "age", *s*, is
days since first infected. The infection at age 0 is governed by the "birth" dynamics,
the infection dynamics, and death due to natural causes is ignored. An infected
individual is assumed to be infectious in an age-dependent way until *T*, when the
individual is either cured or dead, in either case no longer belonging to the
population of the infectives, where *T* is the mean recovery/removal period. The
solution to this partial equation (via the method of characteristics) yields certain
rather general results.
528
529 **Conservation Law**. For $t > T$
530
531 $$R(t) = N(t - T).$$
532 The distribution of the newly recovery/removal follows that of the newly infected
533 with a time delay of *T*. When applied to hospitalizations, it says that those who are

534 admitted to the hospital either recover after a mean hospital stay of T days, or dead
535 after a similar number of days. A more complicated relationship holds for $t < T$.
536
537 Since hospital stay acts like a filter for $R(t)$, the profile for R is slightly wider than
538 that for N. Appendix takes this into account.
539
540 $R(t)$ here actually consists of two parts: the recovered individuals $R_c(t)$ and the
541 removed (dead) individuals, $D(t)$: $R(t) = R_c(t) + D(t)$.
542
543 **Validation after the fact**: This fundamental relationship can be validated
544 statistically with data, provided the data is long enough. This is one of the ways the
545 mean recovery period T is determined statistically from data, but it is not practical
546 in the early phase of the epidemic.
547
Since $R_c(t)$ and $D(t)$ are a subset of $R(t)$, the time lag relationship should also hold.
Additional insight can be gained by looking at them separately. Figures 3, obtained
using the longest data from China, South Korea and Italy during the COVID-19
pandemic, shows that $N(t)$ and $R(t)$ are highly correlated: with correlation
coefficients all over 0.9 when both distributions are smoothed with 3-point boxcar.
The mean time delay of the correlation is denoted by μ, which can be interpreted as
a statistical mean of T. The spread of the cross-correlation is measured by its
standard deviation, σ_c; its value is also given in the figure. The lag time (and
standard deviation) of $R(t)$ for China is 19 days (σ_c=17.7 days) for South Korea is 23
days (σ_c=19.8 days) and for Italy is only 10 days (σ_c=21.2days). Due to their low Case
Fatality Rate (CFR), there is practically no difference between $R(t)$ and $R_c(t)$ for
China and Korea. But for Italy, which has a high CFR, there are differences among
the constituting parts. Why Italy's lag time between $R(t)$ and $N(t)$ is shorter than
China and South Korea needs a deeper examination, and it does not necessarily
mean the shorter the better. The lag time of $D(t)$ for China is 7 days, for Korea is 17
and for Italy is only 3 days. The short survival time for patients in the hospital is an
indication that the Italian hospitals were overwhelmed and allocation of ventilators
was selective to those more likely to survive. This mortality component reduces the
overall time for recovery/removal for Italy to 10 days.
567
568

569
570
571 **Figure 3**. Lagged cross-correlation of $R_c(t)$, $D(t)$ and $R(t)$ with $N(t)$ for China, South
572 Korea and Italy. Collectively or individually, the conservation law and time lag
573 relationship are validated. The separated $R_c(t)$ and $D(t)$ reveal the stress the
574 regional hospitals had experienced.
575

Epidemic curve.
Furthermore, the solution to the model equations (in Appendix) shows that the
time-dependent profile of $N(t)$, sometimes referred to as "the epidemic curve", is
determined by the "age" distribution, and the method of characteristics in the
solution converts that "age" distribution into a time profile. The details of the "age"
distribution depend on the "birth" process and on how long it has been since the
outbreak first begins. We assume that we do not mix data from different epicenters
(the homogeneity assumption). Focusing on a single outbreak, which starts at *t=0*,
we examine the "age" distribution at a time t_B much longer than the initial
incubation period. Then we assume there exists a full spectrum of age with *s*
between 0 and *T* in a Gaussian-like form. To the right of the peak in the "age"
distribution, it is easy to understand that those who are "older" should be less in
number because they were infected during an earlier stage of the epidemic. As the
epidemic grows exponentially, there are more and more "younger" infectives. To the
left of the peak in "age' distribution many infectives are at various stages of
incubation. Those who are newly infected may be much large in number, but since
they have not yet developed symptoms, they are less likely to be tested, hospitalized,

and contribute to the "case" record. For Covid-19, the peak infectiousness occurs in
a period just before and after the onset of symptoms, around 5-6 days since first
infected(23). The above discussion gives an epidemiological justification for a
Gaussian-like distribution of the "age" distribution of the "new cases", more
appropriately called "new hospitalization per day". It is consistent with the
argument given in the Introduction in favor of a Gaussian-like epidemiological curve.
It should be pointed out that historically, for obvious reasons, asymptomatics were
also not included in the "incidence", or "mortality", which were used by Farr to
obtain his law of second ratios or by Brownlee to obtain his normal curves.
602
603   With the possible exception of South Korea, in most countries there could be
604   multiple seeding of the outbreak occurring at slightly different times. These
605   staggered series of outbreaks merge in a continuum in the data for that country,
606   leading to a standard deviation in the new cases that is wider than that from the
607   "age" distribution. This is a complication we need to take into account.
608
Also, in many countries, pressure mounts for policy makers to relax the contact-
reduction measures when case counts pass the peak and are declining. In some
countries where the restrictions are gradually lifted, we should expect a long tail in
the epidemic profile, which is therefore not symmetric with respect to the peak.
This external influence to the original expected progression of the course should be
monitored and adjustment to predictions made in real time. As will be discussed in
subsequent sections, our model has the ability to adapt to these changes in its role
as a monitoring tool. Nevertheless, we find, after the fact, that predictions for the
various turning points are still accurate even without taking into account the
changes. It turns out that, consistent with the above discussion, the relaxation of
contact-reduction measures, which lengthens the standard deviation of the new
cases, is only significant in the later stages of the course, and can be ignored before
the peak, but the prediction on the evolution after the peaks on quantities such as
the end of the epidemic and the total number of infected is likely not accurate unless
the changes are taken into account. (This has not been done in Table 1.)
624
4. A suite of tools for tracking the epidemic.
Based on theoretical considerations, we developed a suite of tools for monitoring
the evolution of the epidemic. They can be used to predict the timing of several
turning points and the number of infectives associated with each. These include t_N,
the peak of the daily new cases; t_R, the peak of daily recovered/removed cases, and
t_p, the turning point where the active infected cases (AIC), or total hospitalizations,
is a maximum. This is the point of maximum strain on the hospital resources. It
needs to be closely monitored to keep the AIC below the hospital capacity. We also
estimate the date when the epidemic ends and the country can be reopened, t_c.
634
A more accurate and robust prediction tool is based on the ratio of $N(t)$ and $R(t)$,
designated as the *NR* ratio. This ratio also alleviates to some extent the problem

related to the data of reported cases being a fraction *p* of the true numbers, as *p*
cancels out in the ratio. Unfortunately, some countries, such as UK and Sweden, do
not keep adequate record of *R(t)*, and many country do not keep a rigorous standard,
which could be detect by the low case recovery rate, indicating the violation, or
leakage, of the conservation law. For these countries a less accurate method, in the
sense of having larger error bars, can still be used when only information on *N(t)* is
available. These are described below. We have used the data from countries that
have the longest records for COVID-19 to verify properties of these methods.
645
646 **4.1 Log of NR ratio**:
647
648 We define the *NR* ratio as
649 $$NR(t)=N(t)/R(t).$$
650 At t_p, *NR=1*.
651
We show in Figure 4, using the data of the epidemic for COVID-19 for the longest
records available in some countries, that the logarithm of *NR(t)* lies on a straight line,
with small scatter, passing through the turning point t_p. And data for various stages
of the epidemic, from the initial exponential growth stage, to near the peak of *AIC*,
and then past the peak, all lie on the same straight line. The intercept with *logNR=0*
yields the turning point. This line, obtained by linear-least-square fit, is little
affected by the rather large artificial spike in the data on 12 February for China,
because of its short duration and the logarithmic value. That reporting problem is
necessarily of short duration because, on the date of definition change, previous
week's cases of infectives according to the new criteria were reported in one day.
After that, the book is cleared, and *N(t)* returned to its normal range.
663

Figure 4. Natural Logarithm of the ratio of daily newly infected to newly
recovered/removed. They lie on straight lines with some small scatter. The straight
line obtained by linear-least squares fit is in dotted line with the same color for each
country. (a) For China, the slopes of the lines are almost the same but with different
intercept; the trend lines cross zero (the black solid line) at different time for
different regions indicating different peaking time for *AIC*. The epicenter Wuhan
(green) has latest turning point than its province Hubei (pink), which has a later
turning point than China as a whole (cyan). As a measure of confidence of the linear-
least square fit, the 95% confidence limit for China is given in the figure. Similar
confidence limits for other regions have been calculated but are not shown for the
sake of clarity of the presentation. (b) Comparing different countries. The time is
aligned by plotting the *NR* ratio only when the recovery/removal case numbers first
exceed 50 (the actual numbers on that day are listed at the top), and T_0 is that
calendar date, listed in the inset. The figure in the bottom panel included the China
cases again, to facilitate comparison with other countries, except the data used were
smoothed by a 3-point boxcar filter in the lower panel.
683
It would be interesting to understand why the empirically determined log *NR(t)* lies
on a straight line, and what determines its slope. See Appendix for a theoretical
support. It is shown that, if the distribution of the new cases is Gaussian-like, the
natural logarithm of the *NR* ratio, should be a linear function of time throughout the
course of the epidemic. The slope of the line is $-\dfrac{T}{\sigma_R^2}$.
σ_R, is the standard deviation of the recovered case distribution and is close to that of
the newly infected case, σ_N, twice of which measures the duration of the epidemic
in that region. This expression is still valid even for a sigma that is changing in time
as a response to changing social contact measures.
693
A comparison of the Logarithm of *NR* ratio for several countries is given in Figure 4
(b). A steeper slope is associated with an early turning point, and also a predictor for
a shorter duration of the epidemic. The shallowest slopes in Figure 4 (b) was for
Italy, where the enormous pressure strained their medical system to the limit,
resulting in the largest σ_R value, and one of the highest case fatality rate in the world,
at more than 12%. Germany and China have similar slopes. For China outside Hubei,
the slope is steepest and the turning point reached 9 days earlier than Wuhan. South
Korea's slope is even steeper due to that countries early action. As a result, Italy
took a full month longer to reach its turning point than Germany and China, and
more than 40 days longer than South Korea.
704
4.2 Predictability
Since the logarithm of *NR* ratio lies on a straight line passing through the turning
point of *AIC*, it would be interesting to explore if the turning point, t_p, can be
predicted by extrapolation using data weeks before it happened. Extrapolating a
straight line is much more practical than other more involved curve fitting

algorithms some other groups have adopted. How far in advance this can be done
appears to be limited by the poor quality of the initial data, when R is small and
highly fluctuating. Figure 5 (a) shows the results of such predictions for China, and 5
(b) for Italy. It is a hind cast since the truth is now known (see Figure 8 for how it is
determined). The horizontal axis indicates the last date of the data used in the
prediction. The beginning date of the data used is 24 January for all experiments for
China. Prior to that day, data quality was poor and the newly recovered number
was zero in some days, giving an infinite NR ratio. For China outside Hubei, the
prediction made on 6 February gives the turning point as 14 February, two days
later than the truth. A prediction made on 8 February already converged to the
truth of 12 February, and stays near the truth, differing by no more than fractions of
a day with more data.
722
The huge data glitch on 12 February in Hubei affected the prediction for Hubei, for
China as whole, and for Hubei-exWuhan. These three curves all show a bump up
starting on 12 February, as the slope of $N(t)$ is artificially lifted. Ironically,
predictions made earlier than 12 February are actually better. For example, for
China as a whole, predictions made on 9 February and 10 February both give 19
February as the turning point, only one day off the truth of 18 February. A
prediction made on 11 February actually gives the correct turning point that would
occur one week later. At the time these predictions are made, the newly infected
cases were rising rapidly, by over 2,000 each day, and later by over 14,000. It would
have been incredulous if one were to announce at that time that the epidemic would
turn the corner a week later. Even with the huge spike for the regions affected by
the Hubei's changing of diagnosis criteria, because of its short duration the artifact
affects the predicted value by no more than 3 days, and the prediction accuracy soon
recovers for China as a whole. For Hubei, the prediction never converges to the true
value, but the over-prediction is only 2 days.
738
Uncertainty associated with prediction using this method for China is shown in
Figure 4(a). The uncertainty is a few days (the 95% confidence level is $\pm$ 5 days)
and is usually somewhat larger than the accuracy. A prediction can be found later to
be accurate but at the time it was made it may be equally likely for the prediction to
be a few days earlier or later. The large uncertainty is again seen to be caused by the
1,4000 bump of new cases in one day on 12 February due to the change in definition.
For applications to other countries and to future epidemics without a change in the
definition of the "infection" to such a large extent, we expect even better prediction
accuracy and smaller uncertainty.
748
749   This can be seen in the prediction for Italy. The error of predicting the turning point
750   3 weeks in advance is only 1 or 2 days. In fact a prediction can be made 6 weeks in
751   advance with an accuracy of 5 days or less.
752
The prediction for US as a whole is less accurate (with errors up to 10 days) because
its data is an aggregate of different epicenters. More accurate predictions can be
made by treating each state separately. This is not done here because although the

data for new cases and deaths are available for each state, recovered data are not
individually available. It is also not accurate for UK because its data for recovered
may be suspect. See Table 1.
759
760
761

762
Figure 5. (a) Prediction of the turning point in *AIC* for different regions of China by
extrapolating the trends in logarithm of *NR*. The horizontal axis indicates the date
the prediction is made using data from 24 January to that date. The vertical axis
gives the dates of the predicted turning point. Dashed horizontal lines indicated the
true dates for the turning point. (b) Prediction of the turning point in *AIC* for Italy.
Data used for all predictions starts on 29 February. The first point shown predicts
the turning point to occur on 19 April (4 days early) 6 weeks in advance using 9
days of data from 29 February to 8 March.
771
4.3. Derivative of Log of *N(t)* and *R(t)*.
Interestingly, the derivative of log *N(t)* or log *R(t)* also lies on a straight line, as
shown in Figure 6 (although the scatter is larger as to be expected for any
differentiation of empirical data)*.* Moreover, the straight line extends without
appreciable change in slope beyond the peak of *N(t)*, suggesting that the distribution
of the newly infected number is approximately Gaussian-like at least up to that point.
For an exponential function, the derivative of its logarithm being a linear function of
time is highly suggestive of a general type of distribution including Gaussian-like or
Rayleigh-like.   The recovery time *T* can be determined as t_R-t_N, where t_R is the peak
of *R(t)* and t_N is the peak of *N(t).* These two peak times can be obtained by extending
the straight line in Figure 6 (a) to intersect the zero line. This predicted result can be
verified statistically after the fact by the lagged correlation of *R(t)* and *N(t)*. If the
distribution is indeed Gaussian or even approximately so, the slope in Figure 6
should be proportional to the reciprocal of the square of its standard deviation, σ,
as:
$$\frac{d\log N(t)}{dt} = \frac{-(t-t_N)}{\sigma_N^2}.$$
788
789 Similarly result holds for the daily number of recovered, *R(t)*.
790

791
Figure 6 (a). The derivative of the logarithm of daily newly infected or recovered for
China and Hubei. Notice the clear separation of the new and recovered cases and
also the subtle difference of their slopes. The zero crossings of the trend line give
the peak dates of the new and recovered case respectively. And the slopes give an
estimate of σ values. (b). Same as (a) but for Italy. Best fit lines for $N(t)$ and $R(t)$ are
shown, and 95% confidence limits are shown for $N(t)$ only. Italy's data appear to
have a 7-day periodic oscillation, probably caused by a reporting issue. AR indicates
accumulated R.
800
801 For countries without an adequate record of $R(t)$, this method can still be used for
802 $N(t)$. We can obtain t_N and σ_N. T and t_p cannot be obtained, but can be estimated
803 roughly as $t_p = t_N + T/2$ (see Theory), using $T \sim 20$ days applicable to countries with
804 similar medical systems.
805
4.4 The number of deaths
Because of public attention, predicting the number of deaths has been the main
emphasis for some models. There is also the belief that death numbers are more
reliable than case numbers, although that is not necessarily true. Although death
number by itself could not satisfy conservation law, death is a subset of the newly
confirmed cases; therefore, the death case distribution should follow the new cases
with a time delay. In fact, this delay time could provide a measure of the efficacy of
the medical system, as explained earlier.
814
815 In Figure 7, we present the cases for Italy, USA and UK.

Figure 7 (a), (b) and (c). Logarithm for death distribution for Italy, USA and UK.
Similar to 4.2, the derivative of logarithm of *D(t)* follows a straight line
approximately, indicating that its distribution is also Gaussian-like, similar to *N(t).*
As a consequence, the peak of *D(t)* occurs at the intersection of that straight line
with zero and could be predicted in advance, although in the three countries shown
this peak has already occurred.

5. Inferred epidemiological characteristics.
Table 1 summarizes predictions for USA, some European countries, South Korea and
China. In some countries, one or both of the critical points have just occurred. Even
so, it is still difficult to know if one is at the peak without using our prediction
procedure. The *TIC*∞ is the Total Infected Cases for the whole epidemic, calculated
as twice the value of *TIC* at t_N , and AIC_{peak} is the maximum of *AIC* at t_p . Appendix
discusses how these quantities are computed/predicted.

	t_p (r)	t_N (r)	t_D (r)	AIC_{peak}--K (r)	$TIC\infty$--K (r)	$AFC\infty$--K (r)	σ_N Days	CFR --%	IPM --Per M	FPM --Per M
China	2-21 (2-17)	2-7 (2-13)	2-17 (2-14)	44.4-54.1 (58.0)	87.7-163 (84.0)	3.8-6.6 (4.6)	8.79	5.53	59.98	3.31
China ex-Hubei	2-11 (2-12)	2-5 (2-4)	2-16 (2-13)	8.99-9.04 (8.94)	15.5-26.1 (15.8)	0.14-0.23 (0.13)	8.49	0.79	11.82	0.09
South Korea	3-15 (3-14)	3-2 (3-1)	3-19 (3-29)	7.08-8.35 (7.49)	8.84-16.2 (10.8)	0.18-0.30 (0.26)	6.31	2.36	209.42	4.94
Italy*	4-23 (4-21)	3-30 (3-21)	3-31 (3-28)	114-115 (108)	203-399 (213+)	24.8-42.1 (29.3+)	12.18	13.76+	3526.7+	485.3+
Ger-many*	4-4 (4-7)	3-31 (3-27)	4-14 (4-15)	50.7-69.5 (70.8)	144-309 (167+)	6.86-14.0 (7.0+)	9.62	4.19+	2014.6+	84.35+
Spain*	4-22 (4-27)	4-6 (3-26)	4-7 (4-1)	114-129 (105)	274-495 (248+)	28.1-48.4 (25.6+)	10.44	10.32+	5316.9+	548.5+
USA*	5-12 (NA)	4-19 (4-9)	4-22 (4-16)	1400-1474 (943+)	1520-2410 (1204+)	90-140 (71.1+)	12.21	5.9+	3649.5+	215.3+
UK*	6-30# (NA)	4-20 (4-10)	4-20 (4-30)	11789# (165+)	249-392 (195+)	33-62 (29.4+)	14.21	15.1+	2932.2+	442.5+

Table 1. Summary of the predictions made on May 5, 2020. **Notes**: Countries with
" *"sign after a country name indicates that the epidemic is still developing, although
some of the particular event might have happened. The number in the parenthesis is
the actual data. A"+" sign indicates the latest value in the particular case that is still
developing and that particular event has not happened yet. The '#' indicated the
result might be questionable, for the lack of recovered case numbers.
IPM= Infection per Million of Population. FPM=Fatality per Million of population.

6. Discussions and Conclusion.
There are actually two modeling approaches: mechanistic and statistical models.
Mechanistic model is based on precisely defined epidemical parameters. Without *a*
priori knowledge of the disease characteristic and existing social contact conditions,
however, many of the parameters used are based on assumptions and educated
guesses. Nevertheless, they are useful for exploring different scenarios for the
policy decision to fight an epidemic prior to the rapid rise of the outbreak. In this
paper, we introduced a statistical model that could also serve an important role
complementary to the mechanistic models, to be used after the outbreak started. It
can provide detailed tracking and prediction of the course of the epidemic. For the
statistical model to work well, good quality data are indispensable. Our study
indicated that with quality data, tracking and prediction of critical events such as the
peak and the turning point could be accurate within days.

Our model is supported by underlying theoretic foundation and validated by the
existing data from the region where the pandemic is waning. Many parameters
characterizing an epidemic can be determined from local-in-time data. Because it is
based on general epidemiological principles we suggest that our approach could be
applied not just to the current Covid-19 epidemic, but also generally to future
epidemics.

Importantly, we made explicit the concept of "suppressed equilibrium" as an end
state of an epidemic in additional to the traditional "herd immunity" state. Based on

the traditional mechanistic model, an epidemic ends after a high percentage of the
population is infected and recovered hence acquiring immunity. This is the so-
called "herd immunity" idea. For COVID-19, which we found to be very contagious,
more so than previously thought, the "herd immunity" end would require almost all
of the population be infected and therefore would bring unthinkable toll in the
number of people sick and dead. A second way for an epidemic to end is with strict
contact-reduction measures, so that although a large pool of susceptible population
still exists, the portion that an infected person comes in contact with is reduced by
the measures adopted, again leading to the Effective Reproductive Number less than
1. Unlike the first end state mentioned above, this "suppressed" state is
"parametrically unstable" in the sense that if the social distancing measures are
relaxed before the epidemic ends or new infection is imported after the first wave
ends, the epidemic will rebound, as a large portion of the population is still
susceptible. For this second state to be a stable equilibrium, the social-distancing
measures, and quarantine of cross-border visitors need to be maintained until it is
clear that the disease has died off. It is this second state that most countries are now
aiming for.
886
The second state will have a much earlier date for the new cases to peak and the
epidemic to end. For the US the peak of the newly infected cases is April 7-11. The
epidemic is estimated here to end in the first week of June, assuming the current
social distancing/stay-at-home measures are maintained till then, and after that
date import of infected from abroad is prevented by strict quarantine of visitors.
These assumptions now do not look like they would hold as states begin to open
their business prematurely. For the US, we predict that the total number of infected
cases is 1.5 to 2.4 million, dependent on the assumption. These are symptomatic
cases that require hospitalization.
896
Since it is the goal of most countries to eventually approach the "suppressed
equilibrium", it is important to note that the slowing growth of the incidence (daily
newly infected) that is observed is not a function of biology, but is a result of
contact-reduction, which is social science. The mechanism of the exhaustion of
susceptibles is not relevant anymore as the number of infected is such a very small
percentage of the susceptible population. Therefore it is not necessary to use a
model, such as SIR or SEIR, to keep track of the change in susceptibles after the start
of the outbreak. If these mechanistic models are to be used to track the progress of
an outbreak, the infection rate a in the models needs to be adjusted constantly, but it
is not clear how to adjust it. In any case, the solution is responding to input of
changing model parameters, and not to the natural biological evolution built into the
model structure, such as recovery and immunity upon recovery.
909
This is a new way to look at the field of epidemiology. The observed slowing of the
growth of incidence and the cresting of the epidemic curve are dominantly the
result of a reduction of "infectivity", as first proposed by John Brownlee one
hundred years ago, except that the change is not due to biology---- from the "loss of
the infecting power on the part of the organism" as he thought, at least not in the

case of COVID-19, but as the consequence of reduced contacts among people in
lowering the Reproduction Number, which is defined as the number of other people
one infected person would infect. This measure is a product of the average number
a typical infected person comes in contact with and the probability that a contacted
person becomes infected. While the study of the latter is biology and medicine, that
for the former is social science and public health.
921
922
923

 Appendices: THEORY

Definition: Let $I(t)$ be the number of active infected at time t. Its change is given

by;

$\dfrac{d}{dt}I = N(t) - R(t),$

where $N(t)$ is the number of newly infected, and $R(t)$ that of the newly recovered or

removed (dead). The term: Acting Infected Case (*AIC*) number is used to denote the

confirmed *I(t)* when we deal with data.

Let t_p, the turning point defined as the peak of the active infected number. At this

point maximum medical resource is needed. This maximum occurs when

$\dfrac{d}{dt}I = 0,$ implying $N(t_p) = R(t_p)$.

There is no need to first find *I(t)* to locate this peak. Figure 8 shows how this is

determined locally. This local-in-time metric avoids the accumulation of poor early

data. After the turning point, the newly recovered starts to exceed the newly

infected. The demand for medical resources, such as hospital beds, isolation wards

and respirators, starts to decrease.

We consider a solitary outbreak. Let $t = 0$ be when the first infection began. For

Wuhan, China, this date is near the end of 2019, perhaps even earlier. Let t_B be the

beginning of the better quality data. This time is beyond the initial incubation

period of the disease and it can be assumed that at that time there is already a

population of infected, some of them asymptomatic but nevertheless infectious.

Let $X(t,s)$ be the number of infected cases at time t, with s being the "age"

distribution, i.e. number of days sick.

The total number of infected is given by:

$I(t) = \int_0^T X(t,s)\,ds$.

After being sick for *T* days, a patient either recovers or is removed (dead). *T* is called

the recovery period (or removal period). It is also called the infectious period if the

patient is infectious during this period. Of course its value varies by patient and by

the efficacy of treatment for each hospital. For the removed it also depends on the

age of the patient and whether there are underlying medical conditions. Only a

mean recovery period is obtainable from data, and so this is in reality a statistical

quantity. We will discuss later how this statistical quantity can be obtained from

data.

Conservation law (see ref(22)):

After first infected and until removed or cured, we have:

$$dX(t,s) = \frac{\partial}{\partial t} X \cdot dt + \frac{\partial}{\partial s} X \cdot ds = 0, \quad 0 < s < T.$$

963 So, since $ds/dt = 1$,

$$\frac{\partial}{\partial t} X + \frac{\partial}{\partial s} X = 0.$$

964 This equation is to be solved using the method of characteristics as

$X(t,s) = $ constant along characteristics defined by $ds/dt = 1$.

Boundary condition: $X(t,0)$, specifies the "birth" process,

965 i.e. how the disease spawns newly infected (with "age" $s = 0$).

Initial condition: $X(0,s) = 0$ for $s > 0$, specifies the initial age distribution at $t = 0$

966 There are two types of characteristics:

967 (i) $s > t$, (ii) $s < t$.

968 The first type of characteristics intersects the $t = 0$ axis, and since the initial
969 condition is zero, we have the solution:
970

971 $X(t,s) \equiv 0$ for $s > t$.

973 That is, there is no infected population who is sick for more days than the lapsed
974 time since the first infection occurred.

976 For the second type of characteristics, $t > s$ the solution is
977

978 $X(t,s) = f(s-t)$

979 with the form of f to be determined by the boundary condition. Even without
980 determining the form of f we have the following general results:

981 For $t > T$, and therefore $t > s$:

$$I(t) = \int_0^T X(t,s)\,ds = \int_0^T f(s-t)\,ds$$

$$= \int_{t-T}^t f(p)\,dp.$$

$$\frac{d}{dt}I = f(t) - f(t-T).$$

Since the rate of increase of $I(t)$ is by definition equal to the newly infected number,
$N(t)$, minus the newly recovered (or removed) number, $R(t)$, we have:

$$N(t) - R(t) = \frac{d}{dt}I = f(t) - f(t-T).$$

For a fatal disease with low fatality rate, where almost all infected cases eventually
recover after a hospital stay of T days, we can identify
$f(t)$ with $N(t)$, and $f(t-T)$ with $R(t)$.

If the disease has a non-negligible fatality rate, we include the dead in $R(t)$.
Main Result: The daily newly recovered/removed number $R(t)$, is related to the
daily newly infected number $N(t)$ as, for $t > T$:
995
996                     $$R(t) = N(t - T).$$
997

998   The second type of characteristics intersects the boundary $s = 0$. The boundary
999   condition itself needs to be solved as a function of t to describe how new infection
1000 (at $s = 0$) occurs. This can be done using a birth model, such as Eq. (1.56) in (23).
1001
1002 *Boundary condition*: Following Murray's Eq. (1.56), the "birth" is assumed to be
1003 proportional to "parents" of suitable "age", with an age-dependent birth rate, $a(s)$.
1004

$$X(t,0) = \int_0^T a(s)X(t,s)ds \ .$$

This equation needs to be solved numerically, except in the case of constant a. The
solution for $X(t,0) = g(t)$ is generally an increasing function of time.

Since $X(t,s) = g(t-s)$, at some time $t_B > T$: $X(t_B,s) = g(t_B - s) \equiv f_0(s)$. That is, an
"age" distribution can be converted into a time profile through the method of
characteristics.
1011
1012 The solution of the integral equation is not presented here. For our purpose here it
1013 suffices to assume that the solution of this model yields a distribution with age that
1014 has a full spectrum $0 < s < T$ of infectives at a time t_B , long after a full incubation
1015 period has passed.

1016 $$X(t_B,s) = f_0(s) = A\exp\{-\frac{(s-s_0)^2}{2b^2}\}; \ A \text{ independent of } s.$$
      $$b = \tfrac{1}{2}T.$$

This distribution is justified as follows:
To the right of the peak in "age" distribution, it is easy to understand that the
numbers for those who are "older" should be less because they were infected earlier
during an earlier stage of the epidemic. As the epidemic grows exponentially, there
are more and more "younger" infectives. To the left of the peak in "age' distribution
many are at various stages of incubation. Those who are newly infected may be
large in number, but they are less infectious and contribute less to the growth of the
subsequent infection, and since they have not developed symptoms, they are less
likely to be hospitalized, tested and contribute to the "case" record. For Covid-19,
the peak infectiousness occurs in a period just before and after the onset of
symptoms, around 5-6 days since first infected (23). The above discussion gives an
epidemiological justification for a Gaussian-like distribution of the "age" distribution
of the "new cases", more appropriately called "new hospitalization per day".
1030

1031　It is important to point out that the assumption of less "cases" to the left of the peak
1032　is the only "filter" that serves to limit our theoretical discussion to the data of
1033　symptomatic cases. If all infected were included, $f_0(s)$ should have a maximum at
1034　$s=0$.
1035　Therefore the solution is, for $t>t_B>0$:

$$X(t,s)=X(t_B,s-t)=f_0(s-t)$$
1036
$$=A\exp\{-\frac{(s-s_0-t)^2}{2b^2}\}.$$

1037

1038
$$N(t)=A\exp\{-\frac{(t-s_0)^2}{2b^2}\}=A\exp\{-\frac{(t-t_N)^2}{2b^2}\}$$

$$R(t)=N(t-T)=A\exp\{-\frac{(t-t_R)^2}{2b^2}\},$$

1039　where t_N is the peak of $N(t)$, and $t_R=t_N+T$ is the peak of $R(t)$.
1040　Both distributions are Gaussians.
1041
1042　For $t_B<t<T$,

$$I(t)=\int_0^t X(t,s)ds+\int_t^T X(t,s)ds$$
1043
$$=\int_0^t f(s-t)ds+\int_t^T 0ds$$
$$=\int_{-t}^0 f(p)dp$$

1044　$$\frac{d}{dt}I=f(-t)=A\exp[-\frac{(t-s_0)^2}{T^2}].$$

1045　$$N(t)=A\exp[-\frac{(t-t_N)^2}{T^2}]$$

$$R(t)=0.$$
1046　Again, *N(t)* is Gaussian, but there is no recovered or removed during this early stage.
1047
1048　**Main Result**: The natural logarithm of the ratio of *N* and *R* is a linear function of
1049　time for $t>T$:
1050

$$NR(t)=\frac{N(t)}{R(t)}=\frac{N(t)}{N(t-T)}$$
1051
$$=\exp\{-\frac{(t-t_N)^2}{2b^2}+\frac{(t-t_N-T)^2}{2b^2}\}=\exp\{\frac{T^2}{2b^2}-\frac{T(t-t_N)}{b^2}\}.$$

$$\log NR=-\frac{T(t-t_N)-\frac{1}{2}T^2}{b^2}$$

1052   a linear function of t.
1053

Heterogeneous Data
The above results are obtained for the case of a single introduction into a region of
infected at $t=0$ and we solve for the subsequent development of the epidemic from
that single source. Consider now a large region consisting of a number of small
regions, and the "seeding" of the infected occurs at different times for different
regions. The large region could be China, and the first infection could be Wuhan,
Hubei and then the regions outside Hubei. Then we may have for the China as a
whole data for the newly infected a sum of several Gaussians staggered in time. As
long as the Gaussians are not separated so much that there are different peaks in the
combined data, the combined data can still be considered as Gaussian, as is the case
in the real data. However, the standard deviation σ of the combined Gaussian is
inevitably larger and is no longer given by b:

1066
$$N(t)=\frac{B}{\sqrt{2\pi}\sigma_N}\exp\{-\frac{(t-t_N)^2}{2\sigma_N^2}\}.$$

1067   We still have $R(t)=N(t-T)$ since this result holds for each sub-region. The result
1068   that $\log NR$ is a linear function of time still holds:

1069
$$\log NR(t)=\log\frac{N(t)}{N(t-T)}=-\frac{T(t-t_N)-\frac{1}{2}T^2}{\sigma_N^2}\;.$$

1070   The slope of the straight line is T/σ_N^2 .
1071

1072   Since the hospital stay can act as a smoothing filter on $N(t)$ to yield $R(t)$, the
1073   standard deviation for $R(t)$ could be slightly wider than that for $N(t)$. So we could
1074   have two different Gaussians (but their integral over all time should be the same):

1075
$$N(t)=\frac{B}{\sqrt{2\pi}\sigma_N}\exp\{-\frac{(t-t_N)^2}{2\sigma_N^2}\};\;\;R(t)=\frac{B}{\sqrt{2\pi}\sigma_R}\exp\{-\frac{(t-t_R)^2}{2\sigma_R^2}\}.$$

1076   Taking this into account, we have, denoting $T=t_R-t_N$:

1077
$$\log NR-\log\frac{\sigma_R}{\sigma_N}=-\{\frac{(t-t_N)^2}{2\sigma_N^2}-\frac{(t-t_N-T)^2}{2\sigma_R^2}\}=-\frac{(\sigma_R^2-\sigma_N^2)(t-t_N)^2+2\sigma_N^2 T(t-t_N)-\sigma_N^2 T^2}{2\sigma_N^2\sigma_R^2}$$

1078

1079   As the values of σ_N and σ_R are very close based on the empirical data, the quadratic
1080   term is always small comparing to the other terms for the length of time we are
1081   considering here. Hence,

1082
$$\log NR(t)=\frac{1}{\sigma_R^2}\{-T(t-t_N)+\frac{1}{2}T^2\}\;,$$

a linear function of time. Its slope is $-\dfrac{T}{\sigma_R^2}.$

A caveat: For widely separated countries, such as China and US, the two Gaussian
peaks are separated in time. These two regions should be treated separately so that

the data for each region has only a single peak. Similarly for Europe and China. See
Appendix. There is also a problem with the aggregate data for the US consisting of
different states with outbreaks separated in time.
1089
1090   **Result**: The natural logarithm of the ratio of two Gaussians of slightly different
1091   standard deviation is approximately a straight line, providing the data used is
1092   homogeneous temporally and spatially.
1093
Sometimes, heterogeneity in one variable could be relaxed in an overall mean. The
case in China is an example. The case could be treated as whole China even though
China included Hubei and Wuhan and the very different region of China outside
Hubei, for the outbreak occurred at the same time. But the cases in China and
European countries and US could not be treated as a whole, for they are separated
both spatially and temporally.   Consequently, it would be impossible to talk about a
single distribution for the global prediction.
1101
Time-varying σ and t_N
The above derivations continue to hold even for σ being a function of time. This
becomes relevant when, for example, after the epidemic passes its peak the policy
makers decide to relax the restrictions on social distancing. Such actions alter the
course of the epidemic and create a longer tail of new cases. *N(t)* is no longer
Gaussian because it does not have fore-aft symmetry.   Since such government
actions are not included in the initial prediction, the prediction scheme needs to
adapt using real-time data after the peak. Our method allows for it.
1110
1111
The peak infected cases:

Writing $N(t_N) = N(t_B)\exp\{\int_{t_B}^{t_N} n(t)dt\}$, and noting $n(t) = \dfrac{d}{dt}\log N(t) = -\dfrac{(t - t_N)}{\sigma_N^2}$, the

exponent is $\int_{t_B}^{t_N} n(t)dt = \dfrac{(t_N - t_B)^2}{2\sigma_N^2} = \dfrac{1}{2}n(t_B)(t_N - t_B)$. Hence the peak infected case number

can be predicted, using the predicted value for t_N starting from a conveniently chosen

time t_B, such as the latest time with data available, as:

$N(t_N) = N(t_B)\exp\{\dfrac{n(t_B)(t_N - t_B)}{2}\}$.

1118
1119   **Accumulated quantities.**
1120   To calculate *I(t)* using reported data, only confirmed cases are used (We call it AIC).
1121   It is given by the accumulated newly confirmed cases minus the accumulated
1122   confirmed recovered. Since the accumulation of early poor data can introduce
1123   errors a more local-in-time formula is given as:
1124

$$I(t) = \int_{-\infty}^{t} N(t)\,dt - \int_{-\infty}^{t} R(t)\,dt = \int_{-\infty}^{t} N(t)\,dt - \int_{-\infty}^{t} N(t-T)\,dt$$

$$= \int_{t-T}^{t} N(t)\,dt.$$

1125

That is, to find I at time t, one only needs to add up the daily newly infected case
numbers for a period of T preceding t. This is an almost local-in-time property even
for this accumulated quantity. For validation, we estimate the peak of the I case
number on 18 February by computing the sum of daily newly infected case numbers
for 15 days, from February 4 to February 18, which yields a peak value for the total
infected cases on 18 February of 54,747. This is within 10% of the reported number
of 57, 805, even after taking into account the deaths (by subtracting the
accumulated deaths of 2,004 from our estimate).
1134
The Total Infected Cases (TIC) is one of the most reported statistics:

$TIC(t) = \int_{0}^{t} N(t)\,dt.$

The total infected cases for the epidemic for a region after it is over is given by
approximately:

$TIC_{\infty} = 2 \cdot TIC(t_N)$,

assuming that $N(t)$ is approximately symmetric about its peak at t_N. In reality $N(t)$
may not be symmetric and likely has a long tail. However, since the number of cases
along the tail is small, the above approximation for the total is still good. Since TIC is
officially available at any time t we will use that reported number, projected forward
to the peak t_N, and then doubling it. Caveat: the official number may include
accumulation of early bad data. Hopefully it is a small percentage of the shear size of
the TIC.
1147
The peak AIC number can be predicted as

$I(t_p) = I(t_B)\exp\{\frac{1}{2}\alpha(t_B)(t_B - t_p)\}$, where t_B is the last available data before the

turning point. It is assumed that $\alpha(t)$ lies on a straight line between t_B and t_p.

This is the maximum load that health services need to plan for, and adopt "flatten the
curve" policies to keep this number under the maximum resources available, such as
hospital beds and isolation wards.
1154

Figure 8. The daily newly infected (in blue) and the daily newly recovered (in red), as a function of time for China as a whole (in solid lines) and Hubei (in dotted red and blue lines). The turning point is determined by when the red and blue curves cross.

 Inset: For China outside Hubei.

The problem with data quality

The performance of all data-driven models depends on the quality of data. In general, there are problems with data from all countries. Underreporting is universal to all countries at least during the initial period of the epidemic. The causes might be different. In China, for example, the data collection was erratic in the early days for lack of awareness of the novel virus. A more common cause was the shortage of test kits to identify the cases. The situation is made more complicate by the existence of asymptomatic cases. However, whether to include those cases or not depends on the definition. In this study, by definition, the cases are limited to confirmed ones with symptoms, for only those cases would tax the medical resources. In fact, the best any country can do is to record the confirmed new case accurately. The cases for death seem to be simple, for the number of deaths has attracted great attention by the government and public alike. Many models are designed to follow only death cases. However, even here, the numbers are not problem-free. For lack of testing, large number of unattributed COVID-19 deaths exists in every country. Finally, the most critical problem is for the recovered case numbers. It is important to have this record, for it is essential for the conservation law discussed above. Furthermore, the new to recovery/removal case ratio (NR Ratio) provides a robust prediction for the turning point of the epidemic. Unfortunately, the recovered case numbers had attracted the least attention. As a result, it is not recorded or reported in some countries, such as UK. Under this condition, many of

1185 the tracking and prediction can still be made as demonstrated in the main text. However,
1186 we strongly urge data for the recovery case to be recorded accurately, for it could not
1187 only provide accurate prediction for medical resource preparation, but also as a measure
1188 of the medical system efficacy.
1189
Finally, a few words on data homogeneity. All models, statistical or mechanistic, require
homogeneous condition. There are actually two types of homogeneity: temporal and
spatial. We believe the temporal homogeneity is more critical. For example, China can
be treated as a whole entity even with part of China, such Hubei and Wuhan under strict
lockdown, and part of China is not. The result still makes sense because of its temporal
homogeneity. Temporal inhomogeneity, however, would render the data nonsensical.
Let us take the global condition as an example. There is a gap in time between the
Chinese cases and the rest of the world (see Figure 9). No model should treat the two
events as one. See Figure 10. On the other hand, the European country and the US could
be treated as the World exChina, for the outbreak in European countries and the US
happened at about the same time, even though they are spatially separated widely.
1201

1202
Figure 9. This figure shows the Reproduction Number in real time starting on 20
January. In this presentation, one can see the pandemic is consisted of a serial
event with severe data inhomogeneity in both space and time. The outbreaks were
almost over in China and South Korea before the earliest cases in Italy and Spain
started to rise. There were 40 days difference between China and South Korea on
one hand, and the cluster of the pandemic flared up in European countries and the
US on the other hand.
1210
1211

Figure 10. China and the rest of the world should be treat separately.

Acknowledgements: NEH and FQ are supported by the National Natural Science Foundation of China under Grant 41821004. KKT's research is supported by the Frederic and Julia Wan Endowed Professorship.
Competing Interests: The authors declare no competing interests.
Data Availability: All data in this study are publicly available from World Health Organization (WHO) at https://www.who.int/emergencies/diseases/novel-coronavirus-2019/situation-reports/
and on the Daily Brief site of the China's National Health Commission at
http://en.nhc.gov.cn/
The Korean data is available at
https://sa.sogou.com/new-weball/page/sgs/epidemic
Coronavirus COVID-19 Global Cases by Johns Hopkins CSSE
https://gisanddata.maps.arcgis.com/apps/opsdashboard/index.html#/bda759474 0fd40299423467b48e9ecf6

https://github.com/CSSEGISandData/COVID-19/tree/master/csse_covid_19_data/csse_covid_19_time_series

References

1238

1.   Cyraboski D. When will the coronavirus outbreak peak? Nature News. 2020.

2.   Adams D. Modelers struggle to predict the future of the COVID-19 pandemic.
The Scientist. 2020.

3.   Kermack W, McKendrick A. A contribution to the mathematical theory of
epidemics. . Proc Roy Soc London 1927;A115:700-21.

4.   Ferguson NM. Impact of non-pharmaceutical interventions (NPIs) to reduce
COVID-19 mortality and healthcare demand. Spiral. 2020.

5.   Flaxman S. Report 13: Estimating the number of infections and the impact of
non-pharmaceutical interventions on COVID-19 in 11 European countries. Spiral.
2020.

6.   Sanche S, Lin YT, Xu C, Romero-Severson E, Hengartner N, Ke R. High
contagiousness and rapid spread of severe acute respiratory syndrom coronavirus 2.
Enmerging Infectious Disease. 2020;26(7).

7.   Bendavid E, Mulaney B, Sood N, Shah S, Ling E, Bromley-Dulfano R, et al.
COVID-19 antibody seroprevalence in Santa Clara county, California. medRxiv. 2020.

8.   Susser M, A. A. Causes of death: epidemic, infectious, and zymoptic diseases.
Vital Statistics: A memorial volume of selections from the reports and writings of
William Farr: Scarecrow Press Inc.; 1975. p. 317-21.

9.   Murray C. Forecasting COVID-19 impact on hospital bed-days, ICU-days,
ventilator-days   and deaths by US state in the next 4 months 2020.

10.   Jewell NP, Lewnard JA, Jewell BL. Caution Warranted: Using the Institute for
Health Metrics and
Evaluation Model for Predicting the Course of the COVID-19 Pandemic. Annuals of
Internal Medicine. 2020;173(3).

11.   Woody S, Tec M, Dahan M, Gaither K, Lachmann M, Fox SJ, et al. Projections
for first-wave COVID-19 deaths
across the US using social-distancing
measures derived from mobile phones. Preprint. 2020.

12.   Fine PEM. John Brownlee and the measurement of infectiousness: An
historical study in epidemic theory. Jornal of Royal Statistical Society A.
1979;142:347-62.

13.   Brownlee J. Statistical studies in immunity. The theory of an epidemic. Proc
Roy Soc Edinburgh. 1907;26:484-521.

14.   Brownlee J. Statistical studies in immunity:the theory of an epidemic. Proc R
Soc Edinburg. 1907;26:484-521.

15.   Huang NE, Qiao F. A data driven time-dependent transmission rate for
tracking an epidemic: a case study of 2019-nCoV. 65, 425-427 , . Sci Bull.
2020;65:425-7.

16.   Delamater PL, Street EJ, Leslie TF, Yang YT, H. JK. Complexity of the Basic
Reproduction Number (R0). Emerging Infectious Diseases. 2019;25.

17.   Wu JT, Leung K, Leung GM. Nowcasting and forecasting the potential
domestic and international spread of the 2019-nCoV outbreak originating in Wuhan,
China: a modelling study. . Lancet. 2020;395(10225):689-97.

18.   Anderson RM, Heesterbeek H, Klinkenberg D, Hollingsworth TD. How will
country-based mitigation measures influence the course of the COVID-19 epidemic?
Lancet. 2020;395:931-4.
19.   Wallinga J, Teunis P. Different epidemic curves for severe acute respiratory
syndrome reveal similar impacts of control measures. Ameri J Epidemiology.
2004;160:509-16.
20.   Fraser C, Donnelly CA, Cauchemez S, Hanage WP, Van Kerkhove MD.
Pandemic potential of a strain of influenza A (H1N1): Early findings. Science.
2009;324:1557-61.
21.   Thakkar N, Burstein R, Hu H, Selvaraj P, Klein D. Social distancing and
mobility reductions have reduced COVID-19 transmission in King County, WA.
Institute for Disease Modeling; 2020.
22.   Murray JD. Mathematical Biology I. Third ed: Springer. 551 p.
23.   Lauer SA, Grantz KH, Bi Q, Jones FK. The incubation period of coronavirus
disease 2019 (COVID-19) from publicly reported confirmed cases: estimation and
application. Annals of Internal Medicine. 2020;172:577-82.
1298