1   **A data-driven tool for tracking and predicting the course of COVID-19**
2   **epidemic as it evolves**
3
4

5   **Authors**:
6   Norden E. Huang[#], Fangli Qiao[#] and Ka-Kit Tung[*]
7
Affiliations:
[#] *Data Analysis Laboratory, FIO, Qingdao* 266061, *China*
[*] *Department of Applied Mathematics, University of Washington, Seattle, WA* 98195
[#] Co-first authors: These authors contribute equally to this work.
FQ: qiaofl@fio.org.cn. NEH: norden@ncu.edu.tw
*Corresponding author: ktung@uw.edu
14

15 **KEYWORDS**
16 Covid-19; Epidemiology; Data-driven approach; prediction of turning point; Existing
17 Infected Cases; peak EIC number; duration of hospital stay; end of epidemic; local-
18 in-time metric.

19
20

21 **ABSTRACT**
22
For an emergent disease, such as Covid-19, with no past epidemiological data
to guide models, modelers struggle to make predictions of the course of the
epidemic (1), and when predictions were made the results would vary widely.
Yet much empirical information is already contained in the data of evolving
epidemiological profiles. We show, for epidemics of low fatality rate, both
empirically with data, and theoretically, how the ratio of daily infected and
recovered cases can be used to track and predict the course of the epidemic.
Ability to predict the turning points and the epidemic's end is of crucial
importance for fighting the epidemic and planning for a return to normalcy.
The accuracy of the prediction of the peaks of the epidemic is validated using
data in different regions in China showing the effects of different levels of
quarantine. The validated tool can be applied to other countries where Covid-
19 has spread, and generally to future epidemics. A preliminary prediction for
South Korea is made with limited data, with end of the epidemic as early as the
second week of April, surprisingly.
38
SIGNIFICANCE: We offer a practical tool, as an alternative to traditional models, for
tracking and predicting the course of an epidemic using the daily data on the
infection and recovery. This data-driven tool can predict the turning points two
weeks in advance, with an accuracy of 2-3 days, validated using data from various
regions in China selected to show the effects of quarantine. It also gives information
on how rapid the rise and fall of the case numbers are.   Although empirical, this
approach has a sound theoretical foundation; the main components of the results
are validated after the epidemic is near an end, as is the case for China, and
therefore generally applicable to future epidemics of low fatality rate.
48
49

Main text:

Introduction.
The current COVID-19 epidemic is caused by a novel corona virus, designated
officially as SARS-CoV-2, spreading from Wuhan, the capital city of Hubei province in
China (*2-4*). The new virus seems to have characteristics different from SARS
(severe acute respiratory syndrome) (*5, 6*): it is less deadly but more virulent (*7-10*).
Modeling the epidemic as it develops has been difficult (*1*). Depending on the model
assumptions, predictions of when it "turns a corner" varies wildly (*11-21*), from now
or until after 650 million people have been infected before peaking in the "worst-
case scenario" (*22*). Now as the epidemic has spread beyond China (*23, 24*), a
reliable prediction of the course of the outbreak in each region is critical for the
management and containment of the epidemic, and reducing public anxiety and
panic. China has instituted some of the strictest quarantine measures around Wuhan
and Hubei, which may or may not be adoptable in other countries (*25-27*). It would
be useful to extract the dependence of the epidemic's evolution on the degree of
quarantine to guide policy decisions, while also to characterize properties of Covid-
19 that are applicable to other countries.

The turning point and the end of the epidemic are the two most watched markers on
its development (*28, 29*). There are various definitions of the turning point. A
common one defines the turning point of the epidemic as the reported daily number
of newly infected reaching a peak and then declining. This is the one touted in the
various news announcements, and also used by some research groups (22). The fact
that the number of newly infected reaching a peak and then declining does not
necessarily imply that the epidemic has "turned a corner", because the total number
of still-infected can still be rising with the associated urgent need for additional
medical resources, such as hospital beds and isolation wards. Furthermore, locating
this peak is highly susceptible to data glitches and change in diagnostic definition.
For example, on 12 February, when Hubei changed its definition of confirmed
infection from the gold standard of nucleic acid gene-sequencing tests to clinical
observations and radiological chest scans, over 14,000 newly infected cases were
added that day, creating a peak that has not been exceeded since. Overwhelmed
doctors in Wuhan pleaded for the change so that they did not have to wait for the
returned tests to confirm the infection. If the definition of the turning point based on
the peak of newly infected were used, it would have given 12 February as the
turning point for Hubei. Outside Hubei, there was no change in definition for the
"infected".

A more meaningful turning point should be based on the number of confirmed
infected individuals, designated as the Existing Infected Cases (*EIC*)(*15*), reaching a
peak and then starting to decline. *EIC* is in theory obtainable from data of the daily
number of newly infected, *N(t)*, and the daily number of newly recovered, *R(t)*, by
subtracting the accumulated sum of *R(t)* from the accumulated sum of *N(t)*. Analysis
of this accumulated quantity is sensitively affected by accumulation of poorer early
data of reported cases, including under-reporting and under-detection of the

number of infected caused by insufficient test kits, in addition to the history of
changing diagnostic criteria. Moreover in practice its peak is often not detected
until several weeks after it has occurred.
99
Since the maximum of *EIC* can be located by the zero of its derivative, we propose
using a local-in-time metric of $N(t_p)=R(t_p)$ at the peak of *EIC*, t_p. We demonstrate
that for the ongoing COVID-19 epidemic, this determination of the turning point is
not sensitive to past data problems, including the rather dramatic increase in *N(t)*,
on 12 February, when Hubei changed its definition of "confirmed infected". Also
since it uses the newest diagnostics, with the testing facilities ramped up, hopefully
the numbers are more accurate.
107
108
109

110
111
112
113
114 **Figure 1**. The daily newly infected (in blue) and the daily newly recovered (in red),
115 as a function of time for China as a whole (in solid lines) and Hubei (in dotted lines).
116 The turning point is determined by when the red and blue curves cross.
117   **Inset:** For China outside Hubei.
118
119
Fig. 1 shows how this turning point is empirically determined using daily time series
of reported *N(t)* and *R(t)*. For China as whole, t_p is found to be February 18; for
Hubei, the province of the epicenter Wuhan, t_p is found to be 19 February, and for
China outside Hubei (China exHubei), 12 February, coincidentally on the same day
as the Hubei data spike. However there is no such bump in the data outside Hubei,
and so is not likely the result of the data artifact. These results, even including that

for Hubei, are not affected by the historical data problems because of our local-in-
time method for determining the turning point.
128
The fact that the turning point for the epidemic in China exHubei occurred earlier
than that for Hubei could reveal the effectiveness of the quarantine of Hubei. In
Wuhan, with hospitals facing the number of infected far exceeding available hospital
beds in the initial period, some infected patients were not adequately isolated.
Secondary and tertiary infections might have played a role in delaying the turning
point. On the other hand, outside Hubei, hospitals were not as overwhelmed
because of the strict quarantine placed on Hubei, which drastically reduced the
import of the disease originating from Hubei. The infected were better isolated,
reducing further spread, and treated in hospitals, resulting in shorter time to
recovery (see Table S1).
139
EIC corresponds to *I(t)* in the traditional SIR (susceptible-infected-recovered)
model(*28*), if deaths are not counted in *R(t).* Most predictions have used models
similar to SIR, though some current ones are much more sophisticated (*12-14, 17,*
21), but they all rely on parameters, such as contact, infection rates, time between
secondary and first infections, and case fatality rates. None of them are known with
any certainty (*1*). Most model predictions of the turning point have the epicenter
Hubei leading the rest of China by 1-2 weeks in its predicted turning point, the
opposite of what the data show. In many SIR types of models, an epidemic would
end after most people are infected and acquire immunity. These models tend to
have the disease run its course sooner the earlier it started.
150
151 Can such a turning point be predicted before it happened, and if so by how many
152 days in advance?
153
154 **Determining the epidemiological characteristics**
155
156 We define the *N* to *R* ratio as
157 $$NR(t)=N(t)/R(t).$$
158 At t_p, $NR=1$.
159
We show in Figure 2, using the data of the epidemic for COVID-19, that the
logarithm of *NR(t)* lies on a straight line, with small scatter, passing through the
turning point $t_p.$ And data for various stages of the epidemic, from the initial
exponential growth stage, to near the peak of *EIC*, and then past the peak, all lie on
the same straight line. The intercept with *logNR=0* yields the turning point. This
line, obtained by linear-least-square fit in the semi-log plot, is little affected by the
rather large artificial spike in the data on 12 February because of its short duration
and the logarithmic value. That reporting problem is necessarily of short duration
because, on the date of definition change, previous week's cases of infected
according to the new criteria were reported in one day. After that, the book is
cleared, and *N(t)* returned to its normal range.
171

172
173
Figure 2. Logarithm of the ratio of daily newly infected to newly recovered. They lie
on straight lines with some small scatter. The straight line obtained by linear-least
squares fit is in dotted line. The slopes of the lines are almost the same but with
different intercept; the trend lines cross zero (the black solid line) at different time
for different regions indicating different peaking time for *EIC*. The epicenter Wuhan
(green) has latest turning point than its province Hubei (pink), which has a later
turning point than China as a whole (cyan).

181
It would be interesting to understand why the empirically determined log *NR(t)* lies
on a straight line, and what determines its slope. See Method for a theoretical
support. For a disease with a low fatality rate, which COVID-19 is (*30*), most newly
infected individuals would eventually recover after a hospital stay of *T* days. So
R(t)~N(t-T). This simple observation lies at the heart of our justification for the
straight line for log(*NR*). In Figures S2 and S4, this relationship is validated using
lagged correlation, at a very high value of 0.95. It is however not assumed in our Fig.
2, which is entirely empirical.

190
The theoretical result in Method suggests that the slope of the linear line is $-T/\sigma_2^2$,
where σ_2 is the standard deviation of the *R(t)* profile. In general, the slope can be
different for different regions with different levels of quarantine and epidemic
characteristics. The hospital treatment efficacy would influence *T* directly, as we
also found. The effect of quarantine would influence the value of σ_1, the standard
deviation of the newly infected, and so indirectly *R(t)* and σ_2. Our empirical result
from Fig. 2 however shows that the slope is the almost the same for different
regions in China, implying that efficacy of treatment and level of quarantine affect *T*
and σ^2 proportionally.
200

Predictability

Since the logarithm of *NR* lies on a straight line passing through the turning point of *EIC*, it would be interesting to explore if the turning point can be predicted by extrapolation using data weeks before it happened (see Figure S1). How far in advance this can be done appears to be limited by the poor quality of the initial data. Fig. 3 shows the results of such predictions (See Method). The horizontal axis indicates the last date of the data used in the prediction. The beginning date of the data used is 24 January for all experiments. Prior to that day, data quality was poor and the newly recovered number was zero in some days, giving an infinite *NR* ratio.

For China outside Hubei, the prediction made on 6 February gives the turning point as 14 February, two days later than the truth. A prediction made on 8 February already converged to the truth of 12 February, and stays near the truth, differing by no more than fractions of a day with more data.

The huge data glitch on 12 February in Hubei affected the prediction for Hubei, for China as whole, and for Hubei-exWuhan. These three curves all show a bump up starting 12 February, as the slope of *N(t)* is artificially lifted. Ironically, predictions made earlier than 12 February are actually better. For example, for China as a whole, predictions made on 9 February and 10 February both give 19 February as the turning point, only one day off the truth of 18 February. A prediction made on 11 February actually gives the correct turning point that would occur one week later. At the time these predictions are made, the newly infected cases were rising rapidly, by over 2,000 each day, and later by over 14,000. It would have been incredulous if one were to announce at that time that the epidemic would turn the corner a week later.

Even with the huge spike for the regions affected by the Hubei's changing of diagnosis criteria, because of its short duration the artifact affects the predicted value by no more than 3 days, and the prediction accuracy soon recovers for China as a whole. For Hubei, the prediction never converges to the true value, but the over-prediction is only 2 days. This smallness of the error is remarkable given that other model predictions differ by weeks or months.

Table S1 lists the mean and standard deviation of the predictions. For applications to other countries and to future epidemics without a change in the definition of the "infection" to such a large extent, we expect even better prediction accuracy.

Figure 3 Prediction of the turning point in *EIC* by extrapolating the trend in
logarithm of *NR* (see Method). The horizontal axis indicates the date the prediction
is made using data prior to that date. The vertical axis gives the dates of the
predicted turning point. Dashed horizontal lines indicated the true dates for the
turning point, as determined from Fig. 1.

Inferring statistical characteristics of the epidemic
Interestingly, the derivative of log *N(t)* or log *R(t)* also lies on a straight line, as
shown in Fig. 4 (although the scatter is larger as to be expected for any
differentiation of empirical data)*.* The positive and negative outliers one day before
and after 12 Feb are caused by the spike up and then down, with little effect on the
fitted linear trend (but increases its variance and therefore uncertainty). Moreover,
the straight line extends without appreciable change in slope beyond the peak of
N(t), suggesting that the distribution of the newly infected number is approximately
Gaussian. For an exponential function, the derivative of its logarithm being a linear
function of time is highly suggestive of a general type of distribution including
Gaussian and Rayleigh.   The recovery time *T* can be determined as t_1-t_0, where t_1 is
the peak of *R(t)* and t_0 is the peak of *N(t).* These two peak times can be obtained by
extending the straight line in Fig. 4 to intersect the zero line. This predicted result
can be verified statistically after the fact by the lagged correlation of *R(t)* and *N(t)*. If
the distribution is indeed Gaussian or even approximately so, the slope in Fig. 4
would be proportional to the reciprocal of the square of its standard deviation, $\boldsymbol{\sigma}$, as:

264
$$\frac{d\log N(t)}{dt} = \frac{-(t-t_0)}{\sigma_1^2}.$$

Similarly result holds for the daily number of recovered, *R(t)*.

The inferred statistical characteristics of the Covid-19 epidemic are summarized in
Table S2 for various regions. The mean recovery time *T,* is about 13 days for China
as a whole. For Wuhan, the city at the epicenter whose hospitals were more
overwhelmed and the patients admitted into hospitals more seriously ill than those
in other provinces, $T \sim 16$ days, while that for Hubei is 14 days. The standard
deviation, σ, is found to be around 8 days, with slight difference between that for
N(t) and for *R(t),* with one exception for Hubei outside Wuhan. Such a fine
subdivision may not be practical for the data quality we have.   The σ tends to be
smaller for China as a whole than Wuhan. One can see that *T* and σ^2 indeed varying
approximately in proportion.
278

279
Figure 4 The derivative of the logarithm of daily newly infected or recovered.
Notice the clear separation of the new and recovered cases and also the subtle
difference of their slopes. The zero crossings of the trend line give the peak dates of
the new and recovered case respectively.   And the slopes give an estimate of σ
values. In this Figure, the following abbreviations are used: C=China; H=Hubei;
N=New Case; R=Recovered.
286
Estimate of "all clear" declaration
We can now estimate a time for a declaration of "all clear". No verification is yet
possible as the predicted date has not occurred. At the turning point, the *EIC* is still
at its peak. For the disease to have run its course, and an "all clear" declaration can
be announced, we require that the newly infected case number to drop to zero, for
prediction practice measured by three standard deviations from the peak of *N(t)*.
Then we wait for two incubation periods, each 14 days, to pass, before we declare

"all clear". Using the inferred disease characteristics in Table S1, our prediction is,
for China outside Hubei: the last week of March. For China as a whole: the first week
of April, barring "imports" of infected from abroad. At this point there may still be
some patients in the hospital who are infected with the virus. The "all clear" call
assumes that these patients are not roaming freely to cause new infections.
299

South Korea
Finally, we apply the present approach the still expanding outbreak in South Korea,
with very limited data. We estimate that the turning point for *EIC* is on March 11.
See Method. An estimate of the end of the epidemic can be given as the second week
of April, using the estimated value for t_0= *3* March, *σ=4.5* days. Remarkably, this date
is around the same time as for Wuhan, China. South Korea owes its quick turning
point and end of the epidemic date to its ability to identity the first infection and the
secondary infections at Shincheonji Church (*31*), where most of the infected were
concentrated. This is reflected in the data: **σ** for South Korea is only half that of
China, with a more rapid rise and fall of the newly infected. Its data for the newly
infected are probably more accurate compared to other countries in similar stage of
the epidemic, due to its massive and speedy (within 6 hours) testing of the
population in its "trace, test and treat" policy.
313

Conclusion.
We offer an alternative data-driven approach to track and predict the course of the
epidemic. Many parameters characterizing an epidemic can be determined from
local-in-time data. Validated by real data, we suggest that our approach could be
applied not just to the current Covid-19 epidemic, but also generally to future
epidemics of low fatality rates. It could also be used as a practical tool for epidemic
management decisions such as quarantine institution and medical resource
planning and allocations (32-35).

327 **METHOD**
328
329 **Theoretical support:**
330 The *NR* ratio is defined as:
331

$$NR(t) = \frac{N(t)}{R(t)} \ .$$

333
334 For an epidemic like COVID-19, where the case fatality rate is low (at around 1%),
335 most of the infected would eventually recover; therefore, we have, as will be verified
336 later:

337 $$R(t)=N(t\text{-}T) \ ; \ NR(t) = \frac{N(t)}{N(t-T)} \ ,$$

338
where T is the hospital stay period before recovery, with its value governed by the
efficacy of the treatment. Using real data, we show that this ratio follows a straight-
line trend. To explain this intriguing feature, we find theoretical support based on
Gaussian distributions for the daily new and recovered case numbers. Gaussian
distribution is a simple and reasonable form for a distribution that has a single peak,
with rapid rise, plateauing near the peak and then declining rapidly. Later, we will
verify using actual data for China that they are indeed very close to Gaussian.
346

$$NR(t) = \frac{N(t)}{N(t-T)} = \frac{exp\left\{-\dfrac{1}{2\sigma^2}(t-t_0)^2\right\}}{exp\left\{-\dfrac{1}{2\sigma^2}(t-t_0-T)^2\right\}} \ ; therefore,$$

$$log\ NR(t) = -\left\{\frac{(t-t_0)^2}{2\sigma^2} - \frac{\left(t-t_0-T\right)^2}{2\sigma^2}\right\} = -\frac{2T(t-t_0)-T^2}{2\sigma^2}$$

348
349 a linear function of t. The intercept with 0 yields $t_p = t_0 + \frac{1}{2}T$.

350
351 In reality, the distribution is only approximately Gaussian, of course. But the
352 approximation is very close for the central part of the distribution near the peak. In
353 fact, central limit theory would favor a Gaussian distribution when the data base is
354 large.
355
Empirically, we find that the σ value for *N(t)* and *R(t)* are close to each other but
slight differences exist, as shown in Table S2. This is to be expected, for even though
the new and recovered case happen in tandem with former leading the latter, the
hospital treatment and stay constitute effectively a smoothing filter on *N(t)* to
produce *R(t)*. The hospital process tends to spread the *R(t)* distribution wider, thus

yield a slightly larger σ values. Given the scatter of the differentiation done for
Figure 4 to infer individual distribution characteristics, the difference may or may
not be significant. More data from various regions under different conditions may
resolve this problem in the future. Taking this difference into account the form of
real *NR* should be modified to be:

366
$$NR(t) = \frac{N(t)}{N(t-T)} = \frac{exp\left\{-\frac{1}{2\sigma_1^2}(t-t_0)^2\right\}}{exp\left\{-\frac{1}{2\sigma_2^2}(t-t_0-T)^2\right\}} \; ; therefore,$$

367
$$log\ NR = -\left\{\frac{(t-t_0)^2}{2\sigma_1^2} - \frac{(t-t_0-T)^2}{2\sigma_2^2}\right\} = -\frac{\left(\sigma_2^2-\sigma_1^2\right)(t-t_0)^2 + 2\sigma_1^2 T(t-t_0) - \sigma_1^2 T^2}{2\sigma_1^2\sigma_2^2}$$

368 As the values of σ_1 and σ_2 are very close based on the empirical data, the quadratic
369 term is always small comparing to the other terms for the length of time we are
370 considering here. Hence.
371

$$logNR(t) = \frac{1}{\sigma_2^2}\left\{-Tt + T^2\right\}: \; a\ linear\ function\ of\ time.$$

$$\therefore \; \frac{dlogNR(t)}{dt} = \frac{-T}{\sigma_2^2} \quad almost\ constant$$

373
374 The turning point is still determined by log *NR*=0, yielding a theoretical value of
375 $t_p=t_0+T/2$. This theoretical value can be used when the data on *R(t)* is not available.
376
377 If the daily data is indeed near Gaussian, then for the daily newly infected cases, we
378 should have approximately,
379

$$N(t) = exp\{-\frac{(t-t_0)^2}{2\sigma_1^2}\} \; ; therefore,$$

380
$$log\ N(t) = \frac{-(t-t_0)^2}{2\sigma_1^2} \quad and \quad \frac{d\ log\ N(t)}{dt} = -\frac{(t-t_0)}{\sigma_1^2}.$$

381 The same is true for the recovered cases, except with t_1 replacing t_0 and σ_2 replacing

382 σ_1.

383
Importantly, the real data indeed validate a near-straight line function for *NR*
throughout all phase of the epidemic, and the near-Gaussian distributions for both
N(t) and *R(t)*. Straight line functions are easy to extend and making predictions easy
and robust. These properties also enable us to infer many of the key statistical

characteristics of the epidemic from empirical data, such as the turning point,
peaking times t_0 and t_1 and the σ of the distributions from the formulas given above.
390
391 There are some subtle points that need to be discussed further. Comparing the *NR*
392 ratio approach and the derivative of individual distribution approach, we can see
393 that the *NR* ratio is much smoother; however, the derivative of individual
394 distribution is richer in information for predicting the 'all clear' time shown later.
395
Validation
a. Lagged correlation
First, we validate statistically using lagged correlation between *N(t)* and *R(t)* the
relationship between the two. Figures S2 and S4 show that they are highly
correlated: with correlation coefficient of 0.95 when both distributions are
smoothed with 5-point box car. The unsmoothed daily data also yield a high
correlation coefficient of 0.80, with *R(t)* lags *N(t)* by $T \sim 15$ days. Both of the
correlation coefficients are statistically significant. The result on *T* is consistent
with that estimated or predicted using the slope of the distribution in Figure 4. The
latter, obtained by the intercept of the straight line, is less accurate because of the
slope is rather shallow.
407
b. Gaussian distribution
A Gaussian distribution is completely characterized by the location of the peak and
the standard deviation. These quantities are determined from the slopes in Figure 4,
and therefore there are no free parameters. Even without the use of disposable
parameters, the fit of Gaussian to the actual distribution is adequate, as can be seen
in Figure S3. The corresponding correlation and Gaussian fits for Hubei province
are given in Figures S4 and S5.
415
416 **c. *EIC***
417 *EIC* is the accumulated newly infected minus the accumulated recovered. Given the
418 result in **a**, a simpler calculation can be performed which avoids the early poor data:
419

$$EIC(t) = \int_{-\infty}^{t} N(t)\,dt - \int_{-\infty}^{t} R(t)\,dt = \int_{-\infty}^{t} N(t)\,dt - \int_{-\infty}^{t} N(t-T)\,dt$$

420
$$= \int_{t-T}^{t} N(t)\,dt.$$

That is, to find *EIC* at time *t*, one only needs to add up the daily newly infected case
numbers for a period of *T* preceding *t*. This is an almost local-in-time property even
for this accumulated quantity. For validation, we estimate the peak of the EIC
number on 18 February by computing the sum of daily newly infected case numbers
for 15 days, from February 4 to February 18, which yields an *EIC* on 18 February of
54,747. This is within 10% of the actual number of 57, 805, even after taking into
account the deaths (by subtracting the accumulated deaths of 2,004 from our
estimate).
429

Estimating the end date of the epidemic:

From the σ and T numbers, one can make predictions on the end of the epidemic as follows. There are two different definitions:

> ***1st End date of the epidemic = t_0+3σ+2* incubation period.***

> ***2nd End date of the epidemic = t_1+3σ+2* incubation period.***

The first one depends on the newly infected case, the second one, on the daily cured cases. If we take the incubation time as 14 days, the end of the epidemic outbreak can be calculated easily from the data given in Table S1. Based on our analysis, Wuhan would come out of the epidemic the latest, long after the rest of the country, at around

> ***1st: February 11 + 3x8 +2x14   or towards the beginning of April.***
> ***2nd: February 24+ 3x8 +2x14   or towards the middle of April.***

The estimate based on the first definition is reported in the main text..

South Korea:
Finally, we will show how this method is applied to the expanding outbreak in South Korea. Figure S6 summarized the available data at the present. The recovered case numbers hovered around 1 and 2 daily up to March 1st. It only picked up toward the end. Starting from 19 February, there seems to be enough new daily infected cases. All these phenomena are not random events, for the South Korea Government has identified that the epic center of the epidemic is at church gathering in the city of Daegu and North Gyeongsang province, where 90% of the cases are found. Specifically, a confirmed COVID-19 patient was reported to have attend the Shincheonji Church of Jesus services twice on February 9th and 16th. Given the incubation period of 7 to 14 days, the initial explosion at February 19th and the first peak value around February 24th are not accidents.

If we use the available daily new cases data, we can get the statistical characteristics of the distribution of the daily new cases from Figure S7, which gives the t_0 as March 3rd and a σ value of 4.5 days. If we further use the turning point as approximately $t_0+T/2$, then the turning point should fall on March 10, assuming T as 14 days based on the over all mean from different regions in China.

For the NR ratio, it is limited by the availability of recovered case number. If we use the limited recovered cases starting from March 1st, we have 7 days of data. The computed the NR ratio together with the trend is given in Figure S8. The turning point, at the zero-crossing of the extended trend line, would occur between March 11th and 12th. This approach does not need to use a value for $T.$

475 It should be pointed out that the Korean data available is only marginal. The
476 predicted date of turning point by *NR* ratio would be between March 11th and 12th;
477 by the derivative of distribution it would be March 10th. The result is not only
478 consistent, but also validated by real data showing the turning point on March 12th,
479 a pleasant surprise.
480
481
482

References

1. David Adam, Modelers Struggle to Predict the Future of the COVID-19 Pandemic. The Scientist, https://www.the-scientist.com/news-opinion/modelers-struggle-to-predict-the-future-of-the-covid-19-pandemic-67261 (2020).

2. WHO, Laboratory testing of human suspected cases of novel coronavirus (nCoV) infection: interim guidance, World Health Organization, Geneva (2020).

3. N. Zhu, D. Zhang, W. Wang, X. Li, B. Yang, J. Song, X. Zhao, B. Huang, W. Shi, R. Lu, P. Niu, F. Zhan, A novel coronavirus from patients with pneumonia in China, 2019. *N. Engl. J. Med.* **382**, 727-733 (2020).

4. R. Lu, X. Zhao, J. Li, P. Niu, B. Yang, H. Wu, W. Wang, H. Song, B. Huang, N. Zhu, Y. Bi, X. Ma, F. Zhan, L. Wang, T. Hu, H. Zhou, Z. Hu, W. Zhou, L. Zhao, …., W. Tan, Genomic characterisation and epidemiology of 2019 novel coronavirus: implications for virus origins and receptor binding. *The Lancet* **395**(10224), 565-574 (2020).

5. Y. Liu, A. A. Gayle, A. Wilder-Smith, J. Rocklöv, The reproductive number of COVID-19 is higher compared to SARS coronavirus. *J. Travel Med.* taaa021 (2020).

6. J. W. Glasser, N. Hupert, M. M. McCauley, R. Hatchett, Modeling and public health emergency responses: Lessons from SARS. *Epidemics* 3: 32-37 (2011), doi:10.1016/j.epidem.2011.01.001.

7. P. Zhou, X. Yang, X. Wang, B. Hu, L. Zhang, W. Zhang, H. Si, Y. Zhu, B. Li, C. Huang, H. Chen, J. Chen, …, Z. Shi, A pneumonia outbreak associated with a new coronavirus of probable bat origin. *Nature* **579**, 270-273 (2020).

8. C. Huang, Y. Wang, X. Li, L. Ren, J. Zhao, Y. Hu, L. Zhang, G. Fan, J. Xu, X. Gu, Z. Cheng, T. Yu, J. Xia, Y. Wei, W. Wu, X. Xie, W. Yin, H. Li, M. Liu, Y. Xiao, H. Gao, L. Guo, J. Xie, G. Wang, R. Jiang, Z. Gao, Q. Jin, J. Wang, B. Cao, Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *The Lancet* **395**(10223), 497-506 (2020).

9. J. F-K. Chan, S. Yuan, K.-H. Kok, K. K.-W. To, H. Chu, J. Yang, F. Xing, J. L. BNurs, C. C.-Y. Yip, R. W.-S. Poon, H.-W. Tsoi, S. S.-F. Lo, K.-H. Chan, V. K.-M. Poon, W.-M. Chan, J. D. Lp, J.-P. Cai, V. C.-C. Cheng, H. Chen, C. K.-M. Hui, K-Y. Yuen, A familial cluster of pneumonia associated with the 2019 novel coronavirus indicating person-to-person transmission: a study of a family cluster. *The Lancet* **395**(10223), 514-523 (2020).

10. X. Xu, P. Chen, J. Wang, J. Feng, H. Zhou, X. Li, W. Zhong, P. Hao, Evolution of the novel coronavirus from the ongoing Wuhan outbreak and modeling of its spike protein for risk of human transmission. *Sci. China Life Sci.* **63**, 457-460 (2020).

11. Z. Chen, W. Zhang, Y. Lu. C. Guo, Z. Guo, C. Liao, X. Zhang, Y. Zhang, X. Han, Q. Li, W. lan Lipkin, J. Lu, From SARS-CoV to Wuhan 2019-nCoV Outbreak: Similarity of Early Epidemic and Prediction of Future Trends. *Biorxiv* preprint (2020), doi: https://doi.org/10.1101/2020.01.24.919241.

12. J. M. Read, J. R. E. Bridgen, D. A. T. Cummings, A. Ho, C. P. Jewell, Novel coronavirus 2019-nCoV: early estimation of epidemiological parameters and epidemic predictions. *medRxiv* preprint (2020), doi: https://doi.org/10.1101/2020.01.23.20018549.

13. J. T. Wu, K. Leung, G. M. Leung, Nowcasting and forecasting the potential domestic and international spread of the 2019-nCoV outbreak originating in Wuhan, China: a modelling study. *The Lancet* **395**(10225), 689-697 (2020).

14. S. Zhao, S. S. Musa, Q. Lin, J. Ran, G. Yang, W. Wang, Y. Lou, L. Yang, D. Gao, D. He, M. S. Wang, Estimating the Unreported Number of Novel Coronavirus (2019-nCoV) Cases in China in the First Half of January 2020: A Data-Driven Modelling Analysis of the Early Outbreak. *J. Clin. Med.* **9**, 388 (2020).

15. N. E. Huang, F. Qiao, A data driven time-dependent transmission rate for tracking an epidemic: a case study of 2019-nCoV. *Sci. Bull.* **65**, 425-427(2020) , https://doi.org/10.1016/j.scib.2020.02.005.

16. Q. Li, W. Feng, Trend and forecasting of the COVID-19 outbreak in China. *J. Infection* arXiv:2002.05866v1, (2020).

17. H. Xiong, H. Yan, Simulating the infected population and spread trend of 2019-nCov under different policy by EIR model. *medRxiv* preprint (2020), doi: https://doi.org/10.1101/2020.02.10.20021519.

18. L. Damon, E. Brooks-Pollock, M. Bailey, M. J. Keeling, A spatial model of CoVID-19 transmission in England and Wales: early spread and peak timing. *medRxiv* preprint (2020), doi: https://doi.org/10.1101/2020.02.12.20022566.

19. H. Sun, Y. Qiu, H. Yan, Y. Huang, Y. Zhu, S. Chen, Tracking and Predicting COVID-19 Epidemic in China Mainland. *Medrxiv* preprint (2020), doi: https://doi.org/10.1101/2020.02.17.20024257.

20. Q. Liu, Z. Liu, D. Li, Z. Gao, J. Zhu, J. Yang, Q. Wang, Assessing the Tendency of 2019-nCoV (COVID-19) Outbreak in China. *medRxiv* preprint (2020), doi: https://doi.org/10.1101/2020.02.09.20021444.

21. L. Peng, W. Yang, D. Zhang, C. Zhuge, L. Hong, Epidemic analysis of COVID-19 in China by dynamical modeling. *arXiv* 2002.06563, (2020).

22. D. Cyranoski, When will the coronavirus outbreak peak? *Nature news* (2020).

23. C. R. MacIntyre, Global spread of COVID-19 and pandemic potential. *Global Biosecurity* **1**(3), (2020).

24. WHO, Coronavirus latest: WHO describes outbreak as pandemic, Nature news (2020), https://www.nature.com/articles/d41586-020-00154-w.

25. K. Kupferschmidt, J. Cohen, Can China's COVID-19 strategy work elsewhere? *Science* **367**(6482), 1061-1062 (2020).

26. J. M. Read, J. R. Bridgen, D. A. Cummings, A. Ho, C. P. Jewell, Novel coronavirus 2019-nCoV: early estimation of epidemiological parameters and epidemic predictions. *medRxiv* (2020), doi:10.1101/2020.01.23.20018549.

27. S. Zhao, Q. Lin, J. Ran, S. S. Musa, G. Yang, W. Wang, Y. Lou, D. Gao, L. Yang, D. He, M. H. Wang, Preliminary estimation of the basic reproduction number of novel coronavirus (2019-nCoV) in China, from 2019 to 2020: A data-driven analysis in the early phase of the outbreak. *Int. J. Infect. Dis.*, 92: 214-217 (2020), https://doi.org/10.1016/j.ijid.2020.01.050.

28. W. Kermack, A. McKendrick, A contribution to the mathematical theory of epidemics. *Proc. Roy. Soc. London A* **115**, 700-721 (1927).

29. D. L. Heymann, N. Shindo, COVID-19: what is next for public health? *Lancet*, 395(10224): 542-545 (2020), https://doi.org/10.1016/S0140-6736(20)30374-3.

30. The Novel Coronavirus Pneumonia Emergency Response Epidemiology Team,
The Epidemiological characteristics of an outbreak of 2019 novel coronavirus
disease (COVID-19)-China, 2020, *China CDC Weekly* (2020).
31. E. Shim, A. Tariq, W. Choi, Y. Lee, G. Chowell, Transmission potential of COVID-19
in   South   Korea.   *medRxiv*   preprint,   (2020),   doi:
https://doi.org/10.1101/2020.02.27.20028829.
32. C. M. Peak, L. M. Childs, Y. H. Grad, C. O. Buckee, Comparing nonpharmaceutical
interventions for containing emerging epidemics. *Proc. Natl. Acad. Sci.* 114(15):
4023-4028 (2017), doi:10.1073/pnas.1616438114.
33. R. S. Dhillon, D. Srikrishna, When is contact tracing not enough to stop an
outbreak?   *Lancet   Infect.   Dis.*, 18: 1302-1304   (2018),
https://doi.org/10.1016/S1473-3099(18)30656-X.
34. X. Pang, Z. Zhu, F. Xu, J. Guo, X. Gong, D. Liu, Z. Liu, D. P. Chin, D. R. Feikin,
Evaluation of control measures implemented in the severe acute respiratory
syndrome outbreak in Beijing, 2003. JAMA, 290(24): 3215-3221 (2003).
35. G. Wang, N. E. Huang, F. Qiao, Quantitative evaluation on control measures for an
epidemic: A case study of COVID-19. *Sci. Bull.* **65** (2020), doi: 10.1360/TB-2020-
0159.

Acknowledgements: NEH and FQ are supported by the National Natural Science
Foundation of China under Grant 41821004. KKT's research is supported by the
Frederic and Julia Wan Endowed Professorship.

Competing Interests: The authors declare no competing interests.
Data Availability: All data in this study are publicly available from World Health
Organization (WHO) at https://www.who.int/emergencies/diseases/novel-
coronavirus-2019/situation-reports/
and on the Daily Brief site of the China's National Health Commission at
http://en.nhc.gov.cn/
The Korean data is available at
https://sa.sogou.com/new-weball/page/sgs/epidemic
Coronavirus COVID-19 Global Cases by Johns Hopkins CSSE
https://gisanddata.maps.arcgis.com/apps/opsdashboard/index.html#/bda759474
0fd40299423467b48e9ecf6
612
613

614 **Supplementary Information:**
615

616
617
Figure S1. Prediction of the turning point of *EIC* using linear least-squares trends
using various data lengths for China exHubei.   All data used start from 24 January.
Different colored straight lines show the linear trend calculated from 24 January to a
particular date. The spread is over a very small range. Then these trends are
extrapolated (extrapolations not shown) to intersect the zero line to yield a
prediction for the turning point. The blue dots are the data.
624

625
626 **Figure S2.** Lagged correlation of *R(t)* with *N(t)* for China as a whole.
627

628
Figure S3. Gaussian fit of *N(t)* and *R(t)*, for China as a whole.

631
Figure S4. Lagged correlation of *R(t)* with *N(t)* for Hubei province.

633
634
635   **Figure S5.** Gaussian fit of *N(t)* and *R(t)*, for Hubei Province.

636
637

	China	Hubei	China-Hubei	Hubei-Wuhan
Truth (data)	18	19	12	15
NR Ratio	20.3±1.6 (Feb 20nd)	22.3±1.0 (Feb 22rd)	12.4±0.9 (Feb 12th)	16.0±1.2 (Feb 16th)

638
639   **Table S1**: Predicted turning point dates. Shown are the mean and standard
640   deviation of the predictions over the prediction period, using the NR ratio method

	Crossing	t_0	t_1	T	Sigma N	Sigma R
China	2/18	2/11	2/24	13	7.5	7.7
Hubei	2/20	2/12	2/26	14	7.9	8.5
Wuhan	2/21	2/14	3/01	16	8.8	8.8
C exHubei	2/12	2/08	2/21	13	7.5	7.1
H exWuhan	2/16	2/13	2/27	14	5.0	8.8

642
643   **Table S2:** Statistical characteristics of the COVID-19 epidemic in different regions in
644   China inferred from data, for N(t), the daily number of newly infected and for R(t),
645   the daily number of recovered.

647
648
Figure S6: The available data from South Korea (as of March 7th). The sporadic
recovered case numbers are mostly in the single digit. If we use the sudden increase
of recovered case matching with the sudden explosive increase of new infected, the
distance is approximately 14 days, a reasonable T value when compared to the
mean value in China. For our data analysis, we used daily newly cases starting
February 19th, for the derivative of individual distribution study; we used data case
from March 1st, for the NR ratio study, in order to have enough recovered cases.
656
657

658
659 **Figure S7**: The derivative of the logarithmic value of daily new infected case
660 distribution.
661
662

663
664

665   **Figure S8**: The *NR* ratio from 7 days of data from March 1^{st} to 7^{th}.   The estimated
666   zero-crossing time would occur between March 11^{th} and 12^{th}, a value consistent
667   with the statistics from the daily new case distribution on March 10^{th}.