A data-driven model for predicting the course of COVID-19 epidemic with applications for China, Korea, Italy, Germany, Spain, UK and USA

(Revised: April 5, 2020)

Authors:
Norden E. Huang[#], Fangli Qiao[#] and Ka-Kit Tung*

Affiliations:
[#] *Data Analysis Laboratory, FIO, Qingdao* 266061*, China*
* *Department of Applied Mathematics, University of Washington, Seattle, WA* 98195
[#] Co-first authors: These authors contribute equally to this work.
FQ: qiaofl@fio.org.cn. NEH: norden@ncu.edu.tw
*Corresponding author: ktung@uw.edu

KEYWORDS
Covid-19; Epidemiology; Covid-19 predictions for Italy, Germany, Spain, UK and USA; Data-driven approach; prediction of turning points; peak active infected number; Covid-19 recovery period; end of Covid-19 epidemic; local-in-time metric for epidemic management.

ABSTRACT

For an emergent disease, such as Covid-19, with no past epidemiological data to guide models, modelers struggle to make predictions of the course of the epidemic (Cyranoski, *Nature News* 18 February 2020). The wildly varying predictions make it difficult to base policy decisions on. On the other hand much empirical information is already contained in data of evolving epidemiological profiles. We offer an additional tool, based on general theoretical principles and validated with data, for tracking the turning points, peak and accumulated case numbers of infected and recovered for an epidemic, and to predict its course. Ability to predict the turning points and the epidemic's end is of crucial importance for fighting the epidemic and planning for a return to normalcy. The accuracy of the prediction of the peaks of the epidemic is validated using data in different regions in China showing the effects of different levels of quarantine. The validated tool can be applied to other countries where Covid-19 has spread, and generally to future epidemics. US is found to have the largest net infection rate, and is predicted to have the largest total infected cases (708K) and will take two weeks longer than Wuhan to reach its turning point, and one week longer than Italy and Germany.

SIGNIFICANCE: We offer a practical tool for tracking and predicting the course of an epidemic using the daily data on the infection and recovery. This data-driven tool can predict the turning points two weeks in advance, with an accuracy of 2-3 days, validated using data from various regions in China selected to show the effects of quarantine. It also gives information on how rapid the rise and fall of the case numbers are, and what the peak and total number of infected are. Although empirical, this approach has a sound theoretical foundation; the main components of the results are validated after the epidemic is near an end, as is the case for China, and therefore is generally applicable to future epidemics.

1. Introduction.

The current COVID-19 epidemic is caused by a novel corona virus, designated officially as SARS-CoV-2, spreading from Wuhan, the capital city of Hubei province in China (*2-4*). The new virus seems to have characteristics different from SARS (severe acute respiratory syndrome) (*5, 6*): it is less deadly but spreads more widely (*7-10*). Modeling the epidemic as it develops has been difficult (*1*). Depending on the model assumptions, predictions of when it "turns a corner" for China varies greatly (*11-21*), up to after 650 million people have been infected before peaking; many have now been shown to be inaccurate (*22*). Now as the epidemic has subsided in China and become a global pandemic (*23, 24*), a reliable forecast of the course of the outbreak in each region is critical for the management and containment of the epidemic, and for balancing the impact from the public health crisis vs the economic crisis. China has instituted some of the strictest quarantine measures around Wuhan and Hubei, which may or may not be adoptable in other countries (*25-27*). It would be useful to extract the dependence of the epidemic's evolution on the degree of quarantine to guide policy decisions, while also to characterize properties of Covid-19 that are applicable to other countries.

Mainstream epidemiological models have their origin in the SIR (Susceptible, Infected, and Recovered or Removed) model (28) and its many variations. We explain in the Supplementary Information why existing model predictions vary so widely by commenting on the assumptions underlying these models.

These SIR-type models, however, serves a critical purpose for long-range policy planning, such as warning policy-decision makers of the gravity of the potential impact and prompting them to take proper actions before it is too late. After the breakout, more information is needed for more detailed planning, such as the arrival of the critical turning points, the number of hospital bed we might need at the peak, and the estimate for when to lift the quarantine, and when to return to normalcy. We offer here an additional tool that has the advantage that it has does not depend on the elusive infection rate or the susceptible population, information needed for most models, but has the disadvantage that it cannot be used when the epidemic first started and the data are inaccurate or incomplete. It is based on daily case numbers (i.e. newly confirmed cases), $N(t)$, and recovered cases, $R(t)$.

Without universal testing, the confirmed case number might be only a subset of the true total infected number, which may never be known unless frequent universal testing is instituted. The asymptomatic infected who are not tested and then recover on their own do not get counted but they also do not tax hospital resources. Nevertheless, they can infect others and some of the latter may develop more serious symptoms that require hospitalization. Then these secondary infections are included in our case data. Our aim is to provide a tool that can be used for the management of medical resources. Since we do not use a model to calculate how the asymptomatic infectives infect others, we do not need to know either the infection rate or the asymptomatic infective numbers. Since those who are admitted to the

99   hospital either recover after a hospital stay of *T* days, or dead after a similar number
100 of days, there should be a delayed relationship between *N(t)* and *R(t)*, which we will
101 explore in the Theory section. Now that the epidemic in China appears to have come
102 to an end, the data from various regions in China can be used to validate the model.
103 After validation we then apply it to other regions in the world.
104
Our estimate of the end date of the epidemic is not based on the number of
susceptibles, *S,* approaching zero as in most models (i.e. most of the population is
infected, hence acquiring immunity), but *N(t)* approaching zero and remaining so
for two incubation periods. The first incubation period is to allow the asymptomatic
infected to show symptoms and the second period to allow those that are infected
by the asymptomatic infected to show symptoms. For prediction purpose, the date
when the *N(t)* is zero is estimated by 3 standard deviations from its peak. These
two quantities can be extracted from the data as the epidemic is developing. Our
estimate of the end of the epidemic is earlier than most model predictions, usually
significantly so, because it does not depend on the herd immunity concept.
115
As is true for all data-driven approaches, our result inevitably depends on the
quality of the data used, and some of the early data of the epidemic are not as good
as the later data, when better diagnostic methods and more complete reporting are
established. However, many of the metrics commonly in use require accumulation
of data from the beginning of the epidemic, and consequently are affected by poor
data or change of diagnostic methods along the way. We try to avoid accumulation
and use local-in-time metrics. Nevertheless, data problems cannot be avoided.
Sensitivity of our conclusion on data problems is extensively discussed in this work.
Figure S1 displays examples of data used in this study. One problem immediately
becomes obvious for the Chinese data: On 12 February, when Hubei changed its
definition of confirmed infection from the gold standard of nucleic acid gene-
sequencing tests to clinical observations and radiological chest scans, over 14,000
newly infected cases were added that day, creating a peak that has not been
exceeded since. Overwhelmed doctors in Wuhan pleaded for the change so that
they did not have to wait for the returned tests to confirm the infection. Outside
Hubei, there was no change in definition for the "infected". How this artifact affects
our conclusion will be discussed.
133
2.Model and its validation using data
Definition: Let $I(t)$ be the number of active infected at time t. Its change is given
by;
$$\frac{d}{dt}I = N(t) - R(t),$$
where $N(t)$ is the number of newly infected, and $R(t)$, designated as removed, is
the sum of the daily recovered and dead. For a disease such as Covid-19 with low
fatality rate, *R(t)* consists of mainly recovered. However even for this disease, the
fatality rate in some regions, such as in Northern Italy, approaches 10%. For these
regions *R(t)* should include the dead as well. Note that for the theory part, *N(t)*

includes both confirmed and unconfirmed cases. The term: Existing Infected Case
(*EIC*) number is used to denote the confirmed *I(t)* when we deal with data.
145

Let t_p , the turning point defined as the peak of the active infected number. At this
point maximum medical resource is needed. This maximum occurs when

$\frac{d}{dt}I = 0,$ implying $N(t_p) = R(t_p).$

This is a local-in-time metric. There is therefore no need to first find *I(t)* to locate
this peak. After the turning point, the newly recovered starts to exceed the newly
infected. The demand for medical resources, such as hospital beds, isolation wards
and respirators, starts to decrease.
153

154   The theoretical foundation for our model is given in Supplementary Information.
155   Here we discuss the main results and offer validation of these results using data
156   from China.
157

158   **Main Result**: The daily newly recovered/removed number *R(t)*, is related to the
159   daily newly infected number *N(t)* as:
160

$$R(t) = N(t - T),$$

for $t>T$, where *T* is the mean recovery period.
This result can be rigorously derived using an age-structured population model (see
reference (36)). It is also common sense: the infected eventually recovered after a
number of days, or dead after a similar number of days. Of course, the number of
days a patient stays in the hospital before discharge depends on the efficacy of
treatment and so varies somewhat, and the time it takes for a patient to die may also
depend on the age and underlying conditions. *T* is therefore a statistical quantity.
Validation: This fundamental relationship can be validated statistically with data.
Figures 1, obtained using data from China during the Covid-19 epidemic, shows that
N(t) and *R(t)* are highly correlated: with correlation coefficient of 0.95 when both
distributions are smoothed with 5-point boxcar. The unsmoothed daily data also
yield a high correlation coefficient of 0.80, with *R(t)* lagging *N(t)* by $T{\sim}15$ days. Both
correlation coefficients are statistically significant. A similar result is found for
Hubei (Figure S2) and other regions (not shown). This is one of the ways the *mean*
recovery period is determined statistically from data, but it is not practical in the
early phase of the epidemic. We will give different methods for the latter purpose.
The result on *T* is consistent with that estimated or predicted later using the slope of
the distribution in Figure 4. The latter, obtained by the intercept of the straight line,
is less accurate because of the slope is rather shallow.
181

182
183 **Figure 1.** Lagged correlation of case numbers $R(t)$ and $N(t)$ for China as a whole.

185 **Main Result:** The natural logarithm of the ratio of N and R is a linear function of
186 time for $t > T$. This relationship is important for the purpose of forecast because it is
187 easy to extrapolate from a straight line into the future.
188

189 **Validation of log NR as a straight line**:
190 From data we use the report newly confirmed case number and the recovered case
191 number to define NR ratio as
192                          $NR(t)=N(t)/R(t)$.
At t_p, $NR=1$.

We show in Figure 2, using the data of the epidemic for COVID-19, that the
logarithm of $NR(t)$ lies on a straight line, with small scatter, passing through the
turning point t_p. And data for various stages of the epidemic, from the initial
exponential growth stage, to near the peak of EIC, and then past the peak, all lie on
the same straight line. The intercept with $logNR=0$ yields the turning point. This
line, obtained by linear-least-square fit in the semi-log plot, is little affected by the
rather large artificial spike in the data on 12 February because of its short duration
and the logarithmic value. That reporting problem is necessarily of short duration
because, on the date of definition change, previous week's cases of infected
according to the new criteria were reported in one day. After that, the book is
cleared, and $N(t)$ returned to its normal range.
206

207
208
Figure 2. Logarithm of the ratio of daily newly infected to newly recovered. They lie
on straight lines with some small scatter. The dotted straight lines are obtained by
linear-least squares fit is. The slopes of the lines are almost the same but with
different intercept; the trend lines cross zero (the black solid line) at different time
for different regions indicating different peaking time for *EIC*. The epicenter Wuhan
(green) has latest turning point than its province Hubei (pink), which has a later
turning point than China as a whole (cyan).
216
The theoretical result in SI suggests that the slope of the linear line is $-T/\sigma_R^2$, where
σ_R is the standard deviation of the $R(t)$ profile. In general, the slope can be different for
different regions with different levels of quarantine and epidemic characteristics.
The hospital treatment efficacy would influence T directly. The effect of quarantine
would influence the value of σ_N, the standard deviation of the newly infected, and so
indirectly $R(t)$ and σ_R. Our empirical result from Fig. 2 however shows that the slope
is the almost the same for different regions in China, implying that efficacy of
treatment and level of quarantine affect T and σ^2 proportionally.
225
Result: The derivative of log $N(t)$ and of log $R(t)$ is each a linear function of time,
with known slope. Their intercept with the zero derivative line yields the time for
their respective peak.

Validation

Empirically, the derivative of log $N(t)$ or log $R(t)$ lies on a straight line, as shown in

Fig. 3 (although the scatter is larger as to be expected for any differentiation of

empirical data)*. The positive and negative outliers one day before and after 12 Feb

are caused by the spike up and then down, with little effect on the fitted linear trend

(but increases its variance and therefore uncertainty). Moreover, the straight line

extends without appreciable change in slope beyond the peak of $N(t)$, suggesting

that the distribution of the newly infected number is approximately Gaussian. The

mean recovery time T can be predicted as $t_R - t_N$, where t_R is the peak of $R(t)$ and t_N

is the peak of $N(t)$. These two peak times can be obtained by extending the straight

line in Fig. 3 to intersect the zero line. This predicted result can be verified

statistically after the fact by the lagged correlation of $R(t)$ and $N(t)$. If the

distribution is indeed Gaussian or even approximately so, the slope in Fig. 3 would

be proportional to the reciprocal of the square of its standard deviation, σ, as (See

SI):

245
$$\frac{d \log N(t)}{dt} = \frac{-(t - t_N)}{\sigma_N^2}.$$

Similarly result holds for the daily number of recovered, $R(t)$.

After the epidemic is nearing the end as is the case in China, fitting the data to a

Gaussian can be done after the fact (see Figures S3 and S4). The fit is satisfactory

even without using any disposal parameters. The parameters used are determined

using slopes of log N and log R (see Table S1)

The inferred statistical characteristics of the Covid-19 epidemic are summarized in

Table S1 for various regions. The mean recovery time T, is about 13 days for China

as a whole. For Wuhan, the city at the epicenter whose hospitals were more

overwhelmed and the patients admitted into hospitals more seriously ill than those

in other provinces, $T \sim 16$ days, while that for Hubei is 14 days. The standard

deviation, σ, is found to be around 8 days, with slight difference between that for

$N(t)$ and for $R(t)$, with one exception for Hubei outside Wuhan. Such a fine

subdivision may not be practical for the data quality we have. The σ tends to be

smaller for China as a whole than Wuhan. One can see that T and σ^2 indeed varying

approximately in proportion.

The peak infected cases:

Writing $N(t_N) = N(t_B) \exp\{\int_{t_B}^{t_N} n(t) dt\}$, and noting $n(t) = \frac{d}{dt} \log N(t) = -\frac{(t - t_N)}{\sigma_N^2}$, the

exponent is $\int_{t_B}^{t_N} n(t)dt = \dfrac{(t_N - t_B)^2}{2\sigma_N^2} = \dfrac{1}{2}n(t_B)(t_N - t_B)$. Hence the peak infected case number

can be predicted, using the predicted value for t_N starting from a conveniently chosen

time t_B , such as the latest time with data available, as:

$N(t_N) = N(t_B)\exp\{\dfrac{n(t_B)(t_N - t_B)}{2}\}$.

Accumulated quantities.
To calculate *I(t)* using reported data, only confirmed cases are used (We call it EIC).
It is given by the accumulated newly confirmed cases minus the accumulated
confirmed recovered. Since the accumulation of early poor data can introduce
errors a more local-in-time formula is given as:

278
$$I(t) = \int_{-\infty}^{t} N(t)\,dt - \int_{-\infty}^{t} R(t)\,dt = \int_{-\infty}^{t} N(t)\,dt - \int_{-\infty}^{t} N(t-T)\,dt$$
$$= \int_{t-T}^{t} N(t)\,dt.$$

That is, to find *I* at time *t*, one only needs to add up the daily newly infected case
numbers for a period of *T* preceding *t*. This is an almost local-in-time property even
for this accumulated quantity. For validation, we estimate the peak of the *I* case
number on 18 February by computing the sum of daily newly infected case numbers
for 15 days, from February 4 to February 18, which yields a peak value for the total
infected cases on 18 February of 54,747. This is within 10% of the reported number
of 57, 805, even after taking into account the deaths (by subtracting the
accumulated deaths of 2,004 from our estimate).
287

288
Figure 3 The derivative of the logarithm of daily newly infected or recovered.
Notice the clear separation of the new and recovered cases and also the subtle
difference of their slopes. The zero crossings of the trend line give the peak dates of
the new and recovered case respectively. And the slopes give an estimate of σ
values. In this Figure, the following abbreviations are used: C=China; H=Hubei;
N=New Case; R=Recovered.
295
3. Predictability
Prediction of the turning point using _NR_ ratio.
We first discuss how the true turning point can be determined from data after it has
occurred. Then we give a method for predicting this true value in advance and
assess the accuracy of the forecasts as function of days in advance when the
prediction is made. A note: after this manuscript was submitted for review the
predictions that we made previously have come to pass. Although consequently the
value of our predictions has greatly diminished, it gives us a chance to compare our
predictions against the truths. This model validation process is important if we are
to apply the same method for prediction to other regions.
306
The turning point and the end of the epidemic are the two most watched markers on
its development (*28, 29*), along with the number of infected at each stage of the
epidemic. There are various definitions of the turning point. A common one defines
the turning point of the epidemic as the reported daily number of newly infected
reaching a peak and then declining. This is the one touted in the various news
announcements, and also used by some research groups (22). The fact that the
number of newly infected reaching a peak and then declining does not necessarily

imply that the epidemic has "turned a corner", because the total number of active
infected can still be rising with the associated urgent need for additional medical
resources, such as hospital beds, isolation wards and ventilators. Furthermore,
locating this peak is highly susceptible to data glitches and change in diagnostic
definition. A more meaningful turning point should be based on the number of
confirmed infected individuals, designated as EIC)(15), reaching a peak and then
starting to decline. EIC is in theory obtainable from data of the daily number of new
confirmed cases, $N(t)$, and the daily number of newly recovered, $R(t)$, by subtracting
the accumulated sum of $R(t)$ from the accumulated sum of $N(t)$. Analysis of this
accumulated quantity is sensitively affected by accumulation of poorer early data of
reported cases, including under-reporting and under-detection of the number of
infected caused by insufficient test kits, in addition to the history of changing
diagnostic criteria. Moreover in practice its peak is often not detected until several
weeks after it has occurred.
328
329 Since the maximum of EIC, can be located by the zero of its derivative, we propose
330 using a local-in-time metric of $N(t_p)=R(t_p)$ at the peak of EIC, t_p.
331
Referring to Figure S1, for China as whole, t_p is found to be February 18; for Hubei,
the province of the epicenter Wuhan, t_p is found to be 19 February, and for China
outside Hubei (China exHubei), 12 February, coincidentally on the same day as the
Hubei data spike. However there is no such bump in the data outside Hubei, and so
is not likely the result of the data artifact. These results, even including that for
Hubei, are not affected by the historical data problems because of our local-in-time
method for determining the turning point.
339
Can such a turning point be predicted before it happened, and if so by how many
days in advance? Since the logarithm of NR lies on a straight line passing through
the turning point of EIC, it would be interesting to explore if the turning point can be
predicted by extrapolation using data weeks before it happened by extrapolation
along the straight line (see Figure S5). How far in advance this can be done appears
to be limited by the poor quality of the initial data. Fig. 4 shows the results of such
predictions. The horizontal axis indicates the last date of the data used in the
prediction. The beginning date of the data used is 24 January for all experiments.
Prior to that day, data quality was poor and the newly recovered number was zero
in some days, giving an infinite NR ratio.
350
351 For China outside Hubei, the prediction made on 6 February gives the turning point
352 as 14 February, two days later than the truth. A prediction made on 8 February
353 already converged to the truth of 12 February, and stays near the truth, differing by
354 no more than fractions of a day with more data.
355
The huge data glitch on 12 February in Hubei affected the prediction for Hubei, for
China as whole, and for Hubei-exWuhan. These three curves all show a bump up
starting 12 February, as the slope of $N(t)$ is artificially lifted. Ironically, predictions
made earlier than 12 February are actually better. For example, for China as a whole,

predictions made on 9 February and 10 February both give 19 February as the
turning point, only one day off the truth of 18 February. A prediction made on 11
February actually gives the correct turning point that would occur one week later.
At the time these predictions are made, the newly infected cases were rising rapidly,
by over 2,000 each day, and later by over 14,000. It would have been incredulous if
one were to announce at that time that the epidemic would turn the corner a week
later.
367
Even with the huge spike for the regions affected by the Hubei's changing of
diagnosis criteria, because of its short duration the artifact affects the predicted
value by no more than 3 days, and the prediction accuracy soon recovers for China
as a whole. For Hubei, the prediction never converges to the true value, but the
over-prediction is only 2 days. This smallness of the error is remarkable given that
other model predictions differ by weeks or months.
374
375   Table S2 lists the mean and standard deviation of the predictions. For applications
376   to other countries and to future epidemics without a change in the definition of the
377   "infection" to such a large extent, we expect even better prediction accuracy.
378
379

380
381
382   **Figure 4** Prediction of the turning point in *EIC* by extrapolating the trend in
383   logarithm of *NR*. The horizontal axis indicates the date the prediction is made using
384   data prior to that date. The vertical axis gives the dates of the predicted turning
385   point. Dashed horizontal lines indicated the true dates for the turning point, as
386   determined from Fig. S1.
387

Estimate of "all clear" declaration

We can now estimate a time for a declaration of "all clear". No verification is yet
possible as the predicted date has not occurred. At the turning point, the *EIC* is still
at its peak. For the disease to have run its course, and an "all clear" declaration can
be announced, we require that the newly infected case number to drop to zero. For
prediction practice this "zero" is measured by three standard deviations from the
peak of *N(t)*. Then we wait for two incubation periods, each 14 days, to pass, before
we declare "all clear". Using the inferred disease characteristics in Table S1, our
prediction is, for China outside Hubei: the last week of March. For China as a whole:
the first week of April, barring "imports" of infected from abroad. At this point there
may still be some patients in the hospital who are infected with the virus. The "all
clear" call assumes that these patients are not roaming freely to cause new
infections.

Prediction for South Korea

Figure S6 summarized the available data for Korea at the present. The recovered
case numbers hovered around 1 and 2 daily up to March 1st. It only picked up
toward the end. Starting from 19 February, there seems to be enough new daily
infected cases. The South Korea Government has identified that the epic center of
the epidemic was at church gatherings in the city of Daegu and North Gyeongsang
province, where 90% of the cases are found. Specifically, a confirmed COVID-19
patient was reported to have attend the Shincheonji Church of Jesus services twice
on February 9th and 16th. Given the incubation period of 7 to 14 days, the initial
explosion at February 19th and the first peak value around February 24th are not
accidents.

If we use the available daily new cases data, we can get the statistical characteristics
of the distribution of the daily new cases from Figure S7, which gives the t_N as March
3rd and a σ_N value of 4.5 days. If we further use the turning point as approximately
$t_N+T/2$, then the turning point should fall on March 10, assuming *T* as 14 days based
on the over all mean from different regions in China.

For the *NR* ratio, it is limited by the availability of recovered case number. If we use
the limited recovered cases starting from March 1st, we have 7 days of data. The
computed the *NR* ratio together with the trend is given in Figure S8. The turning
point, at the zero-crossing of the extended trend line, would occur between March
11th and 12th. This approach does not need to use a value for *T.*

An estimate of the end of the epidemic can be given as the second week of April,
using the estimated value for t_N= *3* March, *σ=4.5 days*. Remarkably, this date is
around the same time as for Wuhan, China. South Korea owes its quick turning point
and end of the epidemic date to its ability to identity the first infection and the
secondary infections at Shincheonji Church (*31*), where most of the infected were
concentrated. This is reflected in the data: *σ* for South Korea is only half that of
China, with a more rapid rise and fall of the newly infected. Its data for the newly

infected are probably more accurate compared to other countries in similar stage of
the epidemic, due to its massive and speedy (within 6 hours) testing of the
population in its "trace, test and treat" policy.
436
437   **4. Effects of quarantine judging from data on the net infection rate.**
438   Following (15), we define a time-dependent *net infection rate* as:

439
$$\alpha(t) = \frac{dI/dt}{I} = \frac{d}{dt}\log I(t).$$

In traditional models, such as the SIR model, there is also a time-dependent
infection rate, which at *t=0* is related to the *Basic Reproductive Number R_0*. If this
number is greater than 1 then an epidemic will ensue, i.e. the infected population
will increase exponentially after the introduction to a susceptible population *S* at *t=0*
some initial infected. That is, from the SIR model equation:

$dI/dt = aSI - bI = bI(aS/b - 1),$

where $aS(t)$ is the infection rate and $b = 1/T$ is the mean recovery rate.

Therefore $\alpha(t) = \dfrac{dI/dt}{I} = b(\dfrac{aS}{b} - 1).$ $\alpha(0) = b(R_0 - 1),$ with $R_0 = \dfrac{aS(0)}{b}.$

Our time-dependent net infection rate generalizes this concept to be independent of
the SIR or other models and be applicable at later times as well: If in the course of an
epidemic, $\alpha(t)$ is positive, the number of infected will grow exponentially, reaching

a peak number of infected when $\alpha(t) = 0$ at $t = t_p$. Then the total number of active

infected will decrease exponentially. One could in analogy to R_0, define a time-

dependent Reproductive Number $R_t = \alpha(t)T - 1$, so that if this number is greater

(less) than 1 the number of infected will grow (decrease) at time t. We will here
use $\alpha(t)$ directly.

456
Since $\alpha(t)$ has a zero and its first derivative near the zero is nonzero (viz negative)

because t_p is the maximum of *I(t)*, it is a linear function of t with negative slope in

that neighborhood. This expectation is verified empirically, using data for the total
existing case numbers. We find that this time dependent infection rate is
approximately a linear function of time in the neighborhood of its zero over a period
of a few weeks.

463
464   This gives another way to predict the turning point t_p which can be used instead of

465   (but is less accurate than) the NR ratio, during the early stage of the epidemic when
466   not enough *R(t)* data is available, as is the case currently for US.

467
The peak *EIC* number can be predicted as

$I(t_p) = I(t_B)\exp\{\frac{1}{2}\alpha(t_B)(t_B - t_p)\},$ where t_B is the last available data before the

turning point. It is assumed that $\alpha(t)$ lies on a straight line between t_B and t_p.

Predicting the peak active infected cases
Since the turning point can be predicted two weeks in advance, the above formula
can be used to predict the peak EIC numbers.
474
Predicting the total infected cases (TIC):

$$TIC(t) = \int_0^t N(t)dt.$$

To predict the total infected cases for the epidemic for a region, we need to do the
above accumulation of $N(t)$ into the future, to the end date of the epidemic. That
total is approximately:

$$TIC_\infty = 2 \cdot TIC(t_N) ,$$

assuming that $N(t)$ is approximately symmetric about its peak at t_N. In reality $N(t)$
may not be symmetric and likely has a long tail. However, since the number of cases
along the tail is small, the above approximation for the total is still good. If the
present time t_B is before t_N, we need a way to predict $TIC(t_N)$. Let the *total infection*
rate be defined as:

$$\beta(t) = \frac{d}{dt}\log TIC(t) = \frac{\frac{d}{dt}TIC(t)}{TIC(t)} = \frac{N(t)}{TIC(t)}.$$

By extrapolate the total infection rate forward in time we can predict:

$$TIC(t_N) = TIC(t_B)\exp\{\int_{t_B}^{t_N} \beta(t)dt\}.$$

USA: t_N is predicted to be 2.2 days from today, April 5. Today's TIC is 308,850. So the
predicted total infected cases for the epidemic when it is over is predicted to be
708,750. This is a very large number, nine times larger than that of China, which has
a much larger population, but is nevertheless much lower than some other
predictions of a few million infected. Its current EIC is 285,000, and is predicted to
peak 12 days from now at 547,000. This is the peak demand for hospital beds.
Germany: Germany has good data. Its t_N was 2 days ago. On that day its TIC was
85,000. So the total infected cases for the epidemic when it is over is predicted to be
2 times that: 170,000. Its current EIC is 68,248. The peak EIC is 69,627, which will
occur two days from now.
Spain: t_N occurred on March 31. On that day its TIC was 94,417. The total TIC when
the epidemic is over is predicted to be twice that, 188,800. Its peak EIC is predicted
to be 84,250, to occur 4 days from now.
UK: Its current TIC is 41,000. Its t_N is 7.5 days from now. Calculating its TIC at that
time and then doubling it yields a total TIC of 134,000 when the epidemic is over.
The peak EIC is predicted to be 83,200, 16 days from now. This is the peak demand
for hospital beds. (It should be noted that the recovered case number for UK is
unusually low, currently at 209, while the dead is much higher, at 4,900. The data
may be doubtful.)
508
Comments on the effects of quarantine

Additionally the net infection rate reveals the effect of measures taken with social
distancing and quarantine. Figure 5 shows the time-dependent net infection rate for
each region starting when the newly confirmed cases exceed 100. This way of
plotting facilitates comparison of different regions at the same stage of the epidemic.
First, China outside Hubei has the lowest time-dependent infection rate after Hubei
was lockdown. Germany and Italy have similar exponential growth rate of the net
infected case numbers, both higher than even Wuhan, the epicenter in China. More
surprisingly, US has the highest exponential net infection rate, higher than Germany
and Italy and China. This can be attributed to the fact that US so far does not have a
nation-wide shutdown, unlike these other countries. Secondly, China outside Hubei
reached its turning point early, in fact 20 days earlier than the epicenter, Wuhan.
We had previously predicted this, which is qualitatively different than many model
predictions, which had the epicenter achieving its turning point 1-2 weeks earlier
than China outside Hubei (13). Italy and Germany are predicted to take a week
longer to reach their turning point, while US will take another week longer than it
will take Germany and Italy.
526
That Italy would take the same amount of time to reach its turning point as Germany
and has approximately the same net infection rate may be due to two reasons: Italy
does not test as widely as Germany and so the case numbers represent a smaller
portion of the infected. Secondly Germany has the lowest fatality rate while Italy one
of the highest. The number of the dead is about the same as the number of cured for
Italy. The dead is included in recovered/removed. If it were not included, *EIC*
would have been higher for Italy. Nevertheless, whether dead or cured, the hospital
bed is vacated.
535
536

537
538
539

540   **Figure 5**. The time-dependent net infection rate (in units of 1/day) as a function of
541   time starting on the date (listed in the inset) when the newly confirmed case
542   number exceeds 100 for each region. To obtain the actual calendar date, add the
543   dates on the horizontal axis to the starting date indicated in the inset. The number of
544   confirmed cases on the starting date is listed at the top.
545
5. Conclusion.
We offer an additional data-driven approach to track and predict the course of the
epidemic. Many parameters characterizing an epidemic can be determined from
local-in-time data. Validated by real data, we suggest that our approach could be
applied not just to the current Covid-19 epidemic, but also generally to future
epidemics. It could also be used as a practical tool for epidemic management
decisions such as quarantine institution and medical resource planning and
allocations (32-35).
554
Two results are of special significance for future policy makers. First the turning
point for the epidemic in China exHubei occurred a week earlier than that for
Wuhan. Second the US will take 2 weeks longer to reach its turning point than even
Wuhan. After its lockdown, Wuhan, with a large susceptible population of 11 million,
enforced straight social distancing, which is more strict than that adopted in the US..
As a consequence, even with a large pool of potential susceptible, the outbreak could
end sooner, as compared to the time it will take US and Europe. For China, the
lockdown of Wuhan and Hubei was the reason why the epidemic outside Hubei was
under control, and the turning point occurred earlier. In Wuhan, with hospitals
facing the number of infected patients far exceeding available hospital beds in the
initial period, some infected patients were not adequately isolated. The infected
were sent home and caused secondary infections among family members. This
might have played a role in delaying the turning point. On the other hand, outside
Hubei, hospitals were not as overwhelmed because of the strict quarantine placed
on Hubei, which drastically reduced the import of the disease originating from
Hubei. The infected were better isolated, reducing further spread, and treated in
hospitals, resulting in shorter time to recovery (see Table S1). This is evidence of the
effectiveness of the city and province-wide lockdown in "flattening the curve"
outside.
574
The additional and surprising finding that the net infection rates in Italy, Germany
and the US are higher than even Wuhan also manifests the effect of the enforcement
of lockdown, stay-at-home and strict social distancing policy in Wuhan, which was
much stricter than those adopted currently in Europe and US. The more lax latitude
and lack of enforcement in the US will lead to a longer period of the epidemic, longer
than even Italy, and the largest number of total infected cases of 710,000 before it is
over.
582
583
584
585

References

1. David Adam, Modelers Struggle to Predict the Future of the COVID-19 Pandemic. The Scientist, https://www.the-scientist.com/news-opinion/modelers-struggle-to-predict-the-future-of-the-covid-19-pandemic-67261 (2020).
2. WHO, Laboratory testing of human suspected cases of novel coronavirus (nCoV) infection: interim guidance, World Health Organization, Geneva (2020).
3. N. Zhu, D. Zhang, W. Wang, X. Li, B. Yang, J. Song, X. Zhao, B. Huang, W. Shi, R. Lu, P. Niu, F. Zhan, A novel coronavirus from patients with pneumonia in China, 2019. *N. Engl. J. Med.* **382**, 727-733 (2020).
4. R. Lu, X. Zhao, J. Li, P. Niu, B. Yang, H. Wu, W. Wang, H. Song, B. Huang, N. Zhu, Y. Bi, X. Ma, F. Zhan, L. Wang, T. Hu, H. Zhou, Z. Hu, W. Zhou, L. Zhao, …, W. Tan, Genomic characterisation and epidemiology of 2019 novel coronavirus: implications for virus origins and receptor binding. *The Lancet* **395**(10224), 565-574 (2020).
5. Y. Liu, A. A. Gayle, A. Wilder-Smith, J. Rocklöv, The reproductive number of COVID-19 is higher compared to SARS coronavirus. *J. Travel Med.* taaa021 (2020).
6. J. W. Glasser, N. Hupert, M. M. McCauley, R. Hatchett, Modeling and public health emergency responses: Lessons from SARS. *Epidemics* 3: 32-37 (2011), doi:10.1016/j.epidem.2011.01.001.
7. P. Zhou, X. Yang, X. Wang, B. Hu, L. Zhang, W. Zhang, H. Si, Y. Zhu, B. Li, C. Huang, H. Chen, J. Chen, …, Z. Shi, A pneumonia outbreak associated with a new coronavirus of probable bat origin. *Nature* **579**, 270-273 (2020).
8. C. Huang, Y. Wang, X. Li, L. Ren, J. Zhao, Y. Hu, L. Zhang, G. Fan, J. Xu, X. Gu, Z. Cheng, T. Yu, J. Xia, Y. Wei, W. Wu, X. Xie, W. Yin, H. Li, M. Liu, Y. Xiao, H. Gao, L. Guo, J. Xie, G. Wang, R. Jiang, Z. Gao, Q. Jin, J. Wang, B. Cao, Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *The Lancet* **395**(10223), 497-506 (2020).
9. J. F-K. Chan, S. Yuan, K.-H. Kok, K. K.-W. To, H. Chu, J. Yang, F. Xing, J. L. BNurs, C. C.-Y. Yip, R. W.-S. Poon, H.-W. Tsoi, S. S.-F. Lo, K.-H. Chan, V. K.-M. Poon, W.-M. Chan, J. D. Lp, J.-P. Cai, V. C.-C. Cheng, H. Chen, C. K.-M. Hui, K-Y. Yuen, A familial cluster of pneumonia associated with the 2019 novel coronavirus indicating person-to-person transmission: a study of a family cluster. *The Lancet* **395**(10223), 514-523 (2020).
10. X. Xu, P. Chen, J. Wang, J. Feng, H. Zhou, X. Li, W. Zhong, P. Hao, Evolution of the novel coronavirus from the ongoing Wuhan outbreak and modeling of its spike protein for risk of human transmission. *Sci. China Life Sci.* **63**, 457-460 (2020).
11. Z. Chen, W. Zhang, Y. Lu. C. Guo, Z. Guo, C. Liao, X. Zhang, Y. Zhang, X. Han, Q. Li, W. lan Lipkin, J. Lu, From SARS-CoV to Wuhan 2019-nCoV Outbreak: Similarity of Early Epidemic and Prediction of Future Trends. *Biorxiv* preprint (2020), doi: https://doi.org/10.1101/2020.01.24.919241.
12. J. M. Read, J. R. E. Bridgen, D. A. T. Cummings, A. Ho, C. P. Jewell, Novel coronavirus 2019-nCoV: early estimation of epidemiological parameters and epidemic predictions. *medRxiv* preprint (2020), doi: https://doi.org/10.1101/2020.01.23.20018549.

13. J. T. Wu, K. Leung, G. M. Leung, Nowcasting and forecasting the potential domestic and international spread of the 2019-nCoV outbreak originating in Wuhan, China: a modelling study. *The Lancet* **395**(10225), 689-697 (2020).

14. S. Zhao, S. S. Musa, Q. Lin, J. Ran, G. Yang, W. Wang, Y. Lou, L. Yang, D. Gao, D. He, M. S. Wang, Estimating the Unreported Number of Novel Coronavirus (2019-nCoV) Cases in China in the First Half of January 2020: A Data-Driven Modelling Analysis of the Early Outbreak. *J. Clin. Med.* **9**, 388 (2020).

15. N. E. Huang, F. Qiao, A data driven time-dependent transmission rate for tracking an epidemic: a case study of 2019-nCoV. *Sci. Bull.* **65**, 425-427(2020) , https://doi.org/10.1016/j.scib.2020.02.005.

16. Q. Li, W. Feng, Trend and forecasting of the COVID-19 outbreak in China. *J. Infection* arXiv:2002.05866v1, (2020).

17. H. Xiong, H. Yan, Simulating the infected population and spread trend of 2019-nCov under different policy by EIR model. *medRxiv* preprint (2020), doi: https://doi.org/10.1101/2020.02.10.20021519.

18. L. Damon, E. Brooks-Pollock, M. Bailey, M. J. Keeling, A spatial model of CoVID-19 transmission in England and Wales: early spread and peak timing. *medRxiv* preprint (2020), doi: https://doi.org/10.1101/2020.02.12.20022566.

19. H. Sun, Y. Qiu, H. Yan, Y. Huang, Y. Zhu, S. Chen, Tracking and Predicting COVID-19 Epidemic in China Mainland. *Medrxiv* preprint (2020), doi: https://doi.org/10.1101/2020.02.17.20024257.

20. Q. Liu, Z. Liu, D. Li, Z. Gao, J. Zhu, J. Yang, Q. Wang, Assessing the Tendency of 2019-nCoV (COVID-19) Outbreak in China. *medRxiv* preprint (2020), doi: https://doi.org/10.1101/2020.02.09.20021444.

21. L. Peng, W. Yang, D. Zhang, C. Zhuge, L. Hong, Epidemic analysis of COVID-19 in China by dynamical modeling. *arXiv* 2002.06563, (2020).

22. D. Cyranoski, When will the coronavirus outbreak peak? *Nature news* (2020).

23. C. R. MacIntyre, Global spread of COVID-19 and pandemic potential. *Global Biosecurity* **1**(3), (2020).

24. WHO, Coronavirus latest: WHO describes outbreak as pandemic, Nature news (2020), https://www.nature.com/articles/d41586-020-00154-w.

25. K. Kupferschmidt, J. Cohen, Can China's COVID-19 strategy work elsewhere? *Science* **367**(6482), 1061-1062 (2020).

26. J. M. Read, J. R. Bridgen, D. A. Cummings, A. Ho, C. P. Jewell, Novel coronavirus 2019-nCoV: early estimation of epidemiological parameters and epidemic predictions. *medRxiv* (2020), doi:10.1101/2020.01.23.20018549.

27. S. Zhao, Q. Lin, J. Ran, S. S. Musa, G. Yang, W. Wang, Y. Lou, D. Gao, L. Yang, D. He, M. H. Wang, Preliminary estimation of the basic reproduction number of novel coronavirus (2019-nCoV) in China, from 2019 to 2020: A data-driven analysis in the early phase of the outbreak. *Int. J. Infect. Dis.*, 92: 214-217 (2020), https://doi.org/10.1016/j.ijid.2020.01.050.

28. W. Kermack, A. McKendrick, A contribution to the mathematical theory of epidemics. *Proc. Roy. Soc. London A* **115**, 700-721 (1927).

29. D. L. Heymann, N. Shindo, COVID-19: what is next for public health? *Lancet*, 395(10224): 542-545 (2020), https://doi.org/10.1016/S0140-6736(20)30374-3.

30. The Novel Coronavirus Pneumonia Emergency Response Epidemiology Team,
The Epidemiological characteristics of an outbreak of 2019 novel coronavirus
disease (COVID-19)-China, 2020, *China CDC Weekly* (2020).
31. E. Shim, A. Tariq, W. Choi, Y. Lee, G. Chowell, Transmission potential of COVID-19
in   South   Korea.   *medRxiv*   preprint,   (2020),   doi:
https://doi.org/10.1101/2020.02.27.20028829.
32. C. M. Peak, L. M. Childs, Y. H. Grad, C. O. Buckee, Comparing nonpharmaceutical
interventions for containing emerging epidemics. *Proc. Natl. Acad. Sci.* 114(15):
4023-4028 (2017), doi:10.1073/pnas.1616438114.
33. R. S. Dhillon, D. Srikrishna, When is contact tracing not enough to stop an
outbreak?   *Lancet   Infect.   Dis.*, 18: 1302-1304   (2018),
https://doi.org/10.1016/S1473-3099(18)30656-X.
34. X. Pang, Z. Zhu, F. Xu, J. Guo, X. Gong, D. Liu, Z. Liu, D. P. Chin, D. R. Feikin,
Evaluation of control measures implemented in the severe acute respiratory
syndrome outbreak in Beijing, 2003. JAMA, 290(24): 3215-3221 (2003).
35. G. Wang, N. E. Huang, F. Qiao, Quantitative evaluation on control measures for an
epidemic: A case study of COVID-19. *Sci. Bull.* **65** (2020), doi: 10.1360/TB-2020-
0159.
36. J.D. Murray: Mathematical Biology I. Third Edition. Springer, 551pp.

Acknowledgements: NEH and FQ are supported by the National Natural Science
Foundation of China under Grant 41821004. KKT's research is supported by the
Frederic and Julia Wan Endowed Professorship.
Competing Interests: The authors declare no competing interests.
Data Availability: All data in this study are publicly available from World Health
Organization (WHO) at https://www.who.int/emergencies/diseases/novel-
coronavirus-2019/situation-reports/
and on the Daily Brief site of the China's National Health Commission at
http://en.nhc.gov.cn/
The Korean data is available at
https://sa.sogou.com/new-weball/page/sgs/epidemic
Coronavirus COVID-19 Global Cases by Johns Hopkins CSSE
https://gisanddata.maps.arcgis.com/apps/opsdashboard/index.html#/bda759474
0fd40299423467b48e9ecf6
716
717

Supplementary Information:

Comments on SIR-type of models

Mainstream epidemiological models have their origin in the SIR (Susceptible, Infected, and Recovered or Removed) model (28) and its many variations. We discuss here why existing model predictions vary so widely by commenting on the assumptions underlying these models, using the classical SIR model as an example.

The number of the infected is increased by the rate of infection, which is assumed to be aIS, and decreased by the rate of recovery/removed, who are assumed to be immune to further infection, bI. ($dI/dt = aIS - bI$). This latter category, R, increases at the rate of bI. ($dR/dt = bI$). Both of the parameters, especially the infection rate a, is largely unknown for an emergent disease (1), and have to be either estimated based on limited statistics the number of contacts a single infected person may have and how many of the contacted people will be infected, or obtained by curve fitting of reported $I(t)$. This approach has problems. First, for Covid-19, there is a population of asymptomatic infected, which may be larger than the reported/confirmed I during the early stages of the epidemic, when wide-scale testing is not available. Since this asymptomatic infected is also infectious, and some of those infected by them could later become confirmed infected, fitting the rate of increase of reported I to estimate the infection rate would inevitably yield a much larger a. Consequently the model predictions using this estimated parameter value may yield a very large peak infection number. Secondly, since $dI/dt = aI(S - b/a)$, the number of infected will increase (approximately exponentially) for $S > b/a$, but for a new virus to which the population has no immunity, the susceptible population could be very large. For China it could be as large as 1.4 billion. For Hubei it is 65 million. The susceptible population should be lower if the quarantine of Wuhan were tight, but it is difficult to estimate what it is under realistic conditions. Thirdly, predictions of the end of the epidemic vary widely because of the basic assumption in some SIR type of models that if most of the population is infected and recovered (and hence acquired immunity) or dead (and hence no longer infectious). The concept of "herd immunity" is rooted in the idea that with enough of the population acquiring immunity this way (or from a vaccine if one is developed in time), the susceptible population is reduced to $S < b/a$, and the rate of increase of I will turn negative. This will require as much as 70%-90% of the population be infected, a huge number. Modern models take into account of the effect of quarantine and isolation in reducing the number of people each infected can contact, thus reducing the infection rate a, so that S does not have to be reduced to such an extent for the epidemic to peak and I starts to decrease. However, such estimates are very dependent on model assumptions.

Of course the modern models of epidemiology are more sophisticated than the SIR model (*12-14, 17, 21*).

762   We offer here an additional tool that has the advantage that it has does not depend
763   on the elusive infection rate or the susceptible population, information needed for
764   most models, but has the disadvantage that it cannot be used when the epidemic
765   first started and the data are inaccurate or incomplete. It is based on daily case
766   numbers (i.e. newly confirmed cases), *N(t)*, and recovered cases, *R(t)*.
767
Our estimate of the end date of the epidemic is not based on the number of
susceptibles, *S,* approaching zero as in most models (i.e. most of the population is
infected, hence acquiring immunity), but *N(t)* approaching zero and remaining so
for two incubation periods. The first incubation period is to allow the asymptomatic
infected to show symptoms and the second period to allow those that are infected
by the asymptomatic infected to show symptoms. For prediction purpose, the date
when the *N(t)* is zero is estimated by 3 standard deviations from its peak. These
two quantities can be extracted from the data as the epidemic is developing. Our
estimate of the end of the epidemic is earlier than most model predictions, usually
significantly so, because it does not depend on the herd immunity concept.
778
779
THEORY:
Definition: Let $I(t)$ be the number of active infected at time t. Its change is given
by;
$$\frac{d}{dt}I = N(t) - R(t),$$
where $N(t)$ is the number of newly infected, and $R(t)$ that of the newly recovered or
removed (dead). Note that for the theory part, *N(t)* includes both confirmed and
unconfirmed cases. The term: Existing Infected Case (*EIC*) number is used to denote
the confirmed *I(t)* when we deal with data*.*
788
Let t_p, the turning point defined as the peak of the active infected number. At this
point maximum medical resource is needed. This maximum occurs when
$$\frac{d}{dt}I = 0, \text{ implying } N(t_p) = R(t_p).$$
There is no need to first find *I(t)* to locate this peak. After the turning point, the
newly recovered starts to exceed the newly infected. The demand for medical
resources, such as hospital beds, isolation wards and respirators, starts to decrease.
795
796   Let $t = 0$ be when the first infection began. For Wuhan, China, this date is near the
797   end of 2019, perhaps even earlier. Let t_B be the beginning of the better quality data.
798   This time is beyond the initial incubation period of the disease and it can be
799   assumed that at that time there is already a population of infected, some of them
800   asymptomatic but nevertheless infectious.
801

Let $X(t,s)$ be the number of infected cases at time t, with s being the "age"
distribution, i.e. number of days sick.
The total number of infected is given by:

$I(t) = \int_0^T X(t,s)ds$.

After being sick for T days, a patient either recovers or is removed (dead). T is called
the recovery period (or removal period). It is also called the infectious time if the
patient is infectious during this period. Of course its value varies by patient and by
the efficacy of treatment for each hospital. For the removed it also depends on the
age of the patient and whether there are underlying medical conditions. Only a
mean recovery period is obtainable from data, and so this is in reality a statistical
quantity. We will discuss later how this statistical quantity can be obtained from
data.
814
815   **Conservation law** (see Murray: Mathematical Biology Part I, Chapter 1 (36)):
816   After first infected and until removed or cured, we have:

    $dX(t,s) = \dfrac{\partial}{\partial t}X \bullet dt + \dfrac{\partial}{\partial s}X \bullet ds = 0, \; 0 < s < T.$

817   So, since $ds/dt = 1,$

    $\dfrac{\partial}{\partial t}X + \dfrac{\partial}{\partial s}X = 0.$

    This equation is to be solved using the method of characteristics as

818     $X(t,s) = $ constant along characteristics defined by $ds/dt = 1.$

    Boundary condition: $X(t,0)$, specifies the "birth" process,

819   i.e. how the disease spawns newly infected (with "age" $s = 0$).

    Initial condition: $X(0,s) = 0$ for $s > 0$, specifies the initial age distribution at $t = 0$

820   There are two types of characteristics:
821   (i) $s > t$, (ii) $s < t$.
822   The first type of characteristics intersects the $t = 0$ axis, and since the initial
823   condition is zero, we have the solution:
824

825     $X(t,s) \equiv 0$ for $s > t.$

826
827   That is, there is no infected population who is sick for more days than the lapsed
828   time since the first infection occurred.

829
830   For the second type of characteristics, $t > s$ the solution is
831

832     $X(t,s) = f(s-t)$
833   with the form of f to be determined by the boundary condition. Even without
834   determining the form of f we have the following general results:
835   For $t > T$, and therefore $t > s$:

836
$$I(t) = \int_0^T X(t,s)\,ds = \int_0^T f(s-t)\,ds$$
$$= \int_{t-T}^t f(p)\,dp.$$

837
$$\frac{d}{dt}I = f(t) - f(t-T).$$

839 Since the rate of increase of confirmed $I(t)$ is by definition equal to the newly
840 confirmed infected number, $N(t)$, minus the newly recovered (or removed) number,
841 $R(t)$, we have:

842
$$N(t) - R(t) = \frac{d}{dt}I = f(t) - f(t-T).$$

For a fatal disease with low fatality rate, where almost all infected cases eventually
recover after a hospital stay of T days, we can identify
$f(t)$ with $N(t)$, and $f(t-T)$ with $R(t)$.
If the disease has a non-negligible fatality rate, we include the dead in $R(t)$.
Main Result: The daily newly recovered/removed number $R(t)$, is related to the
daily newly infected number $N(t)$ as, for $t > T$:

851
$$R(t) = N(t-T).$$

Validation: This fundamental relationship can be validated statistically with data.
Figures 1, obtained using data from China during the Covid-19 epidemic, shows that
$N(t)$ and R(t) are highly correlated: with correlation coefficient of 0.95 when both
distributions are smoothed with 5-point boxcar. The unsmoothed daily data also
yield a high correlation coefficient of 0.80, with $R(t)$ lagging $N(t)$ by $T \sim 15$ days. Both
correlation coefficients are statistically significant. A similar result is found for
Hubei (Figure S2) and other regions (not shown). This is one of the ways the *mean*
recovery period is determined statistically from data, but it is not practical in the
early phase of the epidemic. We will give different methods for the latter purpose.
The result on T is consistent with that estimated or predicted later using the slope of
the distribution in Figure 4. The latter, obtained by the intercept of the straight line,
is less accurate because of the slope is rather shallow. For the regions considered in
Figure 1, the fatality rate is small and so the dead are not included in $R(t)$ for
convenience.
866

The second type of characteristics intersects the boundary $s = 0$. The boundary
condition itself needs to be solved as a function of t to describe how new infection
(at $s = 0$) occurs. This can be done using a birth model, such as Eq. (1.56) in (36).
For our purpose we assume that the solution of this model yields a distribution with
age that has a full spectrum $0 < s < T$ of infectives at a time t_B, long after a full
incubation period has passed.

$$873 \quad X(t_B,s)=f_0(s)=A\exp\{-\frac{(s-s_0)^2}{2b^2}\};\ A\ \text{independent of}\ s.$$

$$b=\tfrac{1}{2}T.$$

875 Therefore the solution is, for $t>t_B>0$:

$$876 \quad \begin{aligned} X(t,s)&=X(t_B,s-t)=f_0(s-t)\\ &=A\exp\{-\frac{(s-s_0-t)^2}{2b^2}\}. \end{aligned}$$

$$878 \quad N(t)=A\exp\{-\frac{(t-s_0)^2}{2b^2}\}=A\exp\{-\frac{(t-t_N)^2}{2b^2}\}$$

$$R(t)=N(t-T)=A\exp\{-\frac{(t-t_R)^2}{2b^2}\},$$

879 where t_N is the peak of $N(t)$, and $t_R=t_N+T$ is the peak of $R(t)$.
880 Both distributions are Gaussians.
881

882 For $t_B<t<T$,

$$883 \quad \begin{aligned} I(t)&=\int_0^t X(t,s)ds+\int_t^T X(t,s)ds\\ &=\int_0^t f(s-t)ds+\int_t^T 0\,ds\\ &=\int_{-t}^0 f(p)dp \end{aligned}$$

$$884 \quad \frac{d}{dt}I=f(-t)=A\exp[-\frac{(t-s_0)^2}{T^2}].$$

$$885 \quad N(t)=A\exp[-\frac{(t-t_N)^2}{T^2}]$$

$$R(t)=0.$$

886 Again, $N(t)$ is Gaussian, but there is no recovered or removed during this early stage.
887

888 **Main Result**: The natural logarithm of the ratio of N and R is a linear function of
889 time for $t>T$:
890

$$891 \quad \begin{aligned} NR(t)&=\frac{N(t)}{R(t)}=\frac{N(t)}{N(t-T)}\\ &=\exp\{-\frac{(t-t_N)^2}{2b^2}+\frac{(t-t_N-T)^2}{2b^2}\}=\exp\{\frac{T^2}{2b^2}-\frac{T(t-t_N)}{b^2}\}. \end{aligned}$$

$$\log NR=-\frac{T(t-t_N)-\tfrac{1}{2}T^2}{b^2}$$

892   a linear function of t. This relationship is important for the purpose of forecast
893   because it is easy to extrapolate from a straight line into the future.
894

895   It intersects 0 at $t - t_N = \frac{1}{2}T$. This yields the turning point, when the NR ratio is 1, and
896   therefore its logarithm is zero.
897

Result: The turning point, defined as the maximum of *I(t)*, is given by $t_p = t_N + \frac{1}{2}T$.

Result: The slope of log NR is equal to $4/T$.

When time is normalized by T, the derivative is given by:

$\dfrac{d\log NR}{d(t/T)} = \dfrac{T^2}{b^2} = 4$, a dimensionless constant.

Heterogeneous Data
The above results are obtained for the case of a single introduction into a region of
infected at *t=0* and we solve for the subsequent development of the epidemic from
that single source. Consider now a large region consisting of a number of small
regions, and the "seeding" of the infected occurs at different times for different
regions. The large region could be China, and the first infection could be Wuhan,
Hubei and then the regions outside Hubei. Then we may have for the China as a
whole data for the newly infected a sum of several Gaussians staggered in time. As
long as the Gaussians are not separated so much that there are different peaks in the
combined data, the combined data can still be considered as Gaussian, as is the case
in the real data. However, the standard deviation σ of the combined Gaussian is
inevitably larger and is no longer given by *b:*

$N(t) = \dfrac{B}{\sqrt{2\pi}\sigma_N} \exp\{-\dfrac{(t - t_N)^2}{\sigma_N^2}\}$.

We still have $R(t) = N(t - T)$ since this result holds for each sub-region. The result
that $\log NR$ is a linear function of time still holds:

$\log NR(t) = \log \dfrac{N(t)}{N(t - T)} = -\dfrac{T(t - t_N) - \frac{1}{2}T^2}{\sigma_N^2}$.

The slope of the straight line is T/σ_N^2 .
921

Since the hospital state can add as a smoothing filter on *N(t)* to yield *R(t)*, the
standard deviation for *R(t)* could be slightly wider than that for *N(t)*. So we could
have two different Gaussians (but their integral over all time should be the same):

$N(t) = \dfrac{B}{\sqrt{2\pi}\sigma_N} \exp\{-\dfrac{(t - t_N)^2}{2\sigma_N^2}\}; \;\; R(t) = \dfrac{B}{\sqrt{2\pi}\sigma_R} \exp\{-\dfrac{(t - t_R)^2}{2\sigma_R^2}\}.$

Taking this into account, we have, denoting $T = t_R - t_N$:

$$\log NR - \log\frac{\sigma_R}{\sigma_N} = -\{\frac{(t-t_N)^2}{2\sigma_N^2} - \frac{(t-t_N-T)^2}{2\sigma_R^2}\} = -\frac{(\sigma_R^2-\sigma_N^2)(t-t_N)^2 + 2\sigma_N^2 T(t-t_N) - \sigma_N^2 T^2}{2\sigma_N^2\sigma_R^2}$$

As the values of σ_N and σ_R are very close based on the empirical data, the quadratic
term is always small comparing to the other terms for the length of time we are
considering here. Hence.

$$\log NR(t) = \frac{1}{\sigma_R^2}\{-T(t-t_N) + \frac{1}{2}T^2\} ,$$

a linear function of time. Its slope is $-\dfrac{T}{\sigma_R^2}$.

Result: The natural logarithm of the ratio of two Gaussians of slightly different
standard deviation is approximately a straight line.

936
Validation of log *NR* as a straight line:
From data we use the report newly confirmed case number and the recovered case
number to define *NR* ratio as
$$NR(t)=N(t)/R(t).$$
At t_p, *NR=1*.

We show in Figure 2, using the data of the epidemic for COVID-19, that the
logarithm of *NR(t)* lies on a straight line, with small scatter, passing through the
turning point t_p. And data for various stages of the epidemic, from the initial
exponential growth stage, to near the peak of *EIC*, and then past the peak, all lie on
the same straight line. The intercept with *logNR=0* yields the turning point. This
line, obtained by linear-least-square fit in the semi-log plot, is little affected by the
rather large artificial spike in the data on 12 February because of its short duration
and the logarithmic value. That reporting problem is necessarily of short duration
because, on the date of definition change, previous week's cases of infected
according to the new criteria were reported in one day. After that, the book is
cleared, and *N(t)* returned to its normal range.

The theoretical result suggests that the slope of the linear line is $-T/\sigma_R^2$, where σ_R is
the standard deviation of the *R(t)* profile. In general, the slope can be different for
different regions with different levels of quarantine and epidemic characteristics.
The hospital treatment efficacy would influence *T* directly. The effect of quarantine
would influence the value of σ_N, the standard deviation of the newly infected, and so
indirectly *R(t)* and σ_R. Our empirical result from Fig. 2 however shows that the slope
is the almost the same for different regions in China, implying that efficacy of
treatment and level of quarantine affect *T* and σ^2 proportionally.

Validation of the slope of log *N(t)* and log *R(t)*
Interestingly, the derivative of log *N(t)* or log *R(t)* also lies on a straight line, as
shown in Fig. 3 (although the scatter is larger as to be expected for any

differentiation of empirical data). The positive and negative outliers one day before
and after 12 Feb are caused by the spike up and then down, with little effect on the
fitted linear trend (but increases its variance and therefore uncertainty). Moreover,
the straight line extends without appreciable change in slope beyond the peak of
$N(t)$, suggesting that the distribution of the newly infected number is approximately
Gaussian. The mean recovery time T can be predicted as $t_R - t_N$, where t_R is the peak
of $R(t)$ and t_N is the peak of $N(t)$. These two peak times can be obtained by extending
the straight line in Fig. 3 to intersect the zero line. This predicted result can be
verified statistically after the fact by the lagged correlation of $R(t)$ and $N(t)$. If the
distribution is indeed Gaussian or even approximately so, the slope in Fig. 3 would
be proportional to the reciprocal of the square of its standard deviation, σ, as:

979
$$\frac{d \log N(t)}{dt} = \frac{-(t - t_N)}{\sigma_N^2}.$$

980
981 Similarly result holds for the daily number of recovered, $R(t)$.

982
983 After the epidemic is nearing the end as is the case in China, fitting the data to a
984 Gaussian can be done after the fact (see Figures S3 and S4). The fit is satisfactory
985 even without using any disposal parameters. The parameters used are determined
986 using slopes of log N and log R (see Table S1)

987
The inferred statistical characteristics of the Covid-19 epidemic are summarized in
Table S1 for various regions. The mean recovery time T, is about 13 days for China
as a whole. For Wuhan, the city at the epicenter whose hospitals were more
overwhelmed and the patients admitted into hospitals more seriously ill than those
in other provinces, $T \sim 16$ days, while that for Hubei is 14 days. The standard
deviation, σ, is found to be around 8 days, with slight difference between that for
$N(t)$ and for $R(t)$, with one exception for Hubei outside Wuhan. Such a fine
subdivision may not be practical for the data quality we have. The σ tends to be
smaller for China as a whole than Wuhan. One can see that T and σ^2 indeed varying
approximately in proportion.
998

999
1000
1001 **Figure S1**. The daily newly infected (in blue) and the daily newly recovered (in red),
1002 as a function of time for China as a whole (in solid lines) and Hubei (in dotted lines).
1003 The turning point is determined by when the red and blue curves cross.
1004 **Inset:** For China outside Hubei.
1005

1006
1007
1008 **Figure S2.** Lagged correlation of $R(t)$ with $N(t)$ for Hubei province.
1009

1010
1011 **Figure S3.** Gaussian fit of *N(t)* and *R(t)*, for China as a whole.
1012

1013
1014 **Figure S4.** Gaussian fit of *N(t)* and *R(t)*, for Hubei Province.
1015

	Crossing	t_0	t_1	T	Sigma N	Sigma R
China	2/18	2/11	2/24	13	7.5	7.7
Hubei	2/20	2/12	2/26	14	7.9	8.5
Wuhan	2/21	2/14	3/01	16	8.8	8.8
C exHubei	2/12	2/08	2/21	13	7.5	7.1
H exWuhan	2/16	2/13	2/27	14	5.0	8.8

1016 **Table S1:** Statistical characteristics of the COVID-19 epidemic in different regions in
1017 China inferred from data, for N(t), the daily number of newly infected and for R(t),
1018 the daily number of recovered.
1019

	China	Hubei	China-Hubei	Hubei-Wuhan
Truth (data)	18	19	12	15
NR Ratio	20.3±1.6 (Feb 20nd)	22.3±1.0 (Feb 22rd)	12.4±0.9 (Feb 12th)	16.0±1.2 (Feb 16th)

1020
1021 **Table S2**: Predicted turning point dates. Shown are the mean and standard
1022 deviation of the predictions over the prediction period, using the NR ratio method
1023

1024
Figure S5. Prediction of the turning point of *EIC* using linear least-squares trends
using various data lengths for China exHubei.   All data used start from 24 January.
Different colored straight lines show the linear trend calculated from 24 January to a
particular date. The spread is over a very small range. Then these trends are
extrapolated (extrapolations not shown) to intersect the zero line to yield a
prediction for the turning point. The blue dots are the data.
1031
1032
1033
1034

1035
1036
Figure S6: The available data from South Korea (as of March 7th). The sporadic
recovered case numbers are mostly in the single digit. If we use the sudden increase
of recovered case matching with the sudden explosive increase of new infected, the
distance is approximately 14 days, a reasonable *T* value when compared to the
mean value in China. For our data analysis, we used daily newly cases starting
February 19th, for the derivative of individual distribution study; we used data case
from March 1st, for the *NR* ratio study, in order to have enough recovered cases.

1044
1045 **Figure S7**: The derivative of the logarithmic value of daily new infected case
1046 distribution.
1047

1048
1049 **Figure S8**: The *NR* ratio from 7 days of data from March 1st to 7th. The estimated
1050 zero-crossing time would occur between March 11th and 12th, a value consistent
1051 with the statistics from the daily new case distribution on March 10th.