Detection of Air and Surface Contamination by Severe Acute Respiratory Syndrome
Coronavirus 2 (SARS-CoV-2) in Hospital Rooms of Infected Patients

Po Ying Chia, MBBS[1,2,3] [*]; Kristen Kelli Coleman, PhD[4] [*]; Yian Kim Tan, PhD[5] [*]; Sean Wei
Xiang Ong[1,2] [*], Marcus Gum, BSc[5]; Sok Kiang Lau, PhD[5]; Stephanie Sutjipto, MBBS[1,2]; Pei
Hua Lee, MBBS[1,2]; Than The Son, DVM[4]; Barnaby Edward Young, MB Bchir[1,2,3]; Donald
K. Milton, MD, DrPH[6]; Gregory C. Gray, MD[4,7,8]; Stephan Schuster, PhD[9]; Timothy
Barkham, MBBS[2,10]; Partha Prathim De, MBBS[2,3]; Shawn Vasoo, MBBS[1,2,3]; Monica Chan,
BMBS[1,2]; Brenda Sze Peng Ang, MBBS, MPH[1,2,3,10]; Boon Huan Tan, PhD[5]; Yee-Sin Leo,
MBBS, MPH[1,2,3,10]; Oon-Tek Ng, MBBS, MPH[1,2,3] [#]; Michelle Su Yen Wong, PhD[5] [#]; and
Kalisvar Marimuthu, MBBS[1,2,10] [#]

[*] These four first authors contributed equally.

[#] These three senior authors contributed equally.

Author Affiliations:
[1] National Centre for Infectious Diseases, Singapore
[2] Tan Tock Seng Hospital, Singapore
[3] Lee Kong Chian School of Medicine, Nanyang Technological University, Singapore
[4] Duke-NUS Medical School, National University of Singapore
[5] DSO National Laboratories, Singapore
[6] Maryland Institute for Applied Environmental Health, University of Maryland School of
Public Health, USA
[7] School of Medicine and Global Health Institute, Duke University, USA
[8] Global Health Research Center, Duke Kunshan University, Kunshan, China
[9] Singapore Centre for Environmental Life Sciences Engineering, Nanyang Technological
University, Singapore
[10] Yong Loo Lin School of Medicine, National University of Singapore

Corresponding Authors:
Kalisvar Marimuthu, National Centre for Infectious Diseases, 16 Jln Tan Tock Seng,
Singapore 308442; (kalisvar_marimuthu@ncid.sg) (+65) 6357 7924)
Oon-Tek Ng, National Centre for Infectious Diseases, 16 Jln Tan Tock Seng, Singapore
308442; (Oon_Ng_Tek@ncid.sg) (+65) 6357 7924)

Running Title: Air and surface contamination by SARS-CoV-2

Word Count: 270 (abstract), 2456 (manuscript)

Abstract:

Background: Understanding the particle size distribution in the air and patterns of environmental contamination of SARS-CoV-2 is essential for infection prevention policies.

Objective: To detect the surface and air contamination by SARS-CoV-2 and study the associated patient-level factors.

Design: Cross-sectional study.

Setting: Airborne infection isolation rooms (AIIRs) at the National Centre for Infectious Diseases, Singapore.

Patients: COVID-19 inpatients with a positive PCR test for SARS-CoV-2 within 72 hours before the environmental sampling.

Measurements: Extent of environmental surface contamination in AIIRs of 30 COVID-19 patients by PCR on environmental swabs. The particle size distribution of SARS-CoV-2 in the air was measured using NIOSH air samplers.

Results: 245 surface samples were collected from 30 rooms of COVID-19 patients, and air sampling was conducted in 3 rooms. 56.7% of the rooms had at least one environmental surface contaminated, with 18.5% of the toilet seats and toilet flush button being contaminated. High touch surface contamination was shown in ten (66.7%) out of 15 patients in the first week of illness, and three (20%) beyond the first week of illness (p = 0.010). Air sampling of two COVID-19 patients (both day 5 of symptoms) detected SARS-CoV-2 PCR-positive particles of sizes >4 µm and 1-4 µm. In a single subject at day 9 of symptoms, no SARS-CoV-2 PCR-positive particles were detected.

Limitations: Viral culture results were not available to assess the viability of the virus contaminating the air and surface.

56 **Conclusion:** Environmental contamination was detected in rooms with COVID-19 patients in

57 early stages of illness, but was significantly less after day 7 of disease. Under AIIR

58 conditions, SARS-CoV-2 respiratory particles can be detected at sizes 1-4 µm and >4 µm in

59 diameter in the air which warrants further studies.

Introduction

Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) causing coronavirus disease 2019 (COVID-19) has spread globally and many countries are experiencing ongoing local transmission despite varying levels of control efforts. Understanding the different transmission routes of SARS-CoV-2 is crucial in planning effective interventions to break the chain of transmission. Although extensive surface contamination with SARS-CoV-2 by a symptomatic patient has been demonstrated (1), little is known about airborne transmission of SARS-CoV-2. It is also unknown if asymptomatic individuals pose the same environmental contamination risk as symptomatic ones, although viral shedding has been demonstrated to continue even after clinical recovery of COVID-19 patients (2). There are multiple reports of asymptomatic patients testing positive for SARS-CoV-2 (3, 4), and the potential transmission of the virus by an asymptomatic person has been described (5). Therefore, viral contamination of the air and surfaces surrounding asymptomatic or recovering COVID-19 patients could have serious implications for outbreak control strategies. This knowledge gap is recognized in the Report of the WHO-China Joint Mission on Coronavirus 2019 (6).

The primary objective of our study was to identify potential patient-level risk factors for environmental contamination by SARS-CoV-2 by sampling the air and surfaces surrounding hospitalized COVID-19 patients at different stages of illness.

Methods

Study design, patient selection and data collection

We conducted this cross-sectional study in airborne infection isolation rooms (AIIRs) at the National Centre for Infectious Diseases, Singapore. These rooms had 12 air exchanges per hour, an average temperature of 23°C, relative humidity of 53 – 59%, and exhaust flow of 579.6 m^3/h.

Patients with a SARS-CoV-2 infection confirmed by a polymerase chain reaction (PCR)-

positive respiratory sample within the prior 72 hours were included. Clinical characteristics,

including the presence of symptoms, day of illness, day of stay in the room, supplemental

oxygen requirement, and baseline characteristics, were collected. One patient from a

previously published pilot study on environmental sampling in the same facility (Patient 30;

Supplemental Table 1) was also included in the current analysis (1).

Cleaning regimen of rooms

Routine environmental cleaning of the rooms was carried out by a trained team of

housekeeping staff. High-touch surfaces (e.g. bed rail, cardiac table, switches) were cleaned

twice daily using 5000 parts per million (ppm) sodium dichloroisocyanurate (NaDCC),

reconstituted using Biospot® Effervescent Chlorine Tablets. The floor was cleaned daily

using 1000ppm NaDCC. All surface sampling was performed in the morning before the first

cleaning cycle for the day.

Air sampling

Six NIOSH BC 251 bioaerosol samplers were placed in each of three AIIRs in the general

ward to collect air samples. Particles collected with the NIOSH sampler are distributed into

three size fractions. Particles >4 μm in diameter are collected in a 15 mL centrifuge tube,

particles 1-4 μm in diameter are collected in a 1.5 mL centrifuge tube, and particles <1 μm in

diameter are collected in a self-assembled filter cassette containing a 37-mm diameter,

polytetrafluoroethylene (PTFE) filter with 3μm pores. All NIOSH samplers were connected

to either SKC AirCheck TOUCH Pumps or SKC Universal air sampling pumps set at a flow-

rate of 3.5 L/min and run for four hours, collecting a total of 5,040 L of air from each

patient's room.

In the room of Patient 1, three NIOSH samplers were attached to each of two tripod stands

and situated at different heights from the ground (1.2m, 0.9m, and 0.7m) near the air exhaust

to capture particles from the unidirectional airflow in the room. Throughout the four-hour

sampling period, Patient 1 was intermittently facing the NIOSH samplers while seated one

meter from the first tripod and 2.1 meters from the second tripod. Four SKC 37mm PTFE

filter (0.3μm pore size) cassettes were also distributed throughout the room and connected to

SKC Universal air sampling pumps set at a flow-rate of 5 L/min, each collecting an

additional 1,200 L of air from the room.

In the rooms of Patients 2 and 3, three NIOSH samplers were attached to each of two tripod

stands and situated at different heights from the ground (1.2m, 0.9m, and 0.7m). Throughout

the four-hour sampling period, Patients 2 and 3 remained in bed within 1 meter from all 6

NIOSH samplers (Supplementary Figure 1). Patient 3 was also talking on the phone for a

significant proportion of time during sampling. Additional SKC pumps with PTFE filter

cassettes were not used in the rooms of Patient 2 and 3.

The 6 NIOSH samples from each room were pooled prior to analysis, but the particle size

fractions remained separated. Each sample pool was representative of 5,040 L air.

Surface sampling

Surface samples were collected with Puritan® EnviroMax Plus pre-moistened macrofoam

sterile swabs (25-88060). Eight to 20 surface samples were collected in each room. Five were

high-touch surfaces, including the cardiac table, entire length of the bed rails including bed

control panel and call bell, bedside locker, electrical switches on top of the beds, and chair in

general ward rooms (Supplemental Figure 1). In ICU rooms, the ventilator and infusion

pumps were sampled instead of the electrical switches on top of the beds and chair

(Supplemental Figure 2). Air exhaust outlets and glass window surfaces were sampled in five

rooms, including the three rooms in which air sampling was performed. Toilet seat and

automatic flush button (one combined swab) were sampled in AIIR rooms in the general

ward.

Sample transfer and processing

All samples were immediately stored at 4°C in the hospital prior to transfer to a BSL-3

laboratory where samples were immediately processed and stored at -80°C unless directly

analyzed. Prior to RNA extraction, NIOSH aerosol sample tubes and filters were processed as

previously described (7), with slight modification due to the pooling of samples.

Laboratory methods

The QIAamp viral RNA mini kit (Qiagen Hilden, Germany) was used for sample RNA

extraction. Real-time PCR assays targeting the envelope (E) genes (8) and an in house orf1ab

assay were used to detect SARS-CoV-2 in the samples (9). All samples were run in duplicate

and with both assays. Positive detection was recorded as long as amplification was observed

in at least 1 assay.

Statistical analysis

Statistical analysis was performed using Stata version 15.1 (StataCorp, College Station,

Texas) and GraphPad Prism 8.0 (GraphPad Software, Inc., San Diego). $P < 0.05$ was

considered statistically significant, and all tests were 2-tailed. For the surface environment,

outcome measures analyzed were any positivity by room and pooled percentage positivity by

day of illness and respiratory viral load (represented by clinical cycle threshold (Ct) value).

We analyzed the factors associated with environmental contamination using the Student t-

test, or the nonparametric Wilcoxon rank-sum test was used for continuous variables

depending on their distribution. The $\chi 2$ or Fisher exact test was used to compare categorical

variables. We plotted the best fit curve by least-square method to study the environmental

contamination distribution across various the days of illness and clinical Ct value.

Results

Environmental sampling was conducted in three AIIRs in the ICU and 27 AIIRs in the

general ward. Air sampling was performed in three of the 27 AIIRs in the general ward. All

patients reported COVID-19 symptoms. Seven patients (23%) were asymptomatic at the time

of environmental sampling. Of the 23 symptomatic patients, 18 (78%) had respiratory

symptoms, one had gastrointestinal symptoms, one had both respiratory and gastrointestinal

symptoms, and three patients (10%) had fever or myalgia only (Supplemental Table 1).

There were no baseline differences between patients with environmental surface

contamination and those without, in terms of age, comorbidities, and positive clinical sample

on the day of sampling. Median cycle threshold (Ct) values of the clinical specimens for

patients with and without environmental surface contamination were 25.69 (IQR 20.37 to

34.48) and 33.04 (28.45 to 35.66) respectively (Table 1).

Of the rooms with environmental contamination, the floor was most likely to be contaminated

(65%), followed by the bed rail (59%), and bedside locker (42%) (Figure 1). Contamination

of toilet seat and automatic toilet flush button was detected in five out of 27 rooms, and all

five occupants had reported gastrointestinal symptoms within the preceding one week of

sampling. We did not detect surface contamination in any of the three ICU rooms.

Presence of environmental surface contamination was higher in week 1 of illness (Figure 2)

and showed association with the clinical cyclical threshold (P=0.06). Surface environment

contamination was not associated with the presence of symptoms or supplementary oxygen

(Table 1). In a subgroup analysis, the presence and extent of high-touch surface

contamination were significantly higher in rooms of patients in their first week of illness

(Figure 2). The best fit curve with the least-squares fit (Figure 3) showed that the extent of

high-touch surface contamination declined with increasing duration of illness and Ct values.

There was also no correlation between the Ct values of clinical samples and the Ct values of

environmental samples across the days of illness (Supplemental Figure 3).

Air samples from two (66.7%) of three AIIRs tested positive for SARS-CoV-2, in particle

sizes >4 µm and 1-4 µm in diameter (Table 1). Total SARS-CoV-2 concentrations in air

ranged from 1.84×10^3 to 3.38×10^3 RNA copies per m^3 air sampled. Rooms with viral

particles detected in the air also had surface contamination detected.

Discussion

Surface sampling revealed that the PCR-positivity high-touch surfaces was associated with

nasopharyngeal viral loads and peaked at approximately day four to five of symptoms. Air

sampling of the AIIR environments of two COVID-19 patients (both day five of illness with

high nasopharyngeal swab viral loads) detected the presence of SARS-CoV-2 particles sized

1-4 µm and > 4 µm. The absence of any detection of SARS-CoV-2 in air samples of the third

patient (day nine of illness with lower nasopharyngeal viral load concentration) suggests that

the presence of SARS-CoV-2 in the air is possibly highest in the first week of illness.

Recent aggregated environmental sampling and laboratory experiments have examined the

particle size distribution of SARS-CoV-2 in the air. A study from Wuhan, China sampled

three different environmental settings and detected aerosol size range particles (10).

Additionally, a recent laboratory study demonstrated the ability of SARS-CoV-2 to remain

viable in aerosols for up to 3 hours (11). While limited in subject numbers, our study

examined this issue at the individual patient-level, thus enabling correlation of particle size

distribution in the air with symptoms duration and nasopharyngeal viral loads. The absence of

aerosol-generating procedures or intranasal oxygen supplementation reduces the possibility

of our current findings being iatrogenic in nature. Larger individual patient-level studies

examining the droplet and aerosolizing potential of SARS-CoV-2 over different distances and

under different patient and environmental conditions are rapidly needed to determine the

generalizability of our current findings.

In the current analysis the presence and concentration of SARS-CoV-2 in air and high-touch

surface samples correlated with the day of illness and nasopharyngeal viral loads of COVID-

19 patients. This finding is supported by multiple observational clinical studies have

demonstrated that SARS-CoV-2 viral loads peak in the first week among COVID-19 patients

(2, 12, 13). This finding could help inform public health and infection prevention measures in

prioritizing resources by risk stratifying COVID-19 patients by their potential to directly or

indirectly transmit the SARS-CoV-2 virus to others.

Our study was limited in that it did not determine the ability of SARS-CoV-2 to be cultured

from the environmental swabs and the differentially-sized air particles which would be vital

to determining the infectiousness of the detected particles. Another study from Nebraska

attempted virus culture on SARS-CoV-2 PCR-positive air samples, however could not isolate

viable virus (14). The difficulty in culturing virus from air samples arises from low virus

concentrations, as well as the compromised integrity of the virus due to air sampling

stressors. Future studies using enhanced virus culture techniques could be considered (15),

and efforts to design a culture method to isolate virus from our samples is underway. Second,

sampling in an AIIR environment may not be representative of community settings and

further work is needed to generalize our current findings. Third, we sampled each room at a

single timepoint during the course of illness and did not track environmental contamination

over the course of illness for individual patients. Fourth, as clinical results were within 72

hours of environmental testing, it is plausible that during the day of testing, viral load was

actually low or negligible, hence limiting environmental contamination.

Current epidemiologic evidence does not seem to point to aerosolization as the key route of

transmission of SARS-CoV-2 (16). Detailed epidemiologic studies of outbreaks, in both

healthcare and non-healthcare settings, should be carried out to determine the relative

contribution of various routes of transmission and their correlation with patient-level factors.

In conclusion, in a limited number of AIIR environments, our current study involving

individual COVID-19 patients not undergoing aerosol-generating procedures or oxygen

supplementation suggest that SARS-CoV-2 can be shed in the air from a patient in particles

sized between 1 to 4 microns. Even though particles in this size range have the potential to

linger longer in the air, more data on viability and infectiousness of the virus would be

required to confirm the potential airborne spread of SARS-CoV-2. Additionally, the

concentrations of SARS-CoV-2 in the air and high-touch surfaces could be highest during the

first week of COVID-19 illness. Further work is urgently needed to examine these findings in

larger numbers and different settings to better understand the factors affecting air and surface

spread of SARS-CoV-2 and inform effective infection prevention policies.

Acknowledgements

We thank Dr Bill Lindsley (National Institute for Occupational Safety and Health, NIOSH)

for loaning us aerosol samplers and guiding us in their use; Dr. Raquel Binder and Emily

Robie from the Duke One Health team for quickly providing us equipment for use in this

study; Dr Hsu Li Yang (Saw Swee Hock School of Public Health, National University of

Singapore) for facilitating the air sampling; and Dr Ding Ying (National Centre for Infectious

Diseases, Singapore) for helping with project coordination. No compensation was received

for their roles in the study.

We further thank the colleagues in DSO National Laboratories in the environmental detection

team and clinical diagnostics team for BSL3 sample processing and analysis as well as the

logistics and repository team for transport of biohazard material, inventory and safekeeping

of received items.

Conflict of Interest Disclosures

None reported.

Funding

This study is funded by NMRC Seed Funding Program (TR19NMR119SD), NMRC COVID-

19 Research Fund (COVID19RF-001) and internal funds from DSO National Laboratories.

Ng OT is supported by NMRC Clinician Scientist Award (MOH-000276). K Marimuthu is

supported by NMRC CS-IRG (CIRG18Nov-0034). Chia PY is supported by NMRC

Research Training Fellowship (NMRC/Fellowship/0056/2018).

Role of Funder/Sponsors

References

1. Ong SWX, Tan YK, Chia PY, Lee TH, Ng OT, Wong MSY, et al. Air, Surface Environmental, and Personal Protective Equipment Contamination by Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) From a Symptomatic Patient. JAMA. 2020.

2. Young BE, Ong SWX, Kalimuddin S, Low JG, Tan SY, Loh J, et al. Epidemiologic Features and Clinical Course of Patients Infected With SARS-CoV-2 in Singapore. JAMA. 2020.

3. Chan JF-W, Yuan S, Kok K-H, To KK-W, Chu H, Yang J, et al. A familial cluster of pneumonia associated with the 2019 novel coronavirus indicating person-to-person transmission: a study of a family cluster. The Lancet. 2020.

4. Pan X, Chen D, Xia Y, Wu X, Li T, Ou X, et al. Asymptomatic cases in a family cluster with SARS-CoV-2 infection. Lancet Infect Dis. 2020.

5. Bai Y, Yao L, Wei T, Tian F, Jin D-Y, Chen L, et al. Presumed Asymptomatic Carrier Transmission of COVID-19. JAMA. 2020.

6. WHO. Report of the WHO-China Joint Mission on Coronavirus Disease 2019 (COVID-19). 2020.

7. Coleman KK, Sigler WV. Airborne Influenza A Virus Exposure in an Elementary School. Scientific Reports. 2020;10(1):1859.

8. Corman VM, Landt O, Kaiser M, Molenkamp R, Meijer A, Chu DK, et al. Detection of 2019 novel coronavirus (2019-nCoV) by real-time RT-PCR. Eurosurveillance. 2020;25(3):2000045.

9. Young BE, Ong SWX, Kalimuddin S, Low JG, Tan SY, Loh J, et al. Epidemiologic Features and Clinical Course of Patients Infected With SARS-CoV-2 in Singapore. JAMA. 2020.

293 10. Liu Y, Ning Z, Chen Y, Guo M, Liu Y, Gali NK, et al. Aerodynamic Characteristics

294   and RNA Concentration of SARS-CoV-2 Aerosol in Wuhan Hospitals during

295   COVID-19 Outbreak. BioRxiv. 2020.

296 11. van Doremalen N, Bushmaker T, Morris DH, Holbrook MG, Gamble A, Williamson

297   BN, et al. Aerosol and Surface Stability of SARS-CoV-2 as Compared with SARS-

298   CoV-1. New England Journal of Medicine. 2020.

299 12. Zou L, Ruan F, Huang M, Liang L, Huang H, Hong Z, et al. SARS-CoV-2 Viral Load

300   in Upper Respiratory Specimens of Infected Patients. N Engl J Med. 2020.

301 13. Wang W, Xu Y, Gao R, Lu R, Han K, Wu G, et al. Detection of SARS-CoV-2 in

302   Different Types of Clinical Specimens. JAMA. 2020.

303 14. Santarpia JL, Rivera DN, Herrera V, Morwitzer MJ, Creager H, Santarpia GW, et al.

304   Transmission Potential of SARS-CoV-2 in Viral Shedding Observed at the University

305   of Nebraska Medical Center. medRxiv. 2020.

306 15. Cao G, Noti J, Blachere F, Lindsley W, Beezhold D. Development of an improved

307   methodology to detect infectious airborne influenza virus using the NIOSH bioaerosol

308   sampler. Journal of Environmental Monitoring. 2011;13(12):3321-8.

309 16. Ng K, Poon BH, Kiat Puar TH, Shan Quah JL, Loh WJ, Wong YJ, et al. COVID-19

310   and the Risk to Health Care Workers: A Case Report. Ann Intern Med. 2020.

312   **Tables & Figures**

313   <u>**Table 1. Airborne severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2)**</u>

314   <u>**detections in hospital rooms of infected patients**</u>

Patient	Day of illness	Symptoms on day of air sampling	Clinical Ct value[*]	Airborne SARS-CoV-2 concentrations (RNA copies m^{-3} air)	Aerosol particle size	Samplers used
1	9	Cough, nausea, dyspnea	33.22	ND ND	-- --	NIOSH SKC Filters
2	5	Cough, dyspnea	18.45	2,000 1,384	>4 μm 1-4 μm	NIOSH
3	5	Asymptomatic[†]	20.11	927 916	>4 μm 1-4 μm	NIOSH

ND = none detected

[*]PCR cycle threshold value from patient's clinical sample

[†]Patient reported fever, cough, and sore throat until the day before the sampling. Patient reported no symptoms on the day of sampling, however was observed to be coughing during sampling

316 **Table 2: Baseline clinical characteristics of COVID-19 patients with environmental**

317 **contamination**

	With surface environment contamination (n=17)	Without surface environment contamination (n=13)	*P*-value
Median age (IQR)	52 (42 to 62)	44 (36 to 55)	0.7535
Male Sex (%)	6 (46%)	8 (47%)	0.961
Median Age Adjusted Charlson's Comorbidity Index (IQR)	1 (0 to 2)	1 (0 to 1)	0.6924
Median day of Illness (IQR)	5 (4 to 9)	13 (5 to 20)	0.1715
Median day of stay in room (IQR)	3 (3 to 8)	4 (2 to 16)	0.9491
Oxygen requirement (%)	0	4 (31)	0.026
Symptomatic (%)	12 (71)	11 (85)	0.427
Respiratory symptoms (%)	11 (65)	7 (54)	0.547
Gastrointestinal symptoms (%)	1 (6)	1 (8)	1.000
Clinical Cycle threshold value, median (IQR)[*]	25.69 (20.37 to 34.48)	33.04 (28.45 to 35.66)	0.056

318 [*]PCR cycle threshold value from patient's clinical sample

320 **Figure 1: Percentage of contaminated swabs from surface samples, in rooms with any**

321 **contamination**

322 All other sites were n=17, except for air exhaust vents where n=5

Figure 2: 2a. Percentage of patients with contamination of high touch surfaces in in the

first week of illness compared with more than first week of illness. 2b. Percentage of

surfaces contaminated across weeks of illness. 2c. Percentage of high-touch surfaces

contaminated across weeks of illness

Figure 3: 3a. Mean percentage of high touch surface contaminated by day of illness with

95% confidence interval with best fit curve. 3b. Percentage of high touch surfaces

contaminated by clinical cycle threshold values with best fit curve. 3c. Mean percentage

of high touch surface contaminated by day of illness with 95% confidence interval

grouped by symptoms

345