

1 **Title: No Evidence for Temperature-Dependence of the COVID-19 Epidemic**

2 **Running Title:** Temperature-independence of COVID-19 Epidemic

3 **Keywords:** COVID-19, epidemic, temperature, exponential rate, R0

4 **Authors:** Tahira Jamil^{1,2}, Intikhab Alam¹, Takashi Gojobori¹, and Carlos M. Duarte^{1,2}

5

6 **Affiliations:**

7 1. Computational Bioscience Research Center (CBRC), King Abdullah University of Science
8 and Technology, Thuwal 23955, Saudi Arabia

9 2. Red Sea Research Centre (RSRC), King Abdullah University of Science and Technology,
10 Thuwal 23955, Saudi Arabia

11

12 **1. Abstract**

13 The pandemic of the COVID-19 disease extended from China across the north-temperate
14 zone, and more recently to the tropics and southern hemisphere. We find no evidence that
15 spread rates decline with temperatures above 20 °C, suggesting that the COVID-19 disease is
16 unlikely to behave as a seasonal respiratory virus.

17

18 **2. Introduction**

19 On 30th January the WHO declared the novel coronavirus (COVID-19) outbreak a
20 public health emergency of international concern (<http://www.euro.who.int/en/home>). The
21 epidemic spread gradually from Wuhan province in China, to other Asian nations, the middle
22 east and Europe. By early March the epidemic was mostly concentrated in territories
23 extending between 30°N and 50 °N (Sajadi et al., 2020), now in late winter, leading to the
24 suggestion, echoed by the global media, that the epidemic is likely to be temperature-
25 dependent. This supported speculation of possible decline in severity with the advent of

26 warmer spring and summer temperatures in north-temperate latitudes (Sajadi et al.,
27 2020; Wang et al., 2020), comparable to many viruses affecting human respiratory systems,
28 including SARS (Tan et al., 2005; Gaunt et al., 2010).

29 However, recent (updated up to April 15, 2020; cf. Methods) data revealed the spread
30 of the epidemic across territories experiencing warm temperatures in the tropics (e.g.
31 Indonesia, Singapore, Brazil) and southern hemisphere as well (e.g. Australia, Argentina).
32 The current distribution of the epidemic challenges, therefore, the inference that SARS-CoV-
33 2 may behave as a seasonal respiratory virus based on previous statistical analyses from
34 earlier realized distributions.

35 Here we examine the relationship between the apparent exponential rate of SARS-
36 CoV-2 spread (γ) and the Effective Reproductive number (R_t) of infection and the average
37 daily temperature (T_{avg}) across nations and Chinese provinces where epidemics, with at least
38 100 case reported, have been reported (data updated up to 15th April, 2020).

39 **3. Methods**

40 *Novel Coronavirus (COVID-19) Cases Data*

41 The Novel Coronavirus (COVID-19) daily data are confirmed cases for affected
42 countries and provinces of China reported between 31st December 2019 to 15th April 2020.
43 The data was collected from the reports released by WHO, European Centre for Disease
44 Prevention and Control (ECDC), and John Hopkin CSSA. Data include confirmed and a
45 cumulative total of COVID-19 cases in affected countries/provinces.

46 *Average ambient temperature*

47 The average temperatures of all the affected countries were collected
48 from <https://www.timeanddate.com/>. The monthly mean temperature of February, March and
49 the two-weeks mean temperature of April of capital cities for the various nations were used
50 as reference temperatures for the country.

51

52 *Statistical Analysis*

53 The number of COVID-19 incidences follows the expected exponential growth,
54 although rates are only robust when cases exceed 100 persons for any country or province.
55 Hence, we fitted the exponential model to each country and each province of China. We
56 calculated exponential rate parameters for the countries where the COVID-19 incident has at
57 least a 10-day growth period, and the total number of cases was at least 100.

$$58 \quad N = ae^{\gamma Days}, \quad \gamma > 0$$

$$59 \quad \log N = \alpha + \gamma Days.$$

60 Where N is the cumulative number of diagnosed persons and Days is the number of days and
61 γ is the exponential rate ($100 \times \gamma = \% \text{ increase per day}$).

62

63 To calculate the effect of temperature on the exponential rate parameter, we first
64 regressed the exponential rate parameters retrieved from the exponential model on $Temp$ and
65 $Temp^2$

$$\gamma \sim Temp + Temp^2$$

66 If the squared term is significant, it provides evidence of nonlinearity.

67 The thermal performance of COVID-19 was characterized by fitting spread rate estimate or
68 growth parameter (γ) and temperature to the Gaussian function;

$$\gamma = ae^{\left[-0.5\left(\frac{Temp-opt}{tol}\right)^2\right]},$$

69 $Temp$ is the average temperature (in °C) that best encompasses the growth period of COVID-
70 19 cases since its first incidence in a country/region of China. Where, a (amplitude) is the
71 coefficient related to maximum of spread rate of countries, the optimum (opt) on the
72 temperature gradient is where the maximum of spread rate is attained and the tolerance (tol)
73 gives the width of the response curve. This model has non-linear form, and the model

74 parameters opt and tol occur nonlinearly in the model function. Parameter of thermal
75 performance curve was estimated by fitting Gaussian model to the growth rate and
76 temperature of infected countries. The initial values for the Gaussian parameters
77 opt , tol and a were obtained directly using maximum-likelihood polynomial regression for
78 the Gaussian function.

79 Estimated the Effective reproductive number (R_t), the average number of infections at
80 time t , per infected case over the course of their infection for COVID-19 for provinces of
81 China and other countries using a discrete γ distribution with a mean of 4.8 days and a
82 standard deviation of 3.5 days for the *serial interval* distribution.

83 All analyses were performed using R statistical computing software.

84

85 **4. Results**

86 Our results show that evidence for a temperature-dependence of the transmission
87 reported in previous papers was likely to be an artifact, reflecting the pathways of spread, and
88 that there is no evidence for thermal dependence of the transmission across the -1 to 36°C
89 T_{avg} range across the affected regions. This suggests little basis to expect evidence for the
90 virus to behave as a seasonal respiratory virus.

91 Epidemiological data consisting in the rate of increase in accumulated diagnosed
92 cases among nations (global) shows γ ranging from $1\% \text{ day}^{-1}$ to $23.8\% \text{ day}^{-1}$ (Figure S1),
93 with an average of $9.82 \pm 0.39\% \text{ day}^{-1}$ (Figure 1, Figure S1A), and apparent R_t of 1.27 ± 0.02
94 (Figure 1, Figure S1A). Surprisingly, γ and R_t across Chinese provinces (mean \pm SE = $1.3 \pm$
95 $0.28\% \text{ day}^{-1}$ and 0.96 ± 0.02) (Figure S1B) were well below those of other nations (mean \pm
96 SE = $19.82 \pm 0.39\% \text{ day}^{-1}$ and 1.27 ± 0.02), possibly because much faster confinement of the
97 Chinese population did not allow for the potential exponential rates under uncontrolled
98 conditions to be realized. The broad variability in realized γ and R_t between nations (global)

99 and provinces (China) largely reflects differences in detection likelihood along with the
100 timing and rigour of adoption of confinement measures.

101 The relationship between γ and R_t and T_{avg} shows no evidence for a reduced spread
102 rate with warming (Figure 1), unlike analyses based on previous data. A number of nations
103 with $T_{avg} > 20$ °C, including subtropical and tropical (Brazil, Cuba, UAE, Saudi Arabia, India
104 and Panama), and southern-hemisphere (Peru, Argentina, Indonesia) nations (Figure 2),
105 support γ and R_t above the median values of 9.6% day⁻¹ and 1.23, respectively (Figure 1).
106 However, the same analysis conducted one weeks ago (15th March), did provide some
107 evidence for low γ and R_t for $T_{avg} > 20$ °C (Figure S2). Our updated results (Figure 1) show,
108 however, that this apparent temperature-dependence was confounded with a prevailing zonal
109 pattern of spread across the north-temperate zone, possibly reflecting the main patterns of
110 human mobility, which delayed arrival of the epidemics to the southern hemisphere and the
111 tropics.

112 5. Discussion

113 These results suggest that, contrary to prior assessments, the spread rate of the
114 COVID-19 pandemic is temperature-independent, it is transmitting in countries with warm
115 weather. This signifying that there is little hope for relief as temperatures in the northern
116 hemisphere increase, and that poor nations with weak health systems in tropical regions, such
117 as African, are at great risk. Therefore, in order to reduce transmission, it's important to
118 employ strong lockdown, social distancing and testing and tracking polices.

119

120 Data sources: The data on COVID-19 is available publicly across many sources; where
121 downloadable data files are updated daily few are listed below;

122 World health organization (<https://www.who.int/emergencies/diseases/novel-coronavirus-2019/situation-reports/>)
123

124

125 Johns Hopkins CSSE (<https://data.humdata.org/dataset/novel-coronavirus-2019-ncov-cases>)

126 [Accessed April 15, 2020]

127

128 European Centre for Disease Prevention and Control

129 ([https://www.ecdc.europa.eu/en/publications-data/download-todays-data-geographic-](https://www.ecdc.europa.eu/en/publications-data/download-todays-data-geographic-distribution-covid-19-cases-worldwide)

130 [distribution-covid-19-cases-worldwide](https://www.ecdc.europa.eu/en/publications-data/download-todays-data-geographic-distribution-covid-19-cases-worldwide)) [Accessed April 15, 2020].

131

132 **Author Contribution**

133

134 CMD and TJ conceived and designed the research, TJ conducted the analysis, TJ and CMD

135 wrote the first draft and all co-authors contributed to improving the paper and approved the

136 submission.

137 **Funding**

138 This research was supported by funding provided by the King Abdullah University of

139 Science and Technology to the CBRC.

140 **References**

141 Gaunt, E.R., Hardie, A., Claas, E.C., Simmonds, P., and Templeton, K.E. (2010).

142 Epidemiology and clinical presentations of the four human coronaviruses 229E, HKU1,
143 NL63, and OC43 detected over 3 years using a novel multiplex real-time PCR method.

144 *Journal of clinical microbiology* 48, 2940-2947.

145 Sajadi, M.M., Habibzadeh, P., Vintzileos, A., Shokouhi, S., Miralles-Wilhelm, F., and

146 Amoroso, A. (2020). Temperature and latitude analysis to predict potential spread and

147 seasonality for COVID-19. Available at SSRN 3550308.

148 Tan, J., Mu, L., Huang, J., Yu, S., Chen, B., and Yin, J. (2005). An initial investigation of the

149 association between the SARS outbreak and weather: with the view of the

150 environmental temperature and its variation. *Journal of Epidemiology & Community*

151 *Health* 59, 186-192.

152 Wang, M., Jiang, A., Gong, L., Luo, L., Guo, W., Li, C., Zheng, J., Li, C., Yang, B., and

153 Zeng, J. (2020). Temperature significant change COVID-19 Transmission in 429

154 cities. medRxiv.

155

156

157

158

159

160

161

162

163

164

165

166

167

168 **Figure legends**

169 **Figure 1.** The relationship between the apparent exponential rate of SARS-CoV-2 spread (β)
170 and the Effective Reproductive number of infection (R_t) and the average daily temperature
171 (T_{avg}) across nations and Chinese provinces where > 100 cases of COVID-19 have been
172 reported (data last accessed April 15, 2020, Figure S1). Green symbols represent provinces in
173 China while red symbols represent other nations. Neither the double exponential function
174 with temperature nor the Gaussian function provided a significant ($p < 0.05$) fit for either γ or
175 R_t with temperature.

176

177 **Figure 2.** Distribution of the apparent exponential rate of SARS-CoV-2 spread (γ) and the
178 Effective Reproductive number of infection (R_t) and the average daily temperature (T_{avg})
179 across nations where > 100 cases of COVID-19 have been reported (data last accessed April
180 15, 2020).

181

182 **Appendix Figure S1.** The apparent average (\pm SE) exponential rate of SARS-CoV-2 spread
183 (γ), the average (and 95% confidence limits) of Effective Reproductive number of infection
184 (R_t) and the average daily temperature (T_{avg}), total case and number of days since the first
185 case reported across nations and Chinese provinces where epidemics, with at least 100 case
186 reported, have been reported (data updated through April 15, 2020).

187

188 **Appendix Figure S2.** The relationship between the apparent exponential rate of SARS-CoV-
189 2 spread (γ) and the Effective Reproductive number (R_t) of infection and the average daily
190 temperature (T_{avg}) across nations and Chinese provinces where > 100 cases of COVID-19
191 have been reported, as of Figure 1, but with data updated only until 15th March. The Gaussian
192 function with temperature provided a significant fit for γ with temperature.



