



26 **ABSTRACT** [275/275 words]

27 **Purpose:** USA Football established five levels of contact (LOC) to guide the intensity of high  
28 school football practices. However, it remains unclear whether head impact exposure differs by  
29 LOC. The purpose of this study was to examine head impact frequency and magnitude by LOC  
30 in the overall sample and three position groups.

31  
32 **Methods:** This longitudinal observational study included 24 high school football players during  
33 all practices and games in the 2019 season. Players wore a sensor-installed mouthguard that  
34 monitored head impact frequency, peak linear acceleration (PLA), and rotational head  
35 acceleration (PRA). Practice/game drills were filmed and categorized into 5 LOCs (*air, bags,*  
36 *control, thud, live*), and head impact data were assigned into 5 LOCs. Player position was  
37 categorized into linemen, hybrid, and skill.

38  
39 **Results:** A total of 6016 head impacts were recorded during 5 LOCs throughout the season. In  
40 the overall sample, total number of impacts, sum of PLA, and PRA per player increased in an  
41 incremental manner (*air<bags<control<thud<live*), with the most head impacts in *live*  
42 ( $113.7 \pm 17.8$  hits/player) and the least head impacts in *air* ( $7.7 \pm 1.9$  hits/player). The linemen and  
43 hybrid groups had consistently higher impact exposure than the skill group. Average head impact  
44 magnitudes by position group were higher during *live* drills (PLA (41.0-45.9g) and PRA (3.3-4.6  
45  $\text{krad/s}^2$ ) per head impact), whereas other LOCs had lower magnitudes (PLA (18.2-23.2g) and  
46 PRA (1.6-2.3 $\text{krad/s}^2$ ) per impact).

47

48 **Conclusion:** Our data suggest that LOC may influence cumulative head impact exposure in high  
49 school football, with players incurring frequent head impacts during *live*, *thud*, and *control*. The  
50 data indicate the importance of considering LOCs to refine practice guidelines and policies to  
51 minimize head impact burden in high school football athletes.

52

## 53 INTRODUCTION

54 The long-term consequence of sport-related head injury is a complex public health issue  
55 with no concrete solution (1, 2). Despite inherent risk of head injury in contact sports (e.g.,  
56 American football, hockey, soccer), participating in these team sports, especially during  
57 developmental age, provides well-documented benefits, including higher levels of physical  
58 activity, improved mental health, and lower likelihood of smoking cigarettes and using illegal  
59 substances (3). In an attempt to promote a safer environment, youth football organizations have  
60 implemented the teaching of proper tackling techniques and reduced the size of the field and  
61 number of players on the field (4). These modifications are further substantiated in high school,  
62 college, and professional football, where kickoffs and touchbacks were adjusted to reduce  
63 injuries (5), well-designed protective helmets are worn, and tackling of defenseless players and  
64 blindside blocking are prohibited (6, 7). Consistent with these adjustments, concussion and  
65 catastrophic injury rates have been reduced (8, 9); however, subconcussive head impact exposure  
66 has proven more complex.

67 Subconcussive head impact is defined as a hit to the head that does not induce overt  
68 concussion symptoms (10). These head impacts are most common in American football, where  
69 athletes can experience several hundred impacts with some exceeding 1,000 head impacts in a  
70 single season (11). Evidence has emerged to indicate that both high school and college football  
71 players with frequent subconcussive head impact exposure exhibit neuronal microstructural  
72 damage (12, 13), abnormal brain activation (14, 15), ocular-motor impairment (16), and  
73 elevation in brain-injury blood biomarkers (17). One line of research suggests that long-term  
74 exposure to these hits is a key factor in developing neurodegenerative disorders later in life (18,  
75 19). Although USA Football (for high school) and the NCAA (for college) have eliminated

76 practicing two times in the same day (two-a-days) to minimize head impact exposure, one study  
77 found that total head impact frequency during a summer camp increased by 26% (20). Similarly,  
78 a policy change to reduce the number of preseason practices from 29 to 25 failed to reduce head  
79 impact exposure in college football players, with one team's cumulative head impacts increasing  
80 up to 35% (21). These mixed results highlight the need to dissect football practices to increase  
81 our understanding of what type of contact drills and intensities cause the greatest exposure to  
82 subconcussive head impacts.

83 USA Football (the national governing body over amateur football) has identified five  
84 levels of contact (LOCs) that define the intensities and structure of football practices nationwide.  
85 The LOCs were designed to guide effective practice schedules through a step-by-step approach  
86 to teach fundamental football skills (22). The National Federation of State High School  
87 Associations implemented the LOCs in their high school football practice guidelines beginning  
88 in 2014. However, it remains unknown whether, and to what extent, different levels of contact  
89 influence head impact exposure in high school football players across an entire season and  
90 between position groups.

91 Therefore, we conducted a longitudinal observational study to examine cumulative head  
92 impact exposure across different drill intensities in high school football players over the course  
93 of a single season. Our primary hypothesis was that there would be an incremental increase in  
94 cumulative head impact kinematics between LOCs in the overall sample with *live* recording the  
95 greatest head impact exposure and air recording the lowest: *air* < *bags* < *control* < *thud* < *live*.  
96 Since the proximity to opponents and nature of contact during nearly every play for linemen  
97 (23), we also hypothesized that linemen would have greater head impact exposure in most LOCs,  
98 compared to the hybrid and skill positions. Our secondary aim was to identify the average head

99 impact magnitudes by LOC and position group. Lastly, we aimed to explore frequency of head  
100 impacts that were within 25-60 g, 60-100 g, or >100 g in each LOC.

101

## 102 **METHODS**

### 103 **Participants**

104 This single-site, observational study included 24 male high school football players at  
105 Bloomington High School-North. The study was conducted during the 2019 football season  
106 including practices and games during the pre-season, in-season, and playoffs. None of the 24  
107 players was diagnosed with a concussion during the study period as confirmed by team athletic  
108 trainer and physician. Inclusion criterion was being an active football team member. Exclusion  
109 criteria included a history of head neck injury (including concussion) in the previous year or  
110 neurological disorders. The Indiana University Institutional Review Board and the Monroe  
111 County Community School Corporation Research Review Board approved the study, and all  
112 participants and their legal guardians gave written informed consent.

113

### 114 **Study Procedures**

115 At the preseason data collection, self-reported demographic information (age,  
116 race/ethnicity, height, weight, number of previously diagnosed concussions, and years of  
117 experience in various contact sports, including tackle football) were obtained. Participants were  
118 custom-fitted with the Vector mouthguard (Athlete Intelligence, Inc.) that measured the number  
119 of hits and magnitude of head linear and rotational acceleration. Participants wore the  
120 mouthguard for all practices and games from pre-season training camp (August 13, 2019) to the  
121 end of the season (November 1, 2019). Head impact data were collected during 34 practices, 11

122 varsity games, and 9 junior varsity games over a 12-week season. Video data were collected  
123 using Hudl (Agile Sports Technologies, Inc.) during the same timeframe as subconcussive head  
124 impact data collection. Participants' playing positions were verified by team coaches and  
125 categorized into three groups as follows: 11 linemen athletes (defensive lineman, offensive  
126 lineman), 7 hybrid athletes (tight end, linebacker, running back), and 6 skill athletes (wide  
127 receiver, defensive back), which is in line with prior literature (24, 25). No quarterbacks were  
128 included in this study. In accordance with USA Football guidelines (26), head impacts were  
129 categorized by LOC: *air*, *bags*, *control*, *thud*, and *live*. See Levels of Contact and Film Review  
130 section for more details and supplemental file A for example video for each LOC.

131

## 132 **Head Impact Measurement**

133 This study used an instrumented Vector mouthguard for measuring frequency of head  
134 impacts as well as linear and rotational head kinematics during impacts, as previously described  
135 (27). The mouthguard employs a triaxial accelerometer (ADXL377, Analog Devices) with 200 g  
136 maximum per axis to sense linear acceleration. For rotational kinematics, a triaxial gyroscope  
137 (L3GD20H, ST Microelectronics) was employed. An impact is detected when a linear acceleration  
138 magnitude exceeds 10.0 g for three consecutive samples (sampling every 0.2 milliseconds). All  
139 impact data with a standard hit duration of 96 milliseconds were transmitted wirelessly through  
140 the antenna transmitter to the sideline antenna and computer, then stored on a secure internet  
141 database. The Vector mouthguard is installed with an in-mouth sensor to ensure that data  
142 acquisition occurs only when the mouthguard is securely fitted in one's mouth. Linear  
143 acceleration data were transformed within the Athlete Intelligence software to the head's center  
144 of gravity based on the 50<sup>th</sup> percentile male. From raw impact data extracted from the server, the

145 number of hits, peak linear acceleration, and peak rotational acceleration were used for  
146 descriptive analyses. Timestamps of all impacts were used to correspond with timeframes of  
147 each LOC.

148

### 149 **Levels of Contact and Film Review**

150 The five LOCs are *air*, *bags*, *control*, *thud*, and *live* with *air* being estimated to have the  
151 lowest intensity and *live* being the highest (26). *Air* is defined as drills being run unopposed and  
152 without contact. *Bags* is defined as drills being run against a bag or soft-contact surface. *Control*  
153 is defined as drills being run at an assigned speed until the moment of contact. It does not  
154 involve tackling, rather contact is above the waist and players stay on their feet. *Thud* is defined  
155 as drills being run at a competitive, fast speed through the moment of contact. It does not involve  
156 full tackling, rather contact is above the waist and players stay on their feet and a quick whistle  
157 ends the drills. *Live* is defined as drills being run in game-like conditions that include live-drill  
158 during practice as well as real games. *Live* should be the only time players are allowed to fully  
159 tackle another player to the ground. All head impacts in *air*, *bags*, *control*, and *thud* were during  
160 practices, whereas *live* occurred in both practices and games.

161 Daily subconcussive head impact exposures were quantified in our full sample of 24  
162 football players. Level of contact was assigned to each of the head impacts by corresponding the  
163 real-time, verified practice plan to the real-time head impact data set. Based on our preliminary  
164 analysis, there was a strong agreement on the head impact occurrence between the Vector  
165 mouthguard and film analysis.

166

### 167 **Statistical Analysis**

168 A series of descriptive analyses was conducted to examine all aims. Our outcome  
169 variables for primary and secondary aims were head impact count, sum of peak linear  
170 acceleration, and sum of peak rotational acceleration. These outcomes were expressed as mean  $\pm$   
171 standard error of means and assessed by 5 LOCs (*air*, *bags*, *control*, *thud*, and *live*) and 3  
172 position groups (lineman, hybrid, and skill). For the exploratory aim, a total number of head  
173 impact within ranges of 25-60 g, 60-100 g, or >100 g in each LOC was assessed in the overall  
174 sample.

175

## 176 **RESULTS**

### 177 *Demographics and overall head impact exposure*

178 A total of 6016 head impacts were recorded during 5 LOCs from 24 high school football  
179 players throughout the season, resulting in an average of 250.7 hits, 5891.0 g, and 528.5 krad/s<sup>2</sup>.  
180 These data are not reflective of head impacts that occurred outside the 5 LOCs, such as walk-  
181 through and pre-practice/game conditioning. Demographics and head impact data in the overall  
182 sample are detailed in Table 1.

183

### 184 *LOC-dependent cumulative head impact exposure*

185 LOCs displayed an influence on head impact exposure in high school football players, as  
186 illustrated by the increasing cumulative exposure corresponding to increased levels of contact.  
187 Specifically, in the overall sample, total number of impacts per player as well as sum of PLA and  
188 PRA increased in an incremental manner (*air* < *bags* < *control* < *thud* < *live*), with the most head  
189 impacts being observed in live ( $113.7 \pm 17.8$  hits,  $2,657.6 \pm 432.0$  g,  $233.9 \pm 40.1$  krad/s<sup>2</sup>) and  
190 the least head impact exposure in air ( $7.7 \pm 1.9$  hits,  $176.9 \pm 42.5$  g,  $16.7 \pm 4.2$  krad/s<sup>2</sup>) (Figure

191 1A-C). Of note, *live* impacts were primarily during games (N=2,512, 92.08%) as opposed to  
192 during practices (N=216, 7.92%).

193 All 3 position groups exhibited similar incremental patterns in head impact kinematics  
194 (*air < bags < control < thud < live*). However, the linemen and hybrid groups had consistently  
195 higher exposure in both frequency and magnitudes than the skill group across all LOCs (Figure  
196 2A-C). Head impact kinematics for each LOC are detailed in Supplemental Table 1.

197

### 198 *Head impact magnitudes by position group and LOC*

199 Average head impact magnitudes by position group were shown to be similar across  
200 LOCs for *air*, *bags*, *control*, and *thud*, ranging between 18.2 and 23.2 g for PLA and 1.6 and 2.3  
201  $\text{krad/s}^2$  for PRA per head impact. However, *live* showed two times as high of PLA (41.0 to 45.9  
202 g: Figure 3A) and PRA (3.3 to 4.6  $\text{krad/s}^2$ : Figure 3B) per head impact as other LOCs. The same  
203 pattern was observed in all 3 groups. Consistent with published papers (13, 23, 28), it was  
204 evident that there was a large number of head impacts across all LOCs that were within 10 to 30  
205 g, which might have diluted the minority of high magnitude head impacts. Our exploratory  
206 analysis revealed that in the overall sample, high magnitude head impacts (60-100 g and >100 g)  
207 were most prevalent in *live*, followed by *thud*, *control*, *bags*, and then *air* (Figure 4A-C).

208

## 209 **DISCUSSION**

210 Subconcussion research is still at its infancy, but it is a rapidly growing area of concern in  
211 sport injury prevention. To contribute to this emerging field, this study examined whether  
212 subconcussive head impact kinematics differed across LOCs and between player position groups.  
213 There were four key findings in this study. First, cumulative head impact exposure increased as

214 the LOC increased, with frequent head impacts observed during *live*, *thud*, and *control* drills.  
215 Second, in line with literature (11), linemen and hybrid groups had consistently higher head  
216 impact exposure compared to the skill position group across all LOCs. Third, the mean head  
217 impact magnitude was similar (18 to 23 g) across *air*, *bags*, *control*, and *thud*, but head impacts  
218 during *live* showed greater magnitudes (40 to 45 g). Lastly, very high impact magnitudes (>100  
219 g) were small in number overall but were more frequent in *thud* and *live* than other LOCs. Taken  
220 together, our data, for the first time, empirically support the USA football's categorization of  
221 LOC while calling for a need to dissect football practice guidelines to make more specific  
222 recommendations for practice and games to minimize cumulative burden on adolescents' brain  
223 health.

224 Owing to the sensor-installed helmets, mouthguards, headbands, and skin patches, our  
225 knowledge of head impact exposure in American football has drastically improved in the past 15  
226 years. These technological advancements allowed researchers to evaluate head impact frequency  
227 and magnitude in various position groups, practice types (e.g., shell-only, full-gear), play types  
228 (e.g., running, passing, special teams), and time-based hit rates (29-34). Previous research has  
229 suggested that overall head impact exposure elevates in relation to increased practice duration,  
230 contact intensity, and time spent in high risk drills (21, 32). However, these variables differ  
231 greatly between players and position groups. For example, previous literature demonstrated that  
232 the differences between individual players accounted for 48% of the variance in head impact  
233 exposure during practice (35). Additionally, recent research suggests different types of plays  
234 (e.g., running, passing, special teams) also have different average head impact magnitudes which  
235 will influence cumulative head impact exposure (34). Despite this high degree of variance  
236 between players and play types, the linemen and hybrid position groups consistently have higher

237 head impact exposure compared to skill positions such as receivers, defensive backs, and  
238 quarterbacks (30, 31). We were able to corroborate these observations in our study in the context  
239 of five different LOC, where frequent head impacts were notable especially during high intensity  
240 drills (*live, thud, control*), with linemen and hybrid players experiencing more head impacts than  
241 skill players.

242 Another major finding from this study was that *live* had the highest average PLA and  
243 PRA compared to *air, bags, control, and thud*, which shared similar impact magnitudes.  
244 Furthermore, strong magnitudes of head impacts (60 – 100 g and >100 g) were frequently  
245 observed during *live* and *thud*. These findings suggest current football guidelines that  
246 recommend limiting full contact (*live & thud*) during practice are appropriate. In fact, the  
247 football team in this study significantly minimized *live* drills during practice (n=216 impacts);  
248 therefore, 92% of head impacts in *live* was recorded during the games (n=2,512 impacts). In  
249 alignment with this idea, there have been attempts by various football governing bodies to  
250 restrict the amount of full contact in practices (20, 21, 26, 31, 36). For example, USA Football's  
251 National Practice Guidelines for Youth Tackle Football suggest that full-contact should not be  
252 done for more than 30 minutes per day and no more than 90 minutes per week during the regular  
253 season (26). By limiting full contact practices from 3 days to 2 days per week, there was an  
254 average decline of 42% in head impact exposure in high school football players, albeit average  
255 magnitude of head impact remained unchanged (31). Some of these positive changes, however,  
256 do not translate into the college setting. Even though the NCAA eliminated two-a-day practices  
257 and reduced practice frequency from 29 to 25 practices during summer camp, to compensate for  
258 the loss of practice times, football teams tended to incorporate more high-intensity, contact-prone  
259 practice drills, leading to increased head impact exposure (20, 21). These data further

260 substantiate the importance of regulating the duration of specific drill types, rather than  
261 restricting practice type and frequency.

262         There are several limitations to this study. Our examination of head impacts in high  
263 school football is limited in that it was conducted on a single high school football team in the  
264 Midwest composed of primarily white males; therefore, the results are not generalizable to the  
265 broader U.S. population of high school football teams. A second limitation of the study is that  
266 the USA Football LOC guidelines are just that, guidelines. They are not legislation, and it is  
267 unknown what percentage of high schools currently utilize the USA Football LOC system; thus,  
268 this further limits the generalizability. Implementation of the LOCs is not a perfectly reliable  
269 variable in that situations occur in practice LOCs that may overlap with other LOCs. Coaches in  
270 this high school attempted to implement the LOCs consistently, but to control up to 22 athletes  
271 participating in a single drill at the same time is not always feasible. These limitations portend a  
272 follow-up research question as to whether regulating the number of repetitions in each LOC is a  
273 more effective approach than teamwide contact restrictions for minimizing the frequency and  
274 magnitude of subconcussive head impacts in individual players (or position groups). Hence, for  
275 future research we suggest examining subject-specific head impact data that incorporate multiple  
276 predictor variables such as the repetitions per player per drill, LOC, position group, and impacts  
277 per event (i.e., drill, practice/game, or week) for head impact kinematic outcomes.

278

## 279 **CONCLUSION**

280         Our data suggest that the LOC may influence cumulative head impact exposure in high  
281 school football players, with players incurring frequent head impacts during *live*, *thud*, and  
282 *control*. Strong magnitudes of head impacts were frequently observed during *live* and *thud*.

283 Many head impact investigations and contact recommendations have focused on limiting the  
284 number of practices (20, 21, 37), time involved in full contact in general (26, 30, 38), or  
285 changing game rules (8). Our data indicate the importance of considering LOCs to refine football  
286 practice guidelines/policies to minimize cumulative head impact burden in high school football  
287 players.

288

289

## 290 REFERENCES

- 291 1. Broglio SP, Eckner JT, Paulson HL, Kutcher JS. Cognitive decline and aging: the role of  
292 concussive and subconcussive impacts. *Exerc Sport Sci Rev.* 2012;40(3):138-44.
- 293 2. Hutchison MG, Di Battista AP, McCoskey J, Watling SE. Systematic review of mental  
294 health measures associated with concussive and subconcussive head trauma in former  
295 athletes. *Int J Psychophysiol.* 2018;132(Pt A):55-61.
- 296 3. Logan K, Cuff S. Organized Sports for Children, Preadolescents, and Adolescents.  
297 *Pediatrics.* 2019;143(6):1-20.
- 298 4. USA Football Website [Internet]. Indianapolis, IN; [cited 2020 3/12]. Available from:  
299 <https://usafootball.com/rookietackle/>.
- 300 5. National Collegiate Athletic Association. New kickoff rule approved for 2018. 2018  
301 [cited 2020 3/12]. Available from: [https://www.ncaa.com/news/football/article/2018-04-](https://www.ncaa.com/news/football/article/2018-04-13/alteration-football-kickoff-rule-approved)  
302 [13/alteration-football-kickoff-rule-approved](https://www.ncaa.com/news/football/article/2018-04-13/alteration-football-kickoff-rule-approved).
- 303 6. National Federation of State High School Associations. New Blocking, Kicking Rules  
304 Address Risk Minimization in High School Football. *NFHS News* [Internet]. 2018 [cited

- 305 2020 3/12]. Available from: [https://www.nfhs.org/articles/new-blocking-kicking-rules-](https://www.nfhs.org/articles/new-blocking-kicking-rules-address-risk-minimization-in-high-school-football/)  
306 [address-risk-minimization-in-high-school-football/](https://www.nfhs.org/articles/new-blocking-kicking-rules-address-risk-minimization-in-high-school-football/).
- 307 7. National Football League [Internet]. New York, NY; [cited 2020 3/12]. Available from:  
308 <https://operations.nfl.com/the-rules/nfl-video-rulebook/defenseless-player/>.
- 309 8. Wiebe DJ, D'Alonzo BA, Harris R, Putukian M, Campbell-McGovern C. Association  
310 Between the Experimental Kickoff Rule and Concussion Rates in Ivy League Football.  
311 *JAMA*. 2018;320(19):2035-6.
- 312 9. McIntosh AS, McCrory P. Preventing head and neck injury. *Br J Sports Med*.  
313 2005;39(6):314-8.
- 314 10. McCrory P, Meeuwisse W, Dvorak J et al. Consensus statement on concussion in sport-  
315 the 5th international conference on concussion in sport held in Berlin, October 2016. *Br J*  
316 *Sports Med*. 2017.
- 317 11. Bailes JE, Petraglia AL, Omalu BI, Nauman E, Talavage T. Role of subconcussion in  
318 repetitive mild traumatic brain injury. *J Neurosurg*. 2013;119(5):1235-45.
- 319 12. Bahrami N, Sharma D, Rosenthal S et al. Subconcussive Head Impact Exposure and  
320 White Matter Tract Changes over a Single Season of Youth Football. *Radiology*.  
321 2016;281(3):919-26.
- 322 13. Hiran AA, Bazarian JJ, Merchant-Borna K et al. A common neural signature of brain  
323 injury in concussion and subconcussion. *Sci Adv*. 2019;5(8):eaau3460.
- 324 14. Breedlove EL, Robinson M, Talavage TM et al. Biomechanical correlates of  
325 symptomatic and asymptomatic neurophysiological impairment in high school football.  
326 *Journal of biomechanics*. 2012;45(7):1265-72.

- 327 15. Talavage TM, Nauman EA, Breedlove EL et al. Functionally-detected cognitive  
328 impairment in high school football players without clinically-diagnosed concussion.  
329 *Journal of neurotrauma*. 2014;31(4):327-38.
- 330 16. Kawata K, Rubin LH, Lee JH et al. Association of Football Subconcussive Head Impacts  
331 With Ocular Near Point of Convergence. *JAMA Ophthalmol*. 2016;134(7):763-9.
- 332 17. Kawata K, Liu CY, Merkel SF, Ramirez SH, Tierney RT, Langford D. Blood Biomarkers  
333 for Brain Injury: What Are We Measuring? *Neuroscience and biobehavioral reviews*.  
334 2016;68:460-73.
- 335 18. McKee AC, Stern RA, Nowinski CJ et al. The spectrum of disease in chronic traumatic  
336 encephalopathy. *Brain : a journal of neurology*. 2013;136(Pt 1):43-64.
- 337 19. Mez J, Daneshvar DH, Kiernan PT et al. Clinicopathological Evaluation of Chronic  
338 Traumatic Encephalopathy in Players of American Football. *Jama*. 2017;318(4):360-70.
- 339 20. Stemper BD, Shah AS, Harezlak J et al. Repetitive Head Impact Exposure in College  
340 Football Following an NCAA Rule Change to Eliminate Two-A-Day Preseason  
341 Practices: A Study from the NCAA-DoD CARE Consortium. *Ann Biomed Eng*.  
342 2019;47(10):2073-85.
- 343 21. Stemper BD, Shah AS, Mihalik JP et al. Head Impact Exposure in College Football  
344 following a Reduction in Preseason Practices. *Med Sci Sports Exerc*. 2020.
- 345 22. USA Football [Internet]. Indianapolis, IN; [cited 2020 3/12]. Available from:  
346 <https://usafootball.com/resources-tools/coach/levels-of-contact/>.
- 347 23. Duma SM, Manoogian SJ, Bussone WR et al. Analysis of real-time head accelerations in  
348 collegiate football players. *Clin J Sport Med*. 2005;15(1):3-8.

- 349 24. Beckwith JG, Greenwald RM, Chu JJ et al. Timing of concussion diagnosis is related to  
350 head impact exposure prior to injury. *Med Sci Sports Exerc.* 2013;45(4):747-54.
- 351 25. Broglio SP, Eckner JT, Martini D, Sosnoff JJ, Kutcher JS, Randolph C. Cumulative head  
352 impact burden in high school football. *Journal of neurotrauma.* 2011;28(10):2069-78.
- 353 26. USA Football [Internet]. Indianapolis, IN; [cited 2020 3/12]. Available from:  
354 <https://assets.usafootball.com/documents/practice-guidelines-youth.pdf>.
- 355 27. Wu LC, Nangia V, Bui K et al. In Vivo Evaluation of Wearable Head Impact Sensors.  
356 *Ann Biomed Eng.* 2015.
- 357 28. Crisco JJ, Fiore R, Beckwith JG et al. Frequency and location of head impact exposures  
358 in individual collegiate football players. *J Athl Train.* 2010;45(6):549-59.
- 359 29. Mihalik JP, Bell DR, Marshall SW, Guskiewicz KM. Measurement of head impacts in  
360 collegiate football players: An investigation of positional and event-type differences. In:  
361 2007, pp. 1229-35.
- 362 30. Broglio SP, Martini D, Kasper L, Eckner JT, Kutcher JS. Estimation of head impact  
363 exposure in high school football: implications for regulating contact practices. *Am J*  
364 *Sports Med.* 2013;41(12):2877-84.
- 365 31. Broglio SP, Williams RM, O'Connor KL, Goldstick J. Football Players' Head-Impact  
366 Exposure After Limiting of Full-Contact Practices. *J Athl Train.* 2016;51(7):511-8.
- 367 32. Asken BM, Brooke ZS, Stevens TC et al. Drill-Specific Head Impacts in Collegiate  
368 Football Practice: Implications for Reducing "Friendly Fire" Exposure. *Ann Biomed Eng.*  
369 2019;47(10):2094-108.
- 370 33. Reynolds BB, Patrie J, Henry EJ et al. Practice type effects on head impact in collegiate  
371 football. *Journal of neurosurgery.* 2015:1-10.

- 372 34. Urban JE, Flood WC, Zimmerman BJ et al. Evaluation of head impact exposure  
373 measured from youth football game plays. *J Neurosurg Pediatr.* 2019;1-10.
- 374 35. Campolettano ET, Rowson S, Duma SM et al. Factors Affecting Head Impact Exposure  
375 in College Football Practices: A Multi-Institutional Study. *Ann Biomed Eng.*  
376 2019;47(10):2086-93.
- 377 36. Cranmer GA, Sanderson J. "Rough Week for Testosterone": Public Commentary around  
378 the Ivy League's Decision to Restrict Tackle Football in Practice. *Western Journal of*  
379 *Communication.* 2018;82(5):631-47.
- 380 37. Broglio K, Matzo M. CE: Acute Pain Management for People with Opioid Use Disorder.  
381 *Am J Nurs.* 2018;118(10):30-8.
- 382 38. Pfaller AY, Brooks MA, Hetzel S, McGuine TA. Effect of a New Rule Limiting Full  
383 Contact Practice on the Incidence of Sport-Related Concussion in High School Football  
384 Players. *Am J Sports Med.* 2019;47(10):2294-9.

385  
386  
387

## 388 **FIGURE LEGENDS**

### 389 **Figure 1: Cumulative head impact kinematics between LOC throughout a season.**

390 Cumulative (A) head impact count, (B) peak linear acceleration, and (C) peak rotational  
391 acceleration per player was influenced by LOC in an incremental manner, with *live* being the  
392 highest and *air* being the lowest.

393

394 **Figure 2: Group-dependent cumulative head impact exposure between LOC.** All 3 groups  
395 shared similar incremental pattern in cumulative (A) head impact count, (B) peak linear  
396 acceleration, and (C) peak rotational acceleration per player, with *live* being the highest and *air*  
397 being the lowest. The linemen and hybrid groups exhibited consistently higher head impact  
398 exposure compared to the skill group.

399  
400 **Figure 3: Average head impact magnitude per impact between LOC and group.** Live  
401 showed nearly times as high head impact magnitude (A. peak linear acceleration; B. peak  
402 rotational acceleration) as 5 other LOC.




403  
404 **Figure 4: Frequency of head impacts within various magnitude range.** A total number of  
405 head impacts from the overall sample throughout a season was categorized into peak linear  
406 acceleration ranging (A) 20 – 60 g, (B) 60-100 g, and (C) > 100 g. Live and thud consistently  
407 showed frequency head impacts in high impact magnitudes.




408  
409  
410  
411  
412  
413  
414  
415  
416


<b>Table 1. Group demographics and head impact kinematics</b>	
<b>Variables</b>	<b>Football</b>
n	24
Sex (%)	24M (100)
Age, y	15.7 ± 1.04
BMI, kg/m <sup>2</sup>	27.3 ± 6.29
No. of previous concussion	
0, n (%)	16 (66.7)
1, n (%)	6 (25.0)
2, n (%)	2 (8.3)
Tackle football experience, y	4.92 ± 2.7
Race, n (%)	
White	21 (88)
Black/African American	0 (0)
Asian	0 (0)
African Indiana/Alaska	1 (4)
Multiracial	2 (8)
Ethnicity, n (%)	
Not Latino/Hispanic	20 (83)
Latino/Hispanic	4 (17)
Playing position, n (%)	
Linemen (OL, DL)	11 (46)
Hybrid (TE, LB, RB)	7 (29)
Skill (WR, DB)	6 (25)
Psychiatric condition	
ADHD	0 (0)
Learning Disability	0 (0)
Major Depressive Disorder	0 (0)
Impact Kinematics for season, median (IQR)	
Impact count	203 (118 - 350)
Peak linear acceleration, g	4310.5 (2,686.3 - 8,616.8)
Peak rotational acceleration, krad/s <sup>2</sup>	415.5 (179.2 - 828.0)
Note: BMI, body mass index. IQR, interquartile range. OL, offensive lineman. DL, defensive lineman. TE, tight end. LB, linebacker. RB, running back. WR, wide receiver. DB, defensive back.	

417

418

**A** Cumulative Impacts / Player**B** Cumulative PLA / Player**C** Cumulative PRA / Player

**A Cumulative Impacts / player by Group****B Cumulative PLA / Player by Group****C Cumulative PRA / Player by Group**

**A Average PLA per Head Impact****B Average PRA per Head Impact**