A territory-wide study of early COVID-19 outbreak in Hong Kong community: A clinical, epidemiological and phylogenomic investigation

Kenneth Siu-Sing LEUNG, Timothy Ting-Leung NG, Alan Ka-Lun WU, Miranda Chong-Yee YAU, Hiu-Yin LAO, Ming-Pan CHOI, Kingsley King-Gee TAM, Lam-Kwong LEE, Barry Kin-Chung WONG, Alex Yat-Man HO, Kam-Tong YIP, Kwok-Cheung LUNG, Raymond Wai-To LIU, Eugene Yuk-Keung TSO, Wai-Shing LEUNG, Man-Chun CHAN, Yuk-Yung NG, Kit-Man SIN, Kitty Sau-Chun FUNG, Sandy Ka-Yee CHAU, Wing-Kin TO, Tak-Lun QUE, David Ho-Keung SHUM, Shea Ping YIP, Wing Cheong YAM, Gilman Kit-Hang SIU[1]

The Hong Kong Polytechnic University, Hong Kong Special Administrative Region, China (T.NG, H-Y.LAO, M-P.CHOI, L-K.LEE, D.SHUM, SP. YIP, G.SIU)

The University of Hong Kong, Hong Kong Special Administrative Region, China (K.LEUNG, K.TAM, W-C.YAM)

Pamela Youde Nethersole Eastern Hospital, Hong Kong Special Administrative Region, China (A.WU, M.YAU, K-C.LUNG)

Ruttonjee Hospital, Hong Kong Special Administrative Region, China (R.LIU)

United Christian Hospital, Hong Kong Special Administrative Region, China (B.WONG, E.TSO, K.FUNG, S.CHAU)

Princess Margaret Hospital, Hong Kong Special Administrative Region, China (A.HO, W-S.LEUNG, M-C.CHAN, W-K.TO)

Tuen Mun Hospital, Hong Kong Special Administrative Region, China (K-T.YIP, Y-Y.NG, K-M.SIN, T-L.QUE)

[1] **Correspondence to: Dr. Gilman KH SIU, Department of Health Technology and Informatics, Hong Kong Polytechnic University, Hong Kong (gilman.siu@polyu.edu.hk)**

Article summary lines: A combined epidemiological and phylogenetic analysis of early COVID-19 outbreak in Hong Kong revealed that a SARS-CoV-2 variant with *ORF3a* G251V mutation accounted for all locally acquired cases, and that asymptomatic carriers could be a huge public health risk for COVID-19 control.

35 **Running title:** Early community outbreak of COVID-19 in Hong Kong

37 **Keywords:** Novel Coronavirus, COVID-19, SARS-CoV-2, community outbreak, whole

38 genome sequencing, clinical demographic, molecular phylogeny, molecular evolutionary

39 analysis

Abstract (148 words)

Initial cases of COVID-19 reported in Hong Kong were mostly imported from China. However, most cases reported in February 2020 were locally-acquired infections, indicating local community transmissions. We extracted the demographic, clinical and epidemiological data from 50 COVID-19 patients, who accounted for 53.8% of the cases in Hong Kong by February 2020. Whole-genome sequencing of the SARS-CoV-2 were conducted to determine the phylogenetic relatedness and transmission dynamics. Only three (6.0%) patients required ICU admission. Phylogenetic analysis identified six transmission clusters. All locally-acquired cases harboured a common mutation *Orf3a* G251V and were clustered in two subclades in global phylogeny of SARS-CoV-2. The estimated time to the most recent common ancestor of local COVID-2019 outbreak was December 24, 2019 with an evolutionary rate of 3.04×10^{-3} substitutions per site per year. The reproduction number value was 1.84. Social distancing and vigilant epidemiological control are crucial to the containment of COVID-19 transmission.

(3452 words)

Introduction

"Coronavirus disease 2019(COVID-19)" refers to a cluster of viral pneumonia cases initially

from Wuhan, Hubei Province, China since December 2019. The aetiology was unknown

during the early stage of the outbreak until Chinese scientists isolated a novel coronavirus,

SARS-CoV-2, on January 7, 2020 and performed genome sequencing(1).

Fever was the main symptom of COVID-19 with about one-third of the patients presenting

acute respiratory distress syndrome(ARDS). About 16% of the patients were in severe

condition on admission, and the estimated mortality rate was 1.4%(2). Sustained human-to-

human transmission was confirmed upon identification of cases clustering among families

and transmission from patients to healthcare workers(3, 4), which triggered China's urgent

public health actions and international concern.

As of February 28, 2020, 78,824 patients have been diagnosed to have COVID-19 in

Mainland China with 2,788 deaths. The disease also spread to 50 other countries (5). The

World Health Organization(WHO) declared COVID-19 pandemic in March 2020. The first

imported case in Hong Kong was on January 23, 2020, who was a Mainland China resident

travelled to Hong Kong from Wuhan via Shenzhen by High-Speed Rail. The first local case

with unknown source, i.e. patient who had no travel record during the 14-day incubation

period, was reported on February 4, 2020(6).

By February 28, 2020, there were 93 COVID-19 cases in Hong Kong with at least 70(75·3%)

cases being local cases and their close contacts(6, 7). Secondary and tertiary transmissions

were observed in some case clusters. As most index cases of these clusters have unknown

sources, hidden transmission chain was believed to have established in the community.

Here, we report the demographic, clinical and epidemiological data of 50 hospitalised

patients, who accounted for 53.8% of COVID-19 cases in Hong Kong at the data cut-off

point (February 28, 2020), including three imported cases and six transmission clusters of

local infections. With the combined use of Nanopore sequencing and Illumina sequencing,

viral genomes of all these cases were characterised. Phylogenetic and molecular evolutionary

analyses were performed to determine the transmission linkage and the evolutionary rate of

COVID-19 cases in the community.

Methods

Cases

For this retrospective, multi-centre study, we intentionally enrolled the positive cases of

COVID-19 that were laboratory-confirmed at four public hospital clusters managed under the

Hospital Authority of Hong Kong, namely Hong Kong East Cluster, Kowloon East Cluster,

Kowloon West Cluster, and New Territories West Cluster, from January 26 to February 28,

2020.

Sputum specimens and throat swabs pooled with nasopharyngeal aspirates were collected

from patients who fulfilled the reporting or enhanced surveillance criteria on admission(8).

Laboratory-confirmed infection was defined when SARS-CoV-2 was detected by real-time

reverse transcription polymerase chain reactions(RT-PCRs), which amplified the envelope(*E*)

gene and RNA-dependent RNA polymerase(*RdRp*) gene(9).

Demographic, clinical, and microbiological data were obtained from patients' medical

records. Epidemiological information was retrieved from the Centre for Health

Protection(CHP) of the Department of Health(6) and the website wars.vote4.hk - Coronavirus

in HK(7). The definitions of clinical symptoms and complications are based on the WHO

guidance(10). We adopted the case numbering system of CHP, which was based on the date

of case confirmation.

This study was approved by the Institutional Review Boards of The Hong Kong Polytechnic

University(RSA20021) and the public hospitals involved(HKECREC-20200014;KCC/KEC-

20200070;KWC-20200040;NTWC-20200038).

Specimen preparation

The respiratory specimens were centrifuged at 16,000g for 2 minutes. Total nucleic acid was

extracted from supernatant using MagNA Pure 96 System(Roche Diagnostics, Germany) or

NucliSENS® easyMAG®(bioMérieux, The Netherlands) according to manufacturers'

instructions. DNase treatment was done by TURBO DNA-free Kit(ThermoFisher Scientific,

USA) to remove residual host DNA.

Reverse Transcription and viral genome amplification using multiplex PCR

DNase-treated RNA was reverse-transcribed using random hexamers and SuperScript IV

reverse transcriptase(Invitrogen, USA) as previously described(11). Viral cDNA was then

amplified using two PCRs containing tiled, multiplexed primers(table S1) described in the

ARTIC network(12). Details of the multiplex PCR are provided in the Supplemental Material.

Nanopore MinION Sequencing

Ligation-based 1D sequencing(SQK-LSK109, ONT, UK) was carried out according to

manufacturer's instructions. Briefly, multiplex PCR amplicons of each sample were

normalised to 1 ng/µL prior to end-repairing and native barcode ligation(EXP-NBD104/114,

ONT, UK). Barcoded samples were pooled and ligated to AMII sequencing adaptor.

Sequencing was performed with Nanopore MinION device using R9.4.1 flow cell for 48

hours.

Illumina MiSeq sequencing

Multiplex PCR amplicons were subjected to library preparation and dual-indexing using

KAPA HyperPrep Kit and Unique Dual-Indexed Adapter Kit(Roche Applied Science, US)

according to manufacturer's instructions. Ligated libraries were enriched by 6-cycle PCR

amplification followed by purification and size selection using AMPure XP beads(Beckman

Coulter, USA). The pooled library was sequenced with MiSeq Reagent Kit V2 Nano on

Illumina MiSeq System.

Bioinformatic analysis

Nanopore sequencing data were analysed using modified Artic Network nCoV-2019 novel

coronavirus bioinformatics protocol(Supplemental Material)(13).

Illumina sequencing reads were mapped with reference to respective draft genome of each

sample constructed from Nanopore data. Variants were called using freebayes(v1.0.0) with

haploid decoding and minimum base quality set at Q30. Consensus genomes were

constructed by GATK(v4.1.4.1) based on the VCF file(14). SPAdes genome

assembler(v3.14.0) and minimap2(v2.17) were used to combine Nanopore and Illumina

sequencing results for *de novo* assembly and identify the sequence of the unmapped gap

regions. The sequences have been submitted to the GenBank with accession no. MT232662-

MT232711.

Genomic and Phylogenetic analysis

To identify the amino acid change caused by each single-nucleotide polymorphism(SNP), consensus genome of each specimen was BLASTed against the reference NC_045512.2 using BlastX. Non-synonymous mutations were identified using custom Python script.

Phylogenetic tree was constructed based on the consensus genomes with PhyML(v3.0) using maximum likelihood algorithm. Best-fitting substitution model was selected by Akaike information criteria, in which general time reversible model with fixed proportion of invariable sites(+I) was selected(15). Bootstrap replicates was set at 1000×, and maximum-likelihood phylogenetic tree was rooted on the earliest published genome (accession no.: NC_045512.2). Transmission cluster was defined by clear epidemiological and onset-time relationship. Meanwhile, an additional 273 SARS-CoV-2 genomes were downloaded from GISAID Severe acute respiratory syndrome coronavirus 2 data hub(16). Together with the genomes from this study, phylogenetic tree was constructed using fast likelihood-based aLRT SH-like method and rooted on SARS-CoV-2 genome NC_045512.2.

Estimation of evolutionary rate and divergence time of transmission

To reconstruct the evolutionary model of COVID-19 cases in Hong Kong, Bayesian interference through Markov Chain Monte Carlo(MCMC) framework was implemented in BEAST(v2.6.2)(17). Bayesian phylodynamic analysis was performed using both strict clock and relaxed clock model with coalescent exponential growth tree priors. MCMC chains were run for 1×10^9 generations and sampled every 500 steps. Bayesian output was analysed after the results were visualised by Tracer(v1.7.1)(18). All parameters had an effective sample size of >200, indicating sufficient sampling.

Results

Fifty COVID-19 patients were included with 54·0% being female and the mean age was 55·2

years (table 1). Three were considered as imported cases as the patients stayed in Wuhan

before travelling to Hong Kong in mid-January. Four travelled to other regions where

COVID-19 active community transmission was not confirmed by that time. These cases were

considered as possibly local infection. The other 43 were defined as local infection with no

recent travel history. Eighteen(36·0%) had chronic illnesses, of which cardiovascular and

cerebrovascular diseases were the most common conditions(table 1). In total, 74·0% of the

patients presented with cough. Fever was presented in 58·0% of the patients on admission,

but gradually developed in 64·0% patients during hospitalisation. Other less common

symptoms include muscle-ache(25·0%), sore throat(24·0%), shortness of breath(24·0%), and

diarrhoea(14·3%)(table 2). Two(4·0%) patients were asymptomatic throughout the study

period. Regarding radiological examination, 27(54·0%) showed bilateral pneumonia,

11(22·0%) unilateral pneumonia, and 17(34·7%) multiple mottling and ground-glass opacity.

None of the patients was co-infected with other respiratory viruses or fungi. Two patients

were culture-positive for *Klebsiella aerogenes* and *Ralstonia pickettii* in their sputum

specimens. Both presented with ARDS and acute respiratory injury accompanied with septic

shock or acute renal injury, and required ICU admission.

Of the 50 cases, 42(84·0%) could be clustered based on their epidemiological linkages(figure

1). Six transmission clusters(Clusters 1–6) were identified. Cluster 1 involved a family of

four members. The father, who travelled to Guangdong, China in late January 2020, was

believed to infect his wife and subsequently their daughter and son-in-law in a family

gathering. Clusters 2 and 3 were family clusters of local infection with unknown source. Both

clusters involved three household members without recent travel history. Cluster 4 was a

super-spreading event(SSE) associated with a barbecue and hotpot party of 19 family

members in late January. Ten had developed symptoms two days after the party. A colleague

of one infected member, who was not present at the party, was also diagnosed to have

COVID-19. Cluster 5 initiated from a resident of a public estate, who was diagnosed on

January 30. Eleven days later, three members of a household living in the same building as,

but 10-storey below from, the index case were also infected. Two household members

attended a family gathering of 29 people at a Chinese seafood restaurant during their

incubation period. Three were diagnosed consecutively around two weeks after the gathering.

Additionally, a Filipino domestic helper of one infected member, who was absent in the

family gathering, was also infected. The earliest reported case of Cluster 6 was a woman who

visited a Buddhist worship hall during Chinese New Year holidays. Later, eight individuals

who visited the same Buddhist worship hall during this period were diagnosed. At the data

cut-off point, at least four other household members who had never been to the worship hall

were also tested positive for SARS-CoV-2.

Consensus genomes of all 50 cases were constructed based on Nanopore sequencing and

refined by Illumina sequencing. On average, 62,387 and 18,747 reads were obtained per

genome with 550× and 132× coverage for Nanopore and Illumina platforms respectively. The

consensus genome size was ~29.9 kbp with GC content ~38%. The genomes were highly

conserved with the first SARS-CoV-2 genome with an average sequence identity of

99·98%(range: 99·94%–100·0%). A total of 64 nonsynonymous substitutions were identified

from all 50 genomes(table 3). G251V in *Orf3a* is the most frequent amino acid substitution

with 44/50(88·0%) of the samples harbouring this mutation, followed by *Orf1ab*

H3233Y(30/50; 60·0%) and *S* gene L8V(27/50; 54·0%).

Genomic-wide SNPs were used to contextualise phylogenetic placement of Hong Kong

strains in SARS-CoV-2global phylogeny. Compared to SARS-CoV-2 strains isolated from

other regions, Hong Kong strains showed limited genomic variability, and tended to

aggregate in two clusters(figure 2). In the first cluster, a total of 13 Hong Kong strains formed

a subclade(yellow box) with the strains from France(n=5), Singapore(n=4), Australia(n=2),

South Korea(n=2), Sweden(n=1) and USA(n=1). The second cluster composed of 31 Hong

Kong strains, which formed another distinct subclade(pink box) with the strains from

Japan(n=3), China(n=2), England(n=2), Australia(n=2) and New Zealand(n=1).

Regarding the phylogenetic relationship among the Hong Kong cases, clustering of samples

was consistent with the epidemiological linkage(figure 3). Four distinctive nodes were

identified. The first node belonged to two imported cases and the cases in Cluster 1 of which

the genomes had very close genetic distance(0–2 SNPs) with the reference genome. The

second node represented cases in Cluster 5, of which four patients lived in different

apartments of the same building in a public housing estate, and were believed to have

transmitted the virus to other members in a subsequent family gathering. The third node was

associated with the cases in Cluster 4 which believed to originate from a family hotpot

gathering. Notably, the genomes of Cluster 4 were highly similar to those in Cluster 3 and

two singleton cases(i.e. Case 23 and Case 43), which shared the same missense mutations at

L8V in the *S* gene, H3233Y in *Orf1ab* and G251V in *Orf3a*. Finally, the fourth node

belonged to the SSE occurring in Buddhist worship hall(Cluster 6), in which *Orf1ab* G295V

and *Orf3a* G251V were identified in all the strains in the cluster.

According to Bayesian time-scaled phylodynamic analysis, estimated tMRCA of COVID-

2019 outbreak in Hong Kong was December 24, 2019(95% Bayesian credible interval [BCI]:

December 11, 2019 to January 5, 2020) with an evolutionary rate 3.04×10^{-3} substitutions per

site per year(95% BCI: 2.04×10^{-3} to 4.09×10^{-3} substitutions per site per year). Based on our

demographic data, time interval from symptom onset to hospital admission was ~8·5 days.

The estimated reproduction number was 1.84(95% BCI: 1.37 to 2.35).

Discussion

This study provides a territory-wide overview of early COVID-19 outbreak in Hong Kong, an

international city with borders connecting to Mainland China, by integrating the demographic,

clinical, epidemiological, phylogenomic, and phylodynamic data.

Fever was reported to be the most common symptom since 81·8%–98·0% of the patients had

fever on admission during the initial outbreak in China(19-22). However, a large-scale study

by Guan *et al* found that fever was only presented in 43·8% of the patients on admission(2).

In this study, fever was identified in only 58·0% of the patients on admission, but developed

in 64.0% after hospitalisation. Afebrile patients might be missed if the surveillance case

definition relied solely on fever. Therefore, laboratory surveillance has been extended to in-

patients and out-patients with respiratory symptoms. ICU admission was relatively

uncommon(6·0%) in our cohort when compared with previous studies(19-21). Two ICU

patients presented with severe complications including ARDS, acute renal damage, and septic

shock. Of note, both patients had bacterial co-infection, which was absent in other cases. In

addition to virulence factors of the pathogens, the host immune status, old age, and presence

of chronic illness might be associated with enhanced disease severity. Immune supportive

treatment and prompt antibiotics administration might reduce complications and mortality.

In Hong Kong, initial cases recorded in January 2020 were mostly imported cases. Since

February 1, majority were local cases and their close contacts, indicating local community

transmissions. Transmission in closed settings especially during family and religious

gatherings is a hallmark of recent cases recorded in Hong Kong. Among six clusters

identified based on epidemiological linkage, three(Clusters 4–6; figure 1) were considered

SSEs with more individuals involved(n=8–13). WGS was performed on all 50 cases to

investigate their phylogenetic relatedness and the transmission linkage.

The SARS-CoV-2 samples in Hong Kong have 99·98% identity to the reference

genome(NC_045512.2), indicating that no major genome modification has occurred since the

initial COVID-19 outbreak in Wuhan. Forty-four(88·0%) of our strains shared a common

mutation *Orf3a* G251V. Except one imported case, all belonged to local cases and their close

contacts without travel records(figure 1). Unlike other regions, such as China and Europe,

where the SARS-CoV-2 genomes scattered across different branches in global phylogeny, the

majority(88·0%) of Hong Kong strains were clustered in two subclades(figure 2). This

suggested that the COVID-19 outbreak in Hong Kong community was mostly arisen from

two ancestors.

Regarding the local phylogenetic analysis, clustering of samples was highly concordant to the

epidemiological linkage. Cluster 1 demonstrated the closest genetic distance to the reference

genome amongst all cases reported in Hong Kong(figures 1 and 3). The index case of Cluster

1(Case 66) was previously defined as possibly local infection as the patient travelled to

Guangdong Province, which was not considered to have active community transmission by

that time. However, our sequencing result showed that the genome of Case 66 was 100%

identical to the Wuhan reference genome, and all cases in Cluster 1 did not harbour *Orf3a*

G251V, which was recognized as a hallmark of the local cases with unknown source in our

community. Therefore, instead of possibly local infections, Cluster 1 was more likely

imported from Mainland China via index case 66.

Cluster 5 originated from a public estate, in which a family of three members(Cases 42, 48,

and 49) was speculated to get infected from a confirmed case(Case 12) who lived in the same

building, but 10-storey above, through a faulty sewage pipe setup. Based on phylogenetic

analysis, viral genomes in Cluster 5 shared a similar genetic distance from the reference

genome and were assigned to the same branch of the tree. This supports a potential

transmission linkage among these cases.

Cluster 4 was a family gathering-associated SSE during Chinese New Year. In concert with

epidemiological information, all 11 cases from Cluster 4 shared three common missense

mutation, namely *S* gene L8V, *Orf1ab* H3233Y and *Orf3a* G251V, with seven sharing

identical genomes. Considering the fast-evolving property of RNA viruses, identical genetic

sequences among the strains implied that transmission occurred in a short period or even in a

single event. Meanwhile, two singleton cases(Case 23 and Case 43) as well as cases from

another local cluster(Case 38, Case 39, and Case 40) shared highly similar genomes to those

of Cluster 4(figures 1 and 3). While no apparent epidemiological linkage were observed, the

high degree of genome similarity suggests that these cases might be originated from a single

source. The speculation was further supported by their geographical distribution of residences

who lived in close proximity to each other, and might share overlapping living circles(figure

4).

Cluster 6 was an SSE occurring in a Buddhist worship hall. Two missense mutations G295V

and L3606F in *Orf1ab* were unique to this cluster. Epidemiological investigation identified a

monk(Case 102; pink box of Cluster 6, figure 1), who was the abbot of the worship hall and

had travelled to Mainland China in early January. He was sent to a quarantine centre in late

February after being found associated with a series of confirmed cases connected with the

worship hall. He was completely asymptomatic throughout the study period. Phylogenetic

analysis showed that this case was closest to the root of the cluster(top portion with strains

shown in red, figure 3), suggesting that Case 102 could be the index patient of the cluster. By

the time of data cut-off, the cluster has already involved 13 patients and the spread was still

ongoing. Here we demonstrated the possibility of "hidden spreader" as a source of COVID-

19 community outbreak. It also highlights the importance of rapid quarantine of the close contacts of confirmed cases regardless of the presence of signs and symptoms in order to halt the COVID-19 community spread.

In the evolutionary clock study, death rate δ(which refers to the duration for the case to become non-contagious) was determined as the lag time between symptom onset date and the hospital admission date(i.e. 8·5 days, equivalent to 366/8.5=45·18/year). While δ was normally calculated based on recovery date(23), admission date was considered for calculation because the transmission link in Hong Kong was practically stopped once the patient was hospitalised. Assuming δ=45·18/year in our calculation, reproduction number within Hong Kong up to February 28, 2020 was estimated at 1.84(95% BCI: 1·37 to 2·35). The value strongly indicated that the outbreak in Hong Kong is still ongoing, but was smaller than the estimated reproduction number of 2·6 in Wuhan(24). The smaller value is a combined outcome of reduced growth rate and increased δ. The former one is attributed to very strong public health awareness among Hong Kong people with greatly reduced social activities and always put on surgical masks during this period,(25, 26) whereas the latter is the result of robust laboratory surveillance and fast quarantine time. In addition, tMRCA for the cases in Hong Kong was determined on December 24, 2019, which is ~25 days since the first symptom onset case(Case 2–the case with the earliest symptom onset, January 18, 2020; figure 1) of our study cohort.

There are several limitations in our study. Despite 53·8% of the cases reported in Hong Kong up to February 28 were included, another 43 cases including two fatal cases were not analysed in this study. Moreover, incubation period of cases with unknown source might vary widely. Studies have shown that incubation period can vary from 4.5-15.8 days,(24) and can be even longer for patients presenting with mild symptoms. However, as the patient might already be infectious during incubation period, resultant reproductive number in this study could still be underestimated. Moreover, current calculation was solely based on

phylodynamic analysis which could be different from those based on epidemiological models.

Finally, gap regions were observed in some consensus genomes. This is mainly because

WGS was performed on respiratory specimens instead of viral culture. The paucity of viral

load in specimens could affect the yield of sequencing libraries. Nevertheless, the uncovered

area only accounted for approximately 1-3% of the entire viral genome while the remaining

mapped regions had an average coverage of >100X, which should provide sufficient and

accurate information for our subsequent analyses.

In conclusion, phylogenomic data were consistent with the epidemiological findings that

transmission in closed settings especially during family and religious gatherings is a hallmark

of COVID-19 outbreak in Hong Kong. Social distancing and vigilant infection control

measures, such as rapid isolation of suspected or confirmed cases and their close contacts, are

crucial to the containment of COVID-19 transmission in the community.

Acknowledgments

This study was supported by Faculty of Health and Social Science and Department of Health Technology and Informatics of The Hong Kong Polytechnic University. We appreciate Oxford Nanopore Technologies Limited, especially Hai WANG and Eva YU, for their supportive service in the delivery of sequencing flow cells and reagents, and the provision of technical advices.

Author Bio

Dr. Kenneth Leung obtained his PhD in the Department of Microbiology, the University of Hong Kong in 2019. He received bioinformatic and phylogenetic analysis training under HKU Pasteur institute. His research focused on molecular diagnosis and epidemiology of emerging infectious disease such as *Mycobacterium tuberculosis* and HIV.

Contributors

KSSL and GKHS designed the study, collected, analysed and interpreted the data, conducted literature search, and drafted and critically reviewed the manuscript; and GKHS was also responsible for securing funding for this study. TTLN, HYL, MPC, KKGT, and LKL conducted experiments and analysed the data. AKLW, MCYY, BKCW, AYMH, KTY, KCL, RWTL, EYKT, WSL, MCC, YYN, KMS, KSCF, SKYC, WKT, and TLQ were responsible for collecting and analysing the data, and finalizing the manuscript. DHKS and SPY analysed the data, and wrote and critically reviewed the manuscript. WCY were involved in data analysis, data interpretation and writing of the manuscript. All authors reviewed and approved the final version of the manuscript.

Declaration of interests

We declare no competing interests.

References

1.   Zhu N, Zhang D, Wang W, Li X, Yang B, Song J, et al. A Novel Coronavirus from
Patients with Pneumonia in China, 2019. N Engl J Med. 2020 Feb 20;382(8):727-33.
2.   Guan WJ, Ni ZY, Hu Y, Liang WH, Ou CQ, He JX, et al. Clinical Characteristics of
Coronavirus Disease 2019 in China. N Engl J Med. 2020 Feb 28.
3.   Chan JF, Yuan S, Kok KH, To KK, Chu H, Yang J, et al. A familial cluster of
pneumonia associated with the 2019 novel coronavirus indicating person-to-person
transmission: a study of a family cluster. Lancet. 2020 Feb 15;395(10223):514-23.
4.   Li Q, Guan X, Wu P, Wang X, Zhou L, Tong Y, et al. Early Transmission Dynamics
in Wuhan, China, of Novel Coronavirus-Infected Pneumonia. N Engl J Med. 2020 Jan 29.
5.   Dong E, Du H, Gardner L. An interactive web-based dashboard to track COVID-19 in
real time. Lancet Infect Dis. 2020 Feb 19.
6.   Centre_for_Health_Protection. Latest local situation of Severe Respiratory Disease
associated with a Novel Infectious Agent. 2020 [cited 2020 24 February]; Available from:
https://www.chp.gov.hk/files/pdf/enhanced_sur_pneumonia_wuhan_eng.pdf
7.   Leung YF, Wong HB, Shing Y, Wong HW, Wong WK. wars.vote4.hk - Coronavirus
in HK. 2020 [cited 2020 25 February]; Available from: https://wars.vote4.hk/en/
8.   Centre_for_Health_Protection. Severe respiratory disease associated with a novel
infectious agent—letters to doctors. 2020 [cited 2020 6 February]; Available from:
https://www.chinadaily.com.cn/a/202001/25/WS5e2bb1b6a31012821727333a.html
9.   Corman VM, Landt O, Kaiser M, Molenkamp R, Meijer A, Chu DK, et al. Detection
of 2019 novel coronavirus (2019-nCoV) by real-time RT-PCR. Euro Surveill. 2020 Jan;25(3).
10.   WHO. Clinical management of severe acute respiratory infection when Novel
coronavirus (nCoV) infection is suspected: interim guidance. 2020 [cited 2020 11 Jan];
Available from: https://www.who.int/internal-publications-detail/clinical-management-of-
severe-acute-respiratory-infection-when-novel-coronavirus-(ncov)-infection-is-suspected
11.   Peiris JS, Lai ST, Poon LL, Guan Y, Yam LY, Lim W, et al. Coronavirus as a
possible cause of severe acute respiratory syndrome. Lancet. 2003 Apr 19;361(9366):1319-25.
12.   Quick J. Artic Network-nCoV 2019 sequencing protocol. 2020 [cited 2020 24
February]; Available from: https://artic.network/ncov-2019
13.   Loman N, Rambaut A. nCoV-2019 novel coronavirus bioinformatics protocol.
ARTIC-nCoV-bioinformaticsSOP-v100 2020 [cited 2020 10 March]; Available from:
https://artic.network/ncov-2019/ncov2019-bioinformatics-sop.html
14.   DePristo MA, Banks E, Poplin R, Garimella KV, Maguire JR, Hartl C, et al. A
framework for variation discovery and genotyping using next-generation DNA sequencing
data. Nat Genet. 2011 May;43(5):491-8.
15.   Guindon S, Dufayard JF, Lefort V, Anisimova M, Hordijk W, Gascuel O. New
algorithms and methods to estimate maximum-likelihood phylogenies: assessing the
performance of PhyML 3.0. Syst Biol. 2010 May;59(3):307-21.
16.   Elbe S, Buckland-Merrett G. Data, disease and diplomacy: GISAID's innovative
contribution to global health. Glob Chall. 2017 Jan;1(1):33-46.
17.   Bouckaert R, Heled J, Kuhnert D, Vaughan T, Wu CH, Xie D, et al. BEAST 2: a
software platform for Bayesian evolutionary analysis. PLoS Comput Biol. 2014
Apr;10(4):e1003537.
18.   Rambaut A, Drummond AJ, Xie D, Baele G, Suchard MA. Posterior Summarization
in Bayesian Phylogenetics Using Tracer 1.7. Syst Biol. 2018 Sep 1;67(5):901-4.
19.   Chen N, Zhou M, Dong X, Qu J, Gong F, Han Y, et al. Epidemiological and clinical
characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a
descriptive study. Lancet. 2020 Feb 15;395(10223):507-13.

20. Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y, et al. Clinical features of patients
infected with 2019 novel coronavirus in Wuhan, China. Lancet. 2020 Feb
15;395(10223):497-506.
21. Wang D, Hu B, Hu C, Zhu F, Liu X, Zhang J, et al. Clinical Characteristics of 138
Hospitalized Patients With 2019 Novel Coronavirus-Infected Pneumonia in Wuhan, China.
JAMA. 2020 Feb 7.
22. Wu J, Liu J, Zhao X, Liu C, Wang W, Wang D, et al. Clinical Characteristics of
Imported Cases of COVID-19 in Jiangsu Province: A Multicenter Descriptive Study. Clin
Infect Dis. 2020 Feb 29.
23. Boskova V, Stadler T, Magnus C. The influence of phylodynamic model
specifications on parameter estimates of the Zika virus epidemic. Virus Evol. 2018
Jan;4(1):vex044.
24. Lai A, Bergna A, Acciarri C, Galli M, Zehender G. Early phylogenetic estimate of the
effective reproduction number of SARS-CoV-2. J Med Virol. 2020 Feb 25.
25. Sung AD, Sung JAM, Thomas S, Hyslop T, Gasparetto C, Long G, et al. Universal
Mask Usage for Reduction of Respiratory Viral Infections After Stem Cell Transplant: A
Prospective Trial. Clin Infect Dis. 2016 Oct 15;63(8):999-1006.
26. Seto WH, Tsang D, Yung RW, Ching TY, Ng TK, Ho M, et al. Effectiveness of
precautions against droplets and contact in prevention of nosocomial transmission of severe
acute respiratory syndrome (SARS). Lancet. 2003 May 3;361(9368):1519-20.

Table 1: Demographics, travel record and baseline medical history of 50 patients
recruited in this study

	Patients (n=50)
Age (yo)	
Mean (SD)	55·2 (19·5)
Range	22–96
≤30	8 (16·0%)
31-40	5 (10·0%)
41-50	6 (12·0%)
51-60	11 (22·0%)
61-70	10 (20·0%)
≥71	10 (20·0%)
Gender	
Female	27 (54·0%)
Male	23 (46·0%)
Travel record within 14 days before symptom onset	7 (14·0%)
Wuhan, Hubei Province, China	3 (6·0%)
Other regions in Mainland China	1 (2·0%)
Regions outside Mainland China	3 (6·0%)
No travel record	43 (86·0%)
Chronic medical illnesses	18 (36·0%)
Cardiovascular and cerebrovascular diseases	14 (28·0%)
Endocrine system diseases	11 (22·0%)
Nervous system diseases	5 (10·0%)
Digestive system diseases	4 (8·0%)
Malignant tumour	1 (2·0%)

    Table 2: Clinical characteristics and outcomes of 50 patients recruited in this study

	Patients (n=50)
Signs and symptoms	
Fever (on admission)	29 (58·0%)
Fever (during hospitalisation)	32 (64·0%)
Cough	37 (74·0%)
Sore throat	12 (24·0%)
Shortness of breath	12 (24·0%)
Muscle ache	12 (25·0%)¶
Diarrhoea	7 (14·3%)#
Rhinorrhoea	4 (8·0%)
Nausea and vomiting	4 (8·2%)#
Confusion	1 (2·0%)
More than one sign or symptom	41 (82·0%)
Asymptomatic	2 (4·0%)
Complications*	
Acute respiratory distress syndrome	2 (4·0%)
Acute respiratory injury	1 (2·0%)
Acute renal injury	5 (10·0%)
Septic shock	1 (2·0%)
More than complications	2 (4·0%)
No complications	45 (90·0%)
Radiological findings	
Unilateral pneumonia	11 (22·0%)
Bilateral pneumonia	27 (54·0%)
Multiple mottling and ground-glass opacity	17 (34·7%)#
No abnormality	4 (8·0%)
Co-infection	
Other viruses	0
Bacteria	2 (4·0%)**
Fungi	0
Clinical outcome†	
In serious condition (Admission to ICU)	3 (6·0%)
Hospitalized; in stable condition	27 (54·0%)
Discharged	20 (40·0%)
Interval from symptom onset to hospital admission (Days)‡	
Average (SD)	8·5 (3·9)
Range	1–26
Length of hospital stay (Days)§	
Average (SD)	17·7 (7·7)
Range	8–35

* The definitions of the complications are provided in Supplementary materials.

† The data cut-off of the clinical outcome of the patients is Feb 28, 2020.

‡ The data from symptomatic patients were excluded.

§ This is calculated based on the 18 patients who had been discharged as of Feb 28, 2020.

¶ Data were missing for two patients.

Data were missing for one patient
** Moderate growth of *Klebsiella aerogenes* and *Ralstonia pickettii* were obtained from sputum specimens collected from case no. 38 and 70 respectively.

Table 3: Amino acid substitutions identified in SARS-CoV-2 isolated from the 50 patients in this study

Amino acid substitutions*	All cases (n=50)	Cases and Transmission Clusters of COVID-19 in the Hong Kong Community						Local Singletons† (n=5)	Imported Cases‡ (n=3)
		Cluster 1 (n=4)	Cluster 2 (n=3)	Cluster 3 (n=3)	Cluster 4 (n=11)	Cluster 5 (n=8)	Cluster 6 (n=13)		
Orf1ab									
G295V	13 (26·0%)	-	-	-	-	-	13 (100·0%)	-	-
P380L	1 (2·0%)	-	-	-	-	1 (12·5%)	-	-	-
E444K	1 (2·0%)	-	-	-	-	-	-	1 (20·0%)	-
A599D	1 (2·0%)	-	-	-	-	1 (12·5%)	-	-	-
V682A	1 (2·0%)	-	1 (33·3%)	-	-	-	-	-	-
A702T	1 (2·0%)	-	-	-	-	1 (12·5%)	-	-	-
Q1001E	1 (2·0%)	-	-	-	-	-	1 (7·7%)	-	-
D1323A	1 (2·0%)	-	-	-	-	1 (12·5%)	-	-	-
V1399E	1 (2·0%)	-	-	-	-	1 (12·5%)	-	-	
T1471I	1 (2·0%)	-	-	-	-	-	1 (7·7%)	-	-
V1483I	1 (2·0%)	-	-	-	-	-	-	1 (20·0%)	-
T1864M	1 (2·0%)	-	-	-	-	1 (12·5%)	-	-	-
Y1868N	1 (2·0%)	-	-	-	-	-	1 (7·7%)	-	-
A1983V	1 (2·0%)	-	-	-	-	-	1 (7·7%)	-	-
F2215S	1 (2·0%)	-	-	-	-	1 (12·5%S)	-	-	-
W2232C	1 (2·0%)	-	-	-	-	1 (12·5%C)	-	-	-
H2799Q	2 (4·0%)	-	-	-	-	1 (12·5%)	1 (7·7%)	-	-
D2833G	1 (2·0%)	-	-	-	-	-	1 (7·7%)	-	-
R2875G	2 (4·0%)	-	-	-	-	-	2 (15·4%)	-	-
S2972P	1 (2·0%)	-	-	-	-	1 (12·5%)	-	-	-
L3116F	1 (2·0%)	-	-	-	-	-	1 (7.7%)	-	-
M3131V	1 (2·0%)	-	-	-	-	-	1 (7·7%)	-	-
H3233Y	30 (60·0%)	-	-	3 (100·0%)	11 (100·0%)	-	12 (92·3%)	4 (80·0%)	-
I3257L	1 (2·0%)	-	-	-	-	-	1 (7·7%)	-	-
Y3500F	1 (2·0%)	-	1 (33·3%)	-	-	-	-	-	-
F3604L	1 (2·0%)	-	-	-	-	-	1 (7·7%)	-	-

L3606F	10 (20·0%)	-	-	-	-	-	10 (76·9%)	-	-
L3715S	1 (2·0%)	-	-	-	-	-	-	-	1 (33·3%)
W4124C	1 (2·0%)	-	-	-	-	-	1 (7·7%)	-	-
T4164I	1 (2·0%)	-	-	-	1 (9·1%)	-	-	-	-
A4247V	1 (2·0%)	-	-	-	-	-	1 (7·7%)	-	-
I4352L	1 (2·0%)	-	-	-	-	-	1 (7·7%)	-	-
Y4379F	2 (4·0%)	-	1 (33·3%)	-	-	-	1 (7·7%)	-	-
D4695Y	1 (2·0%)	-	-	-	-	-	1 (7·7%)	-	-
D4729V	1 (2·0%)	-	-	-	-	1 (12·5%)	-	-	-
G4895V	1 (2·0%)	-	-	-	-	1 (12·5%)	-	-	-
S5360P	1 (2·0%)	-	-	-	-	-	1 (7·7%)	-	-
V5384G	1 (2·0%)	-	-	-	-	-	1 (7·7%)	-	-
I5445V	1 (2·0%)	-	-	-	-	-	1 (7·7%)	-	-
I5547V	1 (2·0%)	-	-	-	-	-	1 (7·7%)	-	-
K6275N	1 (2·0%)	-	-	-	-	-	1 (7·7%)	-	-
P6318S	1 (2·0%)	-	-	-	-	-	1 (7·7%)	-	-
D7091E	2 (4·0%)	-	-	-	-	-	-	1 (20·0%)	1 (33·3%)
S									
L8V	27 (54·0%)	-	-	3 (100·0%)	11 (100·0%)	-	9 (69·2%)	4 (80·0%)	-
D138Y	3 (6·0%)	-	-	-	3 (27·3%)	-	-	-	-
L249F	1 (2·0%)	1 (25·0%)	-	-	-	-	-	-	-
K310E	1 (2·0%)	-	-	-	-	-	-	1 (20·0%)	-
V367F	1 (2·0%)	-	-	-	-	-	-	-	1 (33·3%)
K458N	1 (2·0%)	1 (25·0%)	-	-	-	-	-	-	-
R509K	1 (2·0%)	-	-	-	-	1 (12·5%)	-	-	-
V510L	1 (2·0%)	-	-	-	-	1 (12·5%)	-	-	-
P715H	1 (2·0%)	-	-	-	-	-	1 (7·7%)	-	-
D820N	1 (2·0%)	-	-	-	-	-	1 (7·7%)	-	-
P1263L	1 (2·0%)	-	-	-	-	-	1 (7·7%)	-	-
Orf3a									
G251V	44 (88·0%)	-	3 (100·0%)	3 (100·0%)	11 (100·0%)	8 (100·0%)	13 (100·0%)	5 (100·0%)	1 (33·3%)
M269T	1 (2·0%)	-	-	-	-	-	-	-	1 (33·3%)

M									
D209Y	1 (2·0%)	-	-	-	-	-	1 (7·7%)	-	-
Orf8									
V62L	1 (2·0%)	-	-	-	-	-	-	-	1 (33·3%)
L84S	1 (2·0%)	-	-	-	-	-	-	-	1 (33·3%)
N									
T24N	1 (2·0%)	-	-	-	-	-	1 (7·7%)	-	-
R226S	1 (2·0%)	-	-	1 (33·3%)	-	-	-	-	-
G316E	1 (2·0%)	-	-	-	-	-	-	1 (20·0%)	-
M322I	1 (2·0%)	-	-	-	-	1 (12·5%)	-	-	-
K363M	1 (2·0%)	-	-	-	-	-	1 (7·7%)	-	-

"-" indicates 0 (0·0%).

* The location of amino acid substitutions was based on SARS-CoV-2 reference genome (NC_045512.2).

† The singleton cases did not have recent travel history.

‡ The patients had record of travel to Wuhan City.

Figure Legend

Figure 1. Demographics of patients included in this study
A rectangular box stands for male and eclipse for female. Travel history within the 14-day incubation period prior to symptom onset is highlighted in 1) Cyan for local case without travel history; 2) Pink for travel history to Wuhan within 14-days from symptom onset; 3) Orange for travel history to other regions in Mainland China, and 4) Green for travel history to regions outside Mainland China. The case numbers are those used by the Centre of Health Protection, Department of Health, Hong Kong.[6] Within each cluster, the cases of primary infection (leftmost column) are arranged from top to bottom in the order of the date (dd/m) of symptom onset.

Figure 2. Maximum-likelihood phylogenetic tree constructed based on 50 SARS-CoV-2 genomes in this study and genomes collected from GISAID data hub
A total of 273 worldwide genomes as of February 28, 2020 were randomly selected and downloaded from GISAID Global Cases COVID-19 database. The phylogenetic tree was built with the samples collected in this study based on fast likelihood-based aLRT SH-like method and rooted on the earliest published genome of SARS-CoV-2 (accession no.: NC_045512.2). The Hong Kong strains showed limited genetic variability and tended to aggregate in two subclades, which were highlighted in yellow box and pink box respectively.

Figure 3. Maximum-likelihood phylogenetic tree of the 50 COVID-19 cases in this study
The tree was constructed using all 50 COVID-19 cases included in this study and rooted on the earliest published genome of SARS-CoV-2 (accession no.: NC_045512.2). Bootstrap value was set at 1000×. Samples were colour-coded by epidemiological link as follows: 1) Magenta represents Cluster 1: Imported cluster; 2) Cyan represents Cluster 2: unknown source; 3) Green represents Cluster 3: local; 4) Orange represents Cluster 4: Hotpot party-related Superspreading event (SSE); 5) Blue represents Cluster 5: Public housing estate-related SSE; and 6) Red represents Cluster 6: Buddha worship hall-related SSE. Case 84 and Case 102 were asymptomatic at the time of sample collection, and were marked with asterisks (*) in the diagram. Each case is indicated with a case number (see figure 1) followed by the date (dd/mm/yyyy) of symptom onset.

Figure 4. Geographic distribution of residences of singleton case #23, case #43, and cases in Cluster 3 and 4

Hong Kong

China
South Korea
Japan
Singapore
Thailand
India
Vietnam
Taiwan

England
France
Belgium
Germany
Italy
Switzerland
Sweden
Finland

USA
Canada
Brazil
Maxico

Australia

New Zealand

Cruise

NC045512

0.00050

Singleton cases
CLUSTER 1: Imported cluster
CLUSTER 2: local cluster
CLUSTER 3: local cluster
CLUSTER 4: Hotpot party-related SSE
CLUSTER 5: Public housing estate-related SSE
CLUSTER 6: Buddha Worship hall-related SSE

0.00010

Singleton cases
CLUSTER 3: local cluster
CLUSTER 4: Hotpot party-related SSE
(★: Hotpot party site)

Hotpot party site

Patient #38
Patient #39
Patient #40

Patient #23

Patient #43

Patient #44

Patient #33
Patient #36

Patient #27
Patient #29
Patient #30
Patient #31

Patient #32
Patient #34
Patient #35
Patient #41

5 km