

1                   **Japanese citizens’ behavioral changes and**  
2                   **preparedness against COVID-19:**  
3                   **How effective is Japan’s approach of self-restraint ?**

4  
5                   **Short title: Japanese citizens’ behavioral changes and preparedness against COVID-19**

6  
7                   Kaori Muto, PhD<sup>†\*1</sup>, Isamu Yamamoto, PhD<sup>†2</sup>, Miwako Nagasu, PhD<sup>3</sup>, Mikihiro Tanaka, PhD,<sup>4</sup> Koji

8                   Wada, MD, MSc, PhD<sup>5</sup>

9  
10                  <sup>1</sup> The Institute of Medical Science, The University of Tokyo, Tokyo, Japan

11                  <sup>2</sup> Faculty of Business and Commerce, Keio University, Tokyo, Japan

12                  <sup>3</sup> Faculty of Economics, Keio University, Tokyo, Japan

13                  <sup>4</sup> Faculty of Political Science and Economics, Waseda University, Tokyo, Japan

14                  <sup>5</sup> Graduate School of Medicine and Public Health, International University of Health and Welfare,

15                  Tokyo, Japan

16  
17                  <sup>†</sup> These authors contributed equally.

18                  \* Corresponding author email: [krmt@ims.u-tokyo.ac.jp](mailto:krmt@ims.u-tokyo.ac.jp)

19

## 20 **Abstract**

21

22 The Japanese government instituted countermeasures against COVID-19, a pneumonia caused  
23 by the new coronavirus, in January 2020. Seeking “people’s behavioral changes,” in which the  
24 government called on the public to take precautionary measures or exercise self-restraint, was  
25 one of the important strategies. The purpose of this study is to investigate how and from when  
26 Japanese citizens have changed their precautionary behavior under these circumstances, where  
27 the government has only requested their cooperation. This study uses micro data from a  
28 cross-sectional survey conducted on an online platform of an online research company, based on  
29 quota sampling that is representative of the Japanese population. By the end of March 2020, we  
30 had recruited a total of 11,342 respondents, aged from 20 to 64 years. About 85% reported  
31 practising the social distancing recommended by the government. More females than males and  
32 more older than younger participants are supportive of practicing social distancing. Frequent  
33 handwashing is conducted by 86 percent of all, 92 percent of female and 87.9 percent of  
34 over-40 participants. The most important event influencing these precautionary actions was the  
35 infection aboard the Diamond Princess cruise ship, which occurred in early February 2020  
36 (23%). Information from the central and local governments, received by 60% of the participants,  
37 was deemed trustworthy by 50%. However, the results also showed that about 20% of the  
38 participants were reluctant to implement proper prevention measures. The statistical analysis

39 indicated that the typical characteristics of those people were male, younger (under 30 years  
40 old), unmarried, from lower-income households, with a drinking or smoking habit and a higher  
41 extraversion score. To prevent the spread of infection in Japan, it is imperative to address these  
42 individuals and encourage their behavioral changes using various means to reach and influence  
43 them.

44

45

## 46 **Introduction**

47

## 48 **The new coronavirus in Japan**

49       Pneumonia of unknown cause was detected in China and first officially reported on 31  
50       December 2019. The World Health Organization (WHO) announced a name for the new  
51       coronavirus disease, COVID-19 (coronavirus disease 2019), on 11 February 2020.<sup>[1]</sup> Since then,  
52       COVID-19 has been spreading throughout the world, and a rapid increase in deaths has been  
53       reported in many countries. As of 28 March, a total of 571,678 cases and 26,494 deaths have  
54       been confirmed.<sup>[2]</sup> One study has estimated that there will be a total of 81,114 deaths from  
55       COVID-19 over the next four months in the US alone.<sup>[3]</sup> The number of COVID-19 cases and  
56       deaths in Japan is gradually increasing, with 1,499 cases (including 60 critical cases) and 49  
57       deaths reported as of 28 March.<sup>[4]</sup> Several small clusters of infected groups have been increasing  
58       in urban areas, including those in hospitals and nursing homes, in addition to cases with  
59       unlinked infections. Nevertheless, the total number of deaths and severely ill patients has been  
60       comparatively small, especially relative to the country's population size, and the trend of the  
61       increase is not sharp. The reasons for this mild trend have been questioned outside Japan.<sup>[5]</sup>

62       Over the past few decades, Japan has not experienced any serious damage from new  
63       infectious diseases, such as SARS (severe acute respiratory syndrome), MERS (Middle East  
64       respiratory syndrome) or the Ebola virus. Ironically, this history of escapes might delay the

65 establishment of the emergency operation headquarters in Japan. The implementation of  
66 polymerase chain reaction (PCR) tests, which must be the frontline response to the novel  
67 coronavirus outbreak, has faced time-consuming obstacles. In Japan, a recent revision of the Act  
68 on Special Measures for Pandemic Influenza and New Infectious Diseases Preparedness and  
69 Response allows the Prime Minister to declare a state of emergency for the outbreak, but under  
70 the current legislation, no central or local government can enforce lockdowns such as those  
71 undertaken in other countries. If the Act on the Prevention of Infectious Diseases and Medical  
72 Care for Patients with Infectious Diseases is revised for COVID-19, local governors can restrict  
73 or block the traffic in contaminated places for a specified period of not more than 72 hours for  
74 the purpose of preventing the spread of the coronavirus.

75 Under such limitations, the current goal of the Japanese government is to avoid an  
76 explosive increase in patients that would exceed the limit of intensive or critical care units in  
77 hospitals in urban areas. To meet this goal, the government policy consists of three strategies:  
78 early detection of clusters and rapid response, enhancement of the early diagnosis of patients  
79 and intensive care for severely affected patients, and strengthening of the universal healthcare  
80 system and people's behavioral change.<sup>[6]</sup>

81

## 82 **Three strategies against COVID-19**

83       With regards to the first and second strategies, the Ministry of Health, Labor and Welfare  
84 (MHLW) strongly promotes contact tracing, social distancing and pneumonia surveillance under  
85 the direction of the Patient Cluster Countermeasure Group in the MHLW Headquarters for  
86 Novel Coronavirus Disease Control. Regional public health centers conduct contact tracing,  
87 asking infected persons and their close contacts to maintain social distancing for 14 days and  
88 allocating available hospital beds or hospital wards in designated local communities. In clinical  
89 settings, the large number of computed tomography (CT) scanners in Japan (111.49 per million  
90 population<sup>[7]</sup>) supports physicians with investigating suspicious pneumonia cases in the absence  
91 of conducting massive PCR tests in the population. This policy approach might lead to a  
92 relatively slower increase in the number of cases and deaths.

93       Regarding the third strategy, people's behavioral change, by the middle of February 2020,  
94 the MHLW encouraged the Japanese public to practise frequent handwashing and “coughing  
95 etiquette” (using a handkerchief or sleeve instead of hands to catch a cough or sneeze).  
96 Furthermore, the MHLW has prioritized access to healthcare for elderly people, people suffering  
97 from fatigue or shortness of breath and people with underlying health conditions. The MHLW  
98 has also asked the public not to visit clinics for at least four days if they have experienced cold  
99 symptoms or a fever of 37.5°C or over.<sup>[8]</sup> This restriction might be a shock to Japanese citizens,

100 who are typically allowed to freely access clinics and hospitals.

101 In analyses of contact tracing, it was found that one infected person tended to infect more  
102 than one other person at locations with certain characteristics. On 24 February, the Expert  
103 Meeting on the Control of Novel Coronavirus Infection asked the public to refrain from  
104 attending places involving close face-to-face contact (within an arm's length of each other) in  
105 conversations and similar interactions for more than a given length of time in crowds. Since  
106 then but prior to other similar slogans that have appeared in the world, the government has been  
107 campaigning for avoidance of these situations with the slogan "Avoid overlapping three Ms"  
108 *mippei-kukan* [poorly ventilated closed space], *misshu-basho* [large gathering] and  
109 *missetsu-bamen* [conversations or shouting in close proximity], in addition to regular ventilation  
110 and wiping of shared surfaces (door handles, knobs, and bed fences) and goods with diluted  
111 household chlorine bleach. "Avoid overlapping three Ms" has been the core and unique message  
112 against COVID-19 in Japan.

113

## 114 **Previous studies and our research questions**

115 This study examines three research questions: 1) How do Japanese citizens implement the  
116 government's three Ms precautionary measures? 2) How effective are these requests from the  
117 government? and 3) Who has changed their daily precautionary behavior and who has not?

118           Several previous studies have investigated changes in people’s precautionary behavior  
119 against the coronavirus. For example, an online survey conducted on 29 January of 3,083  
120 mainland Chinese respondents revealed that adults living in urban areas had a better awareness  
121 of the issue than those in rural areas (72.7% vs. 66.1%,  $p<0.001$ ).<sup>[9]</sup> Another online survey  
122 conducted between 23 February and 2 March in the US (N=2,986) and the UK (N=2,988) adult  
123 residents showed that people have a good understanding of the main mode of disease  
124 transmission and common symptoms, although they also have several important misconceptions  
125 and discriminatory attitudes towards people with East Asian ethnicity due to COVID-19’s origin  
126 in China.<sup>[10]</sup> The latest study in Italy clarified the three types of attitude to COVID-19 among  
127 Italian citizens: people who trust authority and choose isolation, fatalists who are keen on social  
128 media and uniformed youth.<sup>[11]</sup> The Gallup International Association also recently conducted a  
129 snap poll in 28 countries (including 1,115 Japanese participants), asking about precautionary  
130 procedures, and their findings indicated that 71 percent of Japanese participants had adopted  
131 more frequent handwashing.<sup>[12]</sup> It is still unclear, however, what the trigger is for behavioral  
132 change around COVID-19 and who is more actively implementing the prevention measures. In  
133 this survey, the response period and sample attribution are also unclear. Furthermore, this survey  
134 is not necessarily informative for policymaking, as it does not reveal who is *not* implementing  
135 prevention measures.

136 Using a large sample of cross-sectional survey data, this study investigates how and at what  
137 point Japanese citizens changed their precautionary behavior in this situation, where the  
138 government has only requested, rather than mandated, their cooperation.

139

## 140 **Materials and Methods**

141

### 142 **Survey design and participants**

143 This study uses micro data from a cross-sectional survey conducted via an online  
144 platform of an online research company, MACROMILL INC, Japan. From a pool of  
145 approximately 1.2 million registered individuals residing in Japan, we recruited a total of 11,342  
146 males and females aged from 20 to 64 years. In the recruitment process for this study, quota  
147 sampling was conducted so that the sample distributions among gender (male or female), age  
148 group (20s, 30s, 40s, 50s or 60s), and employment status (regular employee, non-regular  
149 employee, self-employed or not working) became equal to those of the representative Japanese  
150 population, based on the statistics of the Labor Force Survey (Ministry of Internal Affairs and  
151 Communications). The survey was conducted between 26 and 28 March 2020.

152

### 153 **Questionnaire and analysis**

154 In addition to providing individual characteristics, the participants were asked to

155 answer 11 items rating their prevention measures against novel coronavirus infections, such as  
156 social distancing and coughing etiquette on a scale of 1 to 5. Thus, after summarizing  
157 demographic characteristics based on the total, male and female, and under-40 and over-40  
158 categories, we aggregate and compare a proportion of the participants who have been taking  
159 those prevention measures.

160           The participants were also asked what kind of events caused them to change their  
161 behaviors and rated the frequency and reliability of 10 information sources about the  
162 coronavirus on a scale of 1 to 5. Thus, we calculate and compare the frequency and reliability  
163 depending on the information sources.

164           Next, to detect factors associated with behavioral change, the participants were also  
165 asked about their drinking and smoking habits. Personality traits were measured by the Five  
166 Factor Personality Questionnaire: Ten-Item Personality Inventory (TIPI). We estimate a logit  
167 model, where the dependent variable is a dummy indicating 1 if the participant chose “not at all”  
168 or “not true” to the question “Do you avoid the three overlapping Ms?” and where independent  
169 variables are individual characteristics.

170

## 171 **Data analysis**

172           We analyzed the data using STATA/MP version 16.0 for Mac (StataCorp, College

173 Station, TX, United States).

174

## 175 **Ethical issues**

176 Our survey falls outside the scope of the Japanese government’s Ethical Guidelines for  
177 Medical and Health Research Involving Human Subjects, and there are no national guidelines in  
178 Japan for social and behavioral research. Therefore, our study was carried out in accordance  
179 with the Ethical Principles for Sociological Research of the Japan Sociological Society, which  
180 do not require ethical reviews.

181 All survey participants gave consent to participate in the anonymous online survey by  
182 MACROMILL INC. The authors did not obtain any personal information about the participants.  
183 After being informed about the purposes of the study and their right to quit the survey,  
184 participants agreed to participate. They were provided with the option of “I don’t want to  
185 respond” for all questions. Completion of the entire questionnaire was considered as participant  
186 consent.

187

## 188 **Results**

189

### 190 **Demographic characteristics**

191 The characteristics of the sample, both as a whole and separated by gender (male or

192 female) or age (under or over 40 years old), are summarized in Table 1. The total sample size is  
193 11,342, with almost equal gender distribution. Gender and age distribution are proportional to  
194 that of the Japanese population. University or college graduates constituted about 50–60 percent  
195 of respondents. About half of the total sample is composed of regular employees (usually,  
196 indefinite and full-time employees). About a quarter of respondents had a household income of  
197 4–5 million yen.

198

199

200

**Table 1. Sample characteristics**

	n, (%)									
	All		Male		Female		Under 40 y		Over 40 y	
<b>Total respondents</b>	11,342		5,734		5,608		4,300		7,842	
<b>Female</b>	5,608	(49.44)					2,110	(49.07)	3,498	(49.67)
<b>Age</b>										
20-29 years old	1,964	(17.32)	994	(17.34)	970	(17.30)	1,964	(45.67)	0	(0.00)
30-39 years old	2,336	(20.60)	1,196	(20.86)	1,140	(20.33)	2,336	(54.33)	0	(0.00)
40-49 years old	3,098	(27.31)	1,568	(27.35)	1,530	(27.28)	0	(0.00)	3,098	(43.99)
50-59 years old	2,754	(24.28)	1,373	(23.94)	1,381	(24.63)	0	(0.00)	2,754	(39.11)
60-64 years old	1,190	(10.49)	603	(10.52)	587	(10.47)	0	(0.00)	1,190	(16.90)
<b>Married</b>	6,620	(58.37)	3,241	(56.52)	3,379	(60.25)	1,859	(43.23)	4,761	(67.61)
<b>Parents with children under junior high school</b>	2,972	(26.20)	1,493	(26.04)	1,479	(26.37)	1,464	(34.05)	1,508	(21.41)
<b>University or college graduate</b>	6,278	(55.35)	3,415	(59.56)	2,863	(51.05)	2,478	(57.63)	3,800	(53.96)
<b>Work status</b>										
Regular employee	5,817	(51.29)	3,986	(69.52)	1,831	(32.65)	2,460	(57.21)	3,357	(47.67)
Nonregular employee	2,865	(25.26)	733	(12.78)	2,132	(38.02)	1,010	(23.49)	1,855	(26.34)
Self-employed and others	660	(5.82)	422	(7.36)	238	(4.24)	138	(3.21)	522	(7.41)
Not working	2,000	(17.63)	593	(10.34)	1,407	(25.09)	692	(16.09)	1,308	(18.57)
<b>Household annual income</b>										
Less than 2,000K JPY	646	(7.56)	312	(6.67)	334	(8.63)	250	(8.28)	396	(7.16)
2,000-3,999K JPY	1,939	(22.68)	925	(19.77)	1,014	(26.21)	773	(25.59)	1,166	(21.10)
4,000-5,999K JPY	2,247	(26.29)	1,224	(26.16)	1,023	(26.44)	869	(28.77)	1,378	(24.93)
6,000-6,999K JPY	1,606	(18.79)	929	(19.85)	677	(17.50)	541	(17.91)	1,065	(19.27)
8,000-8,999K JPY	1,052	(12.31)	619	(13.23)	433	(11.19)	322	(10.66)	730	(13.21)
10,000-11,999K JPY	516	(6.04)	318	(6.80)	198	(5.12)	139	(4.60)	377	(6.82)
12,000-14,999K JPY	306	(3.58)	192	(4.10)	114	(2.95)	58	(1.92)	248	(4.49)
15,000-19,999K JPY	151	(1.77)	100	(2.14)	51	(1.32)	42	(1.39)	109	(1.97)
More than 20,000K JPY	85	(0.99)	60	(1.28)	25	(0.65)	27	(0.89)	58	(1.05)

201

202

203

204

205 **To what extent have prevention measures been taken?**

206           In the survey, the participants were first asked to answer to the question “Have you  
207 ever conducted any measures to prevent novel coronavirus infections or outbreaks?” About 76  
208 percent of participants evaluated themselves as having taken some action.

209           To observe the details of those actions, Table 2 shows a variety of prevention measures  
210 taken, aggregating a proportion of the participants who answer “very true” and “true” for each  
211 prevention measure.

212

213

**Table 2. “Have you ever conducted anything to prevent novel coronavirus infections or outbreaks?”**

214

	% of very true and true, (C.I.)				
	All	Male	Female	Under 40 y	Over 40 y
<b>1. Avoid a poorly-ventilated closed space</b>	80.6 (79.8 - 81.3)	75.9 (74.8 - 77.0)	85.3 (84.4 - 86.3)	77.6 (76.3 - 78.8)	82.4 (81.5 - 83.3)
<b>2. Avoid large gatherings</b>	80.5 (79.8 - 81.2)	77.5 (76.4 - 78.6)	83.6 (82.6 - 84.6)	76.3 (75.1 - 77.6)	83.1 (82.2 - 83.9)
<b>3. Avoid conversations or shouting in close proximity</b>	57.0 (56.1 - 57.9)	55.6 (54.3 - 56.8)	58.5 (57.2 - 59.8)	52.3 (50.9 - 53.8)	59.9 (58.7 - 61.0)
<b>4. Avoid places where items 1-3 above overlap</b>	80.6 (79.8 - 81.3)	76.9 (75.8 - 77.9)	84.4 (83.4 - 85.3)	76.7 (75.4 - 77.9)	82.9 (82.1 - 83.8)
<b>5. Do not go to mass gatherings</b>	86.8 (86.2 - 87.5)	82.7 (81.8 - 83.7)	91.0 (90.3 - 91.8)	82.7 (81.6 - 83.8)	89.4 (88.6 - 90.1)
<b>6. Undertake frequent handwashing</b>	86.3 (85.7 - 87.0)	81.9 (80.9 - 82.9)	90.9 (90.2 - 91.7)	83.8 (82.7 - 84.9)	87.9 (87.1 - 88.6)
<b>7. Undertake cough etiquette (use handkerchiefs or sleeves instead of hands)</b>	77.0 (76.2 - 77.8)	72.0 (70.9 - 73.2)	82.1 (81.1 - 83.1)	73.9 (72.6 - 75.2)	78.9 (77.9 - 79.8)
<b>8. Always wear a surgical-style mask when going out</b>	70.1 (69.2 - 70.9)	62.7 (61.4 - 63.9)	77.6 (76.5 - 78.7)	70.0 (68.6 - 71.3)	70.1 (69.1 - 71.2)
<b>9. Avoid going out when you have a cold</b>	76.7 (76.0 - 77.5)	70.9 (69.7 - 72.1)	82.7 (81.7 - 83.7)	72.8 (71.4 - 74.1)	79.2 (78.2 - 80.1)
<b>10. Get sufficient rest and sleep</b>	73.1 (72.3 - 73.9)	68.3 (67.1 - 69.5)	77.9 (76.9 - 79.0)	71.6 (70.2 - 72.9)	74.0 (73.0 - 75.0)
<b>11. Eat a nutritious diet</b>	69.5 (68.6 - 70.3)	64.2 (62.9 - 65.4)	74.9 (73.7 - 76.0)	69.1 (67.7 - 70.5)	69.7 (68.6 - 70.8)
<b>12. Prepare consultation and transportation methods for when you feel ill</b>	41.5 (40.6 - 42.4)	42.7 (41.4 - 44.0)	40.3 (39.0 - 41.6)	41.9 (40.4 - 43.3)	41.3 (40.1 - 42.4)

215

216

217

218

Looking at the first four prevention measures, which have been continuously

219

requested by the Japanese government and the Expert Meeting on Control of Novel Coronavirus

220

Infection, it was found that 80 percent have attempted to avoid the “overlapping three Ms.” Of

221

the total, 57% have attempted avoid conversations or shouting in close proximity, which was a

222

relatively low figure among the three Ms. Looking next at the fifth prevention measure, more

223

than 85 percent of all participants reported practising social distancing by avoiding mass

224 gatherings. Regarding gender and age differences, more females than males and more older than  
225 younger participants are supportive of social distancing, as shown by the differences in the  
226 confidence intervals.

227         Regarding hygiene practices, frequent handwashing is conducted by about 86 percent  
228 of all, about 91 percent of female and about 88 percent of over-40 participants. Coughing  
229 etiquette was implemented by 77 percent of the participants. Many also answered that they have  
230 avoided going out when ill with a cold.

231         As for the measures to strengthen individual immunity, around 70 percent of the  
232 participants reported getting sufficient rest and sleep or eating a nutritious diet. Again, focusing  
233 on gender and age differences, prevention measures are conducted more often by females and  
234 older people.

235         However, regardless of gender and age, about 40 percent of participants have prepared  
236 consultation and transportation methods for if they were to become ill.

237

## 238 **What has caused the behavioral changes?**

239         To explore the triggers of the behavioral changes and preparedness observed above,  
240 the participants were asked “What was the most important event influencing these actions?”; the  
241 responses are summarized in Fig 1. The figure shows that about 23 percent of the participants

242 cited the infection aboard the *Diamond Princess* cruise ship <sup>[13]</sup> that occurred around early  
243 February 2020, when there were still few domestic cases. *Diamond Princess* is a  
244 British-registered cruise ship on which an 80-year-old passenger from Hong Kong tested  
245 positive for COVID-19 on 1 February 2020. Because the ship was in Japanese waters, the ship  
246 was quarantined in February 2020 for nearly a month with about 3,700 passengers and crew on  
247 board. Other participants noted events from the end of February, including the alert from the  
248 Expert Meeting (5.6 percent), the statement of emergency by the governor of Hokkaido  
249 (northern island of Japan) (7.4 percent) and the request by the Prime Minister to not attend mass  
250 gatherings (7.8 percent). The next large trigger was the request by the Prime Minister for  
251 nationwide school closures in Japan on 28 February 2020 (about 14 percent). Finally, worldwide  
252 outbreak around early March (22 percent) also attracted participants' attention.

253         To explore what kind of information affected their behavioral change and  
254 preparedness, the survey asked participants to report the frequency at which they consult certain  
255 sources about the novel coronavirus infection and rate their reliability of the information source.  
256 The results are summarized in Table 3.

257

258

259

**Table 3. “From where do you get and trust information about novel coronavirus infection?”**

260

		%, (C.I.)			
	All	Male	Female	Under 40 y	Over 40 y
<b>1. TV news programs</b>					
Get information: frequently or sometimes	89.0 (88.5 - 89.6)	85.4 (84.5 - 86.3)	92.7 (92.1 - 93.4)	85.2 (84.1 - 86.2)	91.4 (90.8 - 92.1)
Trust information: very much or yes	55.2 (54.3 - 56.1)	51.0 (49.7 - 52.3)	59.5 (58.3 - 60.8)	50.5 (49.0 - 52.0)	58.1 (56.9 - 59.2)
<b>2. TV talk and variety shows</b>					
Get information: frequently or sometimes	69.4 (68.6 - 70.3)	62.4 (61.2 - 63.7)	76.6 (75.5 - 77.7)	67.5 (66.1 - 68.9)	70.6 (69.5 - 71.7)
Trust information: very much or yes	31.4 (30.6 - 32.3)	27.2 (26.0 - 28.3)	35.8 (34.5 - 37.0)	32.5 (31.1 - 33.9)	30.8 (29.7 - 31.9)
<b>3. Newspapers (national and local newspapers)</b>					
Get information: frequently or sometimes	42.0 (41.1 - 42.9)	45.4 (44.1 - 46.7)	38.6 (37.3 - 39.9)	30.0 (28.6 - 31.4)	49.4 (48.2 - 50.5)
Trust information: very much or yes	47.5 (46.6 - 48.5)	45.2 (43.9 - 46.4)	50.0 (48.7 - 51.3)	41.0 (39.6 - 42.5)	51.5 (50.3 - 52.7)
<b>4. Tabloid paper</b>					
Get information: frequently or sometimes	7.9 (7.4 - 8.4)	11.4 (10.5 - 12.2)	4.3 (3.7 - 4.8)	9.6 (8.7 - 10.5)	6.8 (6.2 - 7.4)
Trust information: very much or yes	12.7 (12.1 - 13.3)	13.4 (12.6 - 14.3)	11.9 (11.1 - 12.8)	16.1 (15.0 - 17.2)	10.6 (9.9 - 11.4)
<b>5. Internet news sites</b>					
Get information: frequently or sometimes	86.6 (86.0 - 87.3)	84.7 (83.7 - 85.6)	88.7 (87.8 - 89.5)	84.4 (83.3 - 85.5)	88.0 (87.3 - 88.8)
Trust information: very much or yes	41.8 (40.9 - 42.7)	41.3 (40.0 - 42.5)	42.3 (41.0 - 43.6)	42.1 (40.6 - 43.5)	41.6 (40.5 - 42.8)
<b>6. SNS app news</b>					
Get information: frequently or sometimes	45.6 (44.7 - 46.5)	42.8 (41.5 - 44.1)	48.4 (47.1 - 49.8)	57.5 (56.0 - 58.9)	38.3 (37.2 - 39.5)
Trust information: very much or yes	24.5 (23.7 - 25.3)	23.4 (22.3 - 24.5)	25.7 (24.5 - 26.8)	29.9 (28.5 - 31.3)	21.2 (20.3 - 22.2)
<b>7. Information sent by the Prime Minister</b>					
Get information: frequently or sometimes	66.3 (65.4 - 67.2)	63.6 (62.3 - 64.8)	69.1 (67.9 - 70.3)	59.8 (58.3 - 61.3)	70.3 (69.2 - 71.4)
Trust information: very much or yes	47.5 (46.6 - 48.4)	46.0 (44.8 - 47.3)	48.9 (47.6 - 50.2)	45.7 (44.2 - 47.2)	48.6 (47.4 - 49.7)
<b>8. Information sent by the Ministry of Health, Labor and Welfare</b>					
Get information: frequently or sometimes	63.4 (62.5 - 64.3)	60.8 (59.5 - 62.1)	66.0 (64.8 - 67.2)	58.3 (56.8 - 59.7)	66.5 (65.4 - 67.6)
Trust information: very much or yes	48.8 (47.9 - 49.7)	46.9 (45.7 - 48.2)	50.7 (49.4 - 52.0)	47.6 (46.1 - 49.1)	49.5 (48.4 - 50.7)
<b>9. Information provided by government Expert Meetings</b>					
Get information: frequently or sometimes	56.9 (55.9 - 57.8)	55.2 (54.0 - 56.5)	58.5 (57.2 - 59.8)	50.3 (48.8 - 51.8)	60.8 (59.7 - 62.0)
Trust information: very much or yes	51.4 (50.5 - 52.3)	49.7 (48.4 - 51.0)	53.1 (51.8 - 54.4)	48.8 (47.3 - 50.3)	53.0 (51.8 - 54.1)
<b>10. Information sent by local (prefecture) government</b>					
Get information: frequently or sometimes	58.0 (57.1 - 58.9)	53.7 (52.4 - 55.0)	62.4 (61.1 - 63.6)	54.3 (52.8 - 55.8)	60.2 (59.1 - 61.4)
Trust information: very much or yes	55.6 (54.7 - 56.6)	52.5 (51.2 - 53.8)	58.8 (57.6 - 60.1)	54.0 (52.5 - 55.4)	56.7 (55.5 - 57.8)

261

262

263           Table 3 shows that almost 90 percent receive information from TV news programs and  
264 Internet news sites, and that about 50 percent trust such information. Mainstream scientists are  
265 annoyed about the fear-mongering that happens on TV talk and variety shows; indeed, these  
266 formats are slightly favored, but considered less credible, among the public. Meanwhile,  
267 information from the central and local government (received by 60 percent), including the  
268 Prime Minister and the Expert Meeting, is relatively trusted by the participants (50 percent).  
269 Among official sources, the local government is most trusted. Newspapers (national and local)  
270 are read by only about 42 percent of the participants, and about 48 percent answered that they  
271 trust information from newspapers.

272           Looking at the differences in gender and age, females tend to seek more information  
273 and trust it more than males, except for the information from newspapers. Participants over 40  
274 years old tend to access and trust the information from TV, newspapers and officials more than  
275 those under 40 years old do, while young people often seek and trust news from the Internet and  
276 SNS apps.

277

## 278 **Who does not adhere to social distancing?**

279           As we confirmed in Table 2, more than 80 percent of the participants have been

280 implementing social distancing measures and most Japanese citizens seem to be exhibiting  
281 some behavioral change to prevent coronavirus infections. However, this also means that about  
282 20 percent may not be conducting sufficient prevention measures.

283           To detect what kind of individuals are included in the group not conducting prevention  
284 measure, Table 4 shows the estimation results of the logit model. Like the other tables, Table 4  
285 shows the results based on the total, male and female, and under-40 and over-40 categories. The  
286 number shown in the table is an odds ratio, so the estimates that are significantly higher than 1  
287 indicate a higher tendency to not conduct proper social distancing.

288

289

**Table 4. Estimation results of logit model for not conducting social distancing**

Dependent variable: Avoid places where items 1-3 above overlap in Table 2 -> Not at all or not tue		odds ratio, (C.I.)			
	All	Male	Female	Under 40 y	Over 40 y
<b>Male</b>	1.635*** (1.304 - 2.049)			1.431** (1.039 - 1.970)	1.854*** (1.335 - 2.575)
<b>Age (ref. = 40-49 y)</b>					
20-29 years old	1.671*** (1.274 - 2.192)	1.819*** (1.285 - 2.574)	1.647** (1.053 - 2.574)	1.830*** (1.354 - 2.474)	
30-39 years old	0.910 (0.683 - 1.213)	0.916 (0.639 - 1.314)	0.949 (0.589 - 1.529)		
50-59 years old	0.819 (0.610 - 1.099)	0.915 (0.632 - 1.325)	0.689 (0.423 - 1.123)		0.858 (0.634 - 1.163)
60-64 years old	0.437*** (0.264 - 0.724)	0.631 (0.357 - 1.117)	0.112*** (0.026 - 0.471)		0.448*** (0.265 - 0.758)
<b>Not married</b>	1.445*** (1.117 - 1.870)	1.372* (0.985 - 1.912)	1.514* (0.979 - 2.340)	1.476* (0.954 - 2.282)	1.422** (1.020 - 1.984)
<b>Not having children younger than junior high school age</b>	1.246 (0.935 - 1.661)	1.114 (0.774 - 1.603)	1.485 (0.922 - 2.392)	1.366 (0.868 - 2.151)	1.165 (0.792 - 1.713)
<b>High or junior high school graduate</b>	1.171 (0.957 - 1.434)	1.174 (0.914 - 1.508)	1.152 (0.813 - 1.633)	1.347* (0.999 - 1.815)	1.053 (0.798 - 1.390)
<b>Work status (Ref. = Regular employee)</b>					
Nonregular employee	1.035 (0.795 - 1.349)	0.992 (0.670 - 1.468)	1.119 (0.763 - 1.642)	0.952 (0.654 - 1.387)	1.151 (0.783 - 1.692)
Self-employee and others	0.727 (0.446 - 1.185)	0.958 (0.569 - 1.613)	0.164* (0.022 - 1.197)	0.658 (0.233 - 1.860)	0.745 (0.425 - 1.307)
Not working	0.767 (0.552 - 1.066)	1.089 (0.707 - 1.678)	0.596* (0.343 - 1.035)	0.745 (0.463 - 1.201)	0.807 (0.509 - 1.279)
<b>Household annual income (Ref. = 4,000-5,999K JPY)</b>					
Less than 2,000K JPY	1.441* (0.990 - 2.099)	1.103 (0.667 - 1.824)	1.760* (0.970 - 3.195)	1.883** (1.115 - 3.180)	1.110 (0.638 - 1.931)
2,000-3,999K JPY	1.159 (0.879 - 1.530)	1.064 (0.752 - 1.505)	1.284 (0.797 - 2.068)	1.480* (0.991 - 2.211)	0.936 (0.632 - 1.385)
6,000-6,999K JPY	1.182 (0.880 - 1.589)	1.143 (0.803 - 1.625)	1.234 (0.713 - 2.137)	1.430 (0.908 - 2.254)	1.002 (0.678 - 1.481)
8,000-8,999K JPY	0.984 (0.687 - 1.411)	0.893 (0.577 - 1.383)	1.189 (0.629 - 2.249)	1.119 (0.633 - 1.977)	0.877 (0.550 - 1.401)
10,000-11,999K JPY	0.894 (0.544 - 1.467)	0.919 (0.523 - 1.614)	0.737 (0.253 - 2.144)	1.212 (0.571 - 2.575)	0.694 (0.357 - 1.350)
12,000-14,999K JPY	0.730 (0.362 - 1.471)	0.519 (0.205 - 1.313)	1.366 (0.463 - 4.029)	1.748 (0.653 - 4.677)	0.384* (0.137 - 1.079)
15,000-19,999K JPY	0.460 (0.143 - 1.482)	0.365 (0.087 - 1.525)	0.740 (0.097 - 5.656)	0.415 (0.055 - 3.124)	0.456 (0.108 - 1.922)
More than 20,000K JPY	0.707 (0.216 - 2.314)	0.530 (0.125 - 2.241)	2.026 (0.246 - 16.664)	0.434 (0.056 - 3.340)	0.967 (0.227 - 4.125)
<b>Drinking habit: Drink 3-6 times per week or everyday</b>	1.150 (0.911 - 1.451)	1.278* (0.976 - 1.675)	0.773 (0.476 - 1.256)	1.326 (0.909 - 1.935)	1.053 (0.784 - 1.413)
<b>Smoking habit: Smoke sometimes or everyday</b>	1.077 (0.856 - 1.355)	0.956 (0.729 - 1.253)	1.529* (0.995 - 2.350)	1.124 (0.791 - 1.597)	1.043 (0.767 - 1.419)
<b>Big 5 Personality traits</b>					
Extraversion	1.113*** (1.032 - 1.200)	1.108** (1.008 - 1.219)	1.125* (0.992 - 1.275)	1.151** (1.032 - 1.284)	1.087 (0.978 - 1.208)
Neuroticism	0.986 (0.909 - 1.069)	0.977 (0.880 - 1.085)	1.005 (0.884 - 1.143)	0.988 (0.880 - 1.109)	0.987 (0.880 - 1.107)
Openness	0.976 (0.900 - 1.059)	0.952 (0.861 - 1.054)	1.014 (0.884 - 1.164)	0.917 (0.815 - 1.031)	1.030 (0.919 - 1.154)
Conscientiousness	0.865*** (0.798 - 0.937)	0.865*** (0.782 - 0.957)	0.875* (0.765 - 1.002)	0.909 (0.810 - 1.020)	0.826*** (0.738 - 0.925)
Agreeableness	0.884*** (0.814 - 0.959)	0.906* (0.818 - 1.004)	0.851** (0.742 - 0.976)	0.896* (0.799 - 1.005)	0.874** (0.776 - 0.985)
<b>Constant</b>	0.075*** (0.037 - 0.152)	0.138*** (0.059 - 0.322)	0.055*** (0.017 - 0.182)	0.045*** (0.016 - 0.123)	0.100*** (0.038 - 0.266)
<b>Number of observations</b>	8,548	4,679	3,869	3,021	5,527

291 Note: \*\*\* p<0.01, \*\* p<0.05, \* p<0.1

292            Looking at the estimation results in Table 4, males, people in their 20s and unmarried  
293 people exhibit significantly higher odds ratios, indicating that these groups tend not to conduct  
294 preventive social distancing. Although work status is not generally associated with this  
295 prevention measure, females, regular employees and non-regular employees tended to exhibit  
296 higher odds ratios than self-employed or unemployed people.

297            Regarding household annual income, the lowest group (less than 2,000K JPY) has  
298 significantly higher odds ratio for the total, female and under-40 categories.

299            Higher odds ratios for not conducting social distancing are associated with drinking  
300 for males and smoking for females. Furthermore, those with higher extraversion scores also  
301 tend to exhibit significantly higher odds ratio in many cases, while conscientiousness and  
302 agreeableness are associated with lower odds ratio in most cases.

303

## 304 **Should the government change its policy on mass gatherings?**

305            Before this survey was conducted, the request by the Japanese government for  
306 self-restraint in avoiding mass gatherings had become an issue. For example, on 22 March 2020,  
307 the K-1 Grand Prix, a martial arts event, was held despite the Minister's and local governor's pleas  
308 for restraint, and 6,500 participants were packed into the Saitama Super Arena. On 23 March, more  
309 than 50,000 gathered in Sendai to see the Olympic flame, which had recently arrived from

310 Greece.<sup>[14]</sup> We asked the participants whether they supported this policy approach. About 29  
 311 percent of participants supported the idea that the government should now allow mass  
 312 gatherings. Males tend to support allowing mass gatherings more than females. On the other  
 313 hand, 65 percent supported government limitations of movement, in addition to self-restraint to  
 314 avoid mass gatherings in order to shorten the period of the pandemic. There are no significant  
 315 differences in gender and age category for this question.

316

317 **Table 5. “Do you support the government’s policy?”**

	% of agree and relatively agree, (C.I.)				
	All	Male	Female	Under 40 y	Over 40 y
1. The government should allow mass gatherings now	28.8 (27.9 - 29.6)	31.2 (30.0 - 32.4)	26.3 (25.1 - 27.5)	30.5 (29.1 - 31.8)	27.7 (26.7 - 28.8)
2. The government should limit movement in addition to mass gatherings	64.9 (64.0 - 65.7)	64.1 (62.9 - 65.4)	65.6 (64.4 - 66.9)	63.4 (62.0 - 64.8)	65.8 (64.7 - 66.9)

318

319

## 320 **Discussion**

321

322

323 Under circumstances where there is no enforced ban on mass gathering or travelling  
 324 beyond the home region, our findings indicate that a large portion of Japanese citizens seem to  
 325 be implementing proper prevention measures on their own.

326 First, more than three-quarters of the survey participants have taken some preventive

327 actions, including social distancing, handwashing, coughing etiquette and strengthening  
328 immunity. Because the previous empirical studies did not include developed countries like  
329 Japan,<sup>[15]</sup> there is little scientific evidence that Japanese people prefer cleanliness and tend to  
330 wash their hands relatively more frequently than other countries. In Japanese communities,  
331 water facilities for handwashing with soap and hand sanitizers are normally placed various  
332 public places, such as train stations and supermarkets. Moreover, handwashing became part of  
333 regular practice at home and school through post-war education.<sup>[16]</sup> In general, Japanese people  
334 have developed the discipline of washing their hands before having meals or after using the  
335 toilet. It is also well known that Japanese bow for greetings instead of handshaking, kissing and  
336 hugging. This cultural behavior implies that the frequency of body contact among Japanese  
337 people may be less than for those who greet with handshaking, kissing and hugging. During hay  
338 fever season, Japanese citizens regularly wear surgical-style masks for prevention; it may be  
339 that wearing a mask was a less popular preventive measure than some of the others in this study  
340 due to there being a shortage of these products. These already habitual practices might help  
341 behavioral changes develop among Japanese citizens during these unusual times.

342 In the survey, more than half of the participants had not prepared access to  
343 consultation centers or transportation methods for if they were to become ill, implying that they  
344 had not planned for the possibility of contracting COVID-19. We must advise the public to

345 prepare for such an event, to talk to family and close friends about unexpected advanced care  
346 planning, and to imagine not being able to use ventilators or extracorporeal membrane  
347 oxygenation at the severe stage.

348           Second, one of the main motivations for people's behavioral changes was the infection  
349 aboard the *Diamond Princess* cruise ship in early February 2020. At that time, only a few cases  
350 of domestic infection had been reported in Japan, but news of the quarantine and positive test  
351 results among the passengers was broadcasted daily. This may have contributed to Japanese  
352 citizens changing their mindset and behavior toward precautionary measures earlier than in  
353 Europe and the US. The sudden request by the Prime Minister for nationwide school closures in  
354 the end of February might also have been an effective measure for changing the mindsets of  
355 Japanese citizens toward prevention, even though this move was scientifically questioned and  
356 confusing to the public, especially to single parents and double-income households.

357           Third, information from the Expert Meeting and central/local governments, including  
358 the Prime Minister, are relatively trusted by survey participants. The Expert Meeting and  
359 central/local government have held frequent press conferences, trying to clarify the tentative  
360 scientific risks and advocating for citizens to conduct prevention measures. Such crisis  
361 communication attempts may have caused behavioral changes in Japanese citizens. The most  
362 trusted resource in this study was information from the local government, which was a hopeful

363 result, considering that the countermeasures against the virus are decided and conducted by the  
364 local government. It is also important to support residents with disadvantages to information  
365 and communication in the local communities.

366           However, the results also show that about 20 percent of the participants are reluctant  
367 to implement proper prevention measures. The statistical analysis indicates that typically those  
368 people are male, younger (under 30 years old), unmarried, are in lower income household, have  
369 a drinking or smoking habit and have a higher extraversion score. To prevent the spread of  
370 infection in Japan, it is imperative to address these individuals and encourage their behavioral  
371 change in various ways that will reach and move them. It is notable that approximately 65  
372 percent of the participants support stricter countermeasures, such as limitation of movement. As  
373 we mentioned in the introduction, as of yet, the government has not made stay-at-home orders  
374 mandatory and it has not offered financial aid to those affected by such measures. The current  
375 requests from central/local governments are not legally binding and people/businesses have to  
376 arrange financial compensation independent of the government. We should observe how  
377 effective these measures are in Japan over the long term.

378           There are several limitations to this study. First, the data were self-reported, and  
379 participants' actual behaviors have not been observed. Second, the sample was not collected  
380 based on random sampling. Quota sampling ensured a similar distribution to the Japanese

381 population among demographic groups (gender, age and work status), but the sample within  
382 each group does not necessarily reflect the population. Third, we obtained this dataset at the end  
383 of March 2020, when the infection is not explosively widespread in Japan. This study should be  
384 reanalyzed after the COVID-19 pandemic comes to an end.

385

## 386 **Acknowledgments**

387

388 We would like to thank the participants in our online survey for offering valuable data. This  
389 work was supported by university grants allocated to the Department of Public Policy, Human  
390 Genome Center, The Institute of Medical Sciences, The University of Tokyo. We also appreciate  
391 members of the COVID-PAGE (Public Advisory Group of Experts) for insightful discussions.

392

## 393 **References**

394

395 [1] World Health Organization. WHO characterizes COVID-19 as a pandemic. 11 March 2020.

396 Available from:

397 <https://www.who.int/emergencies/diseases/novel-coronavirus-2019/events-as-they-happen>

398 [2] World Health Organization. WHO situation report as of 29 March 2020. Available from:

399 <https://www.who.int/emergencies/diseases/novel-coronavirus-2019/situation-reports>

- 400 [3] IHME COVID-19 health service utilization forecasting team. Forecasting COVID-19 impact  
401 on hospital bed-days, ICU-days, ventilator days and deaths by US state in the next 4  
402 months. MedRxiv. 26 March 2020. Available from:  
403 [www.healthdata.org/research-article/forecasting-covid-19-impact-hospital-bed-days-icu-da](http://www.healthdata.org/research-article/forecasting-covid-19-impact-hospital-bed-days-icu-days-ventilator-days-and-deaths)  
404 [ys-ventilator-days-and-deaths](http://www.healthdata.org/research-article/forecasting-covid-19-impact-hospital-bed-days-icu-days-ventilator-days-and-deaths)
- 405 [4] Ministry of Health, Labor and Welfare. Coronavirus disease 2019 (COVID-19) situation  
406 within and outside the country as of 29 March 2020. Available from:  
407 [https://www.mhlw.go.jp/stf/seisakunitsuite/bunya/newpage\\_00032.html](https://www.mhlw.go.jp/stf/seisakunitsuite/bunya/newpage_00032.html)
- 408 [5] Masuike H. Japan's virus success has puzzled the world. Is its luck running out? The New  
409 York Times. 2020 March 26. Available from:  
410 [https://www.nytimes.com/2020/03/26/world/asia/japan-coronavirus.html?searchResultPosit](https://www.nytimes.com/2020/03/26/world/asia/japan-coronavirus.html?searchResultPosition=1)  
411 [ion=1](https://www.nytimes.com/2020/03/26/world/asia/japan-coronavirus.html?searchResultPosition=1)
- 412 [6] Expert Meeting on the Novel Coronavirus Disease Control. Views on the novel coronavirus  
413 disease control (in Japanese). 2020 March 9. Available from:  
414 <https://www.mhlw.go.jp/content/10900000/000606000.pdf>
- 415 [7] This number was registered as of 2017 at the OECD Health Statistics in 2019.  
416 [<https://stats.oecd.org/Index.aspx?ThemeTreeId=9>]
- 417 [8] Ministry of Health, Labor and Welfare. Prevention Measures against Coronavirus Disease

- 418 2019 (COVID-19). Available from:
- 419 <https://www.mhlw.go.jp/content/10900000/000607599.pdf>
- 420 [9] Zhan S, Yang YY, Fu C. Public's early response to the novel coronavirus-infected
- 421 pneumonia. *Emerg Microbes Infect.* 2020 Mar 3;9(1):534. Available from:
- 422 <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7067171/>
- 423 [10] Geldsetzer P. Knowledge and perceptions of COVID-19 among the general public in the
- 424 United States and the United Kingdom: a cross-sectional online survey. *Ann Intern Med.*
- 425 2020 Mar 20. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7086377/>
- 426 [11] Bucchi, M, Salacino B. Italian citizens and covid-19. 2020 March 21. In: Public
- 427 Understanding of Science Blog [Internet]. Available from:
- 428 <https://sagepus.blogspot.com/2020/03/italian-citizens-and-covid-19.html>
- 429 [12] Gallup International Association. 2020 March. The coronavirus: a vast scared majority
- 430 around the world. Available from:
- 431 [https://www.gallup-international.com/wp-content/uploads/2020/03/GIA\\_SnapPoll\\_2020\\_C](https://www.gallup-international.com/wp-content/uploads/2020/03/GIA_SnapPoll_2020_COVID_Tables_final.pdf)
- 432 [OVID\\_Tables\\_final.pdf](https://www.gallup-international.com/wp-content/uploads/2020/03/GIA_SnapPoll_2020_COVID_Tables_final.pdf)
- 433 [13] National Institute for Infectious Diseases. Field briefing: Diamond Princess COVID-19
- 434 cases. 2020 Feb. Available from:
- 435 <https://www.niid.go.jp/niid/en/2019-ncov-e/9407-covid-dp-fe-01.html>; and its update

436 available from: <https://www.niid.go.jp/niid/en/2019-ncov-e/9417-covid-dp-fe-02.html>.

437 [14] Denyer S. Japan's social distancing is shrinking as coronavirus fears ease. Too soon?

438 Washington Post. 2020 March 24. Available from:

439 [https://www.washingtonpost.com/world/asia\\_pacific/japans-social-distancing-is-shrinking-](https://www.washingtonpost.com/world/asia_pacific/japans-social-distancing-is-shrinking-)

440 [as-coronavirus-fears-ease-too-soon/2020/03/24/7c816cee-6d0c-11ea-a156-0048b62cdb51\\_](https://www.washingtonpost.com/world/asia_pacific/japans-social-distancing-is-shrinking-as-coronavirus-fears-ease-too-soon/2020/03/24/7c816cee-6d0c-11ea-a156-0048b62cdb51_)

441 [story.html](https://www.washingtonpost.com/world/asia_pacific/japans-social-distancing-is-shrinking-as-coronavirus-fears-ease-too-soon/2020/03/24/7c816cee-6d0c-11ea-a156-0048b62cdb51_story.html)

442 [15] Wolf J, Johnston R, Freeman MC, Ram PK, Slaymaker T, Laurenz E, Prüss-Ustün A.

443 Handwashing with soap after potential faecal contact: global, regional and country

444 estimates. *Int J Epidemiol.* 2019;48(4): 1204–18. Available from:

445 <https://doi.org/10.1093/ije/dyy253>

446 [16] Bay AR. Disciplining shit. *Japan Forum* 2018;30(4):556-582.

447 doi:10.1080/09555803.2019.1594341

448

**Fig 1. "What was the most important event influencing these actions?"**

(%)

