Estimating effects of intervention measures on COVID-19 outbreak in Wuhan taking account of improving diagnostic capabilities using a modelling approach

Jingbo Liang[1], Hsiang-Yu Yuan[1*], Lindsey Wu[2], Dirk U. Pfeiffer[3]

[1]Department of Biomedical Sciences, Jockey Club College of Veterinary Medicine and Life Sciences, City University of Hong Kong, Hong Kong

[2]Department of Infection Biology, Faculty of Infectious and Tropical Diseases, London School of Hygiene & Tropical Medicine, United Kingdom

[3]Centre for Applied One Health Research and Policy Advice, City University of Hong Kong, Hong Kong, China

[*]Correspondence to: Hsiang-Yu Yuan sean.yuan@cityu.edu.hk

Abstract

Background: Although by late February 2020 the COVID-19 epidemic was effectively controlled

in Wuhan, China, the virus has since spread around the world and been declared a pandemic on

March 11. Estimating the effects of interventions, such as transportation restrictions and quarantine

measures, on the early COVID-19 transmission dynamics in Wuhan is critical for guiding future

virus containment strategies. Since the exact number of COVID-19 infected cases is unknown, the

number of documented cases was used by many disease transmission models to infer

epidemiological parameters. However, this means that it would not be possible to adequately

estimate epidemiological parameters and the effects of intervention measures, because the

percentage of all infected cases that were documented changed during the first 2 months of the

epidemic as a consequence of a gradually increasing diagnostic capability.

Methods: To overcome the limitations, we constructed a stochastic susceptible-exposed-infected-

quarantined-recovered (SEIQR) model, accounting for intervention measures and temporal

changes in the proportion of new documented infections out of total new infections, to characterize

the transmission dynamics of COVID-19 in Wuhan across different stages of the outbreak. Pre-

symptomatic transmission was taken into account in our model, and all epidemiological parameters

were estimated using Particle Markov-chain Monte Carlo (PMCMC) method.

Results: Our model captured the local Wuhan epidemic pattern as a two-peak transmission

dynamics, with one peak on February 4 and the other on February 12, 2020. The impact of

intervention measures determined the timing of the first peak, leading to an 86% drop in the R_e

from 3.23 (95% CI, 2.22 to 4.20) to 0.45 (95% CI, 0.20 to 0.69). An improved diagnostic capability

led to the second peak and a higher proportion of documented infections. Our estimated proportion

of new documented infections out of the total new infections increased from 11% (95% CI 1% -

43%) to 28% (95% CI 4% - 62%) after January 26 when more detection kits were released. After

the introduction of a new diagnostic criterion (case definition) on February 12, a higher proportion

of daily infected cases were documented (49% (95% CI 7% - 79%)).

Introduction

Coronavirus disease 2019 (COVID-19), an acute respiratory infection originally identified in the

city of Wuhan in Hubei Province, China, has spread worldwide in 2020[1-2]. Estimating the effects

of intervention measures is still one of the major scientific goals in order to identify proper

prevention measures in the world[3]. The precise estimation of transmissibility R_e is critical for the

identification of appropriate intervention measures to contain the outbreak[1,4,5,6,7,8]. Although many

recent studies have evaluated how intervention measures implemented in Wuhan reduced disease

spreading to regions outside Wuhan[6,9,10,11,12], the investigation of the contribution of interventions

within Wuhan, the epidemic source region itself, has not been done much[13 14], possibly because

that an irregular pattern of transmission dynamics during early February hinders the model fitting

processes, making the precise estimation of the parameters difficult.

To control the virus spreading during the early outbreak stage, the Chinese government

implemented strict travel restrictions on January 23, 2020 in Wuhan [15]. The first epidemic peak

occurred twelve days after the restrictions were implemented. Soon afterwards, the number of new

daily documented cases started to fluctuate for about two weeks around this peak value, with

another extremely high number of cases peaked in the middle, and then finally reduced (Figure

S1). The transmission dynamics with such an irregular and unusual pattern can affect the

estimation of the effects of intervention measures. The high number of documented cases after the

introduction of interventions was generally hypothesized to be mainly caused by improved

diagnostic capability[16], leading to more detected cases rather than caused by the intrinsic growth

of the epidemic. However, most studies have not considered the changes in diagnostic capability

over time, which can affect the number of documented infections and, ultimately, the estimation

of R_e.

Accounting for temporal changes in COVID-19 diagnostic capability is critical for characterizing

transmissibility and understanding the pattern of the local Wuhan epidemic. Recent studies have

shown that the total potential case number has been significantly underestimated, with more than

80% of all infections undocumented during the initial period following the identification of SARS-

CoV-2 as the causative agent[17]. While the number of total new infections is driven by the epidemic

growth, after the introduction of new commercial kits[18] and introduction of more sensitive

diagnostic criteria[16] (Figure 1), diagnostic capacity in Wuhan increased, resulting in a higher

proportion of total new infections been documented. Therefore, it is important to consider the

improvements in diagnostic capacity over time when using the documented data to reconstruct

transmission models for COVID-19 in Wuhan.

A particularly important challenge is to understand the proportion of transmission that occurs prior

to the onset of illness. During the early outbreak, several studies have described the pre-

symptomatic transmission of SARS-CoV-2, including a 20-year-old woman from Wuhan believed

to have passed on the infection to five of her family members[19] and a Chinese individual believed

to have infected her German business partner[20], both in the absence of symptoms. The existence

of pre-symptomatic transmission indicates that COVID-19 infected individuals can be infectious

during the incubation period. However, simple classical susceptible-exposed-infected-recovered

(SEIR) models assume weak or no infectiousness during the incubation period[21] [14] [22], potentially

resulting in an underestimation of overall the infectiousness of COVID-19 cases.

In this study, in order to overcome the difficulties related to describing irregular fluctuations in the

transmission dynamics and the limitation of the simple SEIR model, a stochastic susceptible-

exposed-infected-quarantined-recovered (SEIQR) model was developed to describe the Wuhan

COVID-19 transmission pattern after the initial outbreak stage. This model extends the classic

SEIR model by including pre-symptomatic transmission and quarantined status and allows the

effects of transportation restrictions and quarantine measures on virus transmission patterns to be

estimated while accounting for improvements in the diagnostic capacity over time. After

considering varying diagnostic capabilities, we will show that this model can capture the

transmission dynamics well and can estimate the reduction in R_e precisely.

Methods

Data collection. The daily number of new documented COVID-19 cases from January 11 to

March 10 in Wuhan, Hubei province, China, were collected from the Wuhan Municipal Health

Commission[23] and the National Health Commission of the People's Republic of China[24].

Description of the SEIQR epidemic model. An SEIQR model was developed to estimate the

effect of intervention measures on COVID-19 transmission dynamics in the Wuhan population

(Figure 2). In our model, S, E, I, Q and R represent the number of individuals in susceptible,

exposed, symptomatically infectious, quarantined, and recovered statuses, with the total

population size $N = S + E + I + Q + R$ assumed to be 11 million. Here, we defined susceptible

individuals change to exposed individuals after they have had contact with the virus and become

infected but not yet symptomatic. Exposed individuals were further divided into two groups: E1,

exposed individuals at the latent period who are not able to transmit the disease; E2, exposed

individuals not at the latent period who are at a pre-symptomatic stage (referred to pre-

symptomatically infectious individuals). The proportions of E1 and E2 out of total exposed

individuals were determined using the proportion of the time span of latent period and pre-

symptomatic transmission period within the incubation period. In our model, we assumed that all

exposed individuals become symptomatic cases after incubation period, and both pre-

symptomatically and symptomatically infectious individuals can transmit the disease (Equation 2).

For quarantined status, we assumed that only symptomatically infectious individuals can be

quarantined. The SEIQR equations were derived as follows:

$$S_t = S_{t-1} - \Delta_{E,t}$$

$$E_t = E_{t-1} + \Delta_{E,t} - \Delta_{I,t}$$

$$I_t = I_{t-1} + \Delta_{I,t} - \Delta_{R,t} - \Delta_{Q,t} \tag{1}$$

$$Q_t = Q_{t-1} + \Delta_{Q,t}$$

$$R_t = R_{t-1} + \Delta_{R,t}$$

110   $\Delta_{E,t}$ is defined as the number of newly exposed individuals before symptom onset, during a time

111   interval from t to $t + 1$, $\Delta_{I,t}$ is the number of newly symptomatically infectious cases (new-onset

112   cases), $\Delta_{Q,t}$ is the number of newly quarantined cases, and $\Delta_{R,t}$ is the number of newly recovered

113   individuals. We assumed $\Delta_{E,t}$, $\Delta_{I,t}$, $\Delta_{Q,t}$, and $\Delta_{R,t}$ follow Poisson distributions:

$$\Delta_{E,t} \sim \text{Poisson}\left(\frac{\beta_{t-1}[E2_t + I_{t-1}]S_{t-1}}{N}\right)$$

$$\Delta_{I,t} \sim \text{Poisson}(\sigma \times E_{t-1})$$

$$\Delta_{Q,t} \sim \text{Poisson}(q \times I_{t-1})$$
(2)

$$\Delta_{R,t} \sim \text{Poisson}(\gamma \times I_{t-1})$$

where $E2_t$ is the number of pre-symptomatically infectious individuals (E2) at time t, assumed

determined as $E2_t = \left(\frac{\frac{1}{\sigma} - \eta}{\frac{1}{\sigma}}\right) E_{t-1}$, σ is the rate at which exposed individuals become

symptomatically infectious cases ($1/\sigma$ is the incubation period), η is the latent period, q is the

quarantine rate ($1/q$ the time between symptom onset and quarantine start), γ is the recovery rate,

expressed by $\gamma = 1/(tau - 1/\sigma)$, and tau is the generation time. Here we assumed tau was fixed

to be 10 days considering the period from being infected to recovered was generally longer than

the observed serial interval (e.g. 7.5 days)[1] and the infectious period was estimated to be around

10 days by a virology study[25]. Using a constant value of tau can reduce the model uncertainty. β_t

is the transmission rate on day t. In this model, β_t is assumed to be modulated by the Wuhan

transportation restriction policy, represented as an exponential relationship with a lag effect:

$$\beta_{t+\text{lag1}} = e^{(\alpha \times \text{pol}_t + \log(\beta_0))}$$
(3)

where pol_t is an indicator variable for the daily transportation restriction policy, with $\text{pol}_t = 0$ if

there is no transportation restriction at time t (i.e., before January 23)[15] and $\text{pol}_t = 1$ otherwise. α

is the transportation restriction effect coefficient, β_0 is the basic transmission rate without

transportation restrictions, and lag1 indicates the lag time of the transportation restrictions effect

on the virus transmission rate assumed to be 6 days[13]. Thus, β_t has a constant value throughout the

period before the implementation of transportation restriction and change to a different constant

value after then.

Mapping SEIQR model to observed hospital document cases. Model estimates of new-onset

cases ($\Delta_{I,t}$) can not be compared with observed hospital documented cases directly. This is because

documented data only captures COVID-19 cases who seek hospital care and are successfully

diagnosed, which will only be a proportion of the total number of symptomatically infectious cases

in the population estimated in the model. To address this discordance, we introduced an

observation model to link the SEIQR model simulated symptomatically infectious cases to the

observations. The daily number of hospital documented cases, $(\text{hosp\_document})_{t+\text{lag2}}$, was

assumed followed a normal distribution with the mean defined as the number of newly

symptomatically infectious cases $\Delta_{I,t}$ that were reported (documented) with the delay lag2 of 6

days[13]:

$$(\text{hosp\_document})_{t+\text{lag2}} \sim \text{Normal}\left(\Delta_{I,t} \times p(m|i) \times p(\text{hosp\_diag}|m)_{t+\text{lag2}}, \epsilon^2\right) \qquad (4)$$

where $p(m|i)$, the probability of a symptomatically infectious case seeks medical care, was

assumed to be fixed at 0.8 according to the high motivation of care-seeking behavior in Wuhan[26].

Hospital diagnostic rate, $p(\text{hosp\_diag}|m)_{t+\text{lag2}}$, represents the probability that an infected

outpatient is diagnosed as COVID-19 case by the hospital with the delay of lag2. ϵ^2 is the

distribution variance assumed to be 360000. We also defined $(\text{prop\_doc})_t$, the proportion of

documented cases out of total newly symptomatically infectious cases, could be calculated as

$(\text{prop\_doc})_t = p(m|i) \times p(\text{hosp\_diag}|m)_t$.

Given that the diagnostic capability progressed over time, hospital diagnostic rate

$p(hosp\_diag|m)_t$ was assumed to have three different values during each of the three periods:

$p_1(hosp\_diag|m)$ is the rate for the period prior to January 27 when test kits were limited,

$p_2(hosp\_diag|m)$ is the rate for the period between January 27 and Feburary 11 when test kits

were sufficient but diagnostic criteria was biased without incorporating clinical diagnosis[7], and

$p_3(hosp\_diag|m)$ is the rate for the period after February 12 when test kits were sufficient and

diagnostic criteria became more sensitive based on both clinical diagnosis and laboratory

diagnosis[16]. The values of $p_1(hosp\_diag|m)$, $p_2(hosp\_diag|m)$ and $p_3(hosp\_diag|m)$ were

estimated after fitting the model to the number of daily hospital documented cases. Hospital

documented cases on the specific days of January 27, February 12, and February 13, the dates of

change in testing capacity[7][16](Figure S1), are likely to contain retrospectively documented cases

due to the transition to new diagnostic criteria or test kits[27]. Therefore, we removed the original

values of these three data and re-filled them by using "na.spline" function in R. That is, the

smoothed values of these three dates and the original data of other dates were used during the

model fitting process.

Effective reproductive number R_e. After obtaining the posterior distributions of model

parameters $\beta_t, \sigma, q, \gamma$ and model status S_t, the effective reproductive number $R_e(t)$ before and after

the intervention policy was implemented can be calculated using the next-generation matrix

approach. Following methods previously described by Diekmann et al.[28], the transmission matrices

T and Σ can be calculated. Briefly, each element in T represents the average number of newly

infected cases in the exposed compartment (E) per unit time due to transmission via a single

infected individual in the exposed (E) or infectious group (I), calculated as $\beta_t \left[\left(\frac{\frac{1}{\sigma} - \eta}{\frac{1}{\sigma}} \right) \right] S_t$ or $\beta_t S_t$.

Σ represents the transitions between model states. $R_e(t)$ can be calculated as the first eigenvector

using the following formula:

$$R_e(t) = eig\left((-1) \begin{bmatrix} \dfrac{\beta_t\left[\left(\frac{\frac{1}{\sigma}-\eta}{\frac{1}{\sigma}}\right)\right]S_t}{N} & \dfrac{\beta_t S_t}{N} \\ 0 & 0 \end{bmatrix} \begin{bmatrix} -\sigma & 0 \\ \sigma & -(\gamma+q) \end{bmatrix}^{-1} \right)[1]$$ (6)

where $\beta_t, S_t, \sigma, q, \gamma$, and N are defined as described above.

Model-filters and validations. Since the time-varied true number of individuals in S, E, I, Q and

R statuses were not directly observable, we used Particle Markov-chain Monte Carlo (PMCMC)

method to handle such hidden variables by simultaneously estimating both the parameters and the

hidden variables[29]. Our framework of PMCMC contains two parts: the SEIQR transmission model

that generates the transmission dynamics and the observation model that maps SEIQR model to

observed hospital document cases. All posterior distributions for the epidemiological hidden

variables and parameters were obtained using the PMCMC method, implemented in the Nimble R

library[30].

The priors for the parameters were drawn from the following distributions: for the incubation

period, $1/\sigma \sim U(1,10)$; for the latent period, $\eta \sim U(1,7)$; $1/q \sim U(1,10)$, for the time between

symptom onset and quarantine start; $\beta_0 \sim U(0,1)$ for the basic transmission rate; and $\alpha \sim N(0,1)$,

for transportation control coefficient. In the observation model, the priors for time progressed

hospital diagnostic rates were set as uniform distribution: $p_1(hosp\_diag|m)$ /

$p_2(hosp\_diag|m) \sim U(0,1)$, $p_2(hosp\_diag|m)$ /$p_3(hosp\_diag|m) \sim U(0,1)$,

$p_3(hosp\_diag|m) \sim U(0,1)$.

To assess convergence, three independent chains of the SMC algorithm sets were conducted using

100,000 iterations of 1000 particle samples in each chain. We calculated the effective sample size

(ESS) and Gelman-Rubin convergence diagnostic statistics across the three chains.

Results

Reconstructing disease dynamics. The daily number of documented COVID-19 cases in Wuhan,

increased exponentially up until the first epidemic peak occurring on February 4, and started to

fluctuate around the first peak value for about two weeks. Note that the values of the highest peak

occurring around the end of the second week in two consecutive days in February were ignored in

our study because this peak was primarily caused by the retrospectively documented cases under

new diagnostic criteria, whose actual hospitalization date were diversely distributed and can not

be traced by our model (Figure S1). The irregular fluctuations can be explained by the effects of

interventions and the improved diagnostic capability: the interventions determined the timing of

the first peak and may cause a decline pattern afterward; the improved diagnostic capability led to

an increase in the number of the documented cases. Together, a high number of cases can be

produced for about two weeks. Our stochastic SEIQR model reproduced this irregular pattern by

a two-peak dynamic with the first peak occurring on February 4 and the second peak occurring

shortly on February 12 (Figure 3). Our estimated times and intensities coincide with the observed

epidemic pattern. The estimated incubation period was 5.68 days (95% CI 2.46 - 8.03), and the

estimated latent time was 2.82 days (95% CI 1.10 - 5.40) (Table 1).

Effects of intervention measures. Both transportation restrictions and quarantine measures had

significant impacts on the effective reproductive number R_e. The initial value of R_e was estimated

to be 3.23 (95% CI 2.22 - 4.20) from January 5 to January 28 (Figure 4), but has dropped by 86%

to 0.45 (95% CI 0.20 - 0.69) from January 29 to March 4 after the implementation of transportation

restrictions, calculated based on the estimated values of transmission rate β_t (Figure S2). The

estimated time delay to start quarantine after symptom onset was 5.44 days (95% CI 1.99 - 9.76)

(Table 1). For limiting the outbreak growth, quarantine measures were important but not essential.

Without quarantine measures, the initial value of R_e would increase to 4.54 (95% CI 3.65 - 6.79)

before the implementation of transportation restrictions, and would become 0.63 (95% CI 0.24 -

1.79) after the implementation of the restrictions (Figure 4). Although R_e eventually became less

than one, the high initial value of R_e would have caused a huge burden of the outbreak. We further

tested how the improvements in the diagnostic capacity influenced the estimation of R_e: about

12%-16% overestimation of R_e was found due to a fixed diagnostic capacity (Figure S3); and the

model fitting RMSE (root-mean-square error) was increased to be 278.80, comparing to 243.37

from our model, indicating a more accurate prediction was generated from our model taking

account of improving diagnostic capabilities.

Effects of detection capability. During the epidemic, the detection capability of COVID-19 in

Wuhan has been improved several times through the increased availability of test kits and the

introduction of more sensitive diagnostic criteria (Figure 1). These improvements in the detection

capability greatly affected the proportion of documented infections during three periods. From

January 11 to January 26, the estimated proportion of documented new infections out of total new

infections was 11% (95% CI 1% - 43%), increasing to 28% (95% CI 4% - 62%) following the

increase in test kit production on January 26. Then the proportion further raised to 49% (95% CI

7% - 79%) after February 12 when the diagnostic criteria became more sensitive (Figure 5A). The

estimated potential cumulative number of infections is correlated with but higher than the observed

hospital documented cases in Wuhan, and a sudden surge of hospital documented cases on

February 12 can be explained by the introduction of more sensitive diagnostic criteria (Figure 5B).

Discussion

This is the first study to demonstrate the effects of intervention measures on the transmission

dynamics in Wuhan while taking account of improvements in diagnostic capacity over time. Our

results indicated that the transportation restrictions and quarantine measures together in Wuhan

were able to contain local epidemic growth by substantially reducing R_e by 86%. This proportion

of the reduction in R_e was exactly the same as the proportion of the reduction in the average daily

number of contacts per person (14.6 vs. 2.0) between a baseline period (before the outbreak) and

the outbreak period in a recent study using contact surveys in Wuhan[31]. Since limited studies have

estimated the effects of the transportation restrictions in Wuhan, the reduction of contact rate offers

valuable information to project the possible effects on the reproduction number. Assuming the

transmissibility was proportional to the contact numbers, the reduction ratio of the contact numbers

will be proportional to the reduction ratio in R_e. These results confirm that measuring contact

mixing is an accurate way to estimate the impacts of intervention measures. Furthermore, the

proportion of undocumented infections was estimated to be reduced during the outbreak, as a

consequence of the improvements in diagnostic capability. These findings will help to inform

further analysis aimed at developing prevention strategies and evaluating the effects of public

health interventions.

While most studies assumed a fixed proportion of documented infections over time, the study

presented here estimates an initial proportion of documented infections of 11%, similar to previous

predictions of 14% by Ruiyun et al[17], which progressively increases with the improvement of

diagnostic capability. Our results suggest that the increase in the number of cases during the early

outbreak needs to be interpreted cautiously, given that the proportion of documented infections is

highly dependent on the availability and use of testing kits over time. As detection was enhanced

through improved clinical diagnosis[16], a sharp rise in cumulative cases on February 12 is likely

explained by prior onset cases retrospectively documented under new diagnostic criteria. The

undocumented infections may be of mild illness or insufficiently serious about seeking treatment[17].

Our results show that the estimated proportion of documented new infections out of total new

infections increased to 49% after diagnostic sensitivity was increased.

The estimation of R_e in the study from January 5 to January 28 is consistent with other recent

studies[32] (3.11 by Jonathan et al. [5], 3.15 by Tian et al. [21], 1.4 to 3.9 by Li et al. [1]). Furthermore, our

results demonstrate that both transportation restrictions and quarantine measures were able to

reduce COVID-19 transmission. Transportation restrictions, including stopping all forms of public

transportation, including trains, and air travel, sharply reduced social contacts thereby reducing

virus transmission rates[13][17]. Population behavioral responses (e.g., social distancing, contacts

mixing, wearing facemasks, etc.) could change concurrently with the implementation of

transportation measures[33][31]. Because a gradual increase in documented hospital cases in February

can be partly due to the increased detection capability, the effects of intervention measures

(indicated as the reduction in R_e) was estimated to be larger than previously reported in studies

using fixed detection rates over the course of the epidemic. For example, R_e was estimated to drop

by 55.3% by Kucharski et al[13]. Quarantine of symptomatic infections was also found to be essential

in curbing the epidemic. Our model estimated that the time between symptom onset and quarantine

start was 5.44 days, similar to the estimates previously reported by Tian et al. (5.19 days)[6].

The estimated incubation period was 5.68 days which is also consistent with other recent studies[1]

[21 34 35]. As the estimated latent period is 2.82 days, some transmissions may occur before the

symptom onset. Finding ways to reduce possible contact during the pre-symptomatic transmission

period may be a critical component in containing the spread of the virus. Given the existence of

pre-symptomatic transmission, this study aligns with government recommendations that people

who have had close contact with confirmed cases, regardless of whether they show symptoms or

not, need to be quarantined for 14 days[36].

The current study suggests that although intensive transportation restrictions and quarantine

measures were critical in containing the COVID-19 outbreak in Wuhan, the improvements in

detection capability have to be taken into account in order to evaluate the effectiveness of these

intervention measures more accurately. This will allow more meaningful evaluations of public

health control effects which will be important for decision in relation to which intervention used

in Wuhan should be replicated in other parts of the world in order to effectively control the current

pandemic.

Acknowledgments

This study was supported by grants from the City University of Hong Kong (#7200573 and

#9610416). We thank Dr. Chung Yin (Joey) Leung who has provided invaluable comments. We

thank Prof. Mengsu Yang, Prof. Chih-Ching Huang and Prof. Si Zhao Qin at City University of

Hong Kong for their suggestions and contributions in the preparation of the manuscript.

Author contributions

HY and JL designed the research. JL collected the data, carried out the analysis, wrote the first

draft. HY, LW and DP critically revised the manuscript, and HY gave final approval for

publication.

Declaration of interests

All authors declare no competing interests.

Reference

1.   Li, Q. *et al.* Early Transmission Dynamics in Wuhan, China, of Novel Coronavirus–

Infected Pneumonia. *N. Engl. J. Med.* (2020) doi:10.1056/nejmoa2001316.

2.   Bedford, J. *et al.* COVID-19: towards controlling of a pandemic. *Lancet* **0**, (2020).

3.   Cobey, S. Modeling infectious disease dynamics. *Science (80-.).* eabb5659 (2020)

doi:10.1126/science.abb5659.

4.   Riou, J. & Althaus, C. L. Pattern of early human-to-human transmission of Wuhan 2019

novel coronavirus (2019-nCoV), December 2019 to January 2020. *Eurosurveillance* **25**,

2000058 (2020).

5.   Read, J. M., Bridgen, J. R. E., Cummings, D. A. T., Ho, A. & Jewell, C. P. Novel

coronavirus 2019-nCoV: early estimation of epidemiological parameters and epidemic

predictions. *medRxiv* (2020) doi:10.1101/2020.01.23.20018549.

6.   Tian, H. *et al.* An investigation of transmission control measures during the first 50 days

of the COVID-19 epidemic in China. *Science (80-.).* eabb6105 (2020)

doi:10.1126/science.abb6105.

7.   Cowling, B. J. *et al.* Impact assessment of non-pharmaceutical interventions against

318   coronavirus disease 2019 and influenza in Hong Kong: an observational study. *Lancet*

319   *Public Heal.* (2020) doi:10.1016/S2468-2667(20)30090-6.

320   8.   Zhao, S. *et al.* Preliminary estimation of the basic reproduction number of novel

321   coronavirus (2019-nCoV) in China, from 2019 to 2020: A data-driven analysis in the early

322   phase of the outbreak. *Int. J. Infect. Dis.* **92**, 214–217 (2020).

323   9.   Kraemer, M. U. G. *et al.* The effect of human mobility and control measures on the

324   COVID-19 epidemic in China. *Science (80-.).* eabb4218 (2020)

325   doi:10.1126/science.abb4218.

326   10.   Chinazzi, M. *et al.* The effect of travel restrictions on the spread of the 2019 novel

327   coronavirus (COVID-19) outbreak. *Science* (2020) doi:10.1126/science.aba9757.

328   11.   Du, Z. *et al.* Risk for Transportation of 2019 Novel Coronavirus Disease from Wuhan to

329   Other Cities in China. *Emerg. Infect. Dis.* **26**, (2020).

330   12.   Zhang, J. *et al.* Evolving epidemiology and transmission dynamics of coronavirus disease

331   2019 outside Hubei province, China: a descriptive and modelling study. *Lancet Infect.*

332   *Dis.* **3099**, 1–10 (2020).

333   13.   Kucharski, A. J. *et al.* Articles Early dynamics of transmission and control of COVID-19:

334   a mathematical modelling study. *Lancet Infect. Dis.* (2020) doi:10.1016/S1473-

335   3099(20)30144-4.

336   14.   Lin, Q. *et al.* A conceptual model for the coronavirus disease 2019 (COVID-19) outbreak

337   in Wuhan, China with individual reaction and governmental action. *Int. J. Infect. Dis.* **93**,

338   211–216 (2020).

15.   The State Council_The People's Republic of China. http://www.gov.cn/xinwen/2020-

01/23/content_5471751.htm.

16.   Health Commission of Hubei Province.

http://wjw.hubei.gov.cn/bmdt/ztzl/fkxxgzbdgrfyyq/xxfb/202002/t20200213_2025580.sht

ml.

17.   Li, R. *et al.* Substantial undocumented infection facilitates the rapid dissemination of

novel coronavirus (SARS-CoV2). *Science (80-.).* eabb3221 (2020)

doi:10.1126/science.abb3221.

18.   SFDA Approves New Coronavirus Nucleic Acid Detection Reagent_Chinese government

website. http://www.gov.cn/xinwen/2020-01/27/content_5472368.htm.

19.   Bai, Y. *et al.* Presumed Asymptomatic Carrier Transmission of COVID-19. *JAMA* (2020)

doi:10.1001/jama.2020.2565.

20.   Rothe, C. *et al.* Transmission of 2019-nCoV Infection from an Asymptomatic Contact in

Germany. *N. Engl. J. Med.* (2020) doi:10.1056/nejmc2001468.

21.   Tian, H. *et al.* Early evaluation of the Wuhan City travel restrictions in response to the

2019 novel coronavirus outbreak. *medRxiv* 2020.01.30.20019844 (2020)

doi:10.1101/2020.01.30.20019844.

22.   Tang, B. *et al.* An updated estimation of the risk of transmission of the novel coronavirus

(2019-nCov). *Infect. Dis. Model.* **5**, 248–255 (2020).

23.   Wuhan Municipal Health Commission.

http://wjw.wuhan.gov.cn/ztzl_28/fk/yqtb/index.shtml.

24.   National Health Commission of the People's Republic of China.

http://www.nhc.gov.cn/xcs/xxgzbd/gzbd_index.shtml.

25.   Zou, L. *et al.* SARS-CoV-2 viral load in upper respiratory specimens of infected patients.

New England Journal of Medicine vol. 382 1177–1179 (2020).

26.   The State Council Of The People's Republic Of China, Policy and regulatory documents;

http://www.gov.cn/xinwen/2020-01/24/content_5472017.htm.

27.   Tsang, T. K. *et al.* Effect of changing case definitions for COVID-19 on the epidemic

curve and transmission parameters in mainland China: a modelling study. *Lancet Public*

Heal. (2020) doi:10.1016/S2468-2667(20)30089-X.

28.   Diekmann, O., Heesterbeek, J. A. P. & Roberts, M. G. The construction of next-

generation matrices for compartmental epidemic models. *J. R. Soc. Interface* **7**, 873–885

(2010).

29.   Endo, A., van Leeuwen, E. & Baguelin, M. Introduction to particle Markov-chain Monte

Carlo for disease dynamics modellers. *Epidemics* **29**, 100363 (2019).

30.   NIMBLE – An R package for programming with BUGS models and compiling parts of R.

https://r-nimble.org/.

31.   Zhang, J. *et al.* Changes in contact patterns shape the dynamics of the COVID-19

outbreak in China. *Science (80-.).* eabb8001 (2020) doi:10.1126/science.abb8001.

32.   Park, S. W. *et al.* Reconciling early-outbreak estimates of the basic reproductive number

and its uncertainty: framework and applications to the novel coronavirus (SARS-CoV-2)

outbreak. doi:10.1101/2020.01.30.20019877.

33. Qian, M. *et al.* Psychological responses, behavioral changes and public perceptions during

the early phase of the COVID-19 outbreak in China: a population based cross-sectional

survey. *medRxiv* 2020.02.18.20024448 (2020) doi:10.1101/2020.02.18.20024448.

34. Backer, J. A., Klinkenberg, D. & Wallinga, J. Incubation period of 2019 novel

coronavirus (2019-nCoV) infections among travellers from Wuhan, China, 20–28 January

2020. *Eurosurveillance* **25**, 2000062 (2020).

35. Lauer, S. A. *et al.* The Incubation Period of Coronavirus Disease 2019 (COVID-19) From

Publicly Reported Confirmed Cases: Estimation and Application. *Ann. Intern. Med.*

(2020) doi:10.7326/M20-0504.

36. Jiang, X. *et al.* Is a 14-day quarantine period optimal for effectively controlling

coronavirus disease 2019 (COVID-19)? *medRxiv* 2020.03.15.20036533 (2020)

doi:10.1101/2020.03.15.20036533.

Figure 1. The timeline of improved diagnostic capability and intervention measures implemented

in Wuhan, China. New commercial kits were approved by the State Food and Drug Administration

(SFDA) on January 26[18]; Updated diagnostic criteria, COVID-19 case confirmation should rely on

both clinical diagnosis and laboratory diagnosis, was introduced on February 12[16]; Wuhan

transportation restrictions were implemented on January 23[15].

Figure 2: SEIQR model schema. The population is divided into five compartments:

S (susceptible), E (exposed), I (symptomatically infectious), Q (quarantined), and R (recovered).

E2 is the number of exposed individuals after latent period who are pre-asymptomatically

infectious, β is the transmission rate, σ is the incubation rate, q is the quarantine rate, γ is the

recovery rate. A fraction of newly symptomatic infections seek for medical care and are

eventually documented by hospitals, where $p(m|i)$ is the probability of a symptomatic infectious

case seeks medical care, $p(hosp\_diag|m)_t$ represents the probability that a symptomatic

infectious outpatient is diagnosed as COVID-19 case by the hospital.

Figure 3. Number of daily hospital documented cases in Wuhan. The red lines represent model-estimated cases, grey shadow represents the 95% prediction interval, black points represent the observed documented cases, blue shaded background denotes incrementally increasing proportions of new documented infections out of total new infections in the corresponding period.

Figure 4. Estimation of the effective reproductive number R_e in Wuhan. The red point represents the estimated R_e when quarantine measures were not implemented, the black point represents R_e

when quarantine measures were implemented, and whiskers show the 95% credible intervals.

Figure 5. The prediction of temporal diagnostic capability and potential cumulative infections in

Wuhan. (A) The estimated proportion of new documented infections out of total new onset

infections on different periods with 95% credible intervals. (B) The red line is the predicted

potential total cumulative cases, and the red shadow area represents the 95% prediction interval;

the grey bar is the hospital documented cumulative cases.

Table 1. Parameter estimates of the SEIQR epidemic model. The definitions of the parameters

are described. The mean value and 95% credible interval of the posterior distribution of each of

the parameters are included. Convergence is diagnosed to have occurred when the value of

Gelman-Rubin convergence is close to 1 or the ESS is larger than 200.

Parameters	Definition	Mean	95% CI	Gelman-Rubin convergence	ESS
$1/\sigma$	Incubation period (days)	5.68	(2.46, 8.03)	1.006	261.56
η	Latent period (days)	2.82	(1.10, 5.40)	1.005	309.46
$1/q$	Time between symptom onset and quarantine start (days)	5.44	(1.99, 9.76)	1.003	477.50
α	Transportation restriction coefficient	-1.96	(-2.90, -1.21)	1.003	411.77
β_0	Basic transmission rate without transportation restrictions	0.67	(0.44, 0.97)	1.001	293.01
$p_1(\text{hosp\_diag}|m)$	Hospital diagnostic rate from Jan 11 to Jan 26	0.14	(0.01,0.54)	1.002	396.84
$p_2(\text{hosp\_diag}|m)$	Hospital diagnostic rate from Jan 27 to Feb 11	0.35	(0.05,0.78)	1.008	571.52
$p_3(\text{hosp\_diag}|m)$	Hospital diagnostic rate from Feb 12 to Mar 10	0.61	(0.09,0.98)	1.004	557.22

Supplementary

Figure S1. The original observed hospital documented daily cases without removing values. The

red points indicate the observed number of cases at the dates when many retrospectively

documented cases were counted. Data in these three days were replaced by smoothing values

because they contain many retrospectively documented cases. The black points indicate the

observed number of cases. Blue shaded background denotes incrementally increasing

proportions of new documented infections out of total new infections on the corresponding

period caused by improved diagnostic rates.

Figure S2. Estimation of the transmission rate β_t with 95% credible intervals.

Figure S3. Estimation of the effective reproductive number R_e using a fixed hospital diagnostic

rate in Wuhan. The fixed hospital diagnostic rate was assumed to be equal to the estimated mean

value of the original rate (0.14, see in Table 1) when not considering the improvement of

diagnostic capability. R_e was estimated to be 3.76 (95% CI 2.43 - 4.36) before the transportation

restrictions were implemented and to be 0.56 (95% CI 0.34 - 0.79) after the transportation

restrictions were implemented.

Figure S4. Trace plots of parameter values for the model frame. The three different colours

represent three chains.