The effect of including the Nordic Hamstring exercise on sprint and jump performance in athletes: protocol of a systematic review and meta-analyses

Kasper Krommes[1], Mathias F. Nielsen[1], Laura Krohn[1], Birk M. Grønfeldt[2], Kristian Thorborg[1,3], Per Hölmich[1], Lasse Ishøi[1]

Abstract

The Nordic Hamstring exercise reduces hamstring strain injuries in football and other sports, but the exercise is not well adopted in practice. Barriers from practitioners include fear of performance decrements, due to lack of specificity of the exercise with high speed running. However, in theory, increased eccentric hamstring strength could transfer to faster sprinting due to higher horizontal force production. Studies on the effect of the Nordic Hamstring exercise on performance have been conflicting and no synthesis of the evidence exists. We therefore pose the following question: does including the Nordic Hamstring exercise hamper sprint or jump performance in athletes? We will answer this question by performing a systematic review of the literature, critically appraise relevant studies, and GRADE the evidence across key outcomes and perform meta-analyses, meta-regression and subgroup analyses. In this protocol we outline the planned methods and procedures.

Keywords

Nordic Hamstring Exercise, Eccentric Training, Hamstring Strain Injuries, Performance, Sprint, Jump, Agility, Change of Direction, Soreness, Systematic Review, Meta-analyses,

Progress report: Besides this protocol, our data extraction form and the process of data extraction has been piloted on 3 relevant studies, along with familiarization with the Risk of Bias 2.0 tool. We have also comprised a preliminary search strategy for PubMed.

Supplementary files

- Data Extraction Form (.pdf)
- Populated PRISMA-P checklist (.pdf)

[1] Sports Orthopedic Research Center – Copenhagen, Orthopedic Department, Copenhagen University Hospital, Amager-Hvidovre, Denmark
[2] Clinical Research Centre, Copenhagen University Hospital, Amager-Hvidovre, Denmark
[3] Physical Medicine and Rehabilitation Research - Copenhagen (PMR-C), Department of Physiotherapy, Copenhagen University Hospital, Amager-Hvidovre

Corresponding Author:
Kasper Krommes, Sports Orthopedic Research Center – Copenhagen, Orthopedic Department, Amager-Hvidovre, Copenhagen University Hospital, Denmark, Kettegaard Alle 30, DK-2650. Email: kasper.krommes@regionh.dk Twitter: @krommes

1.0 BACKGROUND

1.1 Introduction

Exercise and sports participation benefits health,[1–3] all-cause mortality[4–6] and disease prevention.[7–10] Football and its various codes are the most popular sports worldwide,[11] but do carry risk of injuries,[12] of which the most common and burdensome are hamstring strain injuries.[13–15] These injuries are preventable by implementation of the Nordic Hamstring exercise in various doses,[16] even in other sports besides football.[17–19] However, even at the elite football level the exercise is not implemented,[20] and hamstring strain injuries continue to top the injury-statistics.[21] Narratives and surveys from the Sports Medicine community suggests that fear of performance-decrements and soreness from doing the exercise are barriers to implementation.[22–25] Recently, randomized controlled trials from elite and amateur football and handball players have shown improvements in sprint performance after Nordic hamstring exercise-interventions,[26–28] although the mechanism responsible has not yet been established.[29] The most relevant performance measure in football and other sports is sprint ability,[30,31] however, potential sprint performance effects of doing the Nordic Hamstring exercise has not yet been synthesized across studies. Such a synthesis address some of the concerns and barriers for implementation of the Nordic Hamstring exercise and supply new data-based arguments regarding impacts on performance, and in consequence likely increase adoption.

1.2 Objective and research questions

In order to provide a data-based estimate on the effect of including the Nordic Hamstring exercise in the team warm-up or in conditioning regiment on performance measures, we aim to answer the following questions:

Primary question:
What is the effect of including the Nordic Hamstring exercise on sprint performance in athletes?

Secondary question:
What is the effect of including the Nordic Hamstring exercise on other performance measures (such as: change of direction/agility, jumping, repeated sprint performance); and does it carry a risk of adverse events (soreness, injuries, and any other reported adverse events)?

2.0 METHODS

The protocol for this review will adhere to the PRISMA-P (Preferred Reporting Items for Systematic reviews and Meta-analyses - Protocol) statement[32] and the proposed search-extension (PRISMA-S)[33] to increase transparency and reproducibility,[34] and we aim to fulfil items on AMSTAR-2 (A MeaSurement Tool to Assess systematic Reviews)[35] to increase study quality, and ROBIS (Risk Of Bias In Systematic reviews)[36] to minimize risk of bias. Additionally, the final report will adhere to the PRISMA statement and the extension for abstracts.[37,38] Along with this pre-printed protocol, we have registered the study in the PROSPERO (international PROSPEctive Register Of systematic

reviews) repository before commencement of data collection to adhere to *a priori* decisions (submitted 31-MAR-
2020, identifier pending).[36,39] After a duration of public peer-review as pre-print, a possible updated version of the
protocol will be uploaded before we commence data collection and perform literature searches, along with an
update to the PROSPERO registration. After publication, all data (populated data extraction forms, search files,
supplementary analyses, statistical code, bias- and GRADE (Grading of Recommendations Assessment,
Development and Evaluation) assessments, populated PRISMA checklist) will be shared as supplementary files with
no restrictions, either on our institutional website, per journal repository or other open access repository (e.g.
figshare.com).

2.1 Outcomes
Modern football and other field-based sports is characterized by large amounts of sprint acceleration efforts,[30] and
consequently exercise interventions to improve this are considered a high priority.[40,41] A decrease of 0.05 s in a 10
meter sprint time by an average player/athlete, may likely result in a 25–30 cm lead during a maximal 10 m sprint.[42]
In e.g. football, such a difference is considered clinically important, and may be crucial to reach a ball before the
opponent player, block a shot, or score a goal.[42] Most sprints in football are under 20 meters, [43] and while sprinting,
team-sport athletes plateau in velocity after approximately 20 meters[44] and thus acceleration will be classified as
sprinting distances <20 meters and maximal velocity as sprinting distances ≥20 meters. To obtain low measurement
error, we will only consider sprint times measured by sensor-based systems, such as high-speed video, laser-sensors,
radar-gun, or force-plates.

Primary outcome
- Change in sprint performance (seconds) during the phase of primarily acceleration (<20 meters)

Secondary outcomes (in prioritized order)
- Change in sprint performance (seconds) during combined acceleration/maximal velocity phase (≥20 meters)
- Changes in maximal velocity, measured as either velocity (m/s or km/h) or split times during maximal
velocity running (between 20 to 40 meter)
- Change in repeated sprint ability
- Change in agility (such as: 'change of direction' test)
- Changes in jump performance (such as countermovement-jump, drop-jump, squat-jump etc.)
- Soreness or other adverse effects related to performing the Nordic Hamstring exercise

Table 1. Split sprint distances and their outcome-designation

	Sprint distance to							
Sprint distance from	**5 m**	**10 m**	**15 m**	**20 m**	**25 m**	**30 m**	**35 m**	**40 m**
0 m	A	A	A	A + MV	A + MV	A + MV	A + MV	A + MV
5 m		A	A	A + MV	A + MV	A + MV	A + MV	A + MV
10 m			A	A + MV	A + MV	A + MV	A + MV	A + MV
15 m				A + MV	A + MV	A + MV	A + MV	A + MV
20 m					MV	MV	MV	MV
25 m						MV	MV	MV
30 m							MV	MV
35 m								MV

A = acceleration phase (primary outcome); MV = Maximal velocity phase

107
108
109 ## 2.2 Criteria for eligibility of studies

111 **Table 2.** Short-form overview of key inclusion and exclusion criteria for eligible studies.

PICO domain	Inclusion criteria	Exclusion criteria
Population	Participants aged 10-40 years (10-17: adolescents, 18-40: adults)	Less than recreationally active participants
Intervention	Prescribed intervention of >3 weeks with minimum dose of 1x3 reps per week	Assisted Nordic Hamstring exercise at below bodyweight-loading
Comparator	Any	
Outcome	Sprinting time, maximal velocity, repeated sprint ability, agility, jumping, soreness or other adverse effects	Measures of sprint performance not based on sensor/gauge/video systems.
Endpoint		Follow-up measures collected >4 weeks after intervention
Design	Between- and within group/subject longitudinal study designs	Cross-sectional studies

112
113
2.2.1 Outcomes of eligible studies

• Change in sprint time during acceleration phase (split times on 0 – <20 meters)

• Change in sprint time during acceleration and maximal velocity phase (split times on 0 - ≥20 meters)

• Change in maximal velocity (velocity or split times on ≥20 – 40 meters)

• Change in repeated sprint ability

• Change in agility

• Change in jump performance

• Soreness or other adverse effects related to performing the Nordic Hamstring Exercise

122
2.2.2 Interventions of eligible studies
Studies prescribing the Nordic Hamstring exercise in any dose exceeding a minimum of 1 set of 3 repetitions
(minimum session-dose in the fifa11+ injury prevention programmes[45]) per week in volume for a minimum of 3
weeks will be considered. The Nordic hamstring exercise should be performed either as originally described by
Mjølsnes *et al*[46] or with arms positioned in front of- or behind the body, or on devices substituting the partner, such
as the 'Nordbord', 'Hamstring Solo' or other equipment. Additional loading during the exercise to maintain the
supramaximal loading of the exercise is allowed, but not assisted variations that decrease the load to below
bodyweight. Such variations will however be allowed if the programme also includes the conventional Nordic
Hamstring exercise. The Nordic hamstring exercise can be performed in isolation as the only intervention, or as add-
on to other types of exercise or conditioning. The Nordic Hamstring exercise and original protocol utilized in the
literature is described in figure 1.[46]
134
135   **Figure 1.** Description of the Nordic Hamstring Exercise and Protocol.

Description of performing the Nordic Hamstring Exercise:

"The Nordic Hamstring exercise is a partner exercise where the player attempts to resist a forward-falling motion using his hamstrings to maximize loading in the eccentric phase. The player is asked to keep their hips fixed in a slightly flexed position throughout the whole range of motion, and to brake the forward fall for as long as possible using their hamstrings, and to try keeping tension in their hamstrings even after they have to "let go". They are asked to use their arms and hands to buffer the fall, let the chest touch the surface, and immediately get back to the starting position by forcefully pushing with their hands to minimize loading in the concentric phase."[46]

Nordic Hamstring Protocol

Week	Session per week	Sets and reps
1	1	2 x 5
2	2	2 x 6
3	3	3 x 6-8
4	3	3 x 8-10
5-10	3	3 x 12, 10 ,8
10+	1	3 x 12, 10 ,8

Reps = repetitions of the exercise. Illustration is copyrighted by the authors©

136
137   ### 2.2.3 Duration of follow-up in eligible studies
138   Studies doing follow-up testing 0 days to 4 weeks after termination of the intervention will be considered. For the
139   acute effect of the Nordic Hamstring exercise, studies testing the same after doing the Nordic Hamstring Exercise
140   will also be considered.
141
2.2.4 Study design of eligible studies
All longitudinal studies implementing the Nordic Hamstring exercise will be considered, including single-group
designs looking at within-group changes only, however, we will perform subgroup analyses for only RCTs and also
for only low risk of bias RCTs.[47] For the acute effect of the Nordic Hamstring exercise on sprint changes, cross-

sectional studies with pre-post testing on the same day (effectively within-day longitudinal studies) will also be
considered. Optimally, we would include only RCTs employing a non-inferiority framework, but given our
knowledge of the field, such an evidence-pool is not available.

149
150 ## 2.2.5 Comparators of eligible studies
151 We will consider any comparators, such as usual training, no training, resistance training, specific sprint training or
152 conditioning specifically for sprint performance enhancement. If studies have more than two groups we will use the
153 Cochrane recommendations for deciding how to include data.[48] For the primary analysis we will prioritize including
154 usual care or similar comparators.

155
156 ## 2.2.6 Populations in eligible studies
157 Studies enrolling adult and adolescent participants (10-40 years) will be considered, whereas studies with children
158 (<10 years) and older adults (>40 years) and the elderly will be excluded. Participants should at least be
159 recreationally active, up to and including elite and professional athletes. We will perform subgroup analyses for
160 adults or adolescents only (18-40 years, and 10-17 years), and for elite/professional athletes only.

161
2.3 Search strategy
We have consulted an expert football strength and conditioning coach at the elite level, and a research librarian for
assistance in choosing terms and refining search strategy for the final search in PubMed, in addition to using the
Word Frequency Analyser tool to suggest possible search terms from relevant literature (Systematic Review
Accelerator, Institute for Evidence-Based Healthcare, Australia).[49] We aim for a search strategy with high sensitivity
and low precision. We have tested all terms individually, and some terms might be added or removed when applying
the same strategy in other databases or platforms. Cochrane Central, Embase, PubMed and SPORTDiscus, will be
sourced for relevant published studies,[50] and the Polyglot Search Translator tool used to translated search strategies
across databases (Systematic Review Accelerator, Institute for Evidence-Based Healthcare, Australia).[51] In addition,
any grey literature or unpublished data we can find through general search-engines (google.com, Google Scholar),
forwards and backwards snowballing, trial-registries (ISRCTN, ICTRP, EU Register, OSF, ANZCTR), or by other
means, will be sourced by hand search.[52] The preliminary search matrix for PubMed is presented below (table 3),
and will be adapted to all utilized databases to fit their respective search hierarchy, thesauruses, and operators. To
our current knowledge of the field, titles and abstract does not always specify population or design fully thus we will
not utilize search terms for these items, in order to not omit relevant studies. No filters or restrictions on dates, study
design, population or language will be applied.[53] If relevant articles with English abstracts but full-text language in
other than Danish, Norwegian, Swedish or English languages are available, translation services will be sought. A
copy of the search-results will be provided as supplementary files. The search process is planned for June 8[th] – June
16[th] 2020. For incomplete literature such as conference abstracts or theses, or insufficient reporting, we will contact
corresponding authors to supply either the full report or information regarding risk of bias and GRADE assessment
items, and to verify values for data extraction.

Table 3. Search matrix and terms for PubMed.

AND →

	Population	Intervention	Comparator	Outcome	Time	Design
OR ↓	Any (not utilized)	nordic hamstring	Any (not utilized)	sprint*	Any (not utilized)	Any (not utilized)
		nordic hamstrings		"high speed running"		
		nordic lowers		((maximal OR maximum) AND (velocit* OR speed)) AND running		
		nordic curls				
		russian curl		Athletic Performance[MeSH]		
		"hamstring lowers"		"performance"		
		(biceps femoris OR semitendinosus OR semimembranosus OR hamstring* OR Hamstring Muscles[MeSH]) AND eccentric*		"hop test"		
				"hop testing"		
				hopping		
				"jump"		
				"jumping"		
				"change of direction"		
				"performance"		
				"change of direction"		
				sore*		
				DOMS*		

*MeSH = medical subject headings. * = truncation, OR and AND are applied as Boolean operators between terms and columns.*

Two reviewers (KK & LI) will independently agree on selection of eligible studies to include using Endnote

(Clarivate Analytics) to remove duplicates and screen records. If a consensus cannot be achieved a third reviewer

(BG) will be involved. A improved PRISMA-style flowchart[54] will be presented, and we will provide a list of studies

excluded on abstract or full-text level as a supplementary. The level of inter-rater agreement at the abstract-level

between reviewers before consensus will be reported as percentage of agreement and unweighted kappa values.

Along with the publication of this protocol as pre-print, we are contacting field experts through social media and

email for suggestion or additional possible relevant studies for inclusion. Citations will be managed in Endnote.

2.4 Data extraction

Duplicate data extraction will be performed by three independent reviewers, with two reviewers per study (MF, LK,

BG),[55] using a piloted data extraction form based on Cochrane templates[56] and Center for Evidence-Based Medicine

guidelines.[57] we have provided the un-populated form as a supplementary (Data Extraction Form.pdf). Data only

presented as visualization will be extracted using a web-based tool (WebPlotDigitizer 4.2, 2019,

https://automeris.io/WebPlotDigitizer). For extracting adverse events we will use a hybrid confirmatory-exploratory

approach, in that we focus on soreness specifically and hereby pre-specified, and will also include any potential

unrecognized or unspecified adverse events.[58] For multiplicity of effect estimates, we will use an reductionist

approach by including one estimate per analysis, per study-group, based on the follow criteria: 1) most recent

(earliest follow-up measure) estimate, 2) intention-to-treat estimates, and 3) for sprint split times, the split closest to

20 meters.[59] In cases of either missing data, or data extraction from visual reporting, we will contact corresponding

authors by email and ask to either provide, or verify/correct data extraction values. After the initial email request, a

reminder after 1 week, and contact by telephone (if telephone number is available and in service),[60] none or the un-

verified data extraction will be utilized. As a supplementary, we will list all studies included for which

corresponding authors could not supply data or verify our visual extractions.

2.5 Critical appraisal

We will use Cochrane Risk of Bias Tool 2.0[41] to evaluate risk of bias in randomized trials.[61] The ROBINS-I tool will be

used for non-randomized trials.[62] Risk of Bias evaluation will be done once per study based on outcomes as

prioritized in the outcome-section. For sprint-outcomes where several eligible sprint-distances are available, only

the ones closest to the 20 m cut-off will be used as to not double-count participant-data. No study will be excluded

based on the results of the critical appraisal, but this information will be used for the quality of cumulative evidence

assessment using the Cochrane Collaborations GRADE criteria[63.] and for subgroup analysis of 'low risk of bias'

studies.[30,64] Rating the quality of individual studies rather than assessing risk of bias is inappropriate and will not be

performed.[65,66] Reviewers will perform duplicate independent bias assessment with the assistance of a machine

learning system[67,68] (MF, LK, & BG), and duplicate GRADE evaluation (KK & LI) and will achieve consensus in case

of conflicting assessments by involving an additional reviewer (KK & BG). The level of inter-tester agreement

between appraisers before consensus will be reported as percentage of agreement and unweighted kappa values. In

sports- and exercise trials, blinding participants and personnel delivering exercise-based interventions are not

feasible and we will therefore not consider domains pertaining to these blinding situations when assigning studies

to "low risk of bias" subgroup analysis. Details from included studies pertaining to reporting of funding, conflict of

interests, transparency, intention-to-treat analyses and pre-registration will be reported alongside general study

characteristics.

2.6 Analyses

Our meta-analyses of individual study results will be performed with a random effects model using the Restricted

Maximum Likelihood (REML) method to estimate a pooled mean of the overall effect with 95% confidence intervals,

and the between study variance of effect sizes (Tau-Squared, T^2). We will report estimated effect size and interpreted

it as a standardized mean difference (SMD) calculated as Cohens' d. However, we will use a Hedges' g correction to

account for small sized studies. Heterogeneity in the meta-analyses will be tested with Cochranes' Q-test (a chi^2-

test).[66] The percentage of variability in the effect estimates which is due to heterogeneity between the studies rather

than chance will be calculated and interpreted using the I^2 statistic.[66] Causes for heterogeneity will be explored

through subgroup analyses and meta-regression analyses of clinical and/or methodological covariates. Categorical

covariates (e.g age categories, playing-level, comparator, duration or volume of the Nordic Hamstring exercise) will

be investigated with stratified meta-analysis and continuous covariates (e.g. intervention dose) will be investigated

with meta-regression. A covariate is considered relevant and to explain heterogeneity in the overall analyses if

inclusion of the covariate reduces the overall Tau-Squared. If a substantial clinical heterogeneity or high risk of bias

is found, combining data in meta-analysis might be inappropriate, hence analysis will only be presented without

total effect estimates, as firm conclusions on these will be inappropriate. Instead, we will present a brief narrative

analysis, and the possible causes for heterogeneity will be debated in the discussion-section. No other qualitative

syntheses are planned. Differences in estimated effects between relevant categorical subgroups will be analysed

using meta-regression. For evaluating the risk of publication- and small study bias, a contour-enhanced funnel plot[69]

will be provided in addition to the Eggers test for Funnel-asymmetry and small study bias[70] and Beggs test for

publication bias.[71] If relevant, reasons for funnel-asymmetry will be debated in the discussion-section. Results of the

two primary outcomes will be presented in forest plots to allow for visual comparisons between studies. To allow for

a more clinically relevant interpretation of the meta-analyses estimates, we will convert the pooled estimate from

SMD to log (Odds Ratio), so we can estimate the odds ratio for becoming faster or slower and calculate the number

needed to treat for 1 athlete to e.g. get faster. Furthermore, we will also convert the pooled estimate from SMD to an

original scale (e.g. seconds), to enhance clinical interpretation of the results. All analyses will be performed in Stata

16 IC (StataCorp LLC, USA). All planned analyses are provided in table 4. Results will be emphasized based on

effect size and confidence intervals, rather than statistical significance.

Table 4. Overview of planned meta-analyses.

Primary analyses	Secondary analyses	Sensitivity and/or meta-regression analyses
<20 meter sprint time (seconds)	≥20 meter sprint time (seconds)	Only studies enrolling sub-elite or elite athletes
	Maximum velocity (velocity and 20-40 m split times)	Only low risk of bias studies
	Repeated sprint ability time	Only RCT studies
	Agility test time pooled	Only adolescents (10-17 years)
	All types of jumps pooled	Only adults (18-40 years)
	Soreness	Only studies examining the Nordic Hamstring exercise without additive interventions
	Acute effect on any sprint distance pooled	Only studies with usual training or no intervention as comparator
		Only studies with specific sprint training as comparator
		Only Nordic volume corresponding to FIFA11+ dose or less weekly
		Only Nordic volume corresponding to more than FIFA 11+ weekly
		Only Nordic intervention of >4 weeks
		Only Nordic intervention of ≤4 weeks

FIFA 11+ = a standardized warm-up program for football, including a progressive dose of the Nordic Hamstring exercise (2x5 reps, 3/week); RCT = Randomized controlled trial.

Potential conflicts of interest

We have no financial interests to declare. Some of the authors (KK, KT, PH, LI) have previously published two trials

within the scope of this review, in which the effect in both trials favours the experimental intervention and is thus

subjected to confirmation bias and self-citation incentives.

Funding and support

No specific funding were given for this study. No funding- or institutional body will have any control or involvement in

data collection, analyses, interpretation, writing the report or decision to publish.

Changelog from version 1.0 to 2.0

•   Added DOI

•   Added soreness to search matrix

•   Added specific trials registries

•   Due to bugs and installment issues, SRA-helper and de-duplikataor tools will not be utilized, and have been
removed from the methods section

•   Due to pandimical issues, the planned dates for carrying out the search has been postponed to June 8-16, 2020

Acknowledgement

278 The authors would like to thank Strength and Conditioning Coach Michael Myhre and Research Librarian Torben
279 Jørgensen with help in choosing search terms and refining the search strategy.

Author contributions

282 All authors contributed to the protocol. Contributions are visualized according to the CRediT framework (the
283 Contributor Roles Taxonomy)[72] in the following table.

	KK	MF	LK	BG	KT	PH	LI
Conceptualization							
Draft of protocol							
Methods: Study design							
Methods: Analyses plan							
Methods: Literature search							
Methods: Critical appraisal and GRADE							
Methods: Data extraction							
Revision and approval of final protocol							

REFERENCES

1. Garber, C. E. *et al.* American College of Sports Medicine position stand. Quantity and quality of exercise for developing and maintaining cardiorespiratory, musculoskeletal, and neuromotor fitness in apparently healthy adults: guidance for prescribing exercise. *Med. Sci. Sports Exerc.* **43**, 1334–1359 (2011).

2. García-Hermoso, A., Ramírez-Campillo, R. & Izquierdo, M. Is Muscular Fitness Associated with Future Health Benefits in Children and Adolescents? A Systematic Review and Meta-Analysis of Longitudinal Studies. *Sports Med.* **49**, 1079–1094 (2019).

3. Oja, P. *et al.* Infographic: Health benefits of specific types of sports. *Br. J. Sports Med.* **51**, 824–824 (2017).

4. Mandsager, K. *et al.* Association of Cardiorespiratory Fitness With Long-term Mortality Among Adults Undergoing Exercise Treadmill Testing. *JAMA Netw. Open* **1**, e183605–e183605 (2018).

5. Saint-Maurice, P. F. *et al.* Association of Leisure-Time Physical Activity Across the Adult Life Course With All-Cause and Cause-Specific Mortality. *JAMA Netw. Open* **2**, e190355–e190355 (2019).

6. Clausen, J. S. R., Marott, J. L., Holtermann, A., Gyntelberg, F. & Jensen, M. T. Midlife Cardiorespiratory Fitness and the Long-Term Risk of Mortality: 46 Years of Follow-Up. *J. Am. Coll. Cardiol.* **72**, 987–995 (2018).

7. Shibata, S. *et al.* The effect of lifelong exercise frequency on arterial stiffness. *J. Physiol.* **596**, 2783–2795 (2018).

8. Friedenreich, C. M., Neilson, H. K., Farris, M. S. & Courneya, K. S. Physical Activity and Cancer Outcomes: A Precision Medicine Approach. *Clin. Cancer Res. Off. J. Am. Assoc. Cancer Res.* **22**, 4766–4775 (2016).

9. Larson, E. B. *et al.* Exercise is associated with reduced risk for incident dementia among persons 65 years of age and older. *Ann. Intern. Med.* **144**, 73–81 (2006).

10. Paley, C. A. & Johnson, M. I. Physical Activity to Reduce Systemic Inflammation Associated

With Chronic Pain and Obesity: A Narrative Review. *Clin. J. Pain* **32**, 365–370 (2016).

11. FIFA.com. FIFA Big Count 2006: 270 million people active in football. *FIFA.com* http://www.fifa.com/media/news/y=2007/m=5/news=fifa-big-count-2006-270-million-people-active-football-529882.html (2007).

12. Ekstrand, J., Hägglund, M. & Waldén, M. Injury incidence and injury patterns in professional football: the UEFA injury study. *Br. J. Sports Med.* **45**, 553–558 (2011).

13. Ekstrand, J., Hägglund, M. & Waldén, M. Epidemiology of muscle injuries in professional football (soccer). *Am. J. Sports Med.* **39**, 1226–1232 (2011).

14. Brooks, J. H. M., Fuller, C. W., Kemp, S. P. T. & Reddin, D. B. Incidence, risk, and prevention of hamstring muscle injuries in professional rugby union. *Am. J. Sports Med.* **34**, 1297–1306 (2006).

15. Petersen, J., Thorborg, K., Nielsen, M. B. & Hölmich, P. Acute hamstring injuries in Danish elite football: a 12-month prospective registration study among 374 players. *Scand. J. Med. Sci. Sports* **20**, 588–592 (2010).

16. Ishøi, L., Krommes, K., Husted, R. S., Juhl, C. B. & Thorborg, K. Diagnosis, prevention and treatment of common lower extremity muscle injuries in sport – grading the evidence: a statement paper commissioned by the Danish Society of Sports Physical Therapy (DSSF). *Br. J. Sports Med.* (2020) doi:10.1136/bjsports-2019-101228.

17. Longo, U. G. *et al.* The FIFA 11+ program is effective in preventing injuries in elite male basketball players: a cluster randomized controlled trial. *Am. J. Sports Med.* **40**, 996–1005 (2012).

18. Sugiura, Y., Sakuma, K., Sakuraba, K. & Sato, Y. Prevention of Hamstring Injuries in Collegiate Sprinters. *Orthop. J. Sports Med.* **5**, (2017).

19. Seagrave, R. A. *et al.* Preventive Effects of Eccentric Training on Acute Hamstring Muscle Injury in Professional Baseball. *Orthop. J. Sports Med.* **2**, 2325967114535351 (2014).

20. Bahr, R., Thorborg, K. & Ekstrand, J. Evidence-based hamstring injury prevention is not adopted by the majority of Champions League or Norwegian Premier League football teams: the Nordic Hamstring survey. *Br. J. Sports Med.* **49**, 1466–1471 (2015).

21. Ekstrand, J., Waldén, M. & Hägglund, M. Hamstring injuries have increased by 4% annually in men's professional football, since 2001: a 13-year longitudinal analysis of the UEFA Elite Club injury study. *Br. J. Sports Med.* **50**, 731–737 (2016).

22. McCall, A. *et al.* Risk factors, testing and preventative strategies for non-contact injuries in professional football: current perceptions and practices of 44 teams from various premier leagues. *Br. J. Sports Med.* **48**, 1352–1357 (2014).

23. Guex, K. & Millet, G. P. Conceptual framework for strengthening exercises to prevent hamstring strains. *Sports Med. Auckl. NZ* **43**, 1207–1215 (2013).

24. Gambetta, V. More on Prevention as the Cause. *Functional Path Training:* *http://www.functionalpathtrainingblog.com/2011/05/more-on-prevention-as-the-cause-.html* http://www.functionalpathtrainingblog.com/2011/05/more-on-prevention-as-the-cause-.html (2016).

25. Oakley, A. J., Jennings, J. & Bishop, C. J. Holistic hamstring health: not just the Nordic hamstring exercise. *Br. J. Sports Med.* **52**, 816–817 (2018).

26. Ishøi, L. *et al.* Effects of the Nordic Hamstring exercise on sprint capacity in male football players: a randomized controlled trial. *J. Sports Sci.* 1–10 (2017) doi:10.1080/02640414.2017.1409609.

27. Krommes, K. *et al.* Sprint and jump performance in elite male soccer players following a 10-week Nordic Hamstring exercise Protocol: a randomised pilot study. *BMC Res. Notes* **10**, 669 (2017).

28. Chaabene, H. *et al.* Effects of an Eccentric Hamstrings Training on Components of Physical Performance in Young Female Handball Players. *Int. J. Sports Physiol.*

Perform. 1–22 (2019) doi:10.1123/ijspp.2019-0005.

29. Suarez-Arrones, L. *et al.* Dissociation between changes in sprinting performance and Nordic hamstring strength in professional male football players. *PloS One* **14**, e0213375 (2019).

30. Barnes, C., Archer, D. T., Hogg, B., Bush, M. & Bradley, P. S. The evolution of physical and technical performance parameters in the English Premier League. *Int. J. Sports Med.* **35**, 1095–1100 (2014).

31. Jeras, N. M. J., Bovend'Eerdt, T. J. H. & McCrum, C. Biomechanical mechanisms of jumping performance in youth elite female soccer players. *J. Sports Sci.* **0**, 1–7 (2019).

32. Shamseer, L. *et al.* Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015: elaboration and explanation. *BMJ* **350**, g7647 (2015).

33. Rethlefsen, M., Ayala, A. P., Kirtley, S., Koffel, J. & Waffenschmidt, S. PRISMA-S Draft. (2019) doi:10.17605/OSF.IO/7NCYS.

34. Lakens, D. *et al. Examining the Reproducibility of Meta-Analyses in Psychology: A Preliminary Report.* https://osf.io/xfbjf (2017) doi:10.31222/osf.io/xfbjf.

35. Shea, B. J. *et al.* AMSTAR 2: a critical appraisal tool for systematic reviews that include randomised or non-randomised studies of healthcare interventions, or both. *BMJ* **358**, j4008 (2017).

36. Whiting, P. *et al.* ROBIS: A new tool to assess risk of bias in systematic reviews was developed. *J. Clin. Epidemiol.* **69**, 225–234 (2016).

37. Moher, D., Liberati, A., Tetzlaff, J., Altman, D. G. & PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *J. Clin. Epidemiol.* **62**, 1006–1012 (2009).

38. Beller, E. M. *et al.* PRISMA for Abstracts: Reporting Systematic Reviews in Journal and Conference Abstracts. *PLOS Med.* **10**, e1001419 (2013).

39. Palpacuer, C. *et al.* Vibration of effects from diverse inclusion/exclusion criteria and analytical choices: 9216 different ways to

perform an indirect comparison meta-analysis. *BMC Med.* **17**, 174 (2019).

40. Haugen, T. Sprint conditioning of elite soccer players: Worth the effort or lets just buy faster players? 3 (2017).

41. Haugen, T. The role and development of sprinting speed in soccer. (University of Agder, Faculty of Health and Sport Science, 2014).

42. Haugen, T. & Buchheit, M. Sprint Running Performance Monitoring: Methodological and Practical Considerations. *Sports Med. Auckl. NZ* **46**, 641–656 (2016).

43. Andrzejewski, M., Chmura, J., Pluta, B., Strzelczyk, R. & Kasprzak, A. Analysis of Sprinting Activities of Professional Soccer Players: *J. Strength Cond. Res.* **27**, 2134–2140 (2013).

44. Clark, K. P., Rieger, R. H., Bruno, R. F. & Stearne, D. J. The National Football League Combine 40-yd Dash: How Important is Maximum Velocity? *J. Strength Cond. Res.* **33**, 1542–1550 (2019).

45. Thorborg, K. *et al.* Effect of specific exercise-based football injury prevention programmes on

the overall injury rate in football: a systematic review and meta-analysis of the FIFA 11 and 11+ programmes. *Br. J. Sports Med.* **51**, 562–571 (2017).

46. Mjølsnes, R., Arnason, A., Østhagen, T., Raastad, T. & Bahr, R. A 10-week randomized trial comparing eccentric vs. concentric hamstring strength training in well-trained soccer players. *Scand. J. Med. Sci. Sports* **14**, 311–317 (2004).

47. Tan, A. C. *et al.* Prevalence of trial registration varies by study characteristics and risk of bias. *J. Clin. Epidemiol.* **113**, 64–74 (2019).

48. 16.5.4 How to include multiple groups from one study. https://handbook-5-1.cochrane.org/chapter_16/16_5_4_how_to_include_multiple_groups_from_one_study.htm.

49. Clark, J. *et al.* A full systematic review was completed in 2 weeks using automation tools: a case study. *J. Clin. Epidemiol.* **121**, 81–90 (2020).

50. Aagaard, T., Lund, H. & Juhl, C. Optimizing literature search in systematic reviews – are MEDLINE, EMBASE and CENTRAL enough

for identifying effect studies within the area of musculoskeletal disorders? *BMC Med. Res. Methodol.* **16**, (2016).

51. Improving the translation of search strategies using the Polyglot Search Translator: a randomised controlled trial. *Bond University Research Portal* https://research.bond.edu.au/en/datasets/improving-the-translation-of-search-strategies-using-the-polyglot.

52. Greenhalgh, T. & Peacock, R. Effectiveness and efficiency of search methods in systematic reviews of complex evidence: audit of primary sources. *BMJ* **331**, 1064–1065 (2005).

53. Beynon, R. *et al.* Search strategies to identify diagnostic accuracy studies in MEDLINE and EMBASE. *Cochrane Database Syst. Rev.* MR000022 (2013) doi:10.1002/14651858.MR000022.pub3.

54. Vu-Ngoc, H. *et al.* Quality of flow diagram in systematic review and/or meta-analysis. *PLOS ONE* **13**, e0195955 (2018).

55. Jian-Yu, E. *et al.* Adjudication rather than experience of data abstraction matters more in reducing errors in abstracting data in systematic reviews. *Res. Synth. Methods* **n/a**,.

56. Data extraction forms. /data-extraction-forms.

57. Data Extraction in Meta-analysis. *CEBM* https://www.cebm.net/2014/06/data-extraction-in-meta-analysis/ (2014).

58. Junqueira, D. 2019 Cochrane Methods Symposium Part 7: Special issues for addressing adverse effects. *Cochrane Training* https://training.cochrane.org/resource/part-7-special-issues-addressing-adverse-effects.

59. López-López, J. A., Page, M. J., Lipsey, M. W. & Higgins, J. P. T. Dealing with effect size multiplicity in systematic reviews and meta-analyses. *Res. Synth. Methods* **9**, 336–351 (2018).

60. Danko, K. J., Dahabreh, I. J., Ivers, N. M., Moher, D. & Grimshaw, J. M. Contacting authors by telephone increased response proportions compared with emailing: results of a randomized study. *J. Clin. Epidemiol.* **115**, 150–159 (2019).

61. Sterne, J. & et al. RoB 2: a revised tool for assessing risk of bias in randomised trials. *BMJ Press*.

62. Sterne, J. A. *et al.* ROBINS-I: a tool for assessing risk of bias in non-randomised studies of interventions. *BMJ* **355**, i4919 (2016).

63. Santesso, N. *et al.* GRADE guidelines 26: informative statements to communicate the findings of systematic reviews of interventions. *J. Clin. Epidemiol.* (2019) doi:10.1016/j.jclinepi.2019.10.014.

64. Schünemann, H. J., Fretheim, A. & Oxman, A. D. Improving the use of research evidence in guideline development: 9. Grading evidence and recommendations. *Health Res. Policy Syst.* **4**, 21 (2006).

65. Boutron, I. & Ravaud, P. Classification systems to improve assessment of risk of bias. *J. Clin. Epidemiol.* **65**, 236–238 (2012).

66. Higgins, J. & Green, S. Cochrane Handbook for Systematic Reviews of Interventions Version 5.1.0 [updated March 2011]. *Cochrane Collab.* **2011**,.

67. Marshall, I. J., Kuiper, J. & Wallace, B. C. RobotReviewer: evaluation of a system for automatically assessing bias in clinical trials. *J. Am. Med. Inform. Assoc. JAMIA* **23**, 193–201 (2016).

68. Viswanathan, M. *et al.* Assessing the Risk of Bias in Systematic Reviews of Health Care Interventions. in *Methods Guide for Effectiveness and Comparative Effectiveness Reviews* (Agency for Healthcare Research and Quality (US), 2008).

69. Sterne, J. A. C. *et al.* Recommendations for examining and interpreting funnel plot asymmetry in meta-analyses of randomised controlled trials. *BMJ* **343**, d4002 (2011).

70. Egger, M., Davey Smith, G., Schneider, M. & Minder, C. Bias in meta-analysis detected by a simple, graphical test. *BMJ* **315**, 629–634 (1997).

71. Begg, C. B. & Mazumdar, M. Operating characteristics of a rank correlation test for publication bias. *Biometrics* **50**, 1088–1101 (1994).

72. Holcombe, A. Farewell authors, hello

    contributors. *Nature* **571**, 147–147 (2019).