Title: Spatial variability in the risk of death from COVID-19

in Italy, 2020

Authors: Kenji Mizumoto[1,2,3]*, Sushma Dahal[3]*, Gerardo Chowell[3,]

*Mizumoto and Dahal contributed equally

Affiliations:

[1] Graduate School of Advanced Integrated Studies in Human Survivability, Kyoto University

Yoshida-Nakaadachi-cho, Sakyo-ku, Kyoto, Japan

[2] Hakubi Center for Advanced Research, Kyoto University, Yoshidahonmachi, Sakyo-

ku,Kyoto, Japan;

[3] Department of Population Health Sciences, School of Public Health, Georgia State

University, Atlanta, Georgia, USA

Correspondence to:

K Mizumoto, Graduate School of Advanced Integrated Studies in Human Survivability,
Kyoto University Yoshida-Nakaadachi-cho, Sakyo-ku, Kyoto, Japan

Tel: +81 75 762 2001 Fax: +81 75 762 2277

E-mail: mizumoto.kenji.5a@kyoto-u.ac.jp
19

20 **Article type:**

21 Original research

23 **Word count:**

24 Abstract: 199   (150-200)
25 Main: 2612   (Max 3500)
26

Abstract

Objectives: Italy is bearing the brunt of the COVID-19 pandemic, as the death toll there has

already surpassed that in Wuhan, the city in China where the coronavirus emerged in

December 2019. Here we employ statistical methods to assess the severity of COVID-19

pandemic across different regions of Italy.

Method: We manually retrieved the daily cumulative numbers of laboratory-confirmed cases

and deaths attributed to COVID-19 stratified by region in Italy. We estimated both the crude

and time-delay adjusted case fatality ratio across five geographic regions of Italy.

Results: The Northwest that includes Lombardy exhibited the highest time-delay-adjusted

CFR at 23.0% followed by the Northeast (14.2%), the Center region (11.6%) and the South

region (12.1%). The Island region has exhibited the lowest CFR (9.6%).

Conclusion: Our estimates in the Northwest and the Northeast are higher than those reported

from most affected areas in China. The CFR in Northwest Italy is 1.9-fold higher than that in

Wuhan. Our finding reflects the need of urgent medical support in the Northwest region and

the appropriate planning and supplies procurement in other regions of Italy focusing on

medical care delivery to those who are at the highest risk of poorer outcomes due to COVID-

19.

Keywords: COVID-19, Italy, time-delay adjusted CFR, 2020

Introduction

In December 2019, Wuhan, the capital of Hubei Province of China reported a cluster of cases

of severe pneumonia of unknown etiology to the China National Health Commission *(1)*, and

on January 7, 2020, a novel coronavirus was isolated from afflicted patients in Wuhan *(1)*.

This novel coronavirus was named as severe acute respiratory syndrome coronavirus 2

(SARS-CoV-2) while the disease it causes was named COVID-19 *(2)*. Since the initial cases

of COVID-19, the virus has rapidly spread throughout China, and subsequently the number

of COVID-19 cases soared in many parts of the world. Considering the explosive spread and

significant severity associated with the novel coronavirus, the WHO declared this public

health emergency a pandemic on March 11, 2020 *(3)*. As of March 29, a total of 634,835

confirmed cases of COVID-19 and 29,957 deaths have been recorded globally with 202

countries/territories/areas reporting variable disease growth rates *(4)*. Moreover, the US has

reported the highest number of cases (16.3%) while Italy has exhibited the highest death toll

(33.4%) *(4)*.

The severity impact of any pandemic situation like COVID-19 depends on the transmission

rate of the disease, the capacity of the health care system, and the spectrum of clinical

severity which is tied to socio-demographic factors (age, gender) and the underlying

prevalence of comorbidities in the population *(5)*. A better understanding of the expected

influx of severe patients to the health care system during the coronavirus pandemic in

different areas of the world is key to anticipate medical resources such as ICU units and

ventilators which are critically needed to save the lives of severely ill patients *(5-7)*.

The Case fatality ratio (CFR) is one of the most important epidemiological metrics to

quantify the clinical severity of emerging infectious disease such as COVID-19. So far,

several studies have attempted to elucidate the case fatality ratios for different population

segments and geographic regions particularly based on epidemiological data from China *(8-*

10). However, there is still a scarcity of studies carefully documenting the severity of the

COVID-19 pandemic in populations outside China. Accumulating epidemiological data

indicates that the CFR varies by geographical location, intensity of transmission,

characteristics of patients such as age, sex, and comorbidity status *(9)*. For example, the time-

delay adjusted CFR for Wuhan was estimated at 12.2% compared to 4.2% for Hubei province

excluding Wuhan and 0.9% in China excluding Hubei province *(11)*. Rough differences in

severity of the pandemic in different countries have been highlighted *(10)*. However, there is

a need to quantify spatial variability in CFR and investigate how this variability is influenced

by population factors and the characteristics of the health care system.

At the time of writing, Italy is exhibiting an alarming death toll from the COVID-19

pandemic but estimates of the case fatality ratio that carefully account for the delay from

onset of symptoms to death are not yet available. In this study we provide estimates of the

COVID-19 CFR for different regions of Italy by relying on publicly available daily series of

confirmed cases and deaths from March 2, 2020 to April 11, 2020.

Methods

Study setting

Italy is located in Southern Europe and is comprised of some islands and the Italian peninsula

that extends into the central Mediterranean Sea. Italy has 20 administrative regions including

5 autonomous regions *(12)*. For our study we divided the country into 5 geographic regions:

Northwest, Northeast, Centre, South, and Island. Lombardia, Piemonte, Liguria, and Valle

d'Aosta were included in Northwest. Emilia Romagna, Friuli-Venezia Giulia, Trentino Alto

Adige, and Veneto were included in Northeast. Lazio, Marche, Toscana, and Umbria were

included in Centre region while Abruzzo, Puglia, Basilicata, Calabria, Compania, and Molise

were included in the South region. The two islands, Sardegna, and Sicilia were categorized as

Islands.

Initial cases of COVID-19 in Italy

The first two confirmed cases of COVID-19 in Italy had a travel history to Wuhan, China and

were reported on 31[st] January 2020. The third case was not confirmed until February 7, 2020.

On February 22, 2020 the cumulative total reached 9 cases and subsequent incident cases

continued to follow a rapidly increasing daily trend *(13)*.

Data sources

The Ministry of Health of Italy releases daily report on COVID-19 cases and deaths *(14)*.

From these reports we manually retrieved the daily cumulative numbers of reported

laboratory confirmed COVID-19 cases and deaths stratified for five Italian regions from

March 2, 2020 to April 11, 2020.

Statistical analysis

The crude CFR is defined as the number of cumulative deaths divided by the number of

cumulative cases at a specific point in time. For the estimation of CFR in real time, we

employed the delay from hospitalization to death, h_s, which is assumed to be given by $h_s =$

$H(s) - H(s-1)$ for s>0 where $H(s)$ is a cumulative density function of the delay from

hospitalization to death and follows a gamma distribution with mean 10.1 days and SD 5.4

days, obtained from the previously published paper *(11)*. Let $\pi_{a,ti}$ be the time-delay adjusted

case fatality ratio on reported day t_i in area a, the likelihood function of the estimate $\pi_{a,ti}$ is

$$L(\pi_{a,t_i}; c_{a,t}) = \prod_{t_i} \begin{pmatrix} \sum_{t=1}^{t_i} c_{a,t} \\ D_{a,t_i} \end{pmatrix} \left(\pi_{a,t_i} \frac{\sum_{t=2}^{t_i} \sum_{s=1}^{t-1} c_{a,t-s} h_s}{\sum_{t=1}^{t_i} c_{a,t}} \right)^{D_{a,t_i}} \left(1 \right.$$

$$\left. - \pi_{a,t_i} \frac{\sum_{t=2}^{t_i} \sum_{s=1}^{t-1} c_{a,t-s} h_s}{\sum_{t=1}^{t_i} c_{a,t}} \right)^{\sum_{t=1}^{ti} c_{a,t} - D_{a,t_i}}$$

where $c_{a,t}$ represents the number of new cases with reported day t in area a, and $D_{a,ti}$ is the

cumulative number of deaths until reported day t_i in area *a (15, 16)*. Among the cumulative

cases with reported day *t* in area a, $D_{a,ti}$ have died and the remainder have survived the

infection. The contribution of those who have died with biased death risk is shown in the

middle parenthetical term and the contribution of survivors is presented in the right

parenthetical term. We assume that $D_{a,ti}$ is the result of the binomial sampling process with

probability $\pi_{a,ti}$.

We estimated model parameters using a Monte Carlo Markov Chain (MCMC) method in a

Bayesian framework. Posterior distributions of the model parameters were estimated by

sampling from the three Markov chains. For each chain, we drew 100,000 samples from the

posterior distribution after a burn-in of 20,000 iterations. Convergence of MCMC chains

were evaluated using the potential scale reduction statistic *(17, 18)*. Estimates and 95%

credibility intervals for these estimates are based on the posterior probability distribution of

each parameter and based on the samples drawn from the posterior distributions.

All statistical analyses were conducted in R version 3.6.1 (R Foundation for Statistical

Computing, Vienna, Austria) using the 'rstan' package.

Results

As of April 11, a total of 152,271 cases and 19,468 deaths due to COVID-19 have been

reported in Italy. Moreover, the Northwest region has reported the highest number of cases at

79,878 (52.5%) and deaths at 12,985 (66.7%). Similarly, the Northeast reported a total of

40,723 (26.7%) of cases and 3,981 (20.4%) deaths. The Island, Center and the South regions

shared less than 20% of the cases and about 10% of the total deaths.

Figure 1 displays the cumulative cases in (A) Northwest, (B) Northeast, (C) Center, (D)

South and (E) Island, and cumulative deaths in (F) Northwest, (G) Northeast (H) Center (I)

South and (J) Island, over time. Cumulative cases and cumulative deaths increased rapidly in

the Northwest region compared to other Italian regions. In contrast, the cumulative number of

cases and deaths in Island region is relatively mild whereas it is moderate in the Northeast,

Center and the South regions.

Figure 2 illustrates observed and model based posterior estimates of the crude case fatality

ratio in (A) Northwest, (B) Northeast, (C) Center, (D) South and (E) Island, and the time-

delay adjusted case fatality ratio in (F) Northwest, (G) Northeast (H) Center (I) South and (J)

Island. Black dots show crude case fatality ratios, and light and dark indicate 95% and 50%

credible intervals for posterior estimates, respectively. Except for the initial days (first 5

days) our model based crude CFR fitted the observed data well. For the time-delay adjusted

CFR, our model based on posterior estimates are much higher than the crude observed CFR.

The differences are much higher for the initial stage and then declined rapidly followed by a

small difference in the later stage of the epidemic across all regions. For the most affected

Northwest region, the time-delay adjusted risk of death was stable at highest point (100%) for

first 7 days (considering March 1st as day 1) and rapidly declined to 50% by day 15 and

thereafter exhibited a gradual decline (about 31% by day 28). For the Northeast, the risk of

death has declined to about 20% and below 16% for the Center and the South regions by day

28. For the Island region, the risk of death is below 12% by day 28 (figure 2). There was an

overall downward trend of CFR for all the five regions, including a period of stable higher

rate for the initial stage in the Northwest and Northeast regions.

A summary of the time delay adjusted case fatality ratio, range of median estimates and crude

CFR of COVID-19 across five regions of Italy are presented in Table 1. The Northwest had

the highest time delay adjusted CFR of 23.0% (95% credible interval: 22.7, 23.4] followed by

the Northeast region (14.2%) [95% CrI: 13.7, 14.6], the Center region (11.6%) [95% CrI:

11.0, 12.1] and the South region (12.1%) [95% CrI: 11.4, 13.0] (Table 1, figure 3). The Island

has exhibited the lowest CFR (9.6%) [95% CrI: 8.4, 10.9]. The crude CFR for Northwest was

16.3% [95% CI: 16.0, 16.5] compared to 9.8% [95% CI: 9.5, 10.1] in Northeast and 6.6%

[95% CI: 5.8, 7.4] in the Island (Table 1).

177   Discussion

In this paper, we have estimated the time delay adjusted case fatality ratio of COVID-19 for

five different areas of Italy. We found that the latest estimate of time-delay adjusted CFR in

Italy varied substantially across regions with the highest value in the Northwest and the

lowest in the Island region. The Northwest includes the regions of Lombardia, which has

reported the highest number of COVID-19 cases and deaths in Italy. The adjusted CFR in the

Northwest was very large (23.0%) which is 1.6-fold higher than the estimate of Northeast

(14.2%), 2.0-fold higher than that of the Center region (11.6%), 1.9-fold higher than that of

the South region (12.1%), and 2.4-fold higher than that estimated for the Island region

(9.6%). Our results call for all-out efforts to mitigate transmission and invest medical

resources to ameliorate the burden on an already overwhelmed health care system.

The adjusted CFR estimates across all the regions in Italy except the Island region are higher

than the previous CFR estimates for Wuhan (12.2%) *(11)* and Korea (1.4%) *(19)*. Wuhan had

the highest time-delay adjusted CFR compared to other regions of China *(11)*. When we

compare the adjusted CFR of the most affected regions in Italy and China, the estimate for

Italy is more than 1.9 times the estimate for China (23.0% vs 12.2%) *(11)*. This difference

may be partly associated with the population structure of the two countries as suggested in a

previous study *(20)*. Italy has an older population compared to China. In 2015 the median age

of total population in Italy was 45.4 years compared to 36.7 years for China *(21)*. In 2019,

approximately 23% of population in Italy was 65 years and older *(20)* compared to 12.6% in

China *(22)*.

Differences in the definition of COVID-19 related deaths and the testing strategies could

affect differences in CFR. Indeed, there is a lack of consistent case definitions of COVID-19

related deaths across countries. In Italy, COVID-19–related deaths include all deaths that

occur in patients that test positive for SARS-CoV-2 via RT-PCR independent of any

preexisting disease that might have caused the death *(20)*. Likewise, in the early phase of

epidemic, there was an extensive testing strategy in Italy that included both symptomatic

cases and their and asymptomatic contacts but later more strict testing policies prioritized

more severe suspected cases requiring hospitalization *(20)*. Further studies are needed to

clarify how differences in case definitions and testing strategies affect CFR estimates in

different countries.

In our study, as the epidemic progressed, we saw a downward trend in the time delay

adjusted CFR for all the five regions. However, for Northwest and Northeast, there was also

an initial phase with steady state at higher CFR which was relatively longer for Northwest

compared to Northeast. A previous study on COVID-19 using data from China has also

found the declining trend of time delay adjusted CFR for Hubei province excluding Wuhan

(11). This trend was also reported for the 2015 MERS outbreak in South Korea in which the

risk of death was significantly associated with older age and underlying health condition *(23)*.

In the early phase of the outbreak of emerging infectious disease like MERS and COVID-19,

the detection rate of mildly symptomatic cases is low and only patients who have serious

conditions are confirmed due to hospitalization as happened in Wuhan *(11, 24)* and South

Korea *(23)*. However, the downward trend of CFR in the later phase of epidemic suggests

both an improvement in epidemiologic surveillance and a decline in the proportion of

vulnerable patients *(11)*. Because of the decline in the proportion of vulnerable patients and

an increased detection of mildly symptomatic cases, the epidemic might get prolonged unless

strict social/physical distancing measures are applied *(11)*.

In Northwest Italy, though the time delay adjusted CFR has been gradually declining, the

latest estimate is still the highest within Italy and globally *(8, 24-27)* and the estimates for the

other regions are also substantially high though not as high as that in the Northwest region.

Our findings reflect the need of urgent medical support in the Northwest region and the

appropriate planning and supplies procurement in the other regions of Italy with a focus on

medical care delivery to those who are at the highest risk of poorer outcomes due to COVID-

19 such as patients categorized as critical, the elderly, and those with multiple comorbidities

including cardiovascular disease, hypertension, and diabetes. Similarly, social distancing

measures are needed to flatten the curve and prevent the health care system from overloading

to a breaking point.

The Italian government put in place extraordinary social distancing measures to reduce

transmission through strict self-isolation, including the restriction of movement in the

Lombardy region of Northwest at the beginning and later expanded at the nationwide level is

praiseworthy *(28)*. However, there are reports from the Lombardy region that indicate an

overwhelming surge in patients requiring ICU admission and overloaded public health

laboratories *(29)*, which underscore the dire need to mitigate transmission to flatten the

incidence curve and reduce the pressure on the healthcare system.

Our study is not exempted from limitations. The preferential ascertainment of severe cases

bias in COVID-19 may have spuriously increased our estimate of CFR *(30)*, which is a

frequent caveat in this type of studies *(31, 32)*. Similarly, for a disease like COVID-19 in

which there is a rapid increase in infection, but the infection-death time is long (ranges from

2 to 8 weeks) *(9)*, our CFR estimate could have been affected by delayed reporting bias *(30,*

33). Similarly, our data on number of cases reflects the date of reporting and not the date of

onset of illness.

In conclusion our latest estimates of the risk of death due to COVID-19 was as high as 23.0%

in the most affected Northwest region of Italy whereas for the least affected Island region the

CFR was estimated at 9.6%. Our estimates in Northeast (23.0%), the Center (11.6%) and the

South (12.1%) are also elevated compared to delay-adjusted CFR estimates from other areas

such as South Korea and China. Our results call for all-out efforts to reduce transmission in

the most severely affected areas of Italy to ameliorate the burden on an already overwhelmed

health care system.

Financial support.

KM acknowledges support from the Japan Society for the Promotion of Science (JSPS)
KAKENHI Grant Number 18K17368 and from the Leading Initiative for Excellent Young
Researchers from the Ministry of Education, Culture, Sport, Science & Technology of Japan.
KK acknowledges support from the JSPS KAKENHI Grant Number 18K19336 and
19H05330. GC acknowledges support from NSF grant 1414374 as part of the joint NSF-
NIH-USDA Ecology and Evolution of Infectious Diseases program.
262
263

Conflict of interest

The authors declare no conflicts of interest.

References

1.   Wang C, Horby PW, Hayden FG, Gao GF. A novel coronavirus outbreak of global

health concern. *Lancet*. **2020**;395(10223):470-3.

2.   WHO. Naming the coronavirus disease (COVID-19) and the virus that causes it 2020.

Available from: https://www.who.int/emergencies/diseases/novel-coronavirus-

2019/technical-guidance/naming-the-coronavirus-disease-(covid-2019)-and-the-virus-that-

causes-it

3.   WHO. Rolling updates on coronavirus disease (COVID-19) 13 March, **2020**

[03/17/2020]. Available from: https://www.who.int/emergencies/diseases/novel-coronavirus-

2019/events-as-they-happen

4.   WHO. Coronavirus disease 2019 (COVID-19): Situation report 69. 29 March 2020.

5.   Lipsitch M, Swerdlow DL, Finelli L. Defining the epidemiology of Covid-19—

studies needed. *N Engl J Med*. **2020**.

6.   Reich NG, Lessler J, Cummings DA, Brookmeyer R. Estimating absolute and relative

case fatality ratios from infectious disease surveillance data. *Biometrics*. **2012**;68(2):598-606.

7.   Wu JT, Leung K, Bushman M, Kishore N, Niehus R, de Salazar PM, et al. Estimating

clinical severity of COVID-19 from the transmission dynamics in Wuhan, China. *Nature*

Med. **2020**. doi: 10.1038/s41591-020-0822-7.

8.   Novel CPERE. The epidemiological characteristics of an outbreak of 2019 novel

coronavirus diseases (COVID-19) in China. Zhonghua liu xing bing xue za zhi= Zhonghua

liuxingbingxue zazhi. **2020**;41(2):145.

9.   WHO. Report of the WHO-China Joint Mission on Coronavirus Disease 2019

(COVID-19). February; **2020**.

10.   Rodriguez-Morales AJ, Cardona-Ospina JA, Gutiérrez-Ocampo E, Villamizar-Peña

R, Holguin-Rivera Y, Escalera-Antezana JP, et al. Clinical, laboratory and imaging features

of COVID-19: A systematic review and meta-analysis. *Travel Med. Infect. Dis*. **2020**:101623.

11.   Mizumoto K, Chowell G. Estimating Risk for Death from 2019 Novel Coronavirus

Disease, China, January-February 2020. *EID*. **2020**;26(6).

12.   The world fact book: Europe- Italy 2020 [03/25/2020]. Available from:

https://www.cia.gov/library/publications/the-world-factbook/geos/print_it.html

13.   WHO. Coronavirus disease 2019 (COVID-19): Situation reports 11, 20, 22, 23, 24.

2020.

14.   Ministry of Health Italy. New Coronavirus Italy: Ministry of Health; 2020. Available

from:

http://www.salute.gov.it/portale/nuovocoronavirus/archivioNotizieNuovoCoronavirus.jsp

15.   Nishiura H, Klinkenberg D, Roberts M, Heesterbeek JA. Early epidemiological

assessment of the virulence of emerging infectious diseases: a case study of an influenza

pandemic. *PLoS One*. **2009**;4(8).

16.   Tsuzuki S, Lee H, Miura F, Chan YH, Jung S-m, Akhmetzhanov AR, et al. Dynamics

of the pneumonic plague epidemic in Madagascar, August to October 2017. *Euro Surveill*.

2017;22(46).

17.   Gamerman D, Lopes HF. Markov chain Monte Carlo: stochastic simulation for

Bayesian inference: CRC Press; **2006**.

18.   Gelman A, Rubin DB. Inference from iterative simulation using multiple sequences.

Stat Sci. **1992**;7(4):457-72.

19.   Shim Eunha, Mizumoto Kenji, Choi Wongyeong, Chowell Gerardo. Estimating the

risk of COVID-19 death during the course of the outbreak in Korea, February- March, 2020.

2020.

20.   Onder G, Rezza G, Brusaferro S. Case-Fatality Rate and Characteristics of Patients

Dying in Relation to COVID-19 in Italy. *JAMA*. **2020**.

21.   World Population Prospects 2019, Online Edition. Rev. 1. United Nations,

Department of Economic and Social Affairs, Population Division. **2019**.

22.   statistica. Population of China in 2019, by broad age group 2019. Available from:

https://www.statista.com/statistics/251524/population-distribution-by-age-group-in-china/

23.   Mizumoto K, Endo A, Chowell G, Miyamatsu Y, Saitoh M, Nishiura H. Real-time

characterization of risks of death associated with the Middle East respiratory syndrome

(MERS) in the Republic of Korea, **2015**. *BMC med*. 2015;13(1):228.

24.   Yang S, Cao P, Du P, Wu Z, Zhuang Z, Yang L, et al. Early estimation of the case

fatality rate of COVID-19 in mainland China: a data-driven analysis. *Ann Transl Med*.

2020;8(4).

25.   Fang Z, Yi F, Wu K, Lai K, Sun X, Zhong N, et al. Clinical Characteristics of 2019

Coronavirus Pneumonia (COVID-19): An Updated Systematic Review. *medRxiv*. **2020**.

26.   Wilson N, Kvalsvig A, Barnard LT, Baker M. Estimating the Case Fatality Risk of

COVID-19 using Cases from Outside China. *medRxiv*. **2020**.

27.   Dehkordi AH, Alizadeh M, Derakhshan P, Babazadeh P, Jahandideh A.

Understanding Epidemic Data and Statistics: A case study of COVID-19. arXiv preprint

arXiv:200306933. **2020**.

28.   Remuzzi A, Remuzzi G. COVID-19 and Italy: what next? *Lancet*. 2020.

29.   Grasselli G, Pesenti A, Cecconi M. Critical Care Utilization for the COVID-19

Outbreak in Lombardy, Italy: Early Experience and Forecast During an Emergency

Response. *JAMA*. **2020**.

30.   Lipsitch M, Donnelly CA, Fraser C, Blake IM, Cori A, Dorigatti I, et al. Potential

biases in estimating absolute and relative case-fatality risks during outbreaks. *PLoS Neglect

Trop D*. **2015**;9(7).

31.   Yu H, Cowling BJ, Feng L, Lau EH, Liao Q, Tsang TK, et al. Human infection with

avian influenza A H7N9 virus: an assessment of clinical severity. *Lancet*.

2013;382(9887):138-45.

32.   Cauchemez S, Fraser C, Van Kerkhove MD, Donnelly CA, Riley S, Rambaut A, et al.

Middle East respiratory syndrome coronavirus: quantification of the extent of the epidemic,

surveillance biases, and transmissibility. *Lancet Infect Dis*. **2014**;14(1):50-6.

33.   Kucharski AJ, Edmunds WJ. Case fatality rate for Ebola virus disease in west Africa.

Lancet. **2014**;384(9950):1260.

Table 1. Summary results of time-delay adjusted case fatality ratio of COVID-19 in the 5

areas in Italy, 2020 (As of April 11, 2020)

Area	Latest estimate	Range of median estimates during the study period	Crude CFR
Northwest	23.0% (95%CrI[§]: 22.7-23.4%)	23.0-99.5%	16.3% (95%CI[¶]:16.0-16.5%) 12985/79878
Northeast	14.2% (95%CrI: 13.7-14.6%)	14.2-98.1%	9.8% (95%CI: 9.5-10.1%) 3981/40723
Center	11.6% (95%CrI: 11.0-12.1%)	11.6-86.6%	8.1% (95%CI: 7.7-8.5%) 1481/18201
South	12.1% (95%CrI: 11.4-13.0%)	12.1-71.7%	7.9% (95%CrI: 7.4-8.5%) 794/10014
Island	9.6% (95%CrI: 8.4-10.9%)	5.1-18.6%	6.6% (95%CrI: 5.8-7.4%) 227/3455

353 [§]CrI: 95% credibility intervals (CrI), [¶]95%CI: 95% confidence interval

357 Figure legend
358

Figure 1: Temporal distribution of cases and deaths by area due to COVID-19, Italy, March-

April 2020.

Cumulative cases in (A) Northwest, (B) Northeast, (C) Center, (D) South and (E) Island, and

cumulative deaths in (F) Northwest, (G) Northeast (H) Center (I) South and (J) Island, over

time. Day 1 corresponds to March 1st in 2020. As the dates of illness onset were not available,

we used dates of reporting.

Figure 2: Temporal variation of risk of death caused by COVID-19, Italy, March-April 2020.

Observed and posterior estimated of crude case fatality ratio in (A) Northwest, (B) Northeast,

(C) Center, (D) South and (E) Island , and time-delay adjusted case fatality ratio in (F)

Northwest, (G) Northeast (H) Center (I) South and (J) Island. Day 1 corresponds to March 1st

in 2020. Black dots shows crude case fatality ratio, and light and dark indicates 95% and 50%

credible intervals for posterior estimates, respectively.

Figure 3. Latest estimates of time-delay adjusted risk of death caused by COVID-19 by area,

2020, Italy.

Distribution of time-delay adjusted case fatality risks derived from the latest estimates (April

11, 2020) are presented. Top to bottom: Northwest, Northeast, Center, South and Island

377
378
379

Crude case fatality ratio, observed and estimated

A Northwest B Northeast C Center D South E Island

Time adjusted case fatality ratio, crude and time adjusted

F Northwest G Northeast H Center I South J Island

Time delay adjusted CFR by area
The latest estimate