

1 **Impact of time and temperature on gut microbiota and SCFA composition in**
2 **stool samples**

3

4 Janet L. Cunningham^a, Ludvig Bramstång^a, Abhijeet Singh^b, Shishanthi Jayarathna^b, Annica J.
5 Rasmusson^a, Ali Moazzami^b, Bettina Müller^b

6

7

8 ^a Department of Neurosciences, Psychiatry, Uppsala University,

9 ^b Swedish University of Agricultural Sciences, Department of Molecular Sciences,
10 BioCentrum, PO Box7015, SE-75007, Uppsala, Sweden

11

12 Janet.Cunningham@neuro.uu.se

13 Abhijeet.Singh@slu.se

14 Shishanthi.jayarathna@slu.se

15 Annica.Rasmusson@medsci.uu.se

16 Ali.Moazzami@slu.se

17 Bettina.Muller@slu.se

18

19 Corresponding author: Bettina Müller

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34 **Abstract**

35

36 Gut dysbiosis has been implicated in the pathophysiology of a growing number of non-
37 communicable diseases. High through-put sequencing technologies and short chain fatty acid
38 (SCFA) profiling enables surveying of the composition and function of the gut microbiota and
39 provide key insights into host-microbiome interactions. However, a methodological problem
40 with analyzing stool samples is that samples are treated and stored differently prior to
41 submission for analysis potentially influencing the composition of the microbiota and its
42 metabolites. In the present study, we simulated the sample acquisition of a large-scale study, in
43 which stool samples were stored for up to two days in the fridge or at room temperature before
44 being handed over to the hospital. To assess the influence of time and temperature on the
45 microbial community and on SCFA composition in a controlled experimental setting, the stool
46 samples of 10 individuals were exposed to room and fridge temperatures for 24 and 48 hours,
47 respectively, and analyzed using 16S rRNA gene amplicon sequencing, qPCR and nuclear
48 magnetic resonance spectroscopy. To best of our knowledge, this is the first study to investigate
49 the influence of storage time and temperature on the absolute abundance of methanogens, and
50 of *Lactobacillus reuteri*. The results indicate that values obtained for methanogens, *L. reuteri*
51 and total bacteria are still representative even after storage for up to 48 hours at RT (20°C) or 4
52 °C. The overall microbial composition and structure appeared to be influenced more by
53 laboratory errors introduced during sample processing than by the actual effects of temperature
54 and time. Although microbial activity was demonstrated by elevated SCFA at both 4 °C and
55 RT, SCFAs ratios were more stable over the different conditions and may be considered as long
56 as samples are come from similar storage conditions.

57

58

59

60

61

62

63

64

65 **Keywords**

66 Storage condition, Methanogens, *Lactobacillus reuteri*, qPCR, microbial composition, 16s
67 rRNA gene amplicon sequencing, SCFA

68 **Introduction**

69 The gut microbiota (GM) has become of increasing interest as gut dysbiosis has been implicated
70 in the pathophysiology or exacerbation of a growing number of non-communicable diseases
71 including diabetes mellitus, obesity, allergies, rheumatoid arthritis, inflammatory bowel
72 disease, liver disease, colorectal cancer, Parkinson's disease, Alzheimer's disease, multiple
73 sclerosis, autism spectrum disorder, depression and anxiety disorders [1-7]. The gut-brain-axis
74 pertains to the bidirectional communication between gut microbiota and the central nervous
75 system through neural, endocrine, immunological and metabolic means. Short-chain fatty acids
76 (SCFAs) are the main metabolic products from bacterial fermentation in the intestine, and have
77 a key role in microbiota-gut-brain cross talk [8]. The human microbiota consists of at least 1
78 000 species of bacteria with varying compositions and densities at different sites, as well as
79 protozoa, viruses and fungi [9]. DNA sequencing technologies including 16S rRNA gene based
80 amplicon sequencing is the main approach used to study microbial diversity, and to understand
81 the role of the gut microbiome in human health and disease. A methodological problem with
82 analyzing stool samples, however, is that the logistics of collecting samples can vary
83 dramatically between the subjects may influence the composition of the microbiota and its
84 metabolic products. The current understanding is that time and temperature appears to have a
85 low, but not negligible impact on bacterial composition and structure in stool samples [10-15].
86 Immediate freezing of samples at -20 °C is considered to be the "gold standard", but might not
87 always be practically feasible. A number of fecal collection methods including immediate
88 addition of preservation solutions such as ethanol or RNA stabilizers or direct application of
89 commercial kits such as the OMNIgene•GUT have been tested and found to be comparable to
90 the reference of instant freezing at least when analyzing more abundant taxa [11-13, 15, 16].
91 Although most of these collection methods are "participant-friendly", but participants are not
92 always able to follow the instructions. Moreover, a vast amount of stool samples may have been
93 collected and stored in repositories without any immediate preservation. In an ongoing large-
94 scale study, we are aiming to explore the role of the gut microbiota and SCFAs in depression
95 and mood disorders. We have, however, observed differences between the patients in terms of
96 storage conditions before the time point for sample submission. To estimate to which degree
97 different management of samples prior to freezing and without preservation might influence the
98 results of future studies, the effect of different temperatures over time on microbial composition
99 and SCFA profile of stool samples needs to be further investigated.

100 The evidence in the literature on this topic is very limited and, to our knowledge, there is no
101 study done regarding this matter relating specifically to low abundant groups such as

102 methanogens and Lactobacilli. *Lactobacillus* spp. are found in the gastrointestinal systems in
103 variable amounts depending on species and age and has gained increasing attention due to their
104 probiotic attributes [17-19]. The role of methanogens in human health and the possible
105 association to infections and diseases are still poorly understood but a mounting body of
106 evidences suggest a significant role [20-24]. Methanogens play an important role in removing
107 excess hydrogen gas from the gut and improving efficiency of microbial fermentation, but also
108 alter the SCFA production [25]. The impact of time and temperature on the SCFA concentration
109 in stool samples, collected without an immediate preservation method, has been insufficiently
110 studied. Recent studies reported rapid changes of the SCFA level in stool samples stored at RT
111 and several human fecal sample collection protocols have been proposed [15, 26, 27]. So far,
112 no conclusions have been drawn for the consideration of SCFA data originating from stool
113 specimens that could not be obtained in accordance with proposed protocols.

114 The primary aim of this methodological study was to analyze the impact of time and
115 temperature on the absolute abundance of methanogens, *Lactobacillus reuteri*, as well as total
116 bacteria, overall microbial composition and the SCFA profile. This would facilitate the
117 determination of reliable thresholds that can be potentially useful in larger studies, where
118 uniform sample collection conducted at home cannot be guaranteed. For doing so, we have
119 chosen times and temperatures for sample storage before being handed-over to the hospital that
120 are within the range of those reported by some participants in our large-scale studies.

121

122 **Material and Methods**

123 **Ethics and subject recruitment**

124 The study includes 10 subjects: five patients and five controls. Samples from patients were
125 collected from the Uppsala Psychiatric Patient Samples (UPP) Framework approved by the
126 Regional Ethics Committee in Uppsala: (Dnr 2012/81, 2012-03-21 and Dnr 2012/81/1, 2012-
127 12-20 and Dnr 2013/219). To facilitate the collection of fresh samples, patients in this study
128 were recruited from a psychiatric ward where they were undergoing treatment for affective
129 disorders. Written informed consent was obtained from patients for material collection. The
130 results from these samples are not linked to individual factors in this study. The controls in this
131 study were predominantly healthy medical students without any specific inclusion or exclusion
132 criteria and formed a comparatively homogenous group. They contributed to this study
133 anonymously and their data cannot be tied to specific subjects. Participation was voluntary and
134 was not compensated. Control subjects gave verbal informed consent for their donated samples
135 to be analyzed anonymously with the purpose of method development and neither samples nor

136 data cannot be traced back to control individuals. The Regional Ethics Committee waived the
137 need for consent in this case in accordance with Swedish law.

138 **Sample management**

139 The subjects were given instructions to store their sample in fridge or cooler immediately upon
140 acquiring and then initiate contact for transportation. Samples were collected within 4 hours
141 and transported at < 4 °C using a cooling bag with ice clamps. After arrival at the laboratory,
142 the samples were mechanically homogenized with a sterile spatula, aliquoted into five Nunc
143 Cryotube Vials™ and entered into the storing protocol including five storage conditions: the
144 first aliquot was frozen immediately at -20 °C (1), the second was frozen after keeping it 24
145 hours at room temperature (20-21°C) (2), the third after 48 hours at room temperature (3), the
146 fourth after 24 hours at 4 °C (4), and the fifth was frozen after 48 hours at 4 °C (5). DNA was
147 extracted from all frozen samples within 48-96 hours after completion of the storing protocol.

148 **DNA purification**

149 DNA was extracted using the QIAamp® Fast DNA Stool Mini Kit including an additional bead-
150 beating step: 200±1 mg stool was weighed in Lysing Matrix E tubes (MPBiomedicals™) and
151 placed on ice. One ml of InhibitEX Buffer was added and the sample was vortexed continuously
152 for 1 min. The tube was placed in a FastPrep Instrument (MPBiomedicals™) for 40 seconds at
153 speed setting 6.0, and centrifuged at 14 000 x g for 10 minutes. Then, 600 µl of the obtained
154 supernatant was transferred into a new microcentrifuge tube and treated as described in the
155 manufacturer's protocol. The DNA concentration was approximated using Qubit® fluorometer
156 dsDNA protocol (Invitrogen). DNA was purified in triplicates from every storage condition
157 except for three samples due to insufficient material. Out of these three, one was purified in
158 duplicates, and two were purified in triplicates but covering only storage condition 1, 4 and 5.

159 **Quantitative PCR (qPCR)**

160 Absolute quantification using qPCR was performed for total bacteria, methanogenic archaea,
161 and *Lactobacillus reuteri*. Methanogens were quantified using the group specific primers
162 Met630f and Met803r [28], *L. reuteri* by using species specific primers [29] and total bacteria
163 by using the primers Eub338 (ACTCCTACGGGAGGCAGCAG) and Eub518
164 (ATTACCGCGGCTGCTGG) [30, 31]. Before quantification, qPCRs were done with DNA
165 dilutions ranging from 1:10, 1:20, 1:100, 1:500, and 1:1 000 including all primer sets in order
166 to determine the dilution factor required for diminishing inhibitory factors, whilst remaining
167 within the range of standard curve. Quantification of all DNA samples was performed in
168 duplicates using two DNA dilutions (determined before as appropriate) using a Bio-Rad iQ5
169 multicolour real-time PCR detection system and the IQ™ SYBR® Green Supermix (Bio-Rad

170 laboratories, Inc.). qPCR reactions were set up to a final volume of 20 μ L containing the
171 following components: 2x IQTM SYBR[®] Green Supermix, 10 μ M each forward and reverse
172 primers, 3 μ L DNA template and 5 μ L milliQ water. Four to six non-template controls were
173 included in each assay. Plasmid-coded partial 16S rRNA gene originating from methanogens
174 (*Methanoculleus bourgensis*), bacteria (*Escherichia coli*) and *L. reuteri*, respectively were used
175 as standard curves and applied in 10^8 to 10^1 copy numbers. The program used was as follows:
176 initial temperature 95 °C for 7 minutes, followed by 40 cycles at 95 °C for 40 s, 60 °C (*L. reuteri*
177 64 °C) for 60 s (*L. reuteri* 30 s) and 72 °C for 40 s (*L. reuteri* 30 s). The specificity of the PCR
178 product was estimated by melting curve analysis, which consisted of 50 gradual denaturation
179 cycles. The temperature range was set from 55 to 95 °C, dwelled 10 s and increased 0.5 °C in
180 each cycle. PCR products were additionally checked by gel electrophoresis. The data generated
181 were collected and analyzed with Bio-Rad iQ5 standard edition optical system software
182 (version 2.0), from which sorted data were exported to Microsoft Excel for further analysis.
183 Final values can be found in the supplementary file.

184 **Preparation of libraries for Illumina amplicon sequencing**

185 16S rRNA amplicon libraries were constructed as triplicates using two consecutive PCR
186 procedures as described in [32]. The first PCR simultaneously targeted the V4 region of both
187 bacteria and archaea, using the primers 515F
188 (ACACTCTTCCCTACACGACGCTCTTCCGATCTNNNGTGBCAGCMGCCGCGAA)
189 and 805R (AGACGTGTGCTCTTCCGATCTGGACTACHVGGGTWTCTAAT) and attaches
190 adaptors to the amplicons [32]. The reaction mixture contained 2x Phusion High-Fidelity DNA
191 Polymerase/dNTP mix (Thermo Fischer Scientific, Hudson, NH, USA), 10 μ M of each primer,
192 and approx. 10 ng DNA template in a final volume of 25 μ L. The condition for amplification
193 was as following: initial denaturing at 98 °C for 30 s, 20 cycles of 10 s at 98 °C, 30 s at 60 °C,
194 4 s at 72 °C, and a final extension at 72 °C for 2 min. The PCR products were checked for size
195 and quality by electrophoresis. Amplicons were purified using Agencourt AMPure XP (Becker
196 Coulter, Brea, CA, USA), using a magnetic particle/DNA volume ratio of 0.8:1. In the second
197 PCR, Illumina-compatible barcodes were added to the amplicons. The PCR reaction contained
198 10 μ L purified amplicon from the first step, 2x Phusion High-Fidelity DNA Polymerase/dNTP
199 mix and 10 μ M each of the primers
200 5'-AATGATACGGCGACCACCAGATCTACACX8ACACTCTTCCCTACACGACG-3'
201 and 5'-
202 CAAGCAGAAGACGGCATAACGAGATX8GTGACTGGAGTTCAGACGTGTGCTCTTCC
203 GATCT-3', where X8 in the primer sequence represented a specific Illumina-compatible

204 barcode (Eurofins Genomics). The following conditions were used for the second PCR step:
205 initial denaturing at 98 °C for 30 s, 8 cycles of 10 s at 98 °C, 30 s at 62 °C, 5 s at 72 °C, and a
206 final extension at 72 °C for 2 min. The PCR products were checked by electrophoresis and
207 purified using Agencourt AMPure XP. The PCR products were then each diluted to a DNA
208 concentration of approx. 30 nM and pooled together. Pair-end sequencing was performed on
209 the MiSeq platform (Illumina, Inc., San Diego, CA, USA) at Eurofins GATC Biotech GmbH
210 (Konstanz, Germany) resulting in an average of 110 000 raw reads per sample. Illumina
211 adapters and primers have been trimmed away using Cutadapt version 2.2 [33]. All reads shorter
212 than 250 base-pairs (bp), longer than 300 bp or untrimmed were discarded. Amplicon sequence
213 variants, abundancies and taxonomic affiliation were determined using the package *dada2*
214 (version 1.6.0) [34] in R (version 3.4.0), which is implemented on the SLUBI computing cluster
215 in Uppsala (running on CentOS Linux release 7.1.1503; module handling by Modules based on
216 Lua: Version 6.0.1; <https://www.slubi.se/>). Trimming and filtering was jointly performed on
217 paired-end reads. Low quality reads were removed by setting the maximum number of expected
218 errors to 2. The remaining sequence reads were denoised, dereplicated, merged and checked for
219 chimeras in package *dada2* according to the DADA2 pipeline tutorial
220 (https://benjjneb.github.io/dada2/tutorial_1_8.html). Taxonomic classification of the 16S
221 ribosomal RNA sequence variants and a phylogenetic tree were obtained by using the Silva
222 taxonomic training dataset v132 formatted for DADA2
223 (https://zenodo.org/record/1172783/files/silva_nr_v132_train_set.fa.gz; https://zenodo.org/record/1172783/files/silva_species_assignment_v132.fa.gz). A phyloseq object was created
224 consisting of the taxonomy table and the OTU table and used for the visualization with package
225 *phyloseq* (version 1.30.0) [35]. The sequence variants were extracted and a phylogenetic tree
226 was generated using default parameters in FastTree (version 2.1.0) [34]. Sample metadata and
227 phylogenetic tree were merged with the phyloseq object and visually analyzed by using the
228 package *phyloseq* and *ggplot2* (version 3.2.1) [35] in R Studio version 3.5.2 ([http://www.r-](http://www.r-project.org)
229 [project.org](http://www.r-project.org)). Weighted Unifrac distances were calculated by function Unifrac in package
230 *phyloseq* and used to plot the ordination in a Principal coordinate analysis plot with the function
231 *cmdscale* in package *stats* (version 3.6.2) in R core packages. Experimental environmental
232 factors were fitted to the ordination plot by using function *envfit* in package *vegan* (version
233 2.5.6) [36]. The sequencing raw data has been submitted to NCBI sequence read archive (SRA)
234 under the BioProject accession number PRJNA609715.

236

237

238 **Analysis of short chain fatty acids (SCFA)**

239 SCFA has been analyzed using nuclear magnetic resonance (NMR) spectroscopy. Between
240 200-250 mg stool sample was diluted with 0.75 mL sodium phosphate buffer (0.4 M, pH 7.0),
241 homogenized by using a vortex and centrifuged at 6300 rpm at 4 °C for 15 min. From the
242 supernatant, 0.75 mL of fecal water has been transferred to a fresh tube and subjected to
243 centrifugation at 20 000 ×g at 4 °C for 15 min. This step was repeated once with 600 µL of
244 supernatant recovered from the previous centrifugation. Finally, 525 µL of the supernatant was
245 mixed with 45 µL D₂O, and 30 µL internal standard. Each sample solution (560 µL) was
246 transferred to a 5 mm NMR tube, and the ¹H NMR spectra were acquired using a Bruker
247 Avance III spectrometer operating at 600 MHz proton frequency and equipped with a
248 cryogenically cooled probe and an auto sampler. Each spectrum was recorded (25 °C, 128
249 transients, acquisition time 1.8 s, relaxation delay 4 s) using a zgpg30 pulse sequence (Bruker
250 Biospin) using excitation sculpting with gradients for suppression of the water resonance. For
251 each spectrum, 65 536 data points were collected over a spectral width of 17 942 Hz [37]. All
252 NMR spectra were processed using Bruker TopSpin 4.1 software. The data were Fourier-
253 transformed after multiplication by a line broadening of 0.3 Hz and referenced to internal
254 standard peak TSP at 0.0 ppm. Baseline and phase were corrected manually. Each spectrum
255 was integrated using Amix 3.7.3 (Bruker BioSpin GmbH, Rheinstetten, Germany) into 0.01-
256 ppm integral regions between 0.5 and 10 ppm, in which areas between 4.52–5.06 ppm were
257 excluded. Each integral region was referenced to the internal standard. The integral regions
258 corresponding to short chain fatty acids were adjusted for mg of stool samples extracted and
259 used for further statistical analysis. Five samples were analyzed using less amount than 250 mg
260 stool respectively. Absolute numbers and calculation can be found in the supplementary file.

261


262 **Results**

263 **Effect of storage condition on the microbial community composition**

264 Sufficient starting material and high quality raw read data were available to compare microbial
265 changes between immediately frozen and storage for 24 h at RT for 8 individuals; changes
266 between frozen and storage for 48 h at RT for 10 individuals, changes between frozen and
267 storage for 24 h at 4 °C for 7 individuals, and changes between frozen and storage for 48 h at 4
268 °C for 9 individuals. In order to evaluate changes in the microbial structure and composition
269 over time both alpha and beta diversity have been analysed. The alpha diversity indices Shannon
270 and Simpson indicated small changes in the community structure for most of the samples
271 (Figure 1). Larger variations were observed for individuals E, F, H. However, the variance in

272 alpha diversity within the triplicates was in part much higher than that observed between
273 aliquots of different storage conditions (Figure S1).

274



275

276 Figure 1: Variation in alpha diversity (Simpson and Shannon) displayed as box plot of
277 individuals A-J. Extended lines indicate variability outside the upper and lower quantity of the
278 box, whereas outliers are plotted as individual points. Blue: immediately frozen samples,
279 orange: 4 °C, red: 20 °C.


280

281

282 Principal coordinate analysis (PCoA) using the weighted UniFrac matrix was used to compare
283 the beta diversity. Most individuals showed variations in the community composition when
284 comparing the differently stored stool samples. Although the majority of them did not form
285 clear clusters with regard to the different storage temperatures, the PCoA revealed little
286 influence of time and temperature on the microbial composition, as indicated by the arrows
287 (Figure 2).

288

289



290

291

292 Figure 2: PCoA plot of weighted UniFrac distance. Principal coordinate analysis displaying
 293 beta diversity of the bacterial communities in the stool samples of 10 individuals. blue: frozen,
 294 orange: 4 °C, red: 20 °C. Ellipses indicate the individuals.

295

296 **Effect of storage condition of absolute abundance of total bacteria, methanogens and**
 297 ***Lactobacillus reuteri***

298 Generally, only marginal variations of the 16S rRNA gene copy numbers were observed in case
 299 of all three targets, independently if frozen immediately or stored at 4 °C or RT for up to 48 h
 300 (Figure 3, 4). The mean average deviations from the immediately frozen sample were between
 301 1.6 and 2.3 % (± 2.5 - 3.2) in case of methanogens, 0.2 and 0.8 % (± 0.9 - 1.8) in case of total
 302 bacteria, and 2.0 and 3.2 % (± 2.2 - 2.8%) in case of *L. reuteri* (Table 1).

303

304 Table 1. Mean average deviation of absolute 16S rRNA gene copies in percent from initial
 305 values obtained from untreated (immediately frozen) samples. Detailed list of values can be
 306 found in Table S1.

%	4°C_24h	4°C_48h	RT_24h	RT_48h
Methanogens	2.0 \pm 3.2	2.3 \pm 3.2	1.6 \pm 2.5	2.3 \pm 2.5
Total bacteria	0.7 \pm 1.8	0.2 \pm 1.6	0.5 \pm 0.9	0.8 \pm 1.5
<i>L. reuteri</i>	2.0 \pm 2.2	2.9 \pm 2.8	2.5 \pm 2.2	3.2 \pm 2.3

307

308



309

310

311 Figure 3: Absolute 16S rRNA gene copy numbers of total methanogens (A) and total bacteria
312 (B) from stool samples retrieved from individuals A-J, stored at the conditions as indicated in
313 the legend. Frozen stands for immediately frozen at -20 °C.

314


315 *L. reuteri* was found in sufficiently high numbers in only four out of ten samples to be above
316 the detection limit of qPCR and to exclude inhibitory effects (Figure 4). No significant effect
317 of storage time or temperature on the absolute abundance of methanogens, total bacteria or *L.*
318 *reuteri* was observed (Figure 3, 4, Table 1).

319

320 Effect of storage condition on the SCFA acetate, propionate and butyrate

321 Five stool samples were subjected to SCFA analysis using NMR. Storage at RT had a dramatic
322 effect on levels of acetate, propionate and butyrate (Figure S2). Already the storage for 24 hours
323 promoted an increase of all three SCFA to more than 100 % in some cases when compared to
324 values obtained from immediately frozen stool sample. Acetate values partly increased by more
325 than 150 %. The mean average deviation from the initial value (immediately frozen) for acetate,
326 propionate and butyrate were between 94 to 105 % (± 35 %-45 %) after 48 h (Table 2).
327 However, also storage at 4 °C led to an increase of all three SCFA in four out of five samples
328 albeit significantly lower (Figure S2). In one sample, the concentration of SCFAs dropped

329 during storage at 4 °C. The mean average deviation for all SCFA from initial values was
 330 between 14 and 20 % ($\pm 10-19$ %) after 24 h at 4 °C considering both increase and decrease
 331 (Table 2).
 332



333
 334
 335 Figure 4: Absolute 16S rRNA gene copy numbers of *L. reuteri* from stool samples retrieved
 336 from individuals B, E, G, and I, stored at the conditions as indicated in the legend. Frozen stands
 337 for immediately frozen at -20 °C.

338
 339 Table 2. Average deviation \pm standard deviation of SCFA in percent from initial values
 340 obtained from immediately frozen samples. Detailed list can be found in the supplementary file.

	frozen -> 4°C_24h	frozen -> 4°C_48h	frozen -> RT_24h	frozen -> RT_48h
Acetate	20 \pm 10	28 \pm 18	99 \pm 67	105 \pm 36
Butyrate	14 \pm 19	18 \pm 9	67 \pm 45	94 \pm 45
Propionate	15 \pm 13	21 \pm 12	76 \pm 30	102 \pm 38
Butyrate/Acetate	19 \pm 8	17 \pm 6	15 \pm 13	19 \pm 7
Butyrate/Propionate	9 \pm 3	12 \pm 10	30 \pm 28	18 \pm 20
Butyrate/(Acetate+Propionate)	18 \pm 7	15 \pm 6	14 \pm 12	16 \pm 6
Acetate/(Acetate+Propionate+Butyrate)	3 \pm 1	3 \pm 2	3 \pm 3	4 \pm 1
Propionate/(Acetate+Propionate+Butyrate)	7 \pm 2	7 \pm 4	10 \pm 8	9 \pm 4
Butyrate/(Acetate+Propionate+Butyrate)	17 \pm 7	15 \pm 6	14 \pm 11	15 \pm 6

341
 342
 343 Storage up to 48 h resulted in slightly further increase of the mean average deviation (Table 2).
 344 However, when calculating SCFA ratios the picture changed (Figure 5): For the majority of the
 345 calculated ratios the mean standard deviation remains below 20 % when compared to the
 346 respective ratios obtained for the immediately frozen sample (Table 2, Figure 5). The average


347 deviation from the initial value was least pronounced in those ratios containing more than one
348 SCFA in the divisor. The ratios acetate/(acetate+propionate+butyrate) and
349 propionate/(acetate+propionate+butyrate) differ by less than 10 % from the ratios obtained from
350 immediately frozen samples (Table 2, Figure 5). Most important and in contrast to the absolute
351 values, the majority of the SCFA ratios indicate a much less pronounced impact of increased
352 temperature or storage time on the SCFA levels. The observed variations in the SCFA ratios
353 remain comparatively independent of temperature and how long a sample was stored before
354 extraction.

355

356 **Discussion**

357 It is generally not recommended to store fecal samples at RT or longer than 12 h at
358 4 °C since metabolism and proliferation of some bacteria might continue, with consequences
359 on microbial structure and SCFA profile. Moreover, the intake of oxygen might damage strictly
360 anaerobic microorganisms [38]. A number of attempts have been made to determine the optimal
361 collection method to preserve microbial community and metabolic composition. Direct freezing
362 at -80 °C is considered the reference method, however is not feasible for many settings. Instead,
363 a -20°C freezer is available in most homes and freezing at -20 °C may also be applied [38, 39]
364 but this requires willingness to place samples in the freezer and resources for shipping the
365 samples in a frozen state. Several studies reported positive effects of preservation media and
366 stabilizers [11-16]. However, even if patients are asked to describe how samples were handled
367 before submission to the laboratory, it is difficult to have complete control over what has
368 happened and samples, for diverse reasons, may have been exposed to room temperature for
369 hours before reaching the laboratory. There are only few studies that looked at the effects of
370 short-term storage conditions on unpreserved microbial community and those came to different
371 conclusions [10, 12, 39-41].

372



373

374

375 Figure 5: Deviation of SCFA ratios in % from those ratios obtained from the immediately frozen
 376 samples displayed as box plots. Blue: samples stored at 4 °C up to 48 h, red: samples stored at
 377 20 °C up to 48 h. Extended lines indicate variability outside the upper and lower quantity of the
 378 box, whereas outliers are plotted as individual points. Ratios are also displayed in Figure S2.
 379 Calculation and numbers can be found in the supplementary file.

380

381 Ott et al reported a significant reduction of both bacterial diversity and total number of
 382 bacteria after already 8 h at both 4 °C and RT and Cardona et al observed alterations in the
 383 relative abundance across most taxa when fecal sample were subjected to RT for 24 h [39, 41].
 384 On the other hand, two studies reported only minor alterations in taxa abundances when stored
 385 at both 4 °C or RT for up to 24 h [10, 12]. Lauber et al. reported stability of the microbiota even
 386 for up to 14 days at 4 °C and 20 °C [40].

387

388 In the present study, we simulated the sample acquisition of a large-scale study, in which
 389 some patients reported storing stool samples for up to two days in the fridge or at room
 390 temperature before handed over to the hospital. To our knowledge, this is the first study
 391 evaluating the impact of storage time and temperature on the absolute abundance of
 392 methanogens, and of *L. reuteri* in particular. Methanogens are extremely oxygen sensitive and
 even trace amount of oxygen causes stress to most of methanogenic species by damaging the

393 cell membrane and proteins [42]. However, we could demonstrate that the absolute abundance
394 of Methanogens appeared stable at both RT and 4 °C for up to 48 h. Only small variation
395 occurred, which did not correlate to time or temperature. The same findings were observed for
396 *Lactobacillus reuteri*, which is facultative anaerobe and might be therefore able to proliferate
397 under certain conditions. Alpha diversity indices and PCoA plots indicated small changes in the
398 overall microbial community, when stored at 4 °C or RT for up to 24 h, which is in agreement
399 with the findings of Ott and Carroll [10, 41]. We also found that even a prolonged storage up to
400 48 h did not significantly alter the microbial composition any further. The variations observed
401 within the triplicates indicates that other factors including differences in sample handling, PCR
402 and sequencing may also account for the observed alterations in the microbial community. This
403 is further supported by the finding that the total number of bacteria remained stable over time.
404 This is contradictory to Ott et al who reported a dramatic reduction of total bacteria, however
405 the different techniques used might not allow a direct comparison of the results [41]. Although
406 the composition and structure of the microbial community has only changed little over time,
407 the community still appeared to be active, especially at RT. The absolute levels of acetate,
408 propionate and butyrate increased dramatically within 24 hours, indicating general metabolic
409 activities. Even storage at 4 °C could not completely suppress metabolic activities, but proved
410 to be clearly beneficial. Interestingly enough, the effect of time and temperature was strongly
411 diminished when looking at ratios instead of absolute values. Obviously, the SCFAs increase
412 almost proportionally with time at both 4 °C and in RT, so that the factors time and temperature
413 became irrelevant. This means that samples collected at then cooled or at RT can be compared
414 to each other, but not with immediately frozen samples and vice versa. The deviations from the
415 immediately frozen samples were only negligible for the ratio
416 acetate/(acetate+propionate+butyrate). A direct comparison with frozen samples should only
417 be considered taking into account the threshold values identified here as mean deviations.
418 Several other studies show that the immediate preservation by e.g. ethanol or RNAlater can
419 replace an immediate freezing of the samples [15, 43]. However, this may not always be
420 feasible. The results found here can thus support those studies that cannot implement immediate
421 freezing or stabilizing methods, but still want to integrate metabolomics.

422

423

424

425

426

427 **Conclusions**

428 By analyzing the impact of storage time and temperature, we conclude that realistic values for
429 the absolute abundance of methanogens, *L. reuteri* and total bacteria can still be obtained even
430 after storage for up to 48 hours at RT or 4 °C. The overall microbial composition appears to be
431 influenced more by laboratory error introduced during sample processing than by the actual
432 effects of storage temperature and time. Although significant microbial activities have been
433 demonstrated at both 4 °C and RT, the SCFAs may be considered as long as these are taken
434 into account as ratios and originated from similar storage conditions.

435

436 **Acknowledgement**

437 We acknowledge the National Genomics Infrastructure (NGI)-SciLifeLab in Uppsala for
438 carrying out Illumina sequencing and the SLU Bioinformatics Infrastructure (SLUBI) for
439 providing the computing cluster for analyzing sequencing data. Funding for this study was
440 obtained from the Ekhaga foundation, The Swedish Society of Medicine; and ALF Funds from
441 Uppsala University Hospital.

442

443 **Supporting information**

444 Data obtained from short chain fatty acid and qPCR analysis can be found in the supplementary
445 file “S1_Appendix”.

446

447 **References**

448

- 449 1. Cong J, Zhang X: **How human microbiome talks to health and disease.** *European*
450 *journal of clinical microbiology & infectious diseases* : official publication of the
451 *European Society of Clinical Microbiology* 2018, **37**(9):1595-1601.
- 452 2. Dwivedi M, Ansarullah, Radichev I, Kemp EH: **Alteration of Immune-Mechanisms**
453 **by Human Microbiota and Development and Prevention of Human Diseases.**
454 *Journal of immunology research* 2017, **2017**:6985256.
- 455 3. Kang DW, Adams JB, Gregory AC, Borody T, Chittick L, Fasano A, Khoruts A, Geis
456 E, Maldonado J, McDonough-Means S *et al*: **Microbiota Transfer Therapy alters**
457 **gut ecosystem and improves gastrointestinal and autism symptoms: an open-label**
458 **study.** *Microbiome* 2017, **5**(1):10.

- 459 4. Kushak RI, Winter HS: **Intestinal microbiota, metabolome and gender**
460 **dimorphism in autism spectrum disorders.** *Research in Autism Spectrum Disorders*
461 2018, **49**:65-74.
- 462 5. Szablewski L: **Human Gut Microbiota in Health and Alzheimer's Disease.** *Journal*
463 *of Alzheimer's disease : JAD* 2018, **62**(2):549-560.
- 464 6. Yang Y, Tian J, Yang B: **Targeting gut microbiome: A novel and potential therapy**
465 **for autism.** *Life sciences* 2018, **194**:111-119.
- 466 7. Zheng P, Zeng B, Zhou C, Liu M, Fang Z, Xu X, Zeng L, Chen J, Fan S, Du X *et al*:
467 **Gut microbiome remodeling induces depressive-like behaviors through a**
468 **pathway mediated by the host's metabolism.** *Molecular psychiatry* 2016, **21**(6):786-
469 796.
- 470 8. Clarke G, Stilling RM, Kennedy PJ, Stanton C, Cryan JF, Dinan TG: **Minireview:**
471 **Gut microbiota: the neglected endocrine organ.** *Molecular endocrinology*
472 *(Baltimore, Md)* 2014, **28**(8):1221-1238.
- 473 9. Gilbert JA, Blaser MJ, Caporaso JG, Jansson JK, Lynch SV, Knight R: **Current**
474 **understanding of the human microbiome.** *Nature medicine* 2018, **24**(4):392-400.
- 475 10. Carroll IM, Ringel-Kulka T, Siddle JP, Klaenhammer TR, Ringel Y:
476 **Characterization of the fecal microbiota using high-throughput sequencing**
477 **reveals a stable microbial community during storage.** *PLoS ONE* 2012,
478 7(10):e46953.
- 479 11. Flores R, Shi J, Yu G, Ma B, Ravel J, Goedert JJ, Sinha R: **Collection media and**
480 **delayed freezing effects on microbial composition of human stool.** *Microbiome*
481 2015, **3**:33.
- 482 12. Penington JS, Penno MAS, Ngui KM, Ajami NJ, Roth-Schulze AJ, Wilcox SA,
483 Bandala-Sanchez E, Wentworth JM, Barry SC, Brown CY *et al*: **Influence of fecal**
484 **collection conditions and 16S rRNA gene sequencing at two centers on human gut**
485 **microbiota analysis.** *Scientific reports* 2018, **8**(1):4386.
- 486 13. Szopinska JW, Gresse R, van der Marel S, Boekhorst J, Lukovac S, van Swam I,
487 Franke B, Timmerman H, Belzer C, Arias Vasquez A: **Reliability of a participant-**
488 **friendly fecal collection method for microbiome analyses: a step towards large**
489 **sample size investigation.** *BMC Microbiol* 2018, **18**(1):110.
- 490 14. Vogtmann E, Chen J, Amir A, Shi J, Abnet CC, Nelson H, Knight R, Chia N, Sinha R:
491 **Comparison of Collection Methods for Fecal Samples in Microbiome Studies.**
492 *American journal of epidemiology* 2017, **185**(2):115-123.

- 493 15. Wang Z, Zolnik CP, Qiu Y, Usyk M, Wang T, Strickler HD, Isasi CR, Kaplan RC,
494 Kurland IJ, Qi Q *et al*: **Comparison of Fecal Collection Methods for Microbiome**
495 **and Metabolomics Studies**. *Frontiers in cellular and infection microbiology* 2018,
496 **8**:301.
- 497 16. Voigt AY, Costea PI, Kultima JR, Li SS, Zeller G, Sunagawa S, Bork P: **Temporal**
498 **and technical variability of human gut metagenomes**. *Genome Biol* 2015, **16**:73.
- 499 17. Duar RM, Lin XB, Zheng J, Martino ME, Grenier T, Perez-Munoz ME, Leulier F,
500 Ganzle M, Walter J: **Lifestyles in transition: evolution and natural history of the**
501 **genus Lactobacillus**. *FEMS Microbiol Rev* 2017, **41**(Supp_1):S27-s48.
- 502 18. Giraffa G, Chanishvili N, Widyastuti Y: **Importance of lactobacilli in food and feed**
503 **biotechnology**. *Research in microbiology* 2010, **161**(6):480-487.
- 504 19. Mu Q, Tavella VJ, Luo XM: **Role of Lactobacillus reuteri in Human Health and**
505 **Diseases**. *Front Microbiol* 2018, **9**:757.
- 506 20. Attaluri A, Jackson M, Valestin J, Rao SS: **Methanogenic flora is associated with**
507 **altered colonic transit but not stool characteristics in constipation without IBS**.
508 *The American journal of gastroenterology* 2010, **105**(6):1407-1411.
- 509 21. Bang C, Weidenbach K, Gutschmann T, Heine H, Schmitz RA: **The intestinal archaea**
510 **Methanosphaera stadtmanae and Methanobrevibacter smithii activate human**
511 **dendritic cells**. *PLoS ONE* 2014, **9**(6):e99411.
- 512 22. Pimentel M, Mayer AG, Park S, Chow EJ, Hasan A, Kong Y: **Methane production**
513 **during lactulose breath test is associated with gastrointestinal disease**
514 **presentation**. *Digestive diseases and sciences* 2003, **48**(1):86-92.
- 515 23. Weaver GA, Krause JA, Miller TL, Wolin MJ: **Incidence of methanogenic bacteria**
516 **in a sigmoidoscopy population: an association of methanogenic bacteria and**
517 **diverticulosis**. *Gut* 1986, **27**(6):698-704.
- 518 24. Zhang H, DiBaise JK, Zuccolo A, Kudrna D, Braidotti M, Yu Y, Parameswaran P,
519 Crowell MD, Wing R, Rittmann BE *et al*: **Human gut microbiota in obesity and**
520 **after gastric bypass**. *Feb 15* 2009, **106**(7):2365-2370.
- 521 25. Abell GCJ, Conlon MA, McOrist AL: **Methanogenic archaea in adult human faecal**
522 **samples are inversely related to butyrate concentration**. *Microbial Ecology in*
523 *Health and Disease* 2006, **18**(3-4):154-160.
- 524 26. Liebisch G, Ecker J, Roth S, Schweizer S, Öttl V, Schött HF, Yoon H, Haller D,
525 Holler E, Burkhardt R *et al*: **Quantification of Fecal Short Chain Fatty Acids by**

- 526 **Liquid Chromatography Tandem Mass Spectrometry-Investigation of Pre-**
527 **Analytic Stability.** *Biomolecules* 2019, **9**(4).
- 528 27. Gratton J, Phetcharaburanin J, Mullish BH, Williams HRT, Thursz M, Nicholson JK,
529 Holmes E, Marchesi JR, Li JV: **Optimized Sample Handling Strategy for**
530 **Metabolic Profiling of Human Feces.** *Analytical chemistry* 2016, **88**(9):4661-4668.
- 531 28. Christophersen CT: **Ph.D.** *University of Western Australia, Perth* 2007.
- 532 29. Matsuda K, Tsuji H, Asahara T, Matsumoto K, Takada T, Nomoto K: **Establishment**
533 **of an analytical system for the human fecal microbiota, based on reverse**
534 **transcription-quantitative PCR targeting of multicopy rRNA molecules.** *Appl*
535 *Environ Microbiol* 2009, **75**(7):1961-1969.
- 536 30. Lane DJ: **16S/23S rRNA sequencing.** *Stackebrandt, E and Goodfellow, M, Eds,*
537 *Nucleic Acid Techniques in Bacterial Systematic, John Wiley and Sons, New York*
538 1991:115-175.
- 539 31. Muyzer G, de Waal EC, Uitterlinden AG: **Profiling of complex microbial**
540 **populations by denaturing gradient gel electrophoresis analysis of polymerase**
541 **chain reaction-amplified genes coding for 16S rRNA.** *Appl Environ Microbiol*
542 1993, **59**(3):695-700.
- 543 32. Hugerth LW, Wefer HA, Lundin S, Jakobsson HE, Lindberg M, Rodin S, Engstrand
544 L, Andersson AF: **DegePrime, a program for degenerate primer design for broad-**
545 **taxonomic-range PCR in microbial ecology studies.** *Appl Environ Microbiol* 2014,
546 **80**(16):5116-5123.
- 547 33. Martin M: **Cutadapt removes adapter sequences from high-throughput**
548 **sequencing reads.** *2011* 2011, **17**(1):3.
- 549 34. Price MN, Dehal PS, Arkin AP: **FastTree 2--approximately maximum-likelihood**
550 **trees for large alignments.** *PLoS ONE* 2010, **5**(3):e9490.
- 551 35. Wickham H: **ggplot2: Elegant Graphics for Data Analysis.** *Journal of Statistical*
552 *Software* 2016.
- 553 36. Jari Oksanen FGB, Michael Friendly, Roeland Kindt, Pierre Legendre, Dan McGlenn,
554 Peter R. Minchin, R. B. O'Hara, Gavin L. Simpson, Peter Solymos, M. Henry H.
555 Stevens, Eduard Szoecs, Helene Wagner: 2018, **vegan: Community Ecology**
556 **Package.**
- 557 37. Rohnisch HE, Eriksson J, Mullner E, Agback P, Sandstrom C, Moazzami AA: **AQuA:**
558 **An Automated Quantification Algorithm for High-Throughput NMR-Based**

559 **Metabolomics and Its Application in Human Plasma.** *Analytical chemistry* 2018,
560 **90(3):2095-2102.**

561 38. Thomas V, Clark J, Dore J: **Fecal microbiota analysis: an overview of sample**
562 **collection methods and sequencing strategies.** *Future microbiology* 2015,
563 **10(9):1485-1504.**

564 39. Cardona S, Eck A, Cassellas M, Gallart M, Alastrue C, Dore J, Azpiroz F, Roca J,
565 Guarner F, Manichanh C: **Storage conditions of intestinal microbiota matter in**
566 **metagenomic analysis.** *BMC Microbiology* 2012, **12(1):158.**


567 40. Lauber CL, Zhou N, Gordon JI, Knight R, Fierer N: **Effect of storage conditions on**
568 **the assessment of bacterial community structure in soil and human-associated**
569 **samples.** *FEMS Microbiol Lett* 2010, **307(1):80-86.**

570 41. Ott SJ, Musfeldt M, Timmis KN, Hampe J, Wenderoth DF, Schreiber S: **In vitro**
571 **alterations of intestinal bacterial microbiota in fecal samples during storage.**
572 *Diagnostic microbiology and infectious disease* 2004, **50(4):237-245.**


573 42. Jarrell KF: **Extreme Oxygen Sensitivity in Methanogenic Archaeobacteria.**
574 *BioScience* 1985, **35(5):298-302.**

575 43. Loftfield E, Vogtmann E, Sampson JN, Moore SC, Nelson H, Knight R, Chia N, Sinha
576 **R: Comparison of Collection Methods for Fecal Samples for Discovery**
577 **Metabolomics in Epidemiologic Studies.** *Cancer epidemiology, biomarkers &*
578 *prevention : a publication of the American Association for Cancer Research,*
579 *cosponsored by the American Society of Preventive Oncology* 2016, **25(11):1483-**
580 **1490.**

581
582
583
584
585
586
587
588
589
590
591
592




593



594

595 Figure S1: Alpha diversity (Simpson and Shannon) of the replicates displayed as box plot at the
 596 different time points and temperatures analyzed. Individuals are indicated by different colors
 597 and capitals A-J. Extended lines indicate variability outside the upper and lower limits of the
 598 box, whereas outliers are plotted as individual points.

599



600

601

602 Figure S2: Effect of time and temperature on SCFA concentrations and SCFA ratios displayed

603 as average deviation from the initial values obtained from immediately frozen samples.

604

605

606

607