

Abstract

Background: Autistic people experience significantly poorer physical and mental health along with reduced life expectancy.

Aim: To identify self-reported barriers to primary care by autistic adults compared to parents of autistic children and non-autistic adults and link these barriers to self-reported adverse health consequences.

Design and Setting: Following consultation with the autistic community at an autistic conference, Autscope, a quantitative and qualitative survey was developed.

Method: The self-report survey was administered online through social media platforms.

Results: The 57-item online survey was completed by 507 autistic adults, 196 parents of autistic children and 157 control subjects. 79.7% of autistic adults, 52.8% of parents and 36.5% of controls reported difficulty visiting a GP. The highest-rated barriers by autistic adults were deciding if symptoms warrant a GP visit (72.2%), difficulty making appointments by telephone (61.9%), not feeling understood (55.8%), difficulty communicating with their doctor (53.1%) and the waiting room environment (50.5%).

Autistic adults reported a preference for online or text based appointment booking, facility to email in advance the reason for consultation, first or last clinic appointment and a quiet place to wait.

Increased adverse health outcomes reported by autistic adults correlated with difficulty attending, including untreated physical and mental health conditions, not attending specialist referral or screening programmes, requiring more extensive treatment or surgery due to late presentations, and untreated potentially life threatening conditions.

Conclusion: Reduction of healthcare inequalities for autistic people requires that healthcare providers understand autistic perspectives and communication needs. Adjustments for autism specific needs are as necessary as ramps for wheelchair users.

How this fits in:

- Adverse health outcomes are common among autistic people and so it is important to understand how we can promote access to primary care.
- This cross sectional study indicates that 79.7% of autistic patients (compared to 36.5% of controls) reported difficulty visiting a GP.
- Common barriers were: deciding if symptoms warrant a GP visit, difficulty using the telephone to book appointments, not feeling understood and difficulty communicating with their doctor.

- Common suggestions to promote access included: online or text based appointment booking facility, emailing in advance the reason for consultation, providing first or last clinic appointment and having a quiet place to wait.

Introduction

Autism is a common neurodevelopmental condition affecting 1-2% of the population[1]. Most autistic people are adult, do not have intellectual disability and are likely to be undiagnosed[2]. Doctors may underestimate of the number of autistic patients under their care[3,4].

Autistic adults have poor physical and mental health compared to the general population[5]. Most medical conditions are more common[6,7], including diabetes, hypertension and obesity[8]. Autistic people experience premature mortality[9,10,11]. Life expectancy is reduced by 16-30years, with increased mortality reported across almost all diagnostic categories[9] and in-hospital mortality is increased[12]. Autistic people are three times more likely to use emergency departments, to require inpatient admission, and to die after attending emergency care[13].

Alongside increased health needs, autistic people report a greater likelihood that their needs are unmet[14]. Pervasive, multifactorial barriers to healthcare access are experienced[15]. Some are shared by other disabled people, but autistic patients experience additional autism-specific barriers[16]. A recent systematic review identified patient-provider communication, sensory sensitivities, and executive functioning/planning difficulties as important barriers to healthcare[17]. Negative experiences with healthcare providers was separately identified as a prominent barrier[18].

In response to primary legislation[19] and statutory guidance[20] The Royal College of General Practitioners (RCGP) developed an Autism Patient Charter[21] that recommended: staff awareness and training; autism friendly environment; reasonable adjustments following disclosure or clinical suspicion of autism; patient-tailored communications; and behaviour-sensitive accommodations. In spite of efforts to champion autism, proposals to formalise autism training[18,22] and specific awareness-raising interventions[21], almost 40% of GPs report no formal training in autism and have limited confidence in managing autistic patients[22]. Greater autism awareness exists where GPs have personal knowledge of autism, either through a relative or friend on the autistic spectrum, or because they themselves are autistic[22].

Communication skills training for health care providers is the most pressing need[4]. GPs[22] and hospital specialists[3] self-report difficulties communicating with autistic patients. Only 25% of primary healthcare providers reported high confidence in communicating with autistic adult patients, or identifying and making necessary accommodations[4].

This study aimed to identify self-reported barriers to primary health care faced by autistic adults with a focus on autism-specific communication, sensory and procedural considerations. Self-reported consequences were captured to add a narrative frame to the existing evidence base around disparities in health outcomes. Since non-autistic parents of

autistic children experience pronounced increase in morbidity and mortality related to cancer and vascular diseases compared to parents of non-autistic children, this paper further illuminates the barriers experienced by that group[23]. This is to our knowledge the largest study of primary healthcare barriers to date and benefits from a high degree of participatory design by the autistic community making a unique and substantial contribution to the current literature.

Methods

An initial survey was undertaken in August 2018 at Autscope, an annual 3-day residential conference organized by and for autistic people[24]. This was undertaken as a quality improvement project for an “Autism Friendly Town” initiative by AslAm, Ireland’s National Autism Charity[25,26] to inform autism training for local healthcare providers. A 16-item questionnaire, entitled “What do you wish your GP knew about autism?” was distributed in paper and online formats containing open-ended questions and open-text response fields, allowing for qualitative answers. Participants at Autscope included autistic attendees, non-autistic family members and carers, who were capable of completing the survey. The majority of Autscope attendees typically have low to moderate support needs. Seventy-five responses were received, reviewed (by MD) and grouped thematically under broad themes. Thematic analysis was pragmatic, based on face validity rather than a formalised qualitative analysis.

Next a 57-item online survey was developed to investigate barriers to primary care in a larger sample of autistic adults and compare with parents of autistic children and a non-autistic adult control group.

The survey was adapted for 3 groups of respondents: 1. Autistic adults, including those who were parents (Autistic); 2. Parents of autistic children, who did not identify as autistic themselves (Parent); 3. Non autistic adults who did not have an autistic child (Control). All were asked to respond specifically about seeking healthcare for themselves, not for their children. Respondents were directed to the appropriate survey however a small number(n=55) answered the incorrect survey. As the questions were the same it was possible to assign their survey responses to the appropriate group.

Question types included yes-no responses, single- and multiple-item selections from a list, Likert scales, and open-ended short answer. Questions explored the specific barriers encountered and the consequences of such barriers, in particular the reasons for delaying or avoiding a visit, difficulties booking, planning and waiting for a GP visit, challenges during a consultation, including communication, sensory, emotional and organisation issues as well as social supports available. Reasonable adjustments to standard care and access were also explored as well as self-reported consequences of failure to access healthcare.

Participants were recruited through social media, primarily Twitter and Facebook, via both personal pages and in autism related groups and shared on the AslAm website.

Survey data were first collected between October 2018 and April 2019. Reminders were given by repeat posting on social media platforms. Preliminary analysis revealed a recurring

theme of total non-engagement with healthcare providers, despite expressed healthcare needs. Consequently the survey was altered to add response options “I do not attend any medical practice?” and “I do not attend any doctor” to the questions “Do you usually attend the same medical practice?” and “Do you usually see the same doctor?”. The amended survey was run again over a two-week period in August 2019. Due to unquantifiable duplication of responses between the first and second data collection periods, only the data from the second collection period are presented here. Analysis revealed good correlation between both sets of results.

Significance of between-group differences on categorical variables was assessed using 2x2 or 2xk Chi-squared test of differences between paired groups: autistic adults with control participants, autistic adults with parents, and parents with control participants. A Wilcoxon-Mann-Whitney U test was similarly employed on ordinal scales.

Results

Participants

860 respondents completed the online survey: 507 autistic adults, 196 non-autistic parents of autistic children and 157 ‘control’ adults (Table 1).

Barriers to access

79.7% of autistic respondents reported difficulty visiting a GP when needed, compared to 52.8% of parents and 36.5% of controls. See Table 2 for specific barriers.

Communication

Factors causing an autistic patient to delay or avoid seeing their doctor when required included: difficulty using the telephone to book an appointment (61.9%); not feeling understood (55.8%); difficulty communicating with the doctor (53.1%) and with reception staff (46.4%). 59% of autistic respondents reported difficulty communicating during a consultation “all the time” or “frequently” compared to 11.3% of parents and 11.5% of controls ($p < 0.001$).

Autistic respondents reported avoiding the telephone (77.9%), voicemail (61.3%) and face-to-face verbal communication (30.0%). 41.0% reported that it is “easier for me to communicate in writing” (Table 3).

Predictability & control

Autistic respondents reported difficulty not knowing the wait duration (70.0%), what would happen during the consultation (63.1%), which doctor they would see (57.6%) and the consultation length (39.6%). Parents had difficulty with unknown waiting time (49.0%) (Supplementary Table 1).

Sensory processing

The waiting room environment was a barrier for 50.5% of autistic respondents. Sensory issues make communication more difficult for 30.8% of the autistic group (Table 2). Only 10.1% of autistic respondents marked “none of the above” to sensory questions compared to 62.2% of parents and 71.3% of controls (Table 4).

Judgement & attitudes

Autistic people experience more anxiety accessing healthcare; only 3.4% stated they didn't feel anxious going to the doctor, compared to 26.6% of parents and 33.1% of controls. Autistic patients report being "concerned I won't be taken seriously when I describe my symptoms" (67.3%); worried about "wasting the doctor's time" (65.7%) and "being considered a hypochondriac" (65.1%); experience difficulty "asking for help" (62.7%) and "discussing mental health" (59.4%) (Supplementary Table 2).

Disclosure of diagnosis

Only 61.5% of autistic patients reported their doctor knew they were autistic, 22.4% were unsure and 16.1% hadn't disclosed their diagnosis. In contrast, 88.3% of parents reported their doctor knew their child's diagnosis.

Planning & Organising

Autistic respondents reported greater difficulties with organisation and planning for healthcare, including difficulties "making an appointment in advance" (59.2%), "prioritising my health issues" (58.4%) and "making changes to my lifestyle or habits" (55.6%). 45.4% reported forgetting a medical appointment and 29.8% attended on the wrong day. Parents (41.8%) reported significantly less difficulty deciding if symptoms warrant a GP visit than autistic (72.2%) and control (65.0%) respondents. (Supplementary Table 3).

Support needs

Autistic adults more frequently reported physical mobility needs (15.9%), and unmet support needs such as "needing a support person to come with me" (20.9%). This extended to secondary care: 17.4% had no one to support hospital visits, collection from hospital (19.5%), or home care following discharge (25.8%). "Not having anyone to look after my child" was a greater barrier for parents (37.2%) than autistic (13.0%) or control (11.5%) groups (Supplementary Table 4).

Adverse consequences

Autistic respondents reported adverse consequences of these barriers more frequently than parents or controls, including untreated mental (69.0%) and physical (62.8%) health conditions or "did not attend referral to a specialist" (47.1%). Notably a large proportion were told they "should have seen a doctor sooner" (59.9%), "required more extensive treatment or surgery" (35.9%), or did not access treatment for a "potentially serious or life threatening condition" (34.2%). Additionally, they were significantly less likely to "attend on schedule for screening programmes" than the other two groups (Figure 1).

Compared to autistic respondents who had no difficulty visiting a doctor, those who experienced difficulty reported more untreated mental and physical health conditions ($p < 0.001$), not attending specialist referral ($p < 0.001$), needing more extensive treatment ($p = 0.009$), untreated life-threatening conditions ($p = 0.006$) and not attending screening ($p = 0.028$). The autistic respondents (4.0%) and parents (2.0%) who do not attend any doctor differed from the controls (4.5%) in two areas: all had difficulty visiting the doctor when needed, compared to 50% of controls ($p = 0.002$). 95% of autistic non-attenders and 75% of parent non-attenders have experienced at least one delayed treatment outcome, compared

to 43% of non-attending controls ($p=0.01$). There were no significant differences in difficulty attending, barriers experienced or adverse outcomes between formally diagnosed autistic respondents and those who self-identified as autistic.

Facilitators

While most respondents reported booking an appointment online would facilitate access, autistic patients specified a need to “email my doctor in advance with a description of the issue I need to discuss” (62.3%), “wait in a quiet place or outside until my turn” (56.0%), and “book an appointment by text” (41.2%). Autistic patients and parents would be helped if they “could book the first or last appointment” (41.4% and 40.3%) and had a “sensory box available in the waiting room” (15.8% and 14.3%) (Supplementary Table 5).

Despite the outlined difficulties of visiting their doctor, autistic patients and parents felt their relationship with their GP was “very important” or “important” significantly more than controls (69.9% autistic and 76.5% parent vs. 56.1% control, $p=0.001$), but only 33.3% of autistic patients reported a good relationship with their doctor ($p<0.001$).

Discussion

Summary

This study describes the results of a large survey that was undertaken with a convenience sample of autistic people and compares their experiences with parents of autistic children and non-autistic adults. The study adds to the literature highlighting the barriers faced by autistic people in accessing and engaging with primary healthcare, which in our study included greater difficulties deciding when to seek care, reluctance to bother their GP, difficulties planning appointments and greater communication difficulties with particular emphasis on use of the telephone. Communication was also impaired by anxiety and sensory issues.

We linked those barriers to self-reported adverse outcomes. Our data indicated that autistic help-seeking occurs later in the natural course of an illness, with self-reported reduced attendance for screening, late presentations, missed opportunities for early detection and more extensive therapy being required. Autistic people delayed or avoided healthcare because they didn't feel understood by their doctors. Furthermore, a significant minority of autistic adults did not disclose their autism diagnosis which further prevents the identification of their autism-specific needs. These barriers have real consequences, as evidenced in reduced life expectancy, and higher levels of physical and mental health conditions amongst autistic people.

Adverse health outcomes have also been highlighted in parents of autistic children[23]. Parents in this survey reported similar experiences to autistic adults, but reasons included prioritising their child's needs and difficulties with childcare. The self-reported barriers and consequences lay between those reported by autistic and control respondents. Parents were more likely to disclose their child's autism diagnosis to their GP in contrast to autistic adults. Autistic adults potentially could benefit from enhanced awareness of the need to disclose this information and the potential benefits to their care.

Limitations

As the study used a convenience sample and self-report survey the generalisability of the data may be limited. Formal qualitative research methods were not used. Respondents required the ability to respond to the survey which means that the perspectives of autistic people with reduced ability to self-report were not represented.

Strengths

The study arose from a community-identified need to develop autism awareness training for healthcare providers and benefits from being driven by an autistic-led research team using participatory methods, therefore it reflected the priorities of the autistic community and increased the likelihood of genuine responses being received. It provided a unique picture of autistic adults' healthcare experiences, in particular highlighting the difficulty with using the telephone which is a distilled, concentrated essence of verbal communication. It also captured those entirely excluded from healthcare due to access barriers. This is the first comparison of the healthcare experiences of autistic adults, parents of autistic children (an under-represented group in the healthcare literature) and adult controls.

Comparison with existing literature

This study confirms the findings of Nicolaidis[14], Raymaker[16] and several recent reviews on the topic, Mason[17], Bradshaw[27] and Walsh[15] which all showed barriers for autistic adults can be divided into three groups: (1)patient-level factors; (2)provider-level factors; and (3)system-level factors.

Our study stratifies individual barriers from the perspective of the autistic person, which addresses healthcare provider training needs identified in Nicolaidis[4]. It couples these barriers to the self-reported adverse consequences, highlighting some of the factors which may lead to excess morbidity and mortality in the autistic population.

Implications for Research

This study suggests a need for personalised healthcare access plans. A prior study investigated using a pre-visit telephone call to identify individualised accommodations[28]. Our data suggest that this could be problematic for autistic adults. The AASPIRE Healthcare Toolkit[29] includes a publicly available online program which generates a computerised report of required healthcare accommodations. Adaptation of such a toolkit for use in NHS General Practice should be considered. Social care interventions and healthcare facilitators in general practice have shown benefit with a vulnerable population[30], similar approaches could benefit an autistic population. The significant difficulties amongst the small number of autistic people not registered with any GP indicate an urgent need for further research into this group.

Implications for Clinical Practice

Figure 2 shows elements of an autism friendly practice that we propose. Such adjustments would minimise anxiety, manage sensory issues, and ensure mutual understanding with clear, unambiguous communication. Autism friendly practices should employ a personalised approach, with a healthcare access needs assessment and, where possible, a specialist liaison nurse or facilitator.

Implications for Policy

Given the known barriers to care and associated impacts, the extension of annual health checks to autistic adults[31, 32] and the recently announced Oliver McGowan Mandatory Training in Learning Disability and Autism[33] are welcome. These will bring important benefits provided they are directly informed by the autistic community and autistic healthcare providers. Autism registries in GP practice have also been recommended[34,35] which would increase recognition of autism. The success of such initiatives will depend on greater awareness by medical practitioners of autistic culture and communication needs. Specific training for GPs in core training and continuing professional development would be beneficial. GPs with a special interest in autism should be facilitated to develop their skills, but management of general health needs and co-occurring conditions should be within the remit of every GP. Implementing existing autism legislation or development where lacking is urgently required to reduce health inequalities for autistic people.

Conclusion

Significant progress towards eliminating healthcare inequalities for autistic people can only be achieved by understanding the healthcare experiences and access barriers for this vulnerable patient group in general practice.

These barriers represent not so much a failure to deliver or to avail of healthcare, but a lack of intersection between the communication patterns of autistic healthcare users and non-autistic providers, a phenomenon described as The Double Empathy Problem[36].

Reasonable accommodations are legally[37] and morally required, and adjustments for communication needs are as necessary for autistic people as ramps for wheelchair users.

Ethical approval

Ethical approval was obtained from SJH/TUH Research Ethics Committee, Tallaght University Hospital, Dublin

Acknowledgements

We acknowledge the input received from the autistic adult community recruited via local groups and online contacts during the development of the online survey. Assistance with content, structure and proofreading of the surveys was received from nine autistic adults in Ireland and the UK. We received assistance from members of peer support group 'Autistic Doctors International'. We also thank Dr David Hillebrandt and Karen Leneh Buckle for their assistance during this project. We are grateful to AsIAM, Ireland's National Autism Charity and Scally's SuperValu, Clonakilty, for the funding to enable open access publication.

References

1. Data CD. Statistics. Autism Spectrum Disorder. Resource Document.[Last accessed on 2020 Oct 30].
2. Royal College of Psychiatrists.The psychiatric management of autism in adults (CR228). [Last Accessed 2020 Oct 30]

3. Zerbo O, Massolo ML, Qian Y, Croen LA. A study of physician knowledge and experience with autism in adults in a large integrated healthcare system. *J Autism Dev Disord*. 2015 Dec 1;45(12):4002-14.
4. Nicolaidis C, Schnider G, Lee J, Raymaker DM, Kapp SK, Croen LA, Urbanowicz A, Maslak J. Development and psychometric testing of the AASPIRE adult autism healthcare provider self-efficacy scale. *Autism*. 2020 Aug 28;1362361320949734.
5. Rydzewska E, Hughes-McCormack LA, Gillberg C, Henderson A, MacIntyre C, Rintoul J, Cooper SA. General health of adults with autism spectrum disorders—A whole country population cross-sectional study. *Res Autism Spectr Disord*. 2019 Apr 1;60:59-66.
6. Croen LA, Zerbo O, Qian Y, Massolo ML, Rich S, Sidney S, Kripke C. The health status of adults on the autism spectrum. *Autism*. 2015 Oct;19(7):814-23.
7. Ming X, Brimacombe M, Chaaban J, Zimmerman-Bier B, Wagner GC. Autism spectrum disorders: concurrent clinical disorders. *J Child Neurol*. 2008 Jan;23(1):6-13.
8. Flygare Wallen, E., Ljunggren, G., Carlsson, A.C., Pettersson, D. and Wändell, P., 2018. High prevalence of diabetes mellitus, hypertension and obesity among persons with a recorded diagnosis of intellectual disability or autism spectrum disorder. *J Intellect Disabil Res*. 62(4), pp.269-280.
9. Hirvikoski T, Mittendorfer-Rutz E, Boman M, Larsson H, Lichtenstein P, Bölte S. Premature mortality in autism spectrum disorder. *Br J Psychiatry*. 2016 Mar;208(3):232-8.
10. Hwang YI, Srasuebku P, Foley KR, Arnold S, Trollor JN. Mortality and cause of death of Australians on the autism spectrum. *Autism Res*. 2019 May;12(5):806-15.
11. Bilder D, Botts EL, Smith KR, Pimentel R, Farley M, Viskochil J, McMahon WM, Block H, Ritvo E, Ritvo RA, Coon H. Excess mortality and causes of death in autism spectrum disorders: a follow up of the 1980s Utah/UCLA autism epidemiologic study. *J Autism Dev Disord*. 2013 May 1;43(5):1196-204
12. Akobirshoev I, Mitra M, Dembo R, Lauer E. In-hospital mortality among adults with autism spectrum disorder in the United States: A retrospective analysis of US hospital discharge data. *Autism*. 2020 Jan;24(1):177-89.
13. Vohra R, Madhavan S, Sambamoorthi U. Emergency department use among adults with autism spectrum disorders (ASD). *J Autism Dev Disord*. 2016 Apr 1;46(4):1441-54.
14. Nicolaidis C, Raymaker D, McDonald K, Dern S, Boisclair WC, Ashkenazy E, Baggs A. Comparison of healthcare experiences in autistic and non-autistic adults: a cross-sectional online survey facilitated by an academic-community partnership. *J Gen Intern Med*. 2013 Jun 1;28(6):761-9.
15. Walsh C, Lydon S, O'Dowd E, O'Connor P. Barriers to Healthcare for Persons with Autism: A Systematic Review of the Literature and Development of A Taxonomy. *Dev Neurorehabil*. 2020 Feb 8:1-8.
16. Raymaker DM, McDonald KE, Ashkenazy E, Gerrity M, Baggs AM, Kripke C, Hourston S, Nicolaidis C. Barriers to healthcare: Instrument development and comparison between autistic adults and adults with and without other disabilities. *Autism*. 2017 Nov;21(8):972-84
17. Mason D, Ingham B, Urbanowicz A, Michael C, Birtles H, Woodbury-Smith M, Brown T, James I, Scarlett C, Nicolaidis C, Parr JR. A systematic review of what barriers and facilitators prevent and enable physical healthcare services access for autistic adults. *J Autism Dev Disord*. 2019 Aug 15;49(8):3387-400.
18. Nicolaidis C, Raymaker DM, Ashkenazy E, McDonald KE, Dern S, Baggs AE, Kapp SK, Weiner M, Boisclair WC. "Respect the way I need to communicate with you": Healthcare experiences of adults on the autism spectrum. *Autism*. 2015 Oct;19(7):824-31
19. Act A. Autism Act UK. Hm Government (ed). 2009.
20. Department of Health, 2010. "Implementing fulfilling and rewarding lives". Statutory guidance for local authorities and NHS organisation to support implementation of the autism strategy. Department of Health, London
21. Buckley C. Making your practice autism friendly. *InnovAiT*. 2017 Jun;10(6):327-31.
22. Unigwe S, Buckley C, Crane L, Kenny L, Remington A, Pellicano E. GPs' confidence in caring for their patients on the autism spectrum: an online self-report study. *Br J Gen Pract*. 2017 Jun 1;67(659):e445-52.
23. Dijkstra-de Neijis L, Leenen PJ, Hays JP, van der Valk ES, Kraaij R, van Rossum EF, Ester WA. Biological Consequences of Psychological Distress in Caregivers of Children with Autism

- Spectrum Disorder and its Potential Relevance to Other Chronic Diseases Including Cancer. *Curr Epidemiol Rep*. 2020 Jul 4:1-0.
24. The Autscope Organisation. Autscope 2018: Exploring Inclusion. Tonbridge, Kent: 2018. Available from: <http://www.autscope.org/2018/> . [Last accessed on 2020 Oct 30].
 25. AsIAM, 2016. A first for Ireland with Clonakilty becoming Ireland's first autism friendly town. Available at <https://asiam.ie/clonakilty-autism-friendly-town/> .[Last accessed on 2020 Oct 30].
 26. AsIAM, 2020. Are you ready to make your Clonakilty commitment for Autism? Available at: <https://asiam.ie/asiam-public-sector-training/autism-friendly-communities/>. [Last accessed on 2020 Oct 30].
 27. Bradshaw P, Pellicano E, van Driel M, Urbanowicz A. How can we support the healthcare needs of autistic adults without intellectual disability?. *Curr Dev Disord Rep*. 2019 Jun 15;6(2):45-56
 28. Saqr Y, Braun E, Porter K, Barnette D, Hanks C. Addressing medical needs of adolescents and adults with autism spectrum disorders in a primary care setting. *Autism*. 2018 Jan;22(1):51-61.
 29. Nicolaidis C, Raymaker D, McDonald K, Kapp S, Weiner M, Ashkenazy E, Gerrity M, Kripke C, Platt L, Baggs A. The development and evaluation of an online healthcare toolkit for autistic adults and their primary care providers. *J Gen Intern Med*. 2016 Oct 1;31(10):1180-9.
 30. Abel J, Kingston H, Scally A, Hartnoll J, Hannam G, Thomson-Moore A, Kellehear A. Reducing emergency hospital admissions: a population health complex intervention of an enhanced model of primary care and compassionate communities. *Br J Gen Pract*. 2018 Nov 1;68(676):e803-10.
 31. National Institute for Health and Care Excellence. Long term plan. [London]: 2019. Chapter 3, Learning Disability and Autism. Available at: <https://www.longtermplan.nhs.uk/online-version/chapter-3-further-progress-on-care-quality-and-outcomes/a-strong-start-in-life-for-children-and-young-people/learning-disability-and-autism/>.
 32. Harper G, Smith E, Parr J, Ingham B, Bisson H, Wilson C, Buckley C, Unigwe S. Autistica action briefing: health checks. Available at: <https://www.autistica.org.uk/downloads/files/Autistica-Action-Briefing-Health-Checks.pdf>. [Last accessed on 2020 Oct 30].
 33. Health Education England . Partners announced to deliver the Oliver McGowan Mandatory Learning Disability and Autism training for all health and social care staff. Health Education England, 2020, <https://www.hee.nhs.uk/news-blogs-events/news/partners-announced-deliver-oliver-mcgowan-mandatory-learning-disability-autism-training-all-health>. [Last accessed on 2020 Oct 30].
 34. National Institute for Health and Care Excellence. The practice establishes and maintains a register of all patients with a diagnosis of autism. [London]: 2017. ([NM153]). Available at: <https://www.nice.org.uk/standards-and-indicators/qofindicators/the-practice-establishes-and-maintains-a-register-of-all-patients-with-a-diagnosis-of-autism>
 35. Westminster Commission on Autism. A Spectrum of Obstacles—An Inquiry into Access to Healthcare for Autistic People.
 36. Milton DE. On the ontological status of autism: the 'double empathy problem'. *Disabil Soc*. 2012 Oct 1;27(6):883-7.
 37. Act E. Equality Act 2010. The Equality Act. 2010.




Figure 1. Adverse Healthcare Outcomes

Late presentation noted for both autistic & parent group, compared to control ($p < 0.001$)

For all other comparisons between autistic group and parents or controls $p < 0.001$.

Table 1: Participant Characteristics

<u>Participants</u>	Autistic	Parent	Control
Participants (n)	507	196	157
<u>Age</u>			
Median (Range)	38 (17 - 73)	42 (25 - 68)*	38 (18 - 70)
<u>Gender</u>			
Male	99 (19.7%)	15 (7.7%)	16 (10.3%)
Female	311 (62.0%)	181 (92.3%)	132 (84.6%)
Non-binary	83 (16.5%)	0 (0.0%)	7 (4.5%)
Prefer not to say	9 (1.8%)	0 (0.0%)	1 (0.6%)
<u>Location</u>			
UK	330 (65.1%)	72 (36.8%)	67 (42.7%)
Ireland	77 (15.2%)	105 (53.6%)	63 (40.1%)
North America	44 (8.7%)	17 (8.7%)	20 (12.8%)
Other	56 (11.0%)	2 (1.0%)	7 (4.4%)
<u>Formal diagnosis of autism</u>			
By psychiatrist	76.60%		
By clinical psychologist	25.40%		
By multidisciplinary team	48.30%		
	26.30%		
<u>Age at diagnosis</u>			
Median (Range)	33 (2 - 67)		

*n=4 parents reported age <15, omitted from median (range)

Table 2: Access Barriers to Healthcare

Access Barriers	Autistic	Parent	Control	Autistic.Control	Autistic.Parent	Parent.Control
Difficulty attending a GP	400 (79.7%)	103 (52.8%)	57 (36.5%)	p<0.001**	p<0.001**	p<0.001**
<u>Which of the following would cause you to delay or avoid seeing your doctor when you need to?</u>						
	Autistic	Parent	Control	Autistic.Control	Autistic.Parent	Parent.Control
Difficulty deciding if symptoms warrant a GP visit	366 (72.2%)	82 (41.8%)	102 (65.0%)	p=0.102 ns	p<0.001**	p<0.001**
Difficulty using the telephone to book appointment	314 (61.9%)	10 (5.1%)	25 (15.9%)	p<0.001**	p<0.001**	p<0.001**
Not feeling understood	283 (55.8%)	29 (14.8%)	21 (13.4%)	p<0.001**	p<0.001**	p=0.821 ns
Difficulty communicating with the doctor during the appointment	269 (53.1%)	13 (6.6%)	10 (6.4%)	p<0.001**	p<0.001**	p=1.000 ns
The waiting room environment	256 (50.5%)	25 (12.8%)	12 (7.6%)	p<0.001**	p<0.001**	p=0.167 ns
Long wait to get an appointment	251 (49.5%)	81 (41.3%)	70 (44.6%)	p=0.324 ns	p=0.062 ns	p=0.612 ns
Difficulty planning an appointment in advance	243 (47.9%)	59 (30.1%)	51 (32.5%)	p<0.001**	p<0.001**	p=0.715 ns
Inability to see a known or preferred doctor	241 (47.5%)	42 (21.4%)	35 (22.3%)	p<0.001**	p<0.001**	p=0.948 ns
Difficulty communicating with the reception staff	235 (46.4%)	11 (5.6%)	13 (8.3%)	p<0.001**	p<0.001**	p=0.437 ns
Not having enough time to visit the doctor	174 (34.3%)	69 (35.2%)	61 (38.9%)	p=0.346 ns	p=0.894 ns	p=0.552 ns
No online booking system	160 (31.6%)	32 (16.3%)	35 (22.3%)	p=0.033*	p<0.001**	p=0.199 ns
Waiting to see the doctor is too difficult	114 (22.5%)	16 (8.2%)	7 (4.5%)	p<0.001**	p<0.001**	p=0.236 ns
Needing a support person to come with me	106 (20.9%)	11 (5.6%)	7 (4.5%)	p<0.001**	p<0.001**	p=0.806 ns
There is an online booking system but it's confusing	102 (20.1%)	11 (5.6%)	9 (5.7%)	p<0.001**	p<0.001**	p=1.000 ns
Not having anyone to look after my child	66 (13.0%)	73 (37.2%)	18 (11.5%)	p=0.708 ns	p<0.001**	p<0.001**
None of the above	6 (1.2%)	16 (8.2%)	17 (10.8%)	p<0.001**	p<0.001**	p=0.502 ns
Regular GP						
	Autistic	Parent	Control	Autistic.Control	Autistic.Parent	Parent.Control
Usually attend the same medical practice	467 (92.7%)	185 (94.4%)	144 (92.3%)	p=0.908	p=0.594	p=0.718
Usually see the same doctor	280 (55.3%)	123 (62.8%)	74 (47.4%)	p=0.223	p=0.138	p=0.012*
I don't attend any doctor	20 (4.0%)	4 (2.0%)	7 (4.5%)	p=0.223	p=0.138	p=0.012*

Table 2: Access Barriers to Healthcare (contd)

<u>Usual reason for visit</u>	Autistic	Parent	Control	Autistic.Control	Autistic.Parent	Parent.Control
Physical condition or illness	435 (85.8%)	173 (88.3%)	145 (92.4%)	p=0.043 *	p=0.462 ns	p=0.272 ns
Mental health difficulties	310 (61.1%)	85 (43.4%)	43 (27.4%)	p<0.001 **	p<0.001 **	p=0.003 **
Issues directly related to autism	113 (22.3%)	41 (20.9%)	0 (0.0%)	p<0.001 **	p=0.770 ns	p<0.001 **

Table 3: Communication Barriers

<u>Reasons to avoid or delay GP visit (Communication)</u>	Autistic	Parent	Control	Autistic.Control	Autistic.Parent	Parent.Control
Difficulty using the telephone to book an appointment	314 (61.9%)	10 (5.1%)	25 (15.9%)	p<0.001**	p<0.001**	p=0.001**
Not feeling understood	283 (55.8%)	29 (14.8%)	21 (13.4%)	p<0.001**	p<0.001**	p=0.821 ns
Difficulty communicating with the doctor during the appointment	269 (53.1%)	13 (6.6%)	10 (6.4%)	p<0.001**	p<0.001**	p=1.000 ns
Difficulty communicating with the reception staff	235 (46.4%)	11 (5.6%)	13 (8.3%)	p<0.001**	p<0.001**	p=0.437 ns
<u>Communication preferences</u>						
Telephone generally avoided where possible	395 (77.9%)	60 (30.6%)	59 (37.6%)	p<0.001**	p<0.001**	p=0.207 ns
Voicemail generally avoided where possible	311 (61.3%)	54 (27.6%)	61 (38.9%)	p<0.001**	p<0.001**	p=0.033*
Verbal, face-to-face communication generally avoided where possible	152 (30.0%)	31 (15.8%)	12 (7.6%)	p<0.001**	p<0.001**	p=0.03*
It is easier for me to communicate in writing	208 (41.0%)	11 (5.6%)	13 (8.3%)	p<0.001**	p<0.001**	p=0.437 ns
<u>Communication challenges</u>						
Anxiety makes it harder to communicate	395 (77.9%)	49 (25.0%)	42 (26.8%)	p<0.001**	p<0.001**	p=0.801 ns
I have difficulty prioritising when describing medical symptoms	333 (65.7%)	38 (19.4%)	34 (21.7%)	p<0.001**	p<0.001**	p=0.695 ns
I need to give the whole story and not leave anything out	332 (65.5%)	30 (15.3%)	18 (11.5%)	p<0.001**	p<0.001**	p=0.373 ns
I need extra time to process what is being said	286 (56.4%)	18 (9.2%)	12 (7.6%)	p<0.001**	p<0.001**	p=0.746 ns
I can't describe my pain or symptoms accurately	272 (53.6%)	26 (13.3%)	36 (22.9%)	p<0.001**	p<0.001**	p=0.026*
Verbal communication is difficult	234 (46.2%)	8 (4.1%)	11 (7.0%)	p<0.001**	p<0.001**	p=0.331 ns
I express emotions differently e.g. I can appear angry when I am afraid or in pain	227 (44.8%)	14 (7.1%)	6 (3.8%)	p<0.001**	p<0.001**	p=0.267 ns
Sensory issues make communication more difficult	156 (30.8%)	10 (5.1%)	2 (1.3%)	p<0.001**	p<0.001**	p=0.094 ns
None of the above (Communication)	11 (2.2%)	59 (30.1%)	58 (36.9%)	p<0.001**	p<0.001**	p=0.214 ns

Table 4: Sensory Barriers

<u>Reasons to avoid or delay GP visit (Sensory)</u>	Autistic	Parent	Control	Autistic.Control	Autistic.Parent	Parent.Control
The waiting room environment	256 (50.5%)	25 (12.8%)	12 (7.6%)	p<0.001 **	p<0.001 **	p=0.167 ns
<u>Specific sensory challenges</u>						
Noise in the waiting room from other patients	319 (62.9%)	37 (18.9%)	19 (12.1%)	p<0.001 **	p<0.001 **	p=0.113 ns
Bright or fluorescent lights	268 (52.9%)	22 (11.2%)	14 (8.9%)	p<0.001 **	p<0.001 **	p=0.593 ns
Crowded waiting area	299 (59.0%)	47 (24.0%)	22 (14.0%)	p<0.001 **	p<0.001 **	p=0.027 *
Uncomfortable furniture	195 (38.5%)	21 (10.7%)	11 (7.0%)	p<0.001 **	p<0.001 **	p=0.308 ns
Unexpected touch	193 (38.1%)	12 (6.1%)	9 (5.7%)	p<0.001 **	p<0.001 **	p=1.000 ns
Music playing in the waiting room	172 (33.9%)	12 (6.1%)	9 (5.7%)	p<0.001 **	p<0.001 **	p=1.000 ns
Smells in the waiting room	171 (33.7%)	26 (13.3%)	8 (5.1%)	p<0.001 **	p<0.001 **	p=0.016 *
Touch, such as during examination	160 (31.6%)	15 (7.7%)	11 (7.0%)	p<0.001 **	p<0.001 **	p=0.979 ns
Noise from the reception desk	140 (27.6%)	11 (5.6%)	4 (2.5%)	p<0.001 **	p<0.001 **	p=0.249 ns
Smells in the doctor's office	104 (20.5%)	10 (5.1%)	6 (3.8%)	p<0.001 **	p<0.001 **	p=0.751 ns
None of the above (Sensory)	51 (10.1%)	122 (62.2%)	112 (71.3%)	p<0.001 **	p<0.001 **	p=0.092 ns

Communication

- allow text-based **e-consulting** **before** the consultation
- provide written instructions or follow-up/worsening advice **after** the consultation
- Use **clear, unambiguous** language

Sensory Processing

- Conduct a **sensory audit** of waiting and consulting spaces (lighting/noise/smell)
- Appropriate **appointment times** (beginning/end of day)

Predictability & Control

- preferred, regular doctor
- instruct on what to expect at appointments
- signpost to clear, directive resources e.g. www.nhs.uk
- advise on expected waiting times

- **online appointment booking**
- allow time to process information

- Education and training for **all** staff
- Individualised access plan

- **Tell the patient** what to expect and what you're doing during examinations

- Consider **longer appointment slots**
- Provide **written information** on procedures & consider enhanced support for screening