

1 **Public perspectives on social distancing and other protective measures in Europe: a cross-sectional**
2 **survey study during the COVID-19 pandemic**

3 *Karien Meier (0000-0003-4855-1576), Toivo Glatz (0000-0002-8994-7860), Mathijs C Guijt (0000-0002-*
4 *9958-8590), Marco Piccininni (0000-0002-1397-0060), Merel van der Meulen (0000-0002-0001-4408),*
5 *Khaled Atmar (0000-0003-1213-3903), Anne-Tess C Jolink (0000-0002-0290-1705), Tobias Kurth (0000-*
6 *0001-7169-2620), Jessica L Rohmann (0000-0003-2420-5716), Amir H Zamanipoor Najafabadi (0000-*
7 *0003-4855-1576), on behalf of the COVID-19 Survey Study group**

8
9 ^a*Karien Meier, psychologist, medical student (karienmeier@gmail.com)*

10 ^b*Toivo Glatz, clinical linguist, research fellow (toivo.glatz@charite.de)*

11 ^a*Mathijs C Guijt, medical student (m.c.guijt.3@umail.leidenuniv.nl)*

12 ^b*Marco Piccininni, biostatistician, research fellow (marco.piccininni@charite.de)*

13 ^a*Merel van der Meulen, medical student (m.van_der_meulen.nech@lumc.nl)*

14 ^a*Khaled Atmar, medical student (k.atmar@umail.leidenuniv.nl)*

15 ^a*Anne-Tess C Jolink, medical student (a.t.c.jolink@umail.leidenuniv.nl)*

16 ^b*Tobias Kurth, professor of public health and epidemiology (tobias.kurth@charite.de)*

17 ^b*Jessica L Rohmann, epidemiologist, research fellow (jessica.rohmann@charite.de)*

18 ^a*Amir H Zamanipoor Najafabadi, epidemiologist in training and MD/PhD student (amir@lumc.nl)*

19 ***COVID-19 Survey Study group**

20 ^a*Layla Andour, medical student (0000-0002-0073-9601) (l.andour@umail.leidenuniv.nl)*

21 ^a*Liesbeth Blommaart, medical student (0000-0002-9838-3825) (l.blommaart@umail.leidenuniv.nl)*

22 ^a*Fleur L Fisher, medical student (0000-0002-7806-8342) (f.l.fisher@lumc.nl)*

23 ^a*Bas Splinter, medical student (0000-0002-5480-3891) (b.splinter@umail.leidenuniv.nl)*

24 ^a*Jasper van de Stadt, medical student (0000-0001-6126-2221) (j.van.de.stadt@umail.leidenuniv.nl)*

25 ^a*Maartje Teunissen, medical student (0000-0003-1702-5215) (m.teunissen@umail.leidenuniv.nl)*

26 ^a*Thijs CM Windmeijer, medical student (0000-0002-5063-4525) (t.c.m.windmeijer@umail.leidenuniv.nl)*

27 ^c*Guido Rohmann, political scientist, PhD student (0000-0001-6758-8812) (rohmann@gsnas.fu-berlin.de)*

28
29 ^aLeiden University Medical Center, Leiden University, Albinusdreef 2, 2333 ZA Leiden, The Netherlands

30 ^bInstitute of Public Health, Charité - Universitätsmedizin Berlin, Charitéplatz 1, 10117 Berlin, Germany

31 ^cJohn-F.-Kennedy-Institut, Freie Universität Berlin, Lansstraße 5-9, 14195 Berlin, Germany

32

33

34 ***Acknowledgements***

35 *Ilaria Vitulano*

36 *Büşra Nur Bozok*

37 *María Alcántara Laguna*

38 *Ariane Scholten*

39 *Renate Meier*

40 *Shabbir R Shahnahpur*

41 *Marcin M Fiszler*

42

43

44 **Correspondence to: Amir H. Zamanipoor Najafabadi, Department of Neurosurgery, Leiden**

45 **University Medical Center, Leiden, The Netherlands. Tel: +31(0)681663780. Email: amir@lumc.nl.**

46

47 Type of article: Research

48 Figures and Tables: 3 Figures and 3 Tables

49 Word count main text: 4050

50 Word count abstract: 399

51 **ABSTRACT**

52 **Objectives:** The extent to which people implement government-issued protective measures is critical in
53 preventing further spread of coronavirus disease 2019 (COVID-19) caused by coronavirus SARS-CoV-2.
54 Our study aimed to evaluate the public belief in the effectiveness of protective measures, the reported
55 implementation of these measures in daily life, and to identify communication channels used to acquire
56 relevant information on COVID-19 in European countries.

57 **Design:** A cross-sectional online survey available in multiple languages was disseminated on social media
58 starting March 19th, 2020. After five days, we computed descriptive statistics for countries with more than
59 500 respondents. Each day, we compiled and categorized community containment measures enacted in
60 each country by stringency (stage I-IV). Response collection continued for one week to explore possible
61 dynamics as containment strategies intensified.

62 **Participants:** In total, 9,796 adults responded, of whom 8,611 resided in the Netherlands (stage III), 604
63 in Germany (stage III), and 581 in Italy (stage IV). An additional 1,365 respondents completed the survey
64 in the following week.

65 **Results:** Participants indicated support for governmental measures related to avoiding social gatherings,
66 selective closure of public places, and hand hygiene and respiratory measures (range for all measures:
67 95.0%-99.7%). Respondents from the Netherlands were less likely to consider a complete social lockdown
68 effective (59.2%), compared to respondents in Germany (76.6%) or Italy (87.2%). Italian residents did not
69 only apply enforced social distancing measures more frequently (range: 90.2%-99.3%, German and Dutch
70 residents: 67.5%-97.0%), but also self-initiated hygienic and social distancing behaviors (range: 36.3%-
71 96.6%, German and Dutch residents: 28.3%-95.7%). Respondents largely reported being sufficiently
72 informed about the COVID-19 outbreak and about behaviors to avoid infection (range across countries:
73 90.2%-91.1%). Information channels most commonly reported included television (range: 53.0%-82.0%),
74 newspapers (range: 31.0%-63.0%), official health websites (range: 39.0%-54.1%), and social media
75 (range: 40.0%-55.8%). We observed no major changes in answers over time.

76 **Conclusions:** In European countries, the degree of public belief in the effectiveness of protective
77 measures was high and residents reported to be sufficiently informed by various communication channels.
78 In March 2020, implementation of enacted and self-initiated measures differed between countries and
79 were highest among Italian respondents, who were subjected to the most elaborate measures of social
80 lockdown and greatest COVID-19 burden in Europe.

81 **Keywords:** COVID-19, SARS-CoV-2, Survey, Public Behaviour, Pandemic, Mitigation

82 INTRODUCTION

83 The recent pandemic of COVID-19 (coronavirus disease 2019) caused by SARS-CoV-2 (Severe
84 Acute Respiratory Syndrome Coronavirus 2) has infected more than 1,000,000 people worldwide in only a
85 few months' time and caused more than 51,000 deaths as of April 2nd, 2020[1]. This rapidly spreading
86 virus imposes a tremendous burden on national healthcare systems, as they lack sufficient material and
87 human resources to respond to the rapidly increasing number of patients requiring intensive care[1,2].
88 Worldwide, public health organizations, as well as national and international government bodies, have
89 suggested systematic implementation of protective, public health measures in an effort to delay the spread
90 of COVID-19[3]. The aim of these measures is to decrease the peak infection rate, while maintaining a
91 high quality of care under finite resources and limited hospital capacities[2,4]. In addition to basic
92 hygienic advice such as regular hand washing, the most important recommendation known to limit and
93 delay the spread of the virus is social (physical) distancing[5,6].

94 In early March 2020, Europe became the epicenter of the COVID-19 pandemic, with more cases
95 and deaths reported than in all other countries (excluding China) combined[1]. Throughout the course of
96 the month, most European countries progressively implemented community isolation measures to increase
97 social distancing, such as imposing work restrictions and the closure of public places and shops. Italy, the
98 first and most severely affected country in Europe, imposed strict community isolation measures on March
99 9th and 11th, 2020, enforcing a nationwide quarantine in response to the alarming increase in the number
100 of cases, which posed a serious threat to the capacity of the Italian healthcare system[7].

101 The extent to which people are informed about and apply the measures advocated by experts and
102 enforced by governments is critical to control the spread of the virus and to optimize patient outcomes
103 during the current COVID-19 pandemic[4,8,9]. The aim of our study was to evaluate public belief in the
104 effectiveness of protective measures, to what extent individuals have implemented these measures in their
105 daily lives, and to identify key communication channels used to acquire information on COVID-19 in
106 European countries. We believe these insights are not only valuable for the ongoing mitigation of the
107 current COVID-19 pandemic, but may also serve to inform governments' and public health organizations'
108 information dissemination and infection control strategies for possible future pandemics.

109 **METHODS**

110 *Design, setting, and participants*

111 The survey instrument used to gather cross-sectional data was compiled by a team of medical
112 students and epidemiologists from the Leiden University Medical Center and the Charité -
113 Universitätsmedizin Berlin. Our initial aim was to collect sufficient data on adults living in Europe, with
114 an emphasis on individuals residing in the Netherlands, Germany, and Italy, however, the survey was also
115 open to residents of other countries. Data collection is still ongoing and is planned to continue for as long
116 as community isolation measures remain in place. In these primary descriptive analyses, we only reported
117 data from countries with at least 500 responses at our first cut-off date, March 23rd, 2020. This date was
118 chosen because several European regional and national governing bodies announced stricter measures
119 around this date. The study was reviewed and granted exempted status from medical ethical approval by
120 the Institutional Review Board of Leiden University Medical Center in The Netherlands (protocol number:
121 N20-037).

122 *Survey instrument*

123 We selected questions from the validated Flu TELEphone Survey Template (FluTEST), which was
124 designed to assess perceptions and behavior during an influenza pandemic [10]. We slightly modified the
125 items to fit the current outbreak context, where necessary and formulated additional questions to assess
126 beliefs in the effectiveness of protective measures [11]. In brief, the survey instrument consisted of 22
127 items in three sections including: 1) five questions regarding beliefs in the effectiveness of public
128 measures to reduce outspread (e.g. selective closure of places and complete social lockdown), 2) 16
129 questions on the personal application of protective measures (e.g. social distancing behaviors and hygienic
130 practices), and 3) one question on the three most frequently used sources to acquire information on the
131 COVID-19 outbreak, and one question on the perception whether or not respondents felt sufficiently
132 informed. The full survey is presented in Supplemental Text 1. To allow for stratified interpretation of the
133 results, additional questions captured sociodemographic information about gender, age, household
134 composition, employment status, educational level, country of residence, being a healthcare provider or
135 (bio)medical student, and prevalent chronic medical conditions. Respondents were able to complete the
136 questionnaire only once per device in an effort to reduce potential repeat responses.

137 The survey was translated into multiple languages by a small panel of native speakers from the
138 original English language version. Due to time constraints, we were unable to formally validate the survey
139 questions in the other languages. The survey went live on March 19th, 2020 in Dutch, English and
140 German. Other languages have been added since initiation (Italian on March 20, 2020; French and Polish

141 on March 21, 2020; Spanish on March 22, 2020; Turkish on March 25, 2020; and Farsi on March 29,
142 2020; see Supplemental Text 1).

143 *Procedures*

144 The full survey was initially piloted on a sample of 50 respondents. After minor modifications to
145 the structure and language, the survey was actively disseminated through (social) media channels, such as
146 WhatsApp, Telegram, Facebook, LinkedIn, Instagram, and Twitter, and in professional networks via
147 electronic mailing lists. The survey was further promoted via a number of local and national news
148 websites and radio stations. On the landing page, participants were briefed about the study and only those
149 providing informed consent for participation were guided to the actual 5-minute survey. On the final page,
150 participants were debriefed about the study, and thanked for their contribution.

151

152 *Assessment of stages of community containment measures*

153 We extracted community containment measures taken by governments in each country included
154 in this study from national governmental announcements and news articles and compiled daily from
155 March 1st, 2020 onwards. Two independent researchers classified stringency of isolation measures by
156 country in four stages based on the Community Containment Measures guideline originally developed by
157 the Centers for Disease Control and Prevention (CDC) during the SARS outbreak in 2003[12]. Any
158 disagreement was resolved by discussion. The CDC guideline describes seven interventions, which we
159 grouped into four stages to create country-specific timelines for the purposes of this study. Guideline
160 interventions 1 (passive monitoring), 2 and 3 (active monitoring without and with activity restrictions,
161 respectively) were grouped together as Stage I (“Low Impact Containment Measures”), since most
162 countries had already implemented these interventions in early March 2020. Guideline interventions 4
163 (working quarantine) and 5 (focused measures) were grouped and classified as Stage II (“Focused
164 Measures to Increase Social Distance”), as many countries applied these interventions simultaneously. We
165 designated intervention 6 as Stage III (“Community-Wide Measures to Increase Social Distance”) and
166 intervention 7 as Stage IV (“Widespread Community Quarantine, Including Cordon Sanitaire”). We
167 detailed and justified the daily stage classification by country in a series of timelines (Supplemental Tables
168 1a-1d; Supplemental Text 2).

169 *Statistical Analyses*

170 Data collected over a five-day period between March 19th, 2020 and March 23th, 2020 at 11:20
171 AM (UTC+0), were used in the primary analyses. We present results of the survey items including

172 sociodemographic characteristics using descriptive summary statistics for the countries having more than
173 500 responses during this primary data collection period (the Netherlands, Germany, and Italy). Nominal
174 variables were described and visualized using frequencies and percentages. We also reported frequencies
175 of missing responses. We present stratified results for the assessed sociodemographic variables only for
176 the Netherlands, as the number of responses was sufficient per individual subgroup. No formal statistical
177 comparisons were made between countries since the primary aim was descriptive and there were no *a*
178 *priori* testable hypotheses.

179 As a secondary analysis, we explored changes in responses for items about the beliefs in the
180 effectiveness of these measures and their implementation over time. As for items about implementation of
181 protective measures, we reported the proportion of positive answers (“Yes”) out of all responses,
182 excluding responses indicating the question was not applicable to their situation. Similarly, for items about
183 the belief in effectiveness of these measures, we considered the proportion of positive answers (“Probably
184 true”). To easily visualize the change over time, we modeled the proportions for each item and for each
185 country separately, using generalized additive models with time as the independent variable, using a
186 shrinkage version of cubic splines with three knots. In addition, we computed and presented visualizations
187 of the differences in proportion between the responses recorded during the primary data collection period
188 and the weeklong extension only for the Netherlands.

189 Data management, analyses and visualizations were conducted using Stata 16.1 (StatCorp LP,
190 College Station, TX) software and R 3.6.3 / RStudio 1.2.

191

192 **RESULTS**

193
194 Between March 19th and 23rd, 2020, a total of 9,796 respondents completed the survey. Three
195 countries met our study inclusion criteria of more than 500 respondents; the Netherlands (n=8,611),
196 Germany (n=604), and Italy (n=581) (see flowchart: Figure 1). During this primary data collection period,
197 the containment measures in the Netherlands and Germany met criteria for Stage III classification
198 (“Community-Wide Measures to Increase Social Distance”), and those in Italy met Stage IV criteria
199 (“Widespread Community Quarantine”).

200 Most respondents opened the survey link via WhatsApp or Facebook (Figure 1). Approximately
201 two-thirds of respondents were female and one-third of respondents were aged 21-30 years old (Table 1).
202 The majority of respondents had a paid job (57.1%) and many had tertiary academic degrees (68.2%)
203 Approximately 18% of respondents were healthcare providers or (bio)medical students. Less than one-
204 fifth of respondents reported suffering from a chronic illness or being in poor medical condition (17.3%).
205 Descriptive sociodemographic characteristics stratified by country of residence are presented in Table 1.

206

207 **Sources used to acquire information on the COVID-19 outbreak**

208
209 Among respondents living in the Netherlands, Germany, or Italy, the most frequently used sources
210 to obtain relevant information included television (e.g. news, range: 53.0%-82.0%), newspapers or news
211 applications (range: 31.0%-63.0%), social media (e.g. Facebook and Twitter, range: 40.0%-55.8%), and
212 official health websites (range: 39.0%-54.1%). Other people (e.g. family, friends and colleagues, range:
213 23.9%-26.7%) and radio were reported less frequently as sources of information (range: 5.3%-26.7%). In
214 all three countries, healthcare professionals (range: 4.3%-7.6%) and official health hotlines (range: 1.2%-
215 1.6%) were the least frequently reported sources of information (Table 1). Almost all respondents living in
216 these three countries reported being sufficiently informed about the current COVID-19 outbreak and what
217 they could do to prevent an infection (range: 90.2%-91.1%; Table 2).

218

219 **Belief in the effectiveness of measures to reduce outspread**

220

221 The majority of respondents believed that avoiding social gatherings, selective closure of public
222 places and locations, hand hygiene measures, and respiratory measures were effective ways to prevent
223 further spread of COVID-19 (range for all measures in all three countries: 95.0%-99.7%; Table 2). Only
224 59.2% of respondents in the Netherlands perceived a complete social lockdown or isolation measures as

225 effective, compared with 76.6% of respondents from Germany and 87.2% from Italy (Figure 2 and Table
226 2).

227

228 **Individual implementation of protective measures**

229

230 For all items, the percentages reported in the text and Table 3 excluded respondents to whom the
231 item did not apply, which was especially important in the interpretation of three items (keeping children at
232 home before any mandates were put in place, range: 41.0%-75.1%; reducing the use of public transport,
233 range: 1.9%-28.5%; and going to school/university/work, range: 2.4%-14.9%). With regard to personal
234 protective behaviors, a high number of respondents from the Netherlands and Germany reported to have
235 washed their hands with soap and water more often than usual (range: 95.0%-95.7%). In general,
236 respondents from Italy reported applying all proposed personal protective behaviors more often than those
237 from the Netherlands or Germany, except for following a healthy diet or using vitamin supplements
238 (36.3%, Netherlands 54.5%, Germany 54.4%) (Figure 3 and Table 3).

239 Behavior related to limiting interactions with people was fairly similar between countries,
240 although respondents from Italy reported more frequently cancelling or postponing social events (98.8%,
241 compared with 94.8% in the Netherlands and 97.0% in Germany) and avoiding crowded places more
242 frequently (99.3%, compared with 92.4% in the Netherlands and 93.8% in Germany). Respondents living
243 in Germany reported avoiding people with cold or flu-like symptoms (81.1%) less frequently than
244 respondents living in Italy (90.2%) or in the Netherlands (89.0%). Regarding behaviors related to avoiding
245 travel, respondents from Italy more often reported to have reduced the amount they went to school or
246 work (94.4%, compared to 88.0% in the Netherlands and 84.9% in Germany, of public transport use
247 (98.6% compared to 89.6% in the Netherlands and 91.3% in Germany), and of going to shops (97.7%,
248 compared to 81.4% in the Netherlands and 67.5% in Germany) (Figure 3 and Table 3). Responses to
249 questions concerning limiting interactions and avoiding traveling may reflect both imposed restrictions
250 and respondents' awareness and willingness to follow protective measures. Therefore, we additionally
251 asked respondents whether they kept children at home prior to formal mandates to assess the percentage of
252 respondents that applied measures on their own accord. Of those indicating the question was applicable to
253 their situation, in Italy, 60.6% kept their children at home before protective measures compared to 53.6%
254 in the Netherlands and 58.1% in Germany.

255

256 **Subgroup analyses among respondents living in the Netherlands**

257

258 Although we conducted no formal comparisons between the sociodemographic subgroups of
259 participants, some patterns were evident (Supplemental Tables 2a-2r). In general, while there were no
260 meaningful differences in the degree of belief in the effectiveness of protective measures among gender
261 groups, females applied these measures most frequently. Among the different age groups, the belief in the
262 effectiveness of a complete social lockdown differed (e.g. ≤ 20 years: 47.5%, 21-40 years: 62.7%), as well
263 as among subgroups with different daily activities (e.g. retired: 55.4%, homemaker/unemployed: 64.7%)
264 and different education levels (primary/secondary: 54.6%, tertiary academic: 61.1%). Chronically ill
265 patients more frequently reported exhibiting protective measures than respondents without any chronic
266 diseases. Different sociodemographic subgroups used different sources of information to obtain
267 information related to the COVID-19 pandemic. With higher age, the percentage of respondents who
268 agreed they felt sufficiently informed was higher (for example: ≤ 20 years: 87.0%, versus > 60 years:
269 94.9%)

270

271

272 **Change in responses over time**

273

274 Immediately following the primary data collection period, we continued collecting data over the
275 next seven days (March 30th, 2020, 11.40 AM, UTC+0) and received responses from 1,588 additional
276 participants, of whom 1,365 reported living in the Netherlands (n=858), in Germany (n=413) and in Italy
277 (n=94). In general, we observed no substantial changes over time (Supplemental Figures 1a-1b), except
278 for a decrease in the belief of effectiveness of a complete social lockdown in Germany (Supplemental
279 Figure 1a). Furthermore, among respondents from the Netherlands, we observed a small increase in the
280 percentages of respondents indicating they believe in the effectiveness of preventive measures (range: 0%-
281 5%) and those indicating they implemented these measures (range: 0%-10%) across both data collection
282 periods (Supplemental Figures 2a-2b).

283 **DISCUSSION**

284 Our findings indicate that in three European countries, the Netherlands, Germany, and Italy, the
285 public belief in the effectiveness and the actual implementation of certain protective measures during the
286 ongoing COVID-19 pandemic in March of 2020 was high. Furthermore, residents reported to be
287 sufficiently informed about the ongoing pandemic using various communication channels.

288 The public belief in the effectiveness of protective measures was highest among respondents
289 residing in Italy, which had the most extensive measures of social lockdown as well as the highest
290 numbers of COVID-19 cases and deaths in Europe in March 2020. Compared to the Netherlands and
291 Germany, respondents living in Italy most often reported not only exhibiting behaviors related to
292 government imposed restrictions but also voluntary hygienic and social measures. Although in general,
293 more than 90% of respondents indicated belief in the effectiveness of imposed measures of social
294 distancing, a complete social lockdown was deemed effective by only 59% of respondents residing in the
295 Netherlands (compared to 77% in Germany and 87% in Italy), where at the time of survey completion,
296 only lighter social distancing measures were enforced. The results of our study suggest that the level of
297 community containment measures implemented by national governments may be rapidly visible in the
298 public beliefs about protective measures, the extent to which people actually exhibit these relevant
299 behaviors, and reflect the severity of the outbreak situation in a given country.

300 To the best of our knowledge, to date, few data on this topic are available. Results from two
301 recently published survey studies conducted in the USA, the UK, and China primarily focus on the
302 respondents' knowledge about COVID-19 and assess understanding pertaining to the disease
303 course[13,14]. Furthermore, another study conducted between January 24th and February 13th, 2020
304 among 1715 Hong Kong residents showed that most respondents obtained information on the COVID-19
305 pandemic from social media and websites[15]. We found that traditional information sources (e.g.
306 television and news) were used most frequently among our respondents. Our study further corroborates
307 and adds to these first findings with similar results regarding beliefs in the effectiveness of hygienic and
308 social distancing measures and the extent to which these measures were exhibited in a European study
309 population.

310 **Social distancing and other behavioral measures**

311 Individuals' adherence to country-specific mitigation measures has the potential to influence the
312 course of the COVID-19 pandemic. Social (physical) distancing has been proposed as one of the most
313 effective measures for mitigating pandemics caused by viruses, including COVID-19[4,7,16]. Large-scale
314 simulation studies have found that closure of borders is only effective to prevent further spread of the

315 virus if they are implemented for more than two weeks and prevent international travel [8]. Moreover, the
316 peak attack rate can be decreased by case isolation, household quarantine, and school, university and work
317 closure[8].

318 As the transmissibility of the SARS-CoV-2 virus is estimated to be similar to or higher than
319 previous coronaviruses such as SARS, social isolation measures are particularly important [17,18]. In the
320 current COVID-19 pandemic, models have shown that the Wuhan quarantine reduced transmission of
321 COVID-19 cases from mainland China to other countries by 77% by early February [9]. Besides social
322 distancing, other behavioral protective measures have also been deemed effective in the mitigation of the
323 current pandemic. For instance, regular hand washing may result in a reduction of peak infection rate up to
324 65% with a delay of 2.7 months and a 29% decrease in total infection rate[19]. Generally, preventive,
325 precautionary behavior is more commonly observed, in females, and in older persons [20–22], which was
326 also reflected in our findings from Europe during the ongoing COVID-19 pandemic.

327 **Provision and acquirement of information during pandemics**

328 Transparent, timely, and easy-to-understand information is essential to increase trust in national
329 governments during pandemics[23]. The increasing use of portable devices and social media is evident in
330 our findings, which indicate frequent use of social media to acquire pandemic-related information (range
331 across countries: 40.0%-55.8%). However, in recent epidemics and pandemics, a substantial amount of
332 online information, especially distributed via social media, was found to be incorrect and
333 misleading[24,25]. Environmental cues to follow behavioral recommendations, favorable attitudes
334 towards prevention measures, and knowledge about the virus were associated with exhibiting protective
335 behavior[5]. Therefore accurate information provision via social media channels is crucial, besides
336 information via traditional information sources.

337 **Study strengths and limitations**

338 Given the evolving pandemic situation, we felt it was important to develop, translate, and
339 disseminate our questionnaire rapidly to capture a snapshot of public perceptions and behaviors in ‘real
340 time’ as the COVID-19 crisis unfolded in Europe and as policy makers enacted formal containment
341 measures in several European countries. Many items in our survey were adapted from an existing
342 validated questionnaire created to assess perceptions and behaviors in response to influenza[10]. We
343 attempted to make our survey accessible to participants of diverse backgrounds by providing the survey in
344 eight languages. These translations could be readily adapted for use in future studies on other viral
345 pandemics.

346 Readers should consider some important limitations when interpreting our findings. First, since
347 the survey was web-based and recruitment was largely through digital channels including social media, we
348 acknowledge the potential for selection bias. We cannot assume that our study population is representative
349 for the individual countries and acknowledge possible over-representation of health-conscious individuals
350 and those more informed or concerned about the outbreak. However, under the exceptional current
351 circumstances, members of the general public who normally would not participate in health-related
352 surveys may be more likely to participate given the media attention, severity, and the outbreak's large
353 impact on many aspects of daily life. Furthermore, with the social (physical) distancing recommendations
354 and enforced measures in place, many confined to their homes have turned to social media and other web-
355 based platforms for social communication, including those belonging to the older age groups.

356 Second, the number of completed survey responses was much higher among residents of the
357 Netherlands compared to Italy or Germany. Since the majority of our research team members are based in
358 the Netherlands and the largest dissemination efforts occurred there, this is not surprising. Third, while the
359 governments of Germany, Italy, and the Netherlands enacted different mitigation measures in each
360 country, our survey was not adapted to country-specific nuances. Hence, we acknowledge that our results
361 might not fully depict whether residents of these countries actually exhibit their country-specific
362 measures. Fourth, we cannot rule out that an individual completed the survey more than once on multiple
363 devices, in another browser, or by clearing cookies; however, repeat submissions from the same device
364 were not accepted.

365 Fifth, with regard to the secondary analysis over the extended data collection period, we observed
366 no major changes in the aggregated answers over time; however, we acknowledge a possible delay
367 between the implementation of formal community isolation measures and the subsequent information
368 uptake and application of these measures by residents. During the extended data collection period,
369 response rates were lower than in the primary collection period, especially from respondents living in
370 Italy.

371 Finally, we emphasize the aim of our study was descriptive, and we caution readers against
372 drawing causal conclusions regarding the observed differences. A formal comparison between countries
373 would require appropriate analytical consideration of variables taking into account sociocultural
374 (including educational systems), political, and structural contexts in each country. In an effort to help the
375 reader better understand each country-specific setting we included information about the stage of
376 containment measures enacted within the included countries in a timeline encapsulating the data collection
377 period.

378

379 **CONCLUSION**

380 The extent to which individuals internalize and respond to (government-mandated) mitigation
381 measures and recommendations is critical to the control of the spread of the SARS-CoV-2 virus and to
382 optimize outcomes during the current COVID-19 pandemic. In our survey study of the general public
383 living in the Netherlands, Germany, and Italy, we found that approval and application of publicly enforced
384 and self-initiated protective measures were highest in Italy, the region with the most extensive measures of
385 social lockdown and highest burden (number of cases and deaths) in Europe, during the study time period
386 in mid-March, 2020. Media channels used to acquire information and the extent to which respondents felt
387 sufficiently informed about the COVID-19 pandemic differed per country and among sociodemographic
388 subgroups in the Netherlands. No substantial changes in the perceived effectiveness of behavioral
389 protective measures and the implementation of these measures in these countries were observed between
390 March 19th and March 30th, 2020, as the COVID-19 pandemic continued to evolve in Europe and formal
391 community isolation measures became stricter. We believe these insights are valuable to inform the
392 information dissemination and infection control strategies of governments and public health organizations
393 during the current crisis and also for future pandemics.

394 **Summary boxes**

395 *What is already known on this topic*

- 396 ● Social distancing and hygienic measures are effective in mitigating pandemics such as the
397 COVID-19 pandemic.
- 398 ● Public beliefs regarding the effectiveness of protective measures, and the extent to which these
399 measures have been applied by European residents during the COVID-19 pandemic remain
400 unknown.

401 *What this study adds*

- 402 ● In three European countries, the degree of public belief in the effectiveness of protective measures
403 was high and respondents living in the Netherlands, Germany and Italy reported feeling
404 sufficiently informed by various communication channels.
- 405 ● Implementation of enacted and self-initiated measures differed between countries and were
406 highest among respondents living in Italy, who were subjected to the most elaborate measures of
407 social lockdown and experienced the greatest outbreak burden.

408 **Contributors:** AHZN conceived this study and AHZN, KM, and JLR developed the study design. All
409 authors contributed to design the survey, translation and verification of the survey items in German, Dutch
410 or Italian, and distribution of the survey in social and professional networks. KM, TG, MP, and RM
411 conducted the statistical analysis and data visualization. MCG and ATCJ classified countries on
412 containment stage. KM, JLR and AHZN drafted the manuscript with support from MM, KA, ATCJ, TG
413 and MP. The COVID-19 study group contributed to translation, pilot testing of the survey, dissemination,
414 and country community isolation measure classification. All authors and study group members helped
415 interpret the results and critically revised the final manuscript. AHZN supervised the project. The
416 corresponding author attests that all listed authors meet authorship criteria and that no others meeting the
417 criteria have been omitted.

418 **Funding:** The authors received no specific funding for this work.

419 **Competing interests:** All authors have completed the ICMJE uniform disclosure form at
420 http://www.icmje.org/coi_disclosure.pdf (available on request from the corresponding author). KM, TG,
421 MCG, MP, MvdM, ATCJ, JLR, AHZN declare no other support from any organisation for the submitted
422 work, no financial relationships with any organisations that might have an interest in the submitted work
423 in the previous three years, no other relationships or activities that could appear to have influenced the

424 submitted work. TK reports outside of the submitted work to having contributed to an advisory board of
425 CoLucid and a research project funded by Amgen, for which the Charité – Universitätsmedizin Berlin
426 received an unrestricted compensation. He further reports having received honoraria from Lilly,
427 Newsenselab, and Total for providing methodological advice, from Novartis and from Daiichi Sankyo for
428 providing a lecture on neuroepidemiology and research methods. He is further a consulting clinical
429 epidemiology editor at The BMJ and has received compensation for editorial services.

430 **Ethical approval:** The study was reviewed and granted exempted status from medical ethical approval by
431 the Institutional Review Board of Leiden University Medical Center in The Netherlands (protocol number:
432 N20-037).

433 **Data sharing:** Requests for data sharing can be directed to AHZN: a.h.zamanipoor@lumc.nl. All survey
434 items are included as supplemental material.

435 **Transparency:** The corresponding author (AHZN) affirms that this manuscript is an honest, accurate, and
436 transparent account of the study being reported; that no important aspects of the study have been omitted;
437 and that any discrepancies from the study as planned have been explained.

438 **Patient and Public Involvement:** The target population, the general public, was actively involved in the
439 dissemination of this survey. No patients were involved in this study, as the target group was the general
440 public.

441 **Dissemination declaration:** Upon request, the data will be available for policy makers and government
442 bodies.

443

444 **REFERENCES**

- 445 1 Dong E, Du H, Gardner L. An interactive web-based dashboard to track COVID-19 in real time.
446 *Lancet Infect Dis* Published Online First: 19 February 2020. doi:10.1016/S1473-3099(20)30120-1
- 447 2 Heymann DL, Shindo N, WHO Scientific and Technical Advisory Group for Infectious Hazards.
448 COVID-19: what is next for public health? *Lancet* 2020;**395**:542–5. doi:10.1016/S0140-
449 6736(20)30374-3
- 450 3 Ebrahim SH, Ahmed QA, Gozzer E, *et al.* Covid-19 and community mitigation strategies in a
451 pandemic. *BMJ* 2020;**368**:m1066. doi:10.1136/bmj.m1066
- 452 4 Anderson RM, Heesterbeek H, Klinkenberg D, *et al.* How will country-based mitigation measures
453 influence the course of the COVID-19 epidemic? *Lancet* 2020;**395**:931–4. doi:10.1016/S0140-
454 6736(20)30567-5
- 455 5 He F, Deng Y, Li W. Coronavirus Disease 2019 (COVID-19): What we know? *Journal of Medical*
456 *Virology*. 2020. doi:10.1002/jmv.25766
- 457 6 Kucharski AJ, Russell TW, Diamond C, *et al.* Early dynamics of transmission and control of COVID-
458 19: a mathematical modelling study. doi:10.1101/2020.01.31.20019901
- 459 7 Saglietto A, D’Ascenzo F, Zoccai GB, *et al.* COVID-19 in Europe: the Italian lesson. *Lancet*
460 Published Online First: 24 March 2020. doi:10.1016/S0140-6736(20)30690-5
- 461 8 Ferguson NM, Cummings DAT, Fraser C, *et al.* Strategies for mitigating an influenza pandemic.
462 *Nature*. 2006;**442**:448–52. doi:10.1038/nature04795
- 463 9 Chinazzi M, Davis JT, Ajelli M, *et al.* The effect of travel restrictions on the spread of the 2019 novel
464 coronavirus (COVID-19) outbreak. *Science* Published Online First: 6 March 2020.
465 doi:10.1126/science.aba9757
- 466 10 Rubin GJ, James Rubin G, Bakhshi S, *et al.* The design of a survey questionnaire to measure
467 perceptions and behaviour during an influenza pandemic: the Flu Telephone Survey Template
468 (FluTEST). *Health Services and Delivery Research*. 2014;**2**:1–126. doi:10.3310/hsdr02410
- 469 11 World Health Organization. Non-pharmaceutical public health measures for mitigating the risk and
470 impact of epidemic and pandemic influenza: annex: report of systematic literature reviews. World
471 Health Organization 2019. <https://apps.who.int/iris/handle/10665/329439> (accessed 2 Apr 2020).
- 472 12 SARS | Guidance | Interventions for Community Containment | CDC.
473 2019.<https://www.cdc.gov/sars/guidance/d-quarantine/app1.html> (accessed 24 Mar 2020).
- 474 13 Geldsetzer P. Knowledge and Perceptions of COVID-19 Among the General Public in the United
475 States and the United Kingdom: A Cross-sectional Online Survey. *Ann Intern Med* Published Online
476 First: 20 March 2020. doi:10.7326/M20-0912
- 477 14 Zhong B-L, Luo W, Li H-M, *et al.* Knowledge, attitudes, and practices towards COVID-19 among

- 478 Chinese residents during the rapid rise period of the COVID-19 outbreak: a quick online cross-
479 sectional survey. *Int J Biol Sci* 2020;**16**:1745–52. doi:10.7150/ijbs.45221
- 480 15 Kwok KO, Li KK, Chan HH, *et al.* Community responses during the early phase of the COVID-19
481 epidemic in Hong Kong: risk perception, information exposure and preventive measures. *medRxiv*
482 Published Online First:
483 2020.<https://www.medrxiv.org/content/10.1101/2020.02.26.20028217v1.abstract>
- 484 16 Hollingsworth TD, Klinkenberg D, Heesterbeek H, *et al.* Mitigation strategies for pandemic influenza
485 A: balancing conflicting policy objectives. *PLoS Comput Biol* 2011;**7**:e1001076.
486 doi:10.1371/journal.pcbi.1001076
- 487 17 Liu Y, Gayle AA, Wilder-Smith A, *et al.* The reproductive number of COVID-19 is higher compared
488 to SARS coronavirus. *J Travel Med* 2020;**27**. doi:10.1093/jtm/taaa021
- 489 18 Bai Y, Yao L, Wei T, *et al.* Presumed Asymptomatic Carrier Transmission of COVID-19. *JAMA*
490 Published Online First: 21 February 2020. doi:10.1001/jama.2020.2565
- 491 19 Vartti AM, Oenema A, Schreck M, *et al.* SARS knowledge, perceptions, and behaviors: a comparison
492 between Finns and the Dutch during the SARS outbreak in 2003. *Int J Behav Med* 2009;**16**:41–8.
493 doi:10.1007/s12529-008-9004-6
- 494 20 Wong CY, Tang CS. Practice of habitual and volitional health behaviors to prevent severe acute
495 respiratory syndrome among Chinese adolescents in Hong Kong. *J Adolesc Health* 2005;**36**:193–200.
496 doi:10.1016/j.jadohealth.2004.02.024
- 497 21 Tang CS, Wong CY. An outbreak of the severe acute respiratory syndrome: predictors of health
498 behaviors and effect of community prevention measures in Hong Kong, China. *Am J Public Health*
499 2003;**93**:1887–8. doi:10.2105/ajph.93.11.1887
- 500 22 Leung GM, Quah S, Ho L-M, *et al.* A Tale of Two Cities: Community Psychobehavioral Surveillance
501 and Related Impact on Outbreak Control in Hong Kong and Singapore During the Severe Acute
502 Respiratory Syndrome Epidemic. *Infection Control & Hospital Epidemiology*. 2004;**25**:1033–41.
503 doi:10.1086/502340
- 504 23 Organization WH, Others. *Communicating risk in public health emergencies: a WHO guideline for*
505 *emergency risk communication (ERC) policy and practice*. World Health Organization 2017.
506 <https://apps.who.int/iris/bitstream/handle/10665/259807/9789241550208-eng.pdf>
- 507 24 Sharma M, Yadav K, Yadav N, *et al.* Zika virus pandemic—analysis of Facebook as a social media
508 health information platform. *Am J Infect Control* 2017;**45**:301–2. doi:10.1016/j.ajic.2016.08.022
- 509 25 Pandey A, Patni N, Singh M, *et al.* YouTube As a Source of Information on the H1N1 Influenza
510 Pandemic. *American Journal of Preventive Medicine*. 2010;**38**:e1–3.
511 doi:10.1016/j.amepre.2009.11.007

512

513 **FIGURE CAPTIONS**

514

515 **Figure 1.** Flow chart of respondents in survey, on March 23rd, 2020.

516

517 **Figure 2.** Being informed about and belief in the effectiveness of policy recommendations during the
518 COVID-19 pandemic on March 23rd, 2020, by country.

519 Note: Response percentages are rounded and may not add up to 100%. Percentages below 5% omitted.

520

521 **Figure 3.** Individual implementation of protective measures in response to COVID-19 pandemic on
522 March 23rd, 2020, by country.

523 Note: Response percentages are rounded and may not add up to 100%. Percentages below 5% omitted.

524

525 Table 1. *Sociodemographic characteristics of respondents on March 23rd, 2020, by country.*

	Netherlands (Stage III)	Germany (Stage III)	Italy (Stage IV)	Total
No.	8,611	604	581	9,796
Gender (%)				
Male	2,492 (28.9)	206 (34.1)	192 (33.1)	2,890 (29.5)
Female	6,095 (70.8)	389 (64.4)	388 (66.8)	6,872 (70.2)
Other	24 (0.3)	9 (1.5)	1 (0.2)	34 (0.4)
Age, years (%)				
≤20 years	675 (7.8)	11 (1.8)	117 (20.1)	803 (8.2)
21-30 years	2,867 (33.3)	222 (36.8)	189 (32.5)	3,278 (33.5)
31-40 years	1,167 (13.6)	223 (36.9)	99 (17.0)	1,489 (15.2)
41-50 years	1,278 (14.8)	70 (11.6)	75 (12.9)	1,423 (14.5)
51-60 years	1,558 (18.2)	47 (7.8)	73 (12.6)	1,678 (17.1)
61-70 years	852 (9.9)	23 (3.8)	26 (4.5)	901 (9.2)
> 70 years	214 (2.5)	8 (1.3)	2 (0.3)	224 (2.3)
Daily activity (%)				
Paid job	4,885 (56.7)	388 (64.2)	321 (55.3)	5,594 (57.1)
Homemaker/Unemployed	573 (6.7)	41 (6.8)	37 (6.4)	651 (6.7)
Student	2,272 (26.4)	131 (21.7)	199 (34.3)	2,602 (26.6)
Retired	557 (6.5)	20 (3.3)	12 (2.1)	589 (6.0)
Other	324 (3.8)	24 (4.0)	12 (2.1)	360 (3.7)
Household composition (%)				
Single	1,391 (16.2)	113 (18.7)	94 (16.2)	1,598 (16.3)
Parent(s) with child(ren)	2,740 (31.8)	174 (28.8)	135 (23.2)	3,049 (31.1)
Living with partner	2,339 (27.2)	189 (31.3)	94 (16.2)	2,622 (26.8)
Shared flat (with roommates)	1,558 (18.1)	112 (18.5)	34 (5.9)	1,704 (17.4)
Other	583 (6.8)	16 (2.7)	224 (38.6)	823 (8.4)
Highest educational qualification (%)				
Primary/Secondary	635 (7.4)	132 (21.9)	301 (51.8)*	1068 (10.9)
Tertiary vocational	1,991 (23.1)	57 (9.4)		2048 (20.9)
Tertiary academic	5,985 (69.5)	415 (68.7)	280 (48.2)	6,680 (68.2)
Healthcare provider/(bio-)medical student (%)				
	1,572 (18.3)	123 (20.4)	48 (8.3)	1,743 (17.8)
Chronic illness or being in poor medical condition (%)				
	1,528 (17.7)	103 (17.1)	64 (11.0)	1,695 (17.3)
Sources used to acquire information on COVID-19 (%)				
Television	6,613 (76.8)	320 (53.0)	476 (82.0)	7,409 (75.6)
Newspaper, mobile news application	5,422 (63.0)	295 (48.8)	180 (31.0)	5,897 (60.2)
Social media	3,441 (40.0)	277 (45.9)	324 (55.8)	4,042 (41.3)
Radio	1,077 (12.5)	161 (26.7)	31 (5.3)	1,269 (13.0)
Official health hotlines	127 (1.5)	7 (1.2)	9 (1.6)	143 (1.5)
Official health websites	3,361 (39.0)	327 (54.1)	268 (46.1)	3,956 (40.4)
Healthcare professionals	381 (4.4)	26 (4.3)	44 (7.6)	451 (4.6)
People I speak to on a daily basis	2,293 (26.6)	161 (26.7)	139 (23.9)	2,593 (26.5)

For full questionnaire and wording, see Supplement 1.

* *Based on differences in the Italian education system, we considered primary, lower secondary school and upper secondary school as “Primary/Secondary” and university degrees as “Tertiary academic”.*

527 Table 2. *Being informed about and belief in the effectiveness of policy recommendations during the*
 528 *COVID-19 pandemic on March 23rd, 2020, by country.*

		Netherlands (Stage III)	Germany (Stage III)	Italy (Stage IV)
No.		8,611	604	581
Have been sufficiently informed (%)	Probably true	7,839 (91.0)	545 (90.2)	529 (91.1)
	Probably false	271 (3.2)	23 (3.8)	29 (5.0)
	Not sure	476 (5.5)	33 (5.5)	21 (3.6)
	No opinion	25 (0.3)	3 (0.5)	2 (0.3)
Belief in effectiveness of recommendations (%)				
Avoid social gatherings	Probably true	8,479 (98.9)	594 (98.5)	576 (99.7)
	Probably false	60 (0.7)	4 (0.7)	2 (0.4)
	Don't know	31 (0.4)	5 (0.8)	0 (0.0)
	Missing	41	1	3
Selective closure of public places/locations	Probably true	8,158 (95.3)	586 (97.2)	568 (98.1)
	Probably false	234 (2.7)	13 (2.2)	11 (1.9)
	Don't know	172 (2.0)	4 (0.7)	0 (0.0)
	Missing	47	1	2
Implementation of hand hygiene measures	Probably true	8,240 (96.0)	589 (98.2)	567 (98.1)
	Probably false	169 (2.0)	4 (0.7)	8 (1.4)
	Don't know	169 (2.0)	7 (1.2)	3 (0.5)
	Missing	33	4	3
Implementation of respiratory measures	Probably true	8,139 (95.0)	583 (97.3)	569 (99.0)
	Probably false	230 (2.7)	9 (1.5)	4 (0.7)
	Don't know	196 (2.3)	7 (1.2)	2 (0.4)
	Missing	46	5	6
Complete social lockdown/isolation	Probably true	5,063 (59.2)	458 (76.6)	495 (87.2)
	Probably false	1,983 (23.2)	72 (12.0)	29 (5.1)
	Don't know	1,500 (17.6)	68 (11.4)	44 (7.8)
	Missing	65	6	13

*For full questionnaire and wording, see Supplement I.
 Response percentages may not add up to 100% due to rounding.*

529 Table 3. Individual implementation of protective measures in response to COVID-19 pandemic on March
 530 23rd, 2020, by country.

No.	Netherlands (Stage III)				Germany (Stage III)				Italy (Stage IV)			
	8,611				604				581			
Personal protective behaviors (%)	Y	N	NS	NA	Y	N	NS	NA	Y	N	NS	NA
Cleaned or disinfected things you might touch more often than usual	4,083 (47.8)	3,753 (43.9)	713 (8.3)	62	292 (48.9)	264 (44.2)	41 (6.9)	7	423 (73.2)	123 (21.3)	32 (5.5)	3
Carried sanitizing hand gel with you when out and about	2,385 (28.3)	5,871 (69.6)	181 (2.2)	174	277 (47.0)	302 (51.2)	11 (1.9)	14	424 (73.5)	147 (25.5)	6 (1.0)	4
Used sanitizing hand gel to clean your hands, more often than usual	4,681 (55.3)	3,601 (42.5)	186 (2.2)	143	424 (71.5)	157 (26.5)	12 (2.0)	11	442 (76.6)	125 (21.7)	10 (1.7)	4
Reduced the amount you touch your eyes, nose and/or mouth	4,208 (49.0)	2,491 (29.0)	1,889 (22.0)	23	374 (62.2)	87 (14.5)	140 (23.3)	3	439 (75.7)	77 (13.3)	64 (11.0)	1
Followed a healthy diet or took vitamin supplements	4,548 (54.5)	3,429 (41.1)	370 (4.4)	264	317 (54.4)	226 (38.8)	40 (6.9)	21	205 (36.3)	308 (54.5)	52 (9.2)	16
Usually carried tissues with you when out and about	4,399 (52.4)	3,833 (45.7)	163 (1.9)	216	400 (69.7)	157 (27.4)	17 (3.0)	30	472 (83.0)	83 (14.6)	14 (2.5)	12
Usually used tissues when sneezing or coughing	4,542 (62.0)	2,551 (34.8)	232 (3.2)	1,286	335 (62.5)	169 (31.5)	32 (6.0)	68	468 (81.8)	82 (14.3)	22 (3.9)	9
Washed your hands with soap and water more often than usual	8,176 (95.0)	371 (4.3)	56 (0.7)	8	576 (95.7)	23 (3.8)	3 (0.5)	2	561 (96.6)	17 (2.9)	3 (0.5)	0
Kept child(ren) at home before any mandates were put in place	1,151 (53.6)	957 (44.6)	38 (1.8)	6,465	111 (58.1)	75 (39.3)	5 (2.6)	413	208 (60.6)	97 (28.3)	38 (11.1)	238
Limiting interactions with people (%)	Y	N	NS	NA	Y	N	NS	NA	Y	N	NS	NA
Cancelled or postponed a social event	8,042 (94.8)	268 (3.2)	178 (2.1)	123	580 (97.0)	11 (1.8)	7 (1.2)	6	572 (98.8)	6 (1.0)	1 (0.2)	2
Kept away from crowded places	7,834 (92.4)	367 (4.3)	277 (3.3)	133	555 (93.8)	22 (3.7)	15 (2.5)	12	575 (99.3)	3 (0.5)	1 (0.2)	2
Tried to avoid people who have the cold or flu-like symptoms	7,523 (89.0)	514 (6.1)	414 (4.9)	160	469 (81.1)	63 (10.9)	46 (8.0)	26	515 (90.2)	31 (5.4)	25 (4.4)	10
Avoiding travel (%)	Y	N	NS	NA	Y	N	NS	NA	Y	N	NS	NA
Reduced the amount of going to school/university/work	6,446 (88.0)	839 (11.5)	43 (0.6)	1,283	478 (84.9)	81 (14.4)	4 (0.7)	41	535 (94.4)	31 (5.5)	1 (0.2)	14
Reduced use of or change the way you use public transport	5,521 (89.6)	511 (8.3)	127 (2.1)	2,452	493 (91.3)	41 (7.6)	6 (1.1)	64	562 (98.6)	5 (0.9)	3 (0.5)	11
Reduced the amount you go to shops	6,888 (81.4)	1,154 (13.6)	421 (5.0)	148	401 (67.5)	134 (22.6)	59 (9.9)	10	563 (97.7)	8 (1.4)	5 (0.9)	5
Other behaviors not mentioned	2,094 (25.6)	5,335 (65.2)	760 (9.3)	422	166 (28.5)	316 (54.2)	101 (17.3)	21	254 (44.0)	274 (47.5)	49 (8.5)	4

Y = Yes; N = No; NS = Not sure; NA = Not applicable. For full questionnaire and wording, see Supplement I.
 Response percentages may not add up to 100% due to rounding.

Opened survey (n=12,018)

Did not give consent (n=248)

Did not finish (n=1,026)

Finished survey (n=10,744)

Respondents from countries with less than 500 respondents (n=948)

Respondents from countries with 500 respondents or more (n=9,796)

Netherlands (n=8,611)

Survey link was opened by...

Facebook (n=3,019)
Twitter (n=470)
Instagram (n=125)
LinkedIn (n=211)
WhatsApp (n=3,988)
News (n=216)
Other (n=582)

Germany (n=604)

Survey link was opened by...

Facebook (n=111)
Twitter (n=26)
Instagram (n=0)
LinkedIn (n=20)
WhatsApp (n=307)
News (n=2)
Other (n=138)

Italy (n=581)


Survey link was opened by...


Facebook (n=88)
Twitter (n=0)
Instagram (n=8)
LinkedIn (n=1)
WhatsApp (n=442)
News (n=4)
Other (n=38)

Netherlands
(N = 8611)

Germany
(N = 604)

Italy
(N = 581)





Response Yes No Not Sure Not Applicable