Exponential phase of covid19 expansion is driven by airport

connections

Marco Túlio Pacheco Coelho[1], João Fabricio Mota Rodrigues[1], Anderson

Matos Medina[2], Paulo Scalco[3], Levi Carina Terribile[4], Bruno Vilela[2], José

Alexandre Felizola Diniz-Filho[1] & Ricardo Dobrovolski[2]

1 Departamento de Ecologia, Universidade Federal de Goiás, CP 131, 74.001-970

Goiânia, Goiás, Brazil.

2 Instituto de Biologia, Universidade Federal da Bahia, Salvador, BA, Brazil

3. Faculdade de Administração, Economia, Ciências Contábeis (FACE), Universidade

Federal de Goiás, Goiânia, Goiás, Brazil.

4. Laboratório de Macroecologia, Universidade Federal de Jataí, Jataí, GO, Brazil

Abstract

The pandemic state of COVID-19 caused by the SARS CoV-2 put the world in

quarantine, led to hundreds of thousands of deaths and is causing an unprecedented

economic crisis. However, COVID-19 is spreading in different rates at different

countries. Here, we tested the effect of three classes of predictors, i.e., socioeconomic,

climatic and transport, on the rate of daily increase of COVID-19. We found that global

connections, represented by countries' importance in the global air transportation

network, is the main explanation for the growth rate of COVID-19 in different

countries. Climate, geographic distance and socioeconomics had a milder effect in this

big picture analysis. Geographic distance and climate were significant barriers in the

past but were surpassed by the human engine that allowed us to colonize most of our

planet land surface. Our results indicate that the current claims that the growth rate of

COVID-19 may be lower in warmer and humid tropical countries should be taken very

carefully, at risk to disturb well-established and effective policy of social isolation that

may help to avoid higher mortality rates due to the collapse of national health systems.

Keywords: COVID-19, SARS CoV-2, Exponential Growth Rates, Network Centrality,

Spatial Analysis, Socioeconomics.

Introduction

With the worldwide spread of the novel Coronavirus Disease 2019 (COVID-19), caused by the SARS-CoV-2 virus, we are experiencing a declared pandemic. One of the largest preoccupations about this new virus regards its notable ability to spread given the absence of any effective treatment, vaccine and immunity in human populations. Epidemiologists quantify the ability of infectious agents to spread by estimating the basic reproduction number (*R0*) statistic [1], which measures the average number of people each contagious person infects. According to the World Health Organization [2], the new coronavirus is transmitting at an *R0* around *1.4-2.5,* which is greater than seasonal influenza viruses that spread every year around the planet (median *R0* of 1.28, [3]). To anticipate the timing and magnitude of public interventions and mitigate the adverse consequences on public health and economy, understanding the factors associated with the survival and transmission of SARS-CoV-2 is urgent.

Because previous experimental [4], epidemiological [5, 6] and modeling [7] studies show the critical role of temperature and humidity on the survival and transmission of viruses, recent studies are testing the effect of environmental variables on SARS-CoV-2 [8, 9] and forecasting monthly scenarios of the spread of the new virus based on climate suitability [10, *but see* 11]. Although temperature and humidity are known to affect the spread and survival of other coronaviruses (i.e., SARS-CoV and MERS-CoV, [12-15] using the current occurrences of SARS-CoV-2 cases to build correlative climatic suitability models without taking into consideration connectivity among different locations and socioeconomic conditions might be inadequate.

Many factors might influence the distribution of diseases at different spatial scales. Climate might affect the spread of viruses given its known effect on biological processes that influences many biogeographical patterns, including the distribution of

diseases and human behavior (e.g., [16]). Geographic distance represents the

geographical space where the disease spread following the distribution of hosts and has

also been found to explain biogeographic patterns [17-19]. Socioeconomic

characteristics of countries could be viewed as a proxy for the ability to identify and

treat infected people and for the governability necessary to make fast political decision

and avoid the spread of new diseases. Finally, the global transportation network might

surpass other factors as it can reduce the relative importance of geographic distance and

facilitate the spread of viruses and their vectors [20,21]. According to the International

Air Transport Association [22] more than 4 billion passengers traveled abroad in 2018.

This amount of travelers reaching most of our planet's surface represents a human niche

construction (i.e. global transportation network; [23]) that facilitates the global spread

of viruses and vectors [20,21] in the same way it facilitated the spread of invasive

species and domesticated animals over modern human history [23].

The spread of SARS-CoV-2 from central China to other locations might be

strongly associated with inter-country connections, which might largely surpass the

effect of climate suitability. Thus, at this point of the pandemic, there is still a

distributional disequilibrium that can generate very biased predictions based on climatic

correlative modeling [24]. Thus, here we used an alternative macroecological approach

[25], based on the geographical patterns of exponential growth rates of the disease at

country level, to investigate variations on the growth rates of SARS-CoV-2. We studied

the effect of environment, socioeconomic and global transportation controlling for

spatial autocorrelation that could bias model significance. By analyzing these factors,

we show that the exponential growth of COVID-19 at global scale is explained mainly

by country's importance in global transportation network (i.e., air transportation).

Material and methods

We collected the number of people infected by the COVID-19 per day from the John Hopkins [26] and European Centre for Disease Prevention and Control (ECDC, [27]). This data is available for 204 countries, for which 65 had more than 100 cases recorded and for which time series had at least 30 days after the 100[th] case. We also performed the analysis considering countries with more than 50 cases, but it did not qualitatively change our results. Thus, we only show the results for countries with more than 100 cases.

In our analysis, we only used the exponential portion of the time series data (i.e. number of people infected per day) and excluded days after stabilization or decrease in total number of cases. We empirically modelled each time series using an exponential growth model for each country and calculated both the intrinsic growth rate (r) and the regression coefficient of the log growth series to be used as the response variable in our models. Because both were highly correlated (Person's r = 0.97), we used only the regression coefficient to represent the growth rate of COVID-19 in our study.

To investigate potential correlates of the virus growth rate, we downloaded climatic and socioeconomic data of each country. We used climatic data represented by monthly average minimum and maximum temperature (°C) and total precipitation (mm) retrieved from the WorldClim database (https://www.worldclim.org) [29]. We used monthly available data for the most recent year available in WorldClim. We extracted climatic data from the months of January, February, March, and December to represent the climatic conditions of the winter season in the Northern Hemisphere and the summer season in the Southern Hemisphere. From these data, we computed the mean value of climatic variables across each country. Finally, minimum and maximum temperatures were combined to estimate monthly mean temperature for December,

January, February, and March, which was used in the model along with total

precipitation for the same months. However, using different combinations of these

variables (i.e., using means of minimum or maximum temperatures, as well as

minimum or maximum for each month) did not qualitatively affect our results.

We extracted socioeconomic data for each country. Human Development Index

(HDI) rank, mean number of school years in 2015, gross national income (GNI) per

capita in 2011, population size in 2015 and average annual population growth rate

between 2010-2015 were used in our study and downloaded from the United Nations

database (http://hdr.undp.org/en/data). We also obtained a mean value of health

investment in each country by averaging the annual health investments between 2005-

2015 obtained from the World Health Organization database

(http://apps.who.int/gho/data/node.home). Due to the strong collinearity among some of

these predictors, HDI rank and mean number of school years were removed from our

final model.

Finally, we also downloaded air transportation data from the OpenFlights [30]

database regarding the airports of the world, which informs where each airport is

located including country location (7,834 airports), and whether there is a direct flight

connecting the airports (67,663 connections). We checked the Openfligths database to

make the airports and connections compatible by including missing or fixing airport

codes and removing six unidentified airport connections resulting in a total of 7,834

airports and 67,657 connections. We used this information to build an air transportation

network that reflects the existence of a direct flight between the airports while

considering the direction of the flight. Thus, the airport network is a unipartite, binary,

and directed graph where airports are nodes and flights are links (Fig 1, Fig S1). In the

following step, we collapsed the airports' network into a country-level network using

the country information to merge all the airports located in a country in a single node

(e.g., United States had 613 airports that were merged in a single vertex representing the

country). The country-level network (Fig 1, Fig S1) is a directed weighted graph where

the links are the number of connections between 226 countries which is collapsed for

the 65 countries that had more than 100 cases and for which time series data had at least

30 days after the 100th case . Afterward, we measured the countries centrality in the

network using the Eigenvector Centrality [31], hereafter centrality, that weights the

importance of a country in the network considering the number of connections with

other countries and how well connected these countries are to other countries – indirect

connections. All networks analyses were generated using package *igraph* [32].

Growth rate of COVID-19

Legend: [0.0344,0.07] (0.07,0.0861] (0.0861,0.105] (0.105,0.123] (0.123,0.179]

Fig 1. Air transportation network among 65 countries that had more than 100 cases and

for which time series data had at least 30 days after the 100[th] case. Different colours

represent modules of countries that are more connected to each other. Different sizes of

each node represent the growth rate of COVID-19 estimated for each country (See

results Fig 2).

We evaluated the relationship between the predictors (climatic, socioeconomic

and transport data) and our growth rate parameter using a standard multiple regression

(OLS) after taking into consideration the distribution of the original predictors as well

as the normality of model residuals. Moreover, OLS residuals were inspected to

evaluate the existence of spatial autocorrelation that could upward bias the significance

of predictor variables on the model using Moran's I correlograms [33]. Prior to the

analysis, we applied logarithmic (mean precipitation, total population size, and network

centrality) and square root (mean health investments) transformations to the data to

approximate a normal distribution.

Results

The models used to estimate COVID-19 growth rate on different countries

showed an average R^2 of 0.91 (SD = 0.04), varying from 0.78 to 0.99, indicating an

overall excellent performance on estimating growth rates. The geographical patterns in

the growth rates of COVID-19 cases do not show a clear trend, at least in terms of

latitudinal variation, that would suggest a climatic effect at macroecological scale (Fig.

2A).

We build one model including only climate and socioeconomic variables, which

explained *only 14%* of the variation on growth rates. This model did not have spatial

autocorrelation in the residuals. When we added country centrality (i.e. country

importance in global transportation network) as a predictor, the R^2 increased to 48.6%.

In this model, annual population growth and precipitation had positive and significant

effects (Table 1, P = 0.036, P =0.041, Fig 2), while health investments had a negative

and significant effect on growth rate (Table 1, P= 0.035, Fig 2). Here, exponential

growth rates increased *strongly* in response to countries importance in the transportation

network which has more than two times the effect size of any significant variable (Table

1). However, it is also important to note that growth rates of COVID-19 weakly

increases with increases of annual population growth and precipitation, and decreases

with higher investments in health (Table 1). Statistical coefficients were not upward

biased by spatial autocorrelation.

Fig 2. Geographical patterns of growth rate of covid-19 in the exponential phase (**A**),

the Eigenvector Centrality that represents countries' importance in global transportation

network (**B**), Annual population growth (**C**), health investments (**D**) and mean

precipitation (**E**). The relationship between growth rate and the log transformed

eigenvector centrality is showed in **F**. **G, H** and **I** are partial plots showing the

relationship between the residuals of growth rates vs log transformed eigenvector

centrality and annual population growth, health investments and mean precipitation.

Table 1. Model statistics for all variables used in the study.

	Standardized Estimate	Estimate	Std Error	t value	P-value
Intercept		0.149	0.026	3.547	< 0.001
Eigenvector Centrality	0.758	0.016	0.002	6.124	**< 0.001**
Gross National Income	0.16	0.000	0.000	1.491	0.141
Population Size	-0.096	-0.004	0.004	-0.953	0.344
Annual population growth	0.306	0.008	0.003	2.139	**0.036**
Heath investment	-0.287	-0.0004	0.000	-2.155	**0.035**
Mean Temperature	-0.127	-0.0003	0.000	-0.908	0.367
Mean Precipitation	0.253	0.007	0.003	2.091	**0.041**

Discussion

The pandemic state of SARS CoV-2 is killing hundreds of thousands of people, put the world in quarantine and is causing an unprecedented economic crisis. The rates of increase of new cases of COVID-19 is faster in some countries than others. To understand why growth rates are different among countries we investigate the effect of climatic, socioeconomical and human transportation variables that could have important roles on the exponential phase of COVID-19. At global scale, temperature, population size and Gross National Income had no significant effect on the exponential phase of COVID-19. However, annual population growth, health investments and precipitation show significant, but weak effects on growth rates. Countries' importance in the global transportation network has a key role on the severity of COVID-19 pandemic in different countries as it is strongly associated with the growth rates of the disease (Fig 2).

The centrality measure is widely used to discover distinguished nodes on many networks, including epidemiological networks (e.g., [34]). Our findings reinforce the importance of propagule pressure on disease dissemination [35, 36]. Aerial transportation is an important predictor of COVID-19 dissemination in China [37], in Brazil [38], and in Mexico [39]. It is quite likely that further phases of COVID-19 spread, in terms of peak of infections and decrease in mortality rates, will be better related to socioeconomics characteristics of each county and their political decisions when secondary transmissions were identified. We can already clearly identify the effects of adopting strong social isolation policies in China (see [37]) and, on the opposite side of this spectrum, in European countries like Italy, Spain and England [40]. Our analyses call attention to the case of Brazil, a well-connected tropical country that

presents one of the highest increase rates of COVID-19 in the tropics in its exponential

phase (Fig 2A). If decision makers take into consideration yet unsupported claims that

growth rates of COVID-19 in its exponential phase might be lower in warmer and

humid countries, we might observe terrible scenarios unrolling in tropical countries,

especially in those with limited health care structure, such as Brazil. As our results also

show, those countries that invested less in health are also the ones with faster growth

rates of covid-19 in its exponential phase (Fig 2, Table 1).

When discussing and modelling the effect of climate on SARS CoV-2 it is

important to remember that modern human society is a complex system composed of

strongly connected societies that are all susceptible to rare events. It is also critical to

consider the negative correlations between climate and local or regional socioeconomic

conditions (i.e., inadequate sanitary conditions and poor nutritional conditions) that

could easily counteract any potential climatic effect at local scales, such as lower

survival rates of viruses exposed to high humidity, temperatures and high UV

irradiation [8, 41] Tropical regions will experience mild climate conditions in a couple

of months. Thus, regardless of the influence of local environmental conditions, tropical

countries could still expect high contagious rates. In addition, our results points to a

positive effect of precipitation on growth rates, which is the contrary of what has been

suggested by climate suitability models. Finally, climatic suitability models might be

ephemeral for very mathematized modelling fields of science such as epidemiology and

virology that developed over time very realistic models that enables the possibility of

learning with parameters of similar viruses (i.e. SARS) that can definitely help and

instruct decision makers to take actions before it is too late.

Here we showed that countries' importance in the global transportation network

has a key role on COVID-19 growth rates in its exponential phase. Our results reinforce

board control measures in international airports [42, 43] during exceedingly early stages

of pandemics to prevent secondary transmissions that could lead to undesired scenarios

of rapid synchronically spread of infectious diseases in different countries. The rapid

international spread of the severe acute respiratory syndrome (SARS) from 2002 to

2003 led to extensively assessing entry screening measures at international borders of

some countries [44, 45]. The 2019-2020 world spread of COVID-19 highlights that

improvements and testing of board control measures (i.e. screening associated with fast

testing and quarantine of infected travellers) might be a cheap solution for humanity in

comparison to health systems breakdowns and unprecedented global economic crises

that the spread of infectious disease can cause. However, it is important to note that

board control of potentially infected travellers and how to effectively identify them is

still a hotly debated topic in epidemiology and there is still no consensus on accurate

methodologies for its application [46].

We do not expect that our results using a macroecological approach at a global

scale would have a definitive effect on decision-making in terms of public health in any

particular country, province, or city. However, we expect that our analyses show that

current claims that growth of COVID-19 pandemics may be lower in developing

tropical countries should be taken very carefully, at risk to disturb well-established and

effective policy of social isolation that may help to avoid higher mortality rates due to

collapse in national health systems.

Acknowledgments

This paper was developed in the context of the human macroecology project on the

National Institute of Science and Technology (INCT) in Ecology, Evolution and

Biodiversity Conservation, supported by CNPq (grant 465610/2014-5) and FAPEG

(grant 201810267000023). JAFDF, RD, LCT are also supported by CNPq productivity

scholarships. We thank Thiago F. Rangel, André Menegotto, Robert D. Morris and

Marcus Cianciaruso for their constructive comments on early version of the manuscript.

References

281 1. Delamater PL, Street EJ, Leslie TF, Yang YT, Jacobsen KH. 2019
282   Complexity of the basic reproduction number (R0). *Emerg. Infect. Dis.* **25**,
283   1–4. (doi:10.3201/eid2501.171901)
284

2. World Health Organization. 2020. Statement on the meeting of the
International Health Regulations (2005) *Emergency Committee regarding*
the outbreak of novel coronavirus (2019-nCoV). Retrieved from
https://www.who.int/news-room/detail/23-01-2020-statement-on-the-
meeting-of-the-international-health-regulations-(2005)-emergency-
committee-regarding-the-outbreak-of-novel-coronavirus-(2019-ncov)
291

292 3. Biggerstaff M, Cauchemez S, Reed C, Gambhir M, Finelli L. 2014 Estimates
293   of the reproduction number for seasonal, pandemic, and zoonotic influenza:
294   A systematic review of the literature. *BMC Infect. Dis.* **14**, 1–20.
295   (doi:10.1186/1471-2334-14-480)
296

297 4. Lowen AC, Mubareka S, Steel J, Palese P. 2007 Influenza virus transmission
298   is dependent on relative humidity and temperature. *PLoS Pathog.* **3**, 1470–
299   1476. (doi:10.1371/journal.ppat.0030151)
300

301 5. Shaman J, Pitzer VE, Viboud C, Grenfell BT, Lipsitch M. 2010 Absolute
302   humidity and the seasonal onset of influenza in the continental United States.
303   *PLoS Biol.* **8**. (doi:10.1371/journal.pbio.1000316)
304

305 6. Barreca AI, Shimshack JP. 2012 Absolute humidity, temperature, and
306   influenza mortality: 30 years of county-level evidence from the united states.
307   *Am. J. Epidemiol.* **176**, 114–122. (doi:10.1093/aje/kws259)
308

309 7. Zuk T, Rakowski F, Radomski JP. 2009 Probabilistic model of influenza
310   virus transmissibility at various temperature and humidity conditions.
311   *Comput. Biol. Chem.* **33**, 339–343.
312   (doi:10.1016/j.compbiolchem.2009.07.005)
313

314 8. Wang J, Tang K, Feng K, Lv W. 2020 High Temperature and High Humidity
315   Reduce the Transmission of COVID-19. *SSRN Electron. J.* , 1–19.
316   (doi:10.2139/ssrn.3551767)

9. Sajadi MM, Habibzadeh P, Vintzileos A, Shokouhi S, Miralles-Wilhelm F, Amoroso A. 2020 Temperature and Latitude Analysis to Predict Potential Spread and Seasonality for COVID-19. *SSRN Electron. J.* , 6–7. (doi:10.2139/ssrn.3550308)

10. Araujo MB, Naimi B. 2020 Spread of SARS-CoV-2 Coronavirus likely to be constrained by climate. *medRxiv* , 2020.03.12.20034728. (doi:10.1101/2020.03.12.20034728)

11. Chipperfield JD. 2020 On the inadequacy of species distribution models for modelling the spread of SARS-CoV-2 : response to Araújo and Naimi. *ecoRxiv*

12. Tan J, Mu L, Huang J, Yu S, Chen B, Yin J. 2005 An initial investigation of the association between the SARS outbreak and weather: With the view of the environmental temperature and its variation. *J. Epidemiol. Community Health* **59**, 186–192. (doi:10.1136/jech.2004.020180)

13. Chan KH, Peiris JSM, Lam SY, Poon LLM, Yuen KY, Seto WH. 2011 The effects of temperature and relative humidity on the viability of the SARS coronavirus. *Adv. Virol.* **2011**. (doi:10.1155/2011/734690)

14. Doremalen NV, Bushmaker T, Munster VJ. 2013 Stability of middle east respiratory syndrome coronavirus (MERS-CoV) under different environmental conditions. *Eurosurveillance* **18**, 1–4. (doi:10.2807/1560-7917.ES2013.18.38.20590)

15. Gaunt ER, Hardie A, Claas ECJ, Simmonds P, Templeton KE. 2010 Epidemiology and clinical presentations of the four human coronaviruses 229E, HKU1, NL63, and OC43 detected over 3 years using a novel multiplex real-time PCR method. *J. Clin. Microbiol.* **48**, 2940–2947. (doi:10.1128/JCM.00636-10)

16. Murray KA, Olivero J, Roche B, Tiedt S, Guégan JF. 2018 Pathogeography: leveraging the biogeography of human infectious diseases for global health management. *Ecography (Cop.).* **41**, 1411–1427. (doi:10.1111/ecog.03625)

17. Poulin R. 2003 The decay of similarity with geographical distance in parasite communities of vertebrate hosts. *J. Biogeogr.* **30**, 1609–1615. (doi:10.1046/j.1365-2699.2003.00949.x)

18. Nekola JC, White PS. 1999 The distance decay of similarity in biogeography and ecology. *J. Biogeogr.* **26**, 867–878. (doi:10.1046/j.1365-2699.1999.00305.x)

19. Warren DL, Cardillo M, Rosauer DF, Bolnick DI. 2014 Mistaking geography for biology: inferring processes from species distributions. *Trends Ecol. {&} Evol.* **29**, 572–580. (doi:10.1016/j.tree.2014.08.003)

20. Brockmann D, Helbing D. 2013 The hidden geometry of complex, network-driven contagion phenomena. *Science (80).* **342**, 1337–1342. (doi:10.1126/science.1245200)

21. Pybus OG, Tatem AJ, Lemey P. 2015 Virus evolution and transmission in an ever more connected world. *Proc. R. Soc. B Biol. Sci.* **282**, 1–10. (doi:10.1098/rspb.2014.2878)

22. International Air Transport Association. 2019. Annual Review(https://www.iata.org/contentassets/c81222d96c9a4e0bb4ff6ced0126f0bb/iata-annual-review-2019.pdf)

23. Boivin NL, Zeder MA, Fuller DQ, Crowther A, Larson G, Erlandson JM, Denhami T, Petraglia MD. 2016 Ecological consequences of human niche construction: Examining long-term anthropogenic shaping of global species distributions. *Proc. Natl. Acad. Sci. U. S. A.* **113**, 6388–6396. (doi:10.1073/pnas.1525200113)

24. De Marco P, Diniz-Filho JAF, Bini LM. 2008 Spatial analysis improves species distribution modelling during range expansion. *Biol. Lett.* **4**, 577–580. (doi:10.1098/rsbl.2008.0210)

25. Burnside WR, Brown JH, Burger O, Hamilton MJ, Moses M, Bettencourt LMA. 2012 Human macroecology: linking pattern and process in big-picture human ecology. *Biol. Rev.* **87**, 194–208. (doi:10.1111/j.1469-185X.2011.00192.x)

26. Dong E, Du H, Gardner L. 2020 An interactive web-based dashboard to track COVID-19 in real time. *Lancet Infect. Dis.* **20**, 533–534. (doi:10.1016/S1473-3099(20)30120-1)

27. European Centre for Disease Prevention and Control (ECDC). COVID-19 - Situation update – worldwide. Stockholm: ECDC; 1 Apr 2020. Available from: https://www.ecdc.europa.eu/en/geographical-distribution-2019-ncov-cases

28. Harris I, Jones PD, Osborn TJ, Lister DH. 2014 Updated high-resolution grids of monthly climatic observations - the CRU TS3.10 Dataset. *Int. J. Climatol.* **34**, 623–642. (doi:10.1002/joc.3711)

29. Fick SE, Hijmans RJ. 2017 WorldClim 2: new 1-km spatial resolution climate surfaces for global land areas. *Int. J. Climatol.* **37**, 4302–4315. (doi:10.1002/joc.5086)

30. Openflights.org database. 2014. http://openflights.org/data.html. Accessed 20 March 2020

31. Bonacich P. 1987 Power and Centrality: A Family of Measures. *Am. J. Sociol.* **92**, 1170–1182. (doi:10.1086/228631)

32. Csardi G, Nepusz T .2006. The igraph software package for complex network research. *InterJournal*, **Complex Systems**, 1695. http://igraph.org.

33. Legendre P, Lengedre L. Numerical Ecology. Elsevier Science.

34. Madotto A, Liu J. 2016 Super-Spreader Identification Using Meta-Centrality. *Sci. Rep.* **6**, 1–10. (doi:10.1038/srep38994)

35. Tian H *et al.* 2017 Increasing airline travel may facilitate co-circulation of multiple dengue virus serotypes in Asia. *PLoS Negl. Trop. Dis.* **11**, 1–15. (doi:10.1371/journal.pntd.0005694)

36. Chinazzi M *et al.* 2020 The effect of travel restrictions on the spread of the 2019 novel coronavirus (COVID-19) outbreak. *Science* **9757**, 1–12. (doi:10.1126/science.aba9757)

37. Kraemer MUG *et al.* 2020 The effect of human mobility and control measures on the COVID-19 epidemic in China. *Science* **497**, 493–497. (doi:10.1126/science.abb4218)

38. Ribeiro SP *et al.* 2020 Severe airport sanitarian control could slow down the spreading of COVID-19 pandemics in Brazil. *medRxiv* , 2020.03.26.20044370. (doi:10.1101/2020.03.26.20044370)

39. Dattilo W, Silva AC, Guevara R, McGregor-Fors I, Pontes S. 2020 COVID-19 most vulnerable Mexican cities lack the public health infrastructure to face the pandemic: a new temporally-explicit model. *medRxiv* , 1–13. (doi:10.1101/2020.04.10.20061192)

40. Enserink, M & Kupferschmidt, K. (2020). Mathematics of life and death: How disease models shape national shutdowns and other pandemic policies. *Science* (https://www.sciencemag.org/news/2020/03/mathematics-life-and-death-how-disease-models-shape-national-shutdowns-and-other#)

41. Duan S-M *et al.* 2003 Stability of SARS coronavirus in human specimens and environment and its sensitivity to heating and UV irradiation. *Biomed. Environ. Sci.* **16**, 246–255.

42. Bitar D. , Goubar A. DJC. 2009 International travels and fever screening during epidemics: a literature review on the effectiveness and potential use of non-contact infrared thermometers. *Euro Surveill* **14**, pii=19115.

43. Nishiura H, Kamiya K. 2011 Fever screening during the influenza (H1N1-2009) pandemic at Narita International Airport, Japan. *BMC Infect. Dis.* **11**, 1–11. (doi:10.1186/1471-2334-11-111)

44. Bell DM *et al.* 2004 Public health interventions and SARS spread, 2003. *Emerg. Infect. Dis.* **10**, 1900–1906. (doi:10.3201/eid1011.040729)

45. John St. RK, King A, De Jong D, Bodie-Collins M, Squires SG, Tam TWS. 2005 Border screening for SARS. *Emerg. Infect. Dis.* **11**, 6–10. (doi:10.3201/eid1101.040835)

46. Sun G., Matsui T., Kirimoto T., Yao Y., Abe S. .2017. Applications of Infrared Thermography for Noncontact and Noninvasive Mass Screening of Febrile International Travelers at Airport Quarantine Stations. In: Ng E., Etehadtavakol M. (eds) Application of Infrared to Biomedical Sciences. Series in BioEngineering. Springer, Singapore

Fig S1. Spatial pattern of the air transportation network among 65 countries that had more than 100 cases and for which time series data had at least 30 days after the 100[th] case. Red colours in different countries represent different growth rates of COVID-19 in each country. Different colours in the nodes represent modules of countries that are more connected to each other