

Social distancing to slow the U.S. COVID-19 epidemic: longitudinal pretest-posttest comparison group study

Mark J. Siedner, MD*^{¶1,2,3}, Guy Harling, ScD*^{3,4,5,6,7}, Zahra Reynolds, MPH¹, Rebecca F. Gilbert, BA¹, Sebastien Haneuse, PhD⁸, Atheendar S. Venkataramani, MD^{9,10}, Alexander C. Tsai, MD^{1,2,7}

Affiliations:

¹ Massachusetts General Hospital, Boston, MA, USA

² Harvard Medical School, Boston, MA, USA

³ Africa Health Research Institute, KwaZulu-Natal, South Africa

⁴ University College London, London, UK

⁵ MRC/Wits Rural Public Health & Health Transitions Research Unit (Agincourt), University of the Witwatersrand, Johannesburg, South Africa

⁶ Department of Epidemiology, Harvard T.H. Chan School of Public Health, Boston, MA, United States

⁷ Harvard Center for Population and Development Studies, Cambridge, MA, United States

⁸ Department of Biostatistics, Harvard T.H. Chan School of Public Health, Boston, MA, United States

⁹ Department of Medical Ethics and Health Policy, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA, United States

¹⁰ Leonard Davis Institute of Health Economics, University of Pennsylvania, Philadelphia, PA, United States

*Contributed equally

¶Corresponding Author and Address:

Mark J. Siedner MD

Medical Practice Evaluation Center

Massachusetts General Hospital

100 Cambridge Street, Suite 1600

Boston, MA 02114

msiedner@mgh.harvard.edu

Word count: 3436 words

Competing interests: ACT receives a stipend as a Specialty Consulting Editor for PLOS Medicine and serves on the journal's editorial board. All other authors declare no competing interests.

Funding Source: Sir Henry Dale Fellowship, Wellcome Trust (UK); Sullivan Family Foundation

Abstract (312 words)

Background: Social distancing measures to address the U.S. coronavirus disease 2019 (COVID-19) epidemic may have notable health and social impacts.

Methods and Findings: We conducted a longitudinal pretest-posttest comparison group study to estimate the change in COVID-19 case growth before versus after implementation of statewide social distancing measures in the U.S. The primary exposure was time before (14 days prior to, and up to 3 days after) versus after (beginning 4 days after, and up to 21 days after) implementation of the first statewide social distancing measures. Statewide restrictions on internal movement were examined as a secondary exposure. The primary outcome was the COVID-19 case growth rate. The secondary outcome was the COVID-19-attributed mortality growth rate. All states initiated social distancing measures between March 10-25, 2020. The mean daily COVID-19 case growth rate decreased beginning four days after implementation of the first statewide social distancing measures, by 0.9% per day (95% confidence interval [CI], -1.3% to -0.4%; $P < 0.001$). We did not estimate a statistically significant difference in the mean daily case growth rate before versus after implementation of statewide restrictions on internal movement (0.1% per day; 95% CI, -0.04% to 0.3%, $P = 0.14$), but there is significant difficulty in disentangling the unique associations with statewide restrictions on internal movement from the unique associations with the first social distancing measures. Beginning seven days after social distancing, the COVID-19-attributed mortality growth rate decreased by 1.7% per day (95% CI, -3.0% to -0.7%; $P < 0.001$). Our analysis is susceptible to potential bias resulting from the aggregate nature of the ecological data, potential confounding by contemporaneous changes

(e.g., increases in testing), and potential underestimation of social distancing due to spillovers across neighboring states.

Conclusions: Statewide social distancing measures were associated with a decrease in the COVID-19 epidemic case growth rate that was statistically significant and a decrease in the COVID-19-attributed mortality growth rate that was not statistically significant.

Author Summary

Why was the study done: There are few empirical data about the population health benefits of imposing statewide social distancing measures to reduce transmission of severe acute respiratory syndrome coronavirus 2, which causes coronavirus disease 2019 (COVID-19).

What did the researchers find: We compared data from each state before vs. after implementation of statewide social distancing measures to estimate changes in mean COVID-19 daily case growth rates. Growth rates declined by approximately 1% per day beginning four days (approximately one incubation period) after statewide social distancing measures were implemented. Stated differently, our model implies that social distancing reduced the total number of COVID-19 cases by approximately 1,600 reported cases at 7 days after implementation, by approximately reported 55,000 cases at 14 days after implementation, and by approximately reported 600,000 cases at 21 days after implementation.

What do these findings mean: Statewide social distancing measures were associated with a reduction in the growth rate of COVID-19 cases in the U.S. However, our analysis is susceptible to potential bias resulting from the aggregate nature of the data, potential confounding by other changes that occurred during the study period (e.g., increases in testing), and potential underestimation of social distancing due to spillovers across neighboring states.

1 **Introduction**

2

3 Modeling estimates suggest that up to 80% of Americans will be infected with severe acute

4 respiratory syndrome coronavirus 2 (SARS CoV-2) if no preventive interventions are

5 implemented [1]. In response, U.S. state governments have applied social distancing measures in

6 an attempt to limit its transmission and reduce morbidity and mortality from coronavirus disease

7 2019 (COVID-19). Such measures have been applied during prior pandemics, with moderate

8 success [2-5], and are predicted to prevent a rapid, overwhelming epidemic in modeling studies

9 [6]. Data on social distancing and its associations with the course of the COVID-19 pandemic are

10 now beginning to emerge, although no studies have examined changes in COVID-19-attributed

11 mortality as an outcome. Recent studies have shown slowed COVID-19 epidemic growth

12 coinciding with reductions in internal movement [7,8], as well as after implementation of shelter-

13 in-place orders [9-11]. Because of the economic and social costs associated with social

14 distancing measures [12,13], there is immense value in using different modalities to quantify the

15 extent to which they have benefits for epidemic control. Herein, we contribute to this body of

16 evidence, using a longitudinal pretest-posttest comparison group study design to compare the

17 daily growth rate of COVID-19 cases, and the daily growth rate of COVID-19-attributed deaths,

18 before and after implementation of social distancing measures in the U.S. Our primary aim was

19 to empirically estimate the public health impact of government mandated non-pharmacological

20 interventions in the period after their initial implementation and prior to their recently staged

21 relaxation.

22

23 **Methods**

24

25 *Data Collection*

26 We searched government websites and third-party sources to identify all statewide social
27 distancing measures implemented between January 21 and May 1, 2020 (see **Supplement Table**
28 **1**). We applied a broad definition of social distancing measures, using a previously published
29 typology [14]: closures of schools, closures of workplaces, cancellations of public events,
30 restrictions on internal movement, and closures of state borders. Restrictions on internal
31 movement, i.e., shelter-in-place orders (often referred to colloquially as “lockdowns”), are
32 generally the most restrictive of these in terms of their impacts on daily movement. To obtain
33 daily state-specific reported COVID-19 cases and deaths, we used the *New York Times* COVID-
34 19 database (<https://github.com/nytimes/covid-19-data>, last accessed May 26, 2020). Reporting
35 of cases and deaths in the *New York Times* database varies by state, but typically includes both
36 laboratory-confirmed and suspected cases, as recommended by the Council of State and
37 Territorial Epidemiologists [15].

38 Our primary outcome was the rate of change in daily COVID-19 cases in each state,
39 calculated as the natural log of cases on each date minus the natural log of cases on the prior
40 date. Analysis was restricted to days in which a state had at least 30 cumulative cases reported, to
41 minimize any effects of volatile rate changes early in the epidemic. The primary exposure of
42 interest was time, measured as a continuous variable and divided into two periods: pre-
43 implementation (14 days prior to, through three days after, implementation of the first statewide
44 social distancing measure) versus post-implementation (beginning 4 days after, and up to 21
45 days, after implementation). We selected this transition point based on previously published
46 estimates of the lower end of the 95% confidence interval of the COVID-19 incubation period
47 [16-18], which is when cases at a population level should be expected to decline in the setting of

48 a structural intervention. The latent period (between infection and the onset of infectiousness) is
49 shorter than the incubation period (between infection and the onset of symptoms), but during the
50 time period covered by this study -- and as of the time of this writing -- active asymptomatic
51 screening programs have not been implemented in the U.S., so case counts would not have been
52 expected to change until individuals became symptomatic and presented for testing. We limited
53 our analysis to 21 days after implementation to prevent any potential diluting effects resulting
54 from relaxation of some social distancing measures, which in some states began less than four
55 weeks after implementation (e.g., Alabama, Alaska, Mississippi, and South Carolina).

56 The secondary outcome was the rate of change in daily COVID-19-attributed deaths in
57 each state, calculated as the natural log of deaths on each date minus the natural log of deaths on
58 the prior date. Analysis was restricted to days in which a state had at least 30 cumulative deaths
59 reported. In the analysis of COVID-19-attributed deaths, there was less certainty about the
60 hypothesized lag between implementation of social distancing and observed changes in daily
61 COVID-19-attributed deaths. The median time from symptom onset to death varies widely in the
62 literature. The COVID-19 Surveillance Group [19] has estimated a median of 8 days from
63 symptom onset to death in Italy, while estimates from China are approximately twice that
64 amount [20-22]. After incorporating a 3-5 incubation period, we hypothesized that a beneficial
65 impact on COVID-19-attributed mortality, if any, would be observed no sooner than 7 days and
66 no later than 14 days after implementation. Therefore, we took a more exploratory approach to
67 modeling the time from enactment of statewide social distancing, selecting time periods for the
68 spline terms (described below) every 3 days during days 4-17 (i.e., 4, 7, 10, 14, and 17 days) and
69 allowed for observation in the post-intervention period up through 30 days.

70 As a secondary exposure, we examined the dates of statewide restrictions on internal
71 movement. For states that did not implement statewide restrictions on internal movement during
72 the study period, we set day 0 as 14 days after the last state implemented a restriction on internal
73 movement (April 21, 2020) and attributed the prior 14 days to the pre-implementation period.

74

75 *Statistical Analysis*

76 We fit mixed effects linear regression models, specifying the log difference in daily cases
77 as the outcome of interest and a random effect for state to allow for within-state correlation of
78 cases over time. Explanatory variables included time in days, implementation period, and a time-
79 by-implementation period product term. This analysis was not conducted as part of a funded
80 study or pre-planned/registered study protocol. However, we followed a clear analysis plan (see
81 **Supplement**), with minor changes made in response to a rapidly changing epidemic and in
82 response to editorial and reviewer feedback. The initial analysis, described in the manuscript we
83 deposited with the *medRxiv* preprint server on April 8, 2020 [23], included data on statewide
84 social distancing measures implemented between January 21 and March 30, 2020, and
85 COVID-19 cases through March 31. Prior to submission of the current manuscript, we updated
86 the dataset and extended the study period to include social distancing measures implemented up
87 to April 8 and COVID-19 cases and deaths up to April 8. In response to editorial and reviewer
88 feedback, we further updated the dataset and extended the study period to include social
89 distancing measures implemented up to May 1 and COVID-19 cases and deaths up to May 26,
90 expanded the analysis to include growth in COVID-19-attributed deaths as a secondary outcome,
91 and added sensitivity analyses investigating a range of plausible incubation periods and an event
92 study specification (see below).

93 To assess the robustness of our findings, we conducted several sensitivity analyses. First,
94 to adjust for potential confounding by population density, we adjusted our estimates by state-
95 level population density [24,25]. To account for weekly periodicity that could also coincide with
96 implementation of social distancing measures, we also adjusted for day of the week [26,27].
97 Second, to assess the extent to which early vs. late implementation of social distancing measures
98 modified the effects of these actions [28], we stratified our estimates by the size of the epidemic
99 in the state. Third, although implementation of social distancing measures is most likely to show
100 population-level effectiveness in reducing coronavirus transmission beginning 3 days later at the
101 lower bound of its estimated incubation period [16], we refitted the regression models specifying
102 a range of different incubation periods.

103 Finally, to assess the extent to which our estimates may have been driven by model
104 specification, we replaced the longitudinal pretest-posttest comparison group study design with a
105 multivariable regression model in which the primary explanatory variables of interest were a
106 series of binary indicators denoting each day before versus after implementation of the first
107 statewide social distancing measures (often described in the econometrics literature as an event
108 study specification [29-31]). This approach compares daily case growth before versus after
109 implementation of the first statewide social distancing measures in states that implemented such
110 measures versus daily case growth in states that did not implement such measures. Our
111 regression model included state fixed effects, to adjust for potential confounding from time-
112 invariant state-level factors or baseline differences in population socioeconomic or health
113 characteristics; and linear and quadratic terms for time (days) to adjust for nationwide secular
114 trends in the outcomes. We computed 95% confidence intervals adjusted for clustering within
115 states, the geographical level at which exposure occurred [32].

116

117 *Ethics Statement*

118 This ecological analysis was based on publicly available data and was exempt from
119 ethical review.

120

121 **Results**

122

123 A complete list of dates of statewide social distancing measures, by type of measure and state, is
124 contained in the **Supplement (Supplement Table 1)**. During March 10–25, all 50 states and the
125 District of Columbia implemented at least one statewide social distancing measure (**Supplement**
126 **Figure 1**). The most widely enacted measures on the first date of implementation were
127 cancellations of public events (34/51 [67%]) and closures of schools (26/51 [51%]). The first
128 social distancing measures were implemented when the median epidemic size was 35 cases
129 (interquartile range [IQR], 17-72).

130 **Figure 1A** shows the mean COVID-19 daily case growth rate mapped against the date of
131 the first statewide distancing measures. At the date of implementation of the first social
132 distancing measure, states had a mean daily case growth rate of 30.8% (95% confidence interval
133 [CI], 29.1-32.6; **Table 1**), corresponding to a doubling of total cases every 3.3 days. From
134 fourteen days prior, through three days after, implementation of the first social distancing
135 measure, the mean daily case growth rate did not change (-0.2% per day; 95% CI, -0.6% to
136 0.3%; $P=0.51$). Beginning four days after implementation of the first statewide social distancing
137 measure, the mean daily case growth rate decreased by an additional 0.9% per day (95% CI,
138 -0.4% to -1.4%; $P<0.001$). This estimate corresponds to a mean daily case growth rate that had

139 declined to 26.5% (doubling of total cases every 3.8 days) by day 7 after enactment of the first
140 statewide social distancing measures, to 19.6% (doubling time of 5.1 days) by day 14, and to
141 12.7% (doubling time of 7.9 days) by day 21.

142 As of May 1, nearly all (45 [90%]) states had implemented a statewide restriction on
143 internal movement. These restrictions on internal movement were implemented a median of 11
144 days after the first statewide social distancing measure (IQR 8-15) was implemented in the
145 respective states, when the median epidemic size was 937 cases (IQR, 225-1414). The mean
146 daily case growth rate had already been declining at a mean rate of -0.8% per day during the 14
147 days prior to implementation of statewide restrictions on internal movement (95% CI, -0.9% to
148 -0.7%; $P<0.001$) (**Table 1, Figure 1B**). There was a drop detected three days after statewide
149 restrictions on internal movement were implemented (-3.1%; 95% CI, -4.7% to -1.5%, $P<0.001$),
150 but no statistically significant difference in the rate of change from before versus after their
151 implementation (0.1% per day; 95% CI, -0.04% to 0.3%, $P=0.14$). As discussed in more detail
152 below, there is substantial difficulty in disentangling the unique associations with statewide
153 restrictions on internal movement from the unique associations with the first social distancing
154 measures.

155
156 In the analysis of the secondary outcome, change in daily COVID-19-attributed deaths,
157 given the uncertainty in the hypothesized lag between implementation of social distancing and
158 observed changes (if any) in daily COVID-19-attributed deaths, we explored a range of
159 thresholds. As shown in **Table 2**, by seven days after implementation of the first statewide social
160 distancing measure, the mean daily growth rate in COVID-19-attributed deaths decreased by
161 2.0% per day (95% CI, -3.0% to -0.9%; $P<0.001$). By 14 days the estimated association was no

162 longer statistically significant (-1.0% per day; 95% CI, -0.2% to 0.1%; $P=0.09$). No additional
163 statistically significant benefit was found after 7 days of implementation of statewide restrictions
164 on internal movement.

165 Sensitivity analyses suggested our estimates were not sensitive to inclusion of additional
166 covariates, did not differ by size of the epidemic at implementation, and were consistent with the
167 known incubation period (**Supplement Tables 2, 3, and 4**). In the event study specification
168 (**Supplement Figure 2**), mean daily case growth was negative by day 4, and the estimates were
169 statistically significant by day 8, consistent with the primary analysis. The event study analysis
170 for change in daily COVID-19-attributed deaths also produced estimates qualitatively similar to
171 the primary analysis, although with slightly larger confidence intervals given the smaller number
172 of events (**Supplement Figure 3**).

173

174 **Discussion**

175

176 In this longitudinal pretest-posttest comparison group study, we found that implementation of
177 social distancing measures was associated with a reduction in the mean daily growth rate of
178 COVID-19 cases and in the mean daily growth rate of COVID-19-attributed deaths. Our
179 estimates imply a more than doubling in the doubling time (from 3.8 days to 8.0 days) by three
180 weeks following the implementation of social distancing measures. Assuming a cumulative
181 epidemic size of 4,125 reported cases (equivalent to the cumulative number of cases in the U.S.
182 at the time of implementation in each state), the reduction in growth rate we estimated
183 corresponds to a difference between 26,281 reported cases with no social distancing versus
184 24,625 reported cases at 7 days after implementation, a difference between 158,518 reported

185 cases with no social distancing versus 102,223 reported cases at 14 days after implementation,
186 and a difference between 904,773 reported cases with no social distancing versus 283,161
187 reported cases at 21 days after implementation. Stated differently, our model implies that social
188 distancing reduced the total number of reported COVID-19 cases by approximately 1,600 cases
189 at 7 days after implementation, by approximately 56,000 reported cases at 14 days after
190 implementation, and by approximately 621,000 reported cases at 21 days after implementation.

191 These results are consistent with both the theoretical effect of social distancing on
192 epidemic spread [6] and the historical benefit observed with the implementation of such
193 interventions during prior epidemics of communicable diseases [28]. They also are largely in
194 keeping with recent data on the impacts of social distancing measures in the U.S. on both
195 reduced mobility [7,8], and case growth rates [9-11], with generally similar effect sizes. Our
196 study extends this literature by further examining COVID-19-attributed mortality as an outcome.
197 The associations between social distancing and case growth rates were most apparent at the
198 lower end of the incubation period that has been estimated based on publicly available data, with
199 some evidence that the change in growth rates may have started even earlier. We suspect that this
200 observation may have resulted from self-imposed social distancing that has reportedly occurred
201 prior to government-issued mandates [33].

202 Our findings should be interpreted with the following limitations in mind. Our estimates
203 would be biased toward the null if: 1) state and local governments had intensified social
204 distancing measures in response to a worsening epidemic, 2) there were substantial violations of
205 the stable unit treatment value assumption (e.g., workplace closures of large employers that had
206 spillover effects across state lines), or 3) surveillance and testing intensified during the study
207 period (thereby resulting in increased case reporting). Moreover, statewide restrictions on

208 internal movement were often implemented after other social distancing measures had already
209 been applied, further biasing our estimate toward the null. Estimates of cases and deaths in our
210 model include those that are both laboratory-confirmed and suspected by health departments, but
211 they are both likely to be under-estimates, due to limitations in testing, the presence of
212 asymptomatic cases, and the occurrence of deaths that are not attributed to COVID-19 [34,35].
213 Nonetheless, our analyses focus on day-to-day changes in the growth rate of cases and deaths, so
214 underreporting would only bias our results if reported versus true outcomes systematically
215 differed from prior to versus after the implementation of social distancing measures. In contrast,
216 our projected estimates of cases prevented are likely to be highly conservative, because they are
217 modeled based on reported cases. While some studies have suggested that cases have been
218 under-reported in the U.S. by 1-2 orders of magnitude [36], deaths appear to be under-counted by
219 approximately 25% [34]. Finally, our estimates of the associations between social distancing and
220 changes in mean daily case growth rates (and the corresponding number of cases averted) cannot
221 be extrapolated linearly beyond the 21-day post-implementation analysis window. At the time of
222 this analysis, states had begun to enact social relaxation measures, which necessarily prevents
223 drawing conclusions about the long-term associational effects of these interventions in isolation.

224 Our study does not ascertain which of the statewide social distancing measures were most
225 effective in reducing mean daily COVID-19 case growth. A variety of social distancing measures
226 were used, often simultaneously, making it difficult to disentangle their independent
227 associations. It is also possible that some state residents changed their behaviors in response to
228 local (e.g., county-level) social distancing measures enacted prior to the statewide measures, or
229 that some state residents changed their behaviors independently of social distancing measures
230 enacted at any level of jurisdiction. The latter phenomenon has been observed empirically in

231 other settings [37,38]. Such behavior change would be consistent with our sensitivity analysis
232 suggesting a signal that extends back to a 0-day incubation period.

233 Finally, our analysis cannot answer questions about the appropriate time for rescinding
234 social distancing measures. While our findings demonstrate that early mitigation efforts have
235 yielded a substantial population health benefit, these benefits should be weighed against their
236 costs. The costs of social distancing are likely to exacerbate the confluence of longstanding
237 economic, social, and health decline that is already occurring in the U.S. [39-41], brought into
238 even sharper relief given emerging data about racial, ethnic, and socioeconomic disparities in the
239 incidence of COVID-19-related burden [42,43]. Moreover, there is currently robust debate about
240 the extent to which legal authority to initiate or rescind social distancing measures resides with
241 the federal or local government [44]. Additional study may be possible in the coming weeks to
242 months if some statewide social distancing measures are relaxed while others are retained, or if
243 measures are relaxed in some local jurisdictions while being retained in others. The use of cross-
244 country data may also be helpful in this regard, albeit limited by any substantial within-country
245 heterogeneity (such as has been observed in the diversity of uncoordinated U.S. federal and state
246 responses to the epidemic).

247 Our finding that implementation of statewide social distancing was associated with a
248 reduction in the mean daily growth rate of COVID-19-attributed deaths should be interpreted
249 with more caution, given the uncertainty in published estimates about the median time from
250 symptom onset to death [19-21]. The strongest associations we estimated, in terms of both
251 statistical significance and magnitude, occurred at seven days after implementation. While this
252 change could be reflective of what is known about time to death for the median hospitalized

253 patient, other plausible thresholds were not associated with a statistically significant reduction in
254 the mean daily growth rate of COVID-19-attributed deaths.

255 The finding that implementation of statewide restrictions on internal movement was
256 associated neither with a statistically significant reduction in the mean daily growth rate of
257 COVID-19 cases nor with a statistically significant reduction in the mean daily growth rate of
258 COVID-19-attributed deaths warrants additional scrutiny. Identification of the unique effect of
259 statewide restrictions on internal movement, separate from the initial social distancing measures,
260 is a difficult undertaking given that they were implemented a median of 10 days later. There
261 were no states in which statewide restrictions on internal movement were implemented without
262 the prior implementation of other social distancing measures (e.g., cancellations of public events
263 and closures of schools). As a result, we could not test hypotheses about the independent causal
264 effects of statewide restrictions on internal movement, but rather could only ascertain the extent
265 to which they were associated with additional reductions in the COVID-19 case growth rate, i.e.,
266 beyond those generated by the initial statewide social distancing measures. Moreover, the null
267 finding should be interpreted in light of the other likely biases toward the null previously
268 discussed.

269 In summary we demonstrate that the U.S. COVID-19 epidemic growth rate began to
270 decline within approximately one incubation period following the initiation of statewide social
271 distancing measures. Exploratory findings suggest that social distancing may also have had
272 lagged mortality benefits. Future work should consider the potential public health benefits of
273 relaxing these measures and, specifically, to what extent and how differential relaxation of
274 certain measures promotes persistent epidemic control.

References

1. Kissler SM, Tedijanto C, Goldstein E, Grad YH, Lipsitch M. Projecting the transmission dynamics of SARS-CoV-2 through the postpandemic period. *Science*. 2020;368(6493):860-8.
2. Markel H, Lipman HB, Navarro JA, Sloan A, Michalsen JR, Stern AM, et al. Nonpharmaceutical interventions implemented by US cities during the 1918-1919 influenza pandemic. *JAMA*. 2007;298(6):644-54.
3. Bootsma MC, Ferguson NM. The effect of public health measures on the 1918 influenza pandemic in U.S. cities. *Proc Natl Acad Sci U S A*. 2007;104(18):7588-93.
4. Ferguson NM, Cummings DA, Cauchemez S, Fraser C, Riley S, Meeyai A, et al. Strategies for containing an emerging influenza pandemic in Southeast Asia. *Nature*. 2005;437(7056):209-14.
5. Taubenberger JK, Morens DM. 1918 Influenza: the mother of all pandemics. *Emerg Infect Dis*. 2006;12(1):15-22.
6. Teslya A, Pham TM, Godijk NG, Kretzschmar ME, Bootsma MCJ, Rozhnova G. Impact of self-imposed prevention measures and short-term government intervention on mitigating and delaying a COVID-19 epidemic [preprint]. medRxiv. Epub 27 Mar 2020. doi:10.1101/2020.03.12.20034827.
7. Andersen M. Early evidence on social distancing in response to COVID-19 in the United States [preprint]. SSRN Working Paper No. 3569368. Amsterdam: Elsevier, 2020. Available at: <https://ssrn.com/abstract=3569368>. Accessed 2 Jun 2020.

8. Abouk R, Heydari B. The immediate effect of COVID-19 policies on social distancing behavior in the United States [preprint]. medRxiv. Epub 28 Apr 2020.
doi:10.1101/2020.04.07.20057356.
9. Courtemanche C, Garuccio J, Le A, Pinkston J, Yelowitz A. Strong social distancing measures in the United States reduced the COVID-19 growth rate. *Health Aff (Millwood)*. 2020;7: Epub 14 May 2020. doi:10.1377/hlthaff.2020.00608.
10. Dave DM, Friedson AI, Matsuzawa K, Sabia JJ. When do shelter-in-place orders fight COVID-19 best? Policy heterogeneity across states and adoption time [preprint]. NBER Working Paper No. 27091. Cambridge: National Bureau of Economic Research, 2020. Available at: <https://www.nber.org/papers/w27091>. Accessed 2 Jun 2020.
11. Hsiang S, Allen D, Annan-Phan S, Bell K, Bolliger I, Chong T, et al. The effect of large-scale anti-contagion policies on the COVID-19 pandemic. *Nature*. 2020: Epub 8 Jun 2020.
doi:10.1038/s41586-020-2404-8.
12. Bayham J, Fenichel EP. Impact of school closures for COVID-19 on the US health-care workforce and net mortality: a modelling study. *Lancet Public Health*. 2020;5(5):e271-78.
13. Eichenbaum MS, Rebelo S, Trabandt M. The macroeconomics of epidemics [preprint]. NBER Working Paper No. 26882. Cambridge: National Bureau of Economic Research, 2020. Available at:
<https://www.nber.org/papers/w26882>.
14. Petherick A, Hale T, Phillips T, Webster S. Variation in government responses to COVID-19 [preprint]. Blavatnik School Working Paper. Oxford: Blavatnik School of Government, University of Oxford, 2020. Available at:
<https://www.bsg.ox.ac.uk/research/publications/variation-government-responses-covid-19>. Accessed 2 Jun 2020.

15. Council of State and Territorial Epidemiologists. Standardized surveillance case definition and national notification for 2019 novel coronavirus disease (COVID-19). Washington, D.C.: Office of the Assistant Secretary for Preparedness and Response, U.S. Department of Health and Human Services, 2020. Available at: <https://asprtracie.hhs.gov/technical-resources/resource/8322/standardized-surveillance-case-definition-and-national-notification-for-2019-novel-coronavirus-disease-covid-19>. Accessed 2 Jun 2020.
16. Lauer SA, Grantz KH, Bi Q, Jones FK, Zheng Q, Meredith HR, et al. The incubation period of coronavirus disease 2019 (COVID-19) from publicly reported confirmed cases: estimation and application. *Ann Intern Med.* 2020;172(9):577-82.
17. Backer JA, Klinkenberg D, Wallinga J. Incubation period of 2019 novel coronavirus (2019-nCoV) infections among travellers from Wuhan, China, 20-28 January 2020. *Euro Surveill.* 2020;25(5).
18. Guan WJ, Ni ZY, Hu Y, Liang WH, Ou CQ, He JX, et al. Clinical characteristics of coronavirus disease 2019 in China. *N Engl J Med.* 2020;382(18):1708-20.
19. COVID-19 Surveillance Group. Characteristics of COVID-19 patients dying in Italy. Report based on available data on March 20th, 2020 [preprint]. Rome: Istituto Superiore di Sanità; 2020. Available from: https://www.epicentro.iss.it/coronavirus/bollettino/Report-COVID-2019_20_marzo_eng.pdf. Accessed 12 Apr 2020.
20. Zhou F, Yu T, Du R, Fan G, Liu Y, Liu Z, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet.* 2020;395(10229):1054-62.

21. Verity R, Okell LC, Dorigatti I, Winskill P, Whittaker C, Imai N, et al. Estimates of the severity of coronavirus disease 2019: a model-based analysis. *Lancet Infect Dis.* 2020;20(6):669-77.
22. Wu JT, Leung K, Bushman M, Kishore N, Niehus R, de Salazar PM, et al. Estimating clinical severity of COVID-19 from the transmission dynamics in Wuhan, China. *Nat Med.* 2020;26(4):506-10.
23. Siedner MJ, Harling G, Reynolds Z, Gilbert RF, Venkataramani A, Tsai AC. Social distancing to slow the U.S. COVID-19 epidemic: an interrupted time-series analysis [preprint]. medRxiv. Epub 8 Apr 2020. doi:10.1101/2020.04.03.20052373.
24. Lee VJ, Aguilera X, Heymann D, Wilder-Smith A, Lancet Infectious Diseases C. Preparedness for emerging epidemic threats: a Lancet Infectious Diseases Commission. *Lancet Infect Dis.* 2020;20(1):17-9.
25. Grassly NC, Fraser C. Mathematical models of infectious disease transmission. *Nature reviews Microbiology.* 2008;6(6):477-87.
26. Lewis MD, Pavlin JA, Mansfield JL, O'Brien S, Boomsma LG, Elbert Y, et al. Disease outbreak detection system using syndromic data in the greater Washington DC area. *Am J Prev Med.* 2002;23(3):180-6.
27. Buckingham-Jeffery E, Morbey R, House T, Elliot AJ, Harcourt S, Smith GE. Correcting for day of the week and public holiday effects: improving a national daily syndromic surveillance service for detecting public health threats. *BMC Public Health.* 2017;17(1):477.
28. Hatchett RJ, Mecher CE, Lipsitch M. Public health interventions and epidemic intensity during the 1918 influenza pandemic. *Proc Natl Acad Sci U S A.* 2007;104(18):7582-7.

29. Sullivan D, von Wachter T. Job displacement and mortality: an analysis using administrative data. *Q J Econ.* 2009;124(3):1265-306.
30. Jacobson L, LaLonde RJ, Sullivan DG. Earnings losses of displaced workers. *Am Econ Rev.* 1993;83(4):685-709.
31. Goodman-Bacon A. Difference-in-differences with variation in treatment timing [preprint]. NBER Working Paper No. 25018. Cambridge: National Bureau of Economic Research, 2018. Available at: <https://www.nber.org/papers/w25018>. Accessed 2 Jun 2020.
32. Bertrand M, Duflo E, Mullainathan S. How much should we trust differences-in-differences estimates? *Q J Econ.* 2004;119(1):249-75.
33. Brzezinski A, Deiana G, Kecht V, Van Dijke D. The COVID-19 pandemic: government vs. community action across the United States [preprint]. INET Oxford Working Paper No. 2020-06. Oxford: Institute for New Economic Thinking at the Oxford Martin School, University of Oxford, 2020. Available at: <https://www.inet.ox.ac.uk/publications/no-2020-06-the-covid-19-pandemic-government-vs-community-action-across-the-united-states/>. Accessed 2 Jun 2020.
34. Weinberger D, Cohen T, Crawford F, Mostashari F, Olson D, Pitzer VE, et al. Estimating the early death toll of COVID-19 in the United States [preprint]. medRxiv. Epub 29 Apr 2020. doi:10.1101/2020.04.15.20066431.
35. Rivera R, Rosenbaum J, Quispe W. Excess mortality in the United States during the peak of the COVID-19 pandemic [preprint]. medRxiv. Epub 28 May 2020. doi:10.1101/2020.05.04.20090324.
36. Lu FS, Nguyen AT, Link N, Santillana M. Estimating the prevalence of COVID-19 in the United States: three complementary approaches [preprint]. medRxiv. Epub 23 Apr 2020. doi:10.1101/2020.04.18.20070821.

37. Cowling BJ, Ali ST, Ng TWY, Tsang TK, Li JCM, Fong MW, et al. Impact assessment of non-pharmaceutical interventions against coronavirus disease 2019 and influenza in Hong Kong: an observational study. *Lancet Public Health*. 2020;5(5):e279-88.
38. Lewnard JA, Liu VX, Jackson ML, Schmidt MA, Jewell BL, Flores JP, et al. Incidence, clinical outcomes, and transmission dynamics of severe coronavirus disease 2019 in California and Washington: prospective cohort study. *BMJ*. 2020;369:m1923.
39. Case A, Deaton A. *Deaths of despair and the future of capitalism*. Princeton: Princeton University Press; 2020.
40. Venkataramani AS, Bair EF, O'Brien RL, Tsai AC. Association between automotive assembly plant closures and opioid overdose mortality in the United States: a difference-in-differences analysis. *JAMA Internal Medicine*. 2020;180(2):254-62.
41. Muennig PA, Reynolds M, Fink DS, Zafari Z, Geronimus AT. America's declining well-being, health, and life expectancy: not just a white problem. *Am J Public Health*. 2018;108(12):1626-31.
42. Chin T, Kahn R, Li R, Chen JT, Krieger N, Buckee CO, et al. U.S. county-level characteristics to inform equitable COVID-19 response [preprint]. *medRxiv*. Epub 11 Apr 2020. doi:10.1101/2020.04.08.20058248.
43. Unequal protection. *The Economist*. April 18-24, 2020:17-8.
44. Fritze J. Authority to reopen U.S. for business in question. *USA Today*. April 14, 2020: E2.

Figure 1A-B. Scatter plots and predictive margins with 95% confidence interval derived from regression models of the daily COVID-19 growth rate pre vs. post-implementation of the first statewide social distancing measures (1A) and statewide restrictions on internal movement (1B). The red line indicates the date of implementation in each state. The green dashed line is 4 days after implementation of the social distancing measure.




Table 1. Linear regression models for the growth in mean daily COVID-19 cases before vs. after implementation of the first statewide social distancing measure and statewide restrictions on internal movement

	First statewide social distancing measure			Statewide restriction on internal movement		
	b	95% CI	P-value	b	95% CI	P-value
Constant	0.309	0.291, 0.326	<0.001	0.183	0.164, 0.201	<0.001
Time (days relative to intervention)	-0.002	-0.006, 0.003	0.44	-0.008	-0.009, -0.006	<0.001
Post-intervention period	0.006	-0.017, 0.029	0.59	-0.031	-0.047, -0.015	<0.001
Time × post-intervention period	-0.009	-0.013, -0.004	<0.001	0.001	-0.00, 0.003	0.14

Notes: b, estimated regression coefficient. CI, confidence interval. The post-intervention period begins four days after social distancing measures were implemented to account for the disease incubation period.

Table 2. Linear regression models for growth in mean daily COVID-19-attributed deaths before vs. after implementation of the first statewide social distancing measure and statewide restrictions on internal movement, assuming a range of days between symptom onset and death

Time at which post-intervention period begins	First statewide social distancing measure			Statewide restriction on internal movement		
	b	95% CI	P-value	b	95% CI	P-value
4 days	-0.018	-0.033, -0.0-2	0.02	-0.005	-0.001, -0.002	0.005
7 days	-0.020	-0.030, -0.009	<0.001	-0.002	-0.06, 0.002	0.29
10 days	-0.010	-0.021, 0.015	0.09	-0.000	-0.002, 0.002	0.98
14 days	0.001	-0.004, 0.007	0.78	0.002	0.000, 0.004	0.02
17 days	0.001	-0.003, 0.006	0.53	0.003	0.001, 0.006	0.001

Notes: b, estimated regression coefficient; CI, confidence interval. Each row corresponds to a separate regression model fitted to the data that includes 3 variables: time in days, implementation period, and a time-by-implementation period product term. The post-intervention period specified in the regression models varies from 4-17 days after social distancing measures were implemented to account for the disease incubation period as well as uncertainty in the expected time from symptom onset to death.