Fast spread of COVID-19 in Europe and the US and its implications: even modest public health goals require comprehensive intervention

Authors: Ruian Ke, PhD[1,*], Steven Sanche, PhD[1,2], Ethan Romero-Severson, PhD[1], Nick Hengartner, PhD[1]

Affiliations:

[1]T-6 Theoretical Biology and Biophysics, Theoretical Division, Los Alamos National Laboratory, NM87545, USA.

[2]T-CNLS Center for Nonlinear Studies, Los Alamos National Laboratory, NM87545, USA.

[*]Correspondences should be addressed to:

Ruian Ke

Email: rke@lanl.gov

Phone: 1-505-667-7135

Mail: Mail Stop K710,

T-6 Theoretical Biology and Biophysics,

Los Alamos National Laboratory,

NM87545, USA.

Word counts:

<u>Abstract</u>: 247   <u>Main text</u>: 3,425

Summary

Background The COVID-19 pandemic caused more than 800,000 infections and 40,000 deaths by the end of March 2020. However, some of the basic epidemiological parameters, such as the exponential epidemic growth rate, are not well estimated, partially because of confounding factors in data collection during a rapidly growing outbreak, such as underreporting, delays in case confirmation and changes in surveillance intensities.

Methods We developed an inference approach using a mathematical model to control for these confounding factors. We fitted the model to both infection incidence and death count data collected from eight European countries and the US. Public health implications of empirical estimates were examined using simulations.

Findings In all countries, the early epidemic period was characterized by exponential growth with rates between 0·19-0·29/day (epidemic doubling times between 2·4-3·7 days). However, the proportion of cases that had been detected was low (less than 20% of cases detected) except for Germany (23·1%; CI: 5-85%). With such high epidemic growth rates, moderate intervention efforts will have little impact on the public health outcome; high levels of efforts to achieve greater than 77-86% reduction in transmission are needed, no matter the goal is to slowdown the growth to protect a large fraction of population from infection within 18 months or to reverse the growth all together.

Interpretation The extremely fast spread of COVID-19 in Europe and the US suggest a highly infectious virus with a high R_0. Early, strong and comprehensive intervention efforts are necessary, whether the aim is mitigation or containment.

Introduction

COVID-19 originated in Wuhan China in Dec, 2019[1]. It has spread rapidly and caused a global pandemic within a short period of time. As of March 31, 2020, the global pandemic lead to more than 800,000 total confirmed cases and 40,000 deaths. Estimation of the rate of early epidemic spread in Wuhan, China, lead to different conclusions. Initially, it was suggested that the epidemic grew at 0·1-0·14/day, leading to an epidemic doubling time of 5-7 days[2-5]. However, using domestic travel data and two distinct approaches, we estimated that the epidemic in Wuhan grew much faster than initially estimated, and the growth rate is likely to be between 0·21-0·3/day, translating to a doubling time between 2·3 to 3·3 days, and an R_0 approximately at 5·7 with a large confidence interval[6]. A fast epidemic spread is consistent with multiple other lines of evidence, such as the rapid increase of the epidemic curve by symptom onset published by China CDC[7] and the growth in the number of death cases in Hubei, China during late January 2020[6]. However, it was not clear whether COVID-19 can spread in other countries as fast as in Wuhan, China.

Accurate estimation of the rate of epidemic growth is important for many practical aspects. First, it is crucial for forecasting the epidemic trajectory, the burden on health care systems and potential health and economic damage, so that appropriate and timely responses can be prepared. Second, it sets the baseline for evaluation of effectiveness of intervention strategies. Third, it is important for accurate estimation of the basic reproductive number, R_0, which in turn is used for many control measures, including evaluation of the vaccine/herd immunity threshold needed to stop transmission[6,8]. However, a major challenge to the inference of the growth of COVID-19 is that as a result of a fast-growing outbreak and a sizable infected population with no or mild-to-moderate symptoms[9,10], case confirmation data is influenced by many factors in addition to the true epidemic growth, including substantial underreporting[11], i.e. low detection rate, changes in surveillance intensity and delays in case confirmation. Simply fitting an exponential curve to case confirmation data may lead to erroneous conclusions when confounding factors are not taken into account or estimated from other sources of data.

Here, we argue that because death and the cause of death are usually recorded reliably and are less affected by surveillance intensity changes or delay in confirmation than case counts, the time series of death counts reflects the growth of an epidemic reliably, with a delay in onset determined by the time between infection to death. Based on this idea, we designed a simple methodology to disentangle the epidemic growth from confounding factors, such as underreporting, delays in case confirmation and changes in surveillance intensity. We fit models to both case incidence data and death count data collected from eight European countries and the US in March 2020. We show that in most countries, the detection rate of infected individuals is in general low, and COVID-19 spreads very fast in these countries. For such a fast-epidemic growth, our results suggest that very strong and active control measures need to be implemented as early as possible regardless of the public health goal (e.g. mitigation versus containment), and that moderate control measures will not achieve measurable public health benefit.

Methods

Data

We collected daily case confirmation and death count data for the US and 8 countries in Europe from the John Hopkins CSSE (Center for Systems Science and Engineering) database

101 (). The data is accessed and extracted on March
102 31, 2020. The data consists of time series of numbers of case confirmations and deaths by
103 country (cumulative). Daily incidences were derived from the cumulative counts. We used data
104 from the following countries: France (FR), Italy (IT), Spain (SP), Germany (GR), Belgium (BE),
105 Switzerland (SW), Netherlands (NT), United Kingdom (UK) and the US (US).
106
We included a subset of case confirmation and death count data for inference based on the two
following criteria. First, to minimize the impact of stochasticity and uncertainty in early data
collection, we used case confirmation incidence data starting from the date when the cumulative
number of cases was greater than 100, and used daily new death count data starting from the date
when the cumulative death count is greater than 20 in each country. Second, to estimate the early
outbreak growth in each country before control measures were implemented and at the same time
maximize the power of inference, we allowed a maximum of 15 days of data points for the two
types of data, leading to a maximum of 30 data points for each country. Note that the end date of
incidence data used for inference is at or close to the date when strong control measures were
implemented in each country (Table S1). We tested the sensitivity of model predictions when
only 10, 13 days of data points are included for inference. The results are robust to this variation
(Table S1), suggesting that the choice of 15 days is reasonable and that the data shows consistent
exponential growth during this period.
120
121 **Model**
122 We construct a SEIR type model using ordinary differential equations (ODEs; see
123 Supplementary materials). We consider the exponentially growing phase of the outbreak and
124 thus make the common assumption that the susceptible population is constant over time. Then,
125 the total number of infected individuals $I^*(t) = E(t) + I(t)$ can be expressed as:

$$I^*(t) = I_0^* e^{rt} \qquad (1)$$

126 where r is the exponential growth rate of the epidemic (the growth rate for short below), and I_0^*
127 is the number of total infected individual at time 0, set as January 20, 2020. Note the choice of
128 the date of time 0 does not affect our estimation.
129
130 We solve the ODE model and derive the following expressions for the key quantities for model
131 inference (see Supplementary material). The descriptions and values used for the parameters in
132 the ODE model are summarized in Table 1.
133
134 The true daily incidence of infected individuals, $\Omega(t)$, is:

$$\Omega(t) = \frac{\beta(k + r)}{r(k + r + \beta)} I_0^* \left(e^{rt} - e^{r(t-1)} \right) \qquad (2)$$

135 where β and $1/k$ are the transmission potential of the virus and the latent period of infection,
136 respectively.
137
138 The daily new confirmed case count, $\Psi(t)$, is related to the true daily incidence, $\Omega(t)$ as:

$$\Psi(t) = \theta(t) \frac{g}{(g + r)} \Omega(t) \qquad (3)$$

139 where $\theta(t)$ is the detection rate, i.e. the fraction of newly individuals at time t who are later
140 detected by surveillance later on. $1/g$ is the average duration between infection and case
141 confirmation.

143 The daily new death count, $\Phi(t)$, is related to the true daily incidence, $\Omega(t)$ as:

$$\Phi(t) = \left(\frac{nd}{r+nd}\right)^n X \frac{k}{k+r} \Omega(t) \qquad (4)$$

where X is the infection fatality ratio (sometimes referred as case fatality ratio, CFR, depending
on definition). We assumed a realistic (Erlang) distribution for the period between onset of
infectiousness and death[6], where $1/d$ and n are the mean and the shape parameter for the
distribution.

The expressions above clearly establish that during the exponential growth of an epidemic, the
ratio between death counts Φ and the confirmed cases Ψ (i.e. two widely reported numbers in
public databases, publications and news reports) is not only dependent on infection fatality ratio
and the detection rate, but also a highly nonlinear relationship between the distribution of the
period from infection to death and the growth rate, r, as indicated by the $\left(\frac{nd}{r+nd}\right)^n$ term. Failure to
take this into account may lead erroneous conclusions.

We tested three different scenarios for surveillance intensity changes over time, modeled as the
detection rate, $\theta(t)$:
1) θ is a constant, i.e. no change over time;
2) $\theta(t) = p_{min} + (p_{max} - p_{min})\frac{t^m}{t^m + K^m}$, i.e. θ is a Hill-type function of t;
3) $\theta(t)$ is equal to p_{min} before t_1, increases linearly to p_{max} between t_1 and t_2 and stay
constant at p_{max} after t_2, i.e. θ is a semi-linear function of t.
Note that $1/g$ can be a time dependent function as we and others shown previously[6,12]. To keep
the model simple, we implicitly assume that the time dependent changes in g can be included in
the estimation of $\theta(t)$.

See supplementary materials for details of data collection, modeling analysis, parameter choice
and estimation and uncertainty quantifications.

Results
Estimation of the epidemic growth rate and surveillance intensity
We constructed an SEIR type model and fitted the model to both the incidence (case
confirmation) data and the daily new death count data from eight countries in Europe and the US.
We selected data from a period during early outbreak before or a few days after strong control
measures, such as school and work closure, and locking down cities etc., were implemented in
these countries (see Methods for details and Table S1). There are clear decreases in the rate of
exponential growth of infection incidence after the end dates of data selection in most of these
countries (Fig. 1), and this is likely to reflect the impact of the strong control measures
implemented[13].

We estimated that the exponential growth rate of early outbreaks, r, ranges between 0·19 and
0·29/day in the nine countries, translating to doubling times between 2·4-3·7 days (Fig. 1). The
two countries with the highest point estimates of the exponential rate are Spain and the US, at
0·29 and 0·28/day, respectively; whereas Switzerland and Netherlands have lower point
estimates at 0·19/day. Accounting for uncertainties in the parameter values (see Methods), we

found that the epidemic growth rates in France, Germany, Italy, Spain and the US are mostly
likely higher than 0·2/day (Fig. 2A).

By explicitly considering surveillance intensity, our model enables us to estimate the detection
rate, i.e. the probability of an infected person being identified by surveillance. Assuming a
constant detection rate during the period when data used for inference are collected, we
estimated that Germany has a point estimate of the detection rate at 23·1%. In 8 other countries
examined, the detection rates were relatively low, ranging between 2·2-9·3%. This provides a
natural explanation of the high number of reported cases compared to the relative low number of
deaths in March 2020.

We caution that unlike the epidemic growth rate which is constrained by data, the estimation of
detection rate is highly dependent on the fixed parameter values, such as the infection fatality
ratio assumed in the model. If the infection fatality ratio is lower than we assumed, we would
estimate even lower detection rates. To fully assess the uncertainties in the estimation, we
performed sensitivity analysis varying parameter values within ranges based on the best current
knowledge (see Supplementary Material). Taking into account these uncertainties, we estimate
that Germany has a detection rate between 5% and 86%, and Switzerland has a detection rate
between 2% and 26% (Fig. 2). The detection rates in other countries are likely between 1-10%.
Although large uncertainties exist, we find that the relative detection rates among countries are
robust to parameter uncertainties, i.e. the detection rate in Germany is much higher than the rates
in the other countries. Overall, we find that even in countries with well-developed medical and
public health infrastructures, the detection rate for COVID-19 is in general low, likely due to the
high percentage of infected individuals with no or mild-to-moderate symptoms[9,10]. Our results
emphasize the importance of wearing personal protective equipment to prevent transmission
from the large population of unidentified individuals, and more aggressive testing of infection
and contact tracing are needed to identify most infected individuals.

We further tested the possibility of changes in surveillance intensity, and found no statistical
support (Table S2). We emphasize that this conclusion only applies to the period when the
incidence data used for estimation were collected (Table S1). It is likely that the surveillance
intensity was different during other periods of the outbreaks. As shown in Fig. 1, in Belgium,
France, Italy and Netherland, and UK, the red open circles, i.e. data that were not used for
inference, are mostly below the red band predicted by our model during very early outbreak.
This indicates that in these countries, the detection rates were even lower than we estimated such
that there were very few cases detected, although thousands of infected individuals were already
infected. Some of the red open circles in the US seems during late March are above the red band,
suggesting increases in surveillance intensity.

Implications to intervention strategies - 'hit hard, hit early'
Using our empirical estimates of the growth rates, we explored the implications for public health
efforts needed to control the COVID-19 outbreak. We considered an outbreak scenario in a city
with a population size of 10 million. Because it may take at least one and a half years for an
effective vaccine to be developed and deployed, below we compared outbreak outcomes under
different scenarios at 18 months.

We first calculated the length of time for the epidemic to reach epidemic peak, assuming only
one infected individual at day 0. When the growth rate is higher than 0·027/day, the epidemic
peak will occur in less than 18 months and a large fraction of the population (>80%) will be
infected. If our goal is that the total fraction of infected individuals is less than 10% at the end of
18 months, the growth rate has to be less than 0·025/day (i.e. an extremely slow growth rate with
a doubling time of more than 28 days; Fig. 3B). This suggests that moderate social distancing
efforts will be insufficient to delay the epidemic peak beyond 18 months. On the other hand, if
these targeted growth rates are achieved through very strong public health interventions, a little
more effort would lead to enormous public health benefit, i.e. the total infected fraction
decreases exponentially when r decrease beyond 0·023/day as shown in Fig. 3B.
241
To further corroborates our results, we calculated the efforts needed to achieve three goals in 18
months: 1) virus containment, i.e. epidemic stops growing, 2) the total infected population is
10%, and 3) the total infected population is 1%. We found that the efforts needed are similar,
especially when the population of infected individuals is already more than 100 (Fig. 3C), i.e. a
scenario that many cities around the globe are facing now. For example, when an outbreak grows
at rate 0·29/day, the levels of efforts needed to achieve the three goals are between 84% and 86%
reduction in transmission; whereas when the growth rate is 0·19 /day, the levels of effort needed
are between 77% and 80% reduction. These high levels of reduction needed argue for very
strong and comprehensive intervention efforts implemented as soon as possible, no matter
whether the goal is containment or mitigation - a strategy reminiscent of the 'hit hard, hit early'
paradigm in treating HIV infection in a patient[16].
253
Discussion
The epidemic growth rate for disease spread depends on many factors, including biological[17],
demographic, and social factors. In this work, we report high COVID-19 epidemic growth rates
between 0·19-0·29/day and short doubling times between 2·4-3·7 days across the eight most
affected countries in Europe and in the US (as of March 31, 2020). This is consistent with our
previous estimate for the early COVID-19 outbreak in Wuhan, China[6]. Altogether, these results
demonstrate COVID-19 to be a highly transmissible disease in the absence of strong control
measurements irrespective of heterogeneities in geographic and social settings. We also find that
most of infected individuals are not identified/detected, similar to findings in Wuhan, China[6,11].
This has important implications to both pharmaceutical and non-pharmaceutical interventions as
we discuss below.
265
First, with the global efforts to develop vaccines for COVID-19, it is important to have an
accurate measure of the basic reproductive number, R_0, to set the threshold level of herd
immunity needed to prevent transmission. Previously, R_0 for COVID-19 were estimated to be
between 2-3[2-5] and were widely reported in official documents[18] and public media. These
estimates are mostly based on an epidemic growth rate between 0·1 and 0·14/day[2-5], which we
now know to be inconsistent with many new lines of evidence and data as discussed in this paper
and a previous work[6]. The growth rates that we estimated in this paper are simply not consistent
with a low R_0. Previously we showed that with a mean serial interval, defined as the duration of
time between onset of symptoms in an index case and a secondary case, of 6-9 days[3,19] and an
epidemic growth rate between 0·19-0·29/day, the value of R_0 must be higher than 3[6]. Using the
same framework as in Ref. 6, we find that when the growth rate is 0·19/day, the median R_0 is

estimated to be 3·9 (95% CI: 3·1 and 5·0); whereas when the growth rate is 0·29/day, the median
R_0 is estimated to be 7·1 (95% CI: 5·1 and 9·6)[6]. Although shorter serial intervals are reported in
the literature[20,21], it was noted that this is likely due to strong intervention efforts[19,20]. Given the
potentially long infectious period in individuals with either mild or severe symptoms[22], the mean
serial interval during early outbreak in the absence of strong intervention (a likely scenario in
most countries examined here) is unlikely less than 6 days. Overall, our results imply that a large
fraction of the population needs to be vaccinated if an effective vaccine is to prevent the spread
of the virus. In addition, if the virus is allowed to spread through the population, a large fraction
of the population (>74%) will be infected even if the growth curve is flattened by control efforts.

Second, the awareness of the extraordinary high rates of COVID-19 spread during the current
outbreak is critically important for epidemic preparedness. This is because the short doubling
time means that health care systems can be overwhelmed in a couple of weeks rather than several
months in the absence of control. A recent report shows that the number of COVID-19 patients
admitted to intensive care units (ICUs) in Italy grew at a rate of approximately 0·25/day[23],
remarkably consistent with our estimates. With such a high growth rate, there was only a very
short window period for preparation[23]. Of course, heterogeneities in the growth rate may exist
among different areas within each a country. We note that our inference is largely driven by data
collected from highly populated areas, such as Wuhan in China, Lombardy in Italy, and New
York city in the US. Further work is needed to assess heterogeneity in the rate of spread across
areas with different population densities.

Third, the high epidemic growth rates suggest that moderate control efforts will not sufficiently
slow the virus spread to achieve measurable public health benefits. This may explain the
continuous growth of the outbreak in some countries despite measures, such as work and school
closures, were in place. We found that similarly high levels of efforts to reduce overall
transmission by 77-86% are needed to delay the epidemic peak and protect a large fraction of the
population from infection in 18 months (mitigation) or to reduce R_0 below 1 (containment).

Lastly, our finding that the majority of infected individuals are not identified suggests that in
most infected individuals, the symptoms are likely to be mild, and many of them may not be
aware of their infection status. Part of the higher detection probability in Germany is possibly
due to their early use of contact tracing, which will find asymptomatic and mild cases that might
otherwise not be detected. This argues for extensive, universally available testing to identify and
isolate most infected individuals as well as the use of personal protective equipment to prevent
potential transmission from individuals with no or mild symptoms.

Overall, in the absence of very strong control, the virus can cause high mortality and
morbidity[14,15] due to the high number of expected infections, which places an extremely heavy
burden on even the most advanced health care systems[15,24]. Thus, with COVID-19, half-
measures will not be effective in meeting public health goals. Even for more modest goals such
as "flattening the curve", we probably need all feasible tools available, i.e. extensive testing,
isolation and quarantine, use of personal protective equipment, coupled with comprehensive
social distancing. This is a strategy reminiscent of the 'hit hard, hit early' paradigm in treating
HIV infected individuals[16]. China, South Korea, and Singapore have proven that containment is
possible with appropriate measures. Because there will be extensive economic impacts of the

global COVID-19 pandemic regardless of how principled our response is, the question is not
balancing public health with damage to the economy, but rather how many lives can we save.
325
326
Acknowledgments
We would like to thank Alan Perelson for suggestions and critical reading of the manuscript. The
work is partially funded by the Laboratory Directed Research and Development (LDRD) Rapid
Response Program through the Center for Nonlinear Studies at Los Alamos National Laboratory.
RK and SS would like to acknowledge funding from DARPA (HR0011938513), the Center for
Nonlinear Studies. ERS was funded though NIH grant (R01AI135946).
333
334 **Declaration of interests**
335 The authors declare no competing interests.
336
337 **Author contributions**
338 RK and NH conceived the project; RK performed literature search and designed the study; SS
339 collected data; RK, SS and ERS performed analyses; RK, SS, ERS and NH wrote and edited the
340 manuscript.

References:

1.   WHO. Pneumonia of unknown cause – China (https://wwwwhoint/csr/don/05-january-2020-pneumonia-of-unkown-cause-china/en/; accessed January 30, 2020) 2020.

2.   Kucharski AJ, Russell TW, Diamond C, et al. Early dynamics of transmission and control of COVID-19: a mathematical modelling study. *The Lancet Infectious Diseases* 2020.

3.   Li Q, Guan X, Wu P, et al. Early Transmission Dynamics in Wuhan, China, of Novel Coronavirus-Infected Pneumonia. *N Engl J Med* 2020.

4.   Riou J, Althaus CL. Pattern of early human-to-human transmission of Wuhan 2019 novel coronavirus (2019-nCoV), December 2019 to January 2020. *Euro Surveill* 2020; **25**(4).

5.   Wu JT, Leung K, Leung GM. Nowcasting and forecasting the potential domestic and international spread of the 2019-nCoV outbreak originating in Wuhan, China: a modelling study. *The Lancet* 2020.

6.   Sanche S, Lin Y, Xu C, Romero-Severson E, Hengartner N, Ke R. The novel coronavirus, SARA-nCoV-2, is highly contagious and more infectious than initially estimated. *Emerging Infectious Diseases* 2020; (accepted).

7.   The Novel Coronavirus Pneumonia Emergency Response Epidemiology Team. The Epidemiological Characteristics of an Outbreak of 2019 Novel Coronavirus Diseases (COVID-19) - China, 2020. *China CDC Weekly* 2020; **2**(8): 113-22.

8.   Lipsitch M, Cohen T, Cooper B, et al. Transmission dynamics and control of severe acute respiratory syndrome. *Science* 2003; **300**(5627): 1966-70.

9.   Mizumoto K, Kagaya K, Zarebski A, Chowell G. Estimating the asymptomatic proportion of coronavirus disease 2019 (COVID-19) cases on board the Diamond Princess cruise ship, Yokohama, Japan, 2020. *Euro Surveill* 2020; **25**(10).

10.   Zou L, Ruan F, Huang M, et al. SARS-CoV-2 Viral Load in Upper Respiratory Specimens of Infected Patients. *N Engl J Med* 2020.

11.   Li R, Pei S, Chen B, et al. Substantial undocumented infection facilitates the rapid dissemination of novel coronavirus (SARS-CoV2). *Science* 2020.

12.   Ng Y, Li Z, Chua YX, et al. Evaluation of the Effectiveness of Surveillance and Containment Measures for the First 100 Patients with COVID-19 in Singapore — January 2–February 29, 2020. *MMWR Morb Mortal Wkly Rep* 2020.

13.   Flaxman S, Mishra S, Gandy A, et al. Estimating the number of infections and the impact of nonpharmaceutical interventions on COVID-19 in 11 European countries. (https://wwwimperialacuk/media/imperial-college/medicine/sph/ide/gida-fellowships/Imperial-College-COVID19-Europe-estimates-and-NPI-impact-30-03-2020pdf; accessed Mar 30, 2020) 2020.

14.   Dorigatti I, Okell L, Cori A, et al. Severity of 2019-novel coronavirus (nCoV). (https://wwwimperialacuk/media/imperial-college/medicine/sph/ide/gida-fellowships/Imperial-College-COVID19-severity-10-02-2020pdf; accessed Mar 30, 2020) 2020.

15.   Wu JT, Leung K, Bushman M, et al. Estimating clinical severity of COVID-19 from the transmission dynamics in Wuhan, China. *Nature Medicine* 2020.

16.   Ho DD. Time to hit HIV, early and hard. *N Engl J Med* 1995; **333**(7): 450-1.

17.   Wrapp D, Wang N, Corbett KS, et al. Cryo-EM structure of the 2019-nCoV spike in the prefusion conformation. *Science* 2020; **367**(6483): 1260-3.

18.   WHO. Report of the WHO-China Joint Mission on Coronavirus Disease 2019 (COVID-
19) (https://wwwwhoint/docs/default-source/coronaviruse/who-china-joint-mission-on-covid-19-
final-reportpdf; *accessed March 30, 2020)* 2020.
19.   Bi Q, Wu Y, Mei S, et al. Epidemiology and Transmission of COVID-19 in Shenzhen
China: Analysis of 391 cases and 1,286 of their close contacts. *medRxiv* 2020:
2020.03.03.20028423.
20.   Du Z, Xu X, Wu Y, Wang L, Cowling B, Ancel Meyers L. Serial interval of COVID-19
among publicly reported confirmed cases. *Emerging Infectious Diseases*. 2020.
21.   Nishiura H, Linton NM, Akhmetzhanov AR. Serial interval of novel coronavirus
(COVID-19) infections. *Int J Infect Dis* 2020; **93**: 284-6.
22.   Zhou F, Yu T, Du R, et al. Clinical course and risk factors for mortality of adult
inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet* 2020.
23.   Grasselli G, Pesenti A, Cecconi M. Critical Care Utilization for the COVID-19 Outbreak
in Lombardy, Italy: Early Experience and Forecast During an Emergency Response. *JAMA* 2020.
24.   Walker PG, Whittaker C, Watson O, et al. The Global Impact of COVID-19 and
Strategies for Mitigation and Suppression. (https://wwwimperialacuk/media/imperial-
college/medicine/sph/ide/gida-fellowships/Imperial-College-COVID19-Global-Impact-26-03-
2020v2pdf; *accessed March 30, 2020)* 2020.
25.   Lauer SA, Grantz KH, Bi Q, et al. The Incubation Period of Coronavirus Disease 2019
(COVID-19) From Publicly Reported Confirmed Cases: Estimation and Application. *Ann Intern*
Med 2020.
26.   Zhou F, Yu T, Du R, et al. Clinical course and risk factors for mortality of adult
inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet* 2020;
395(10229): 1054-62.
409
410
411

Figure captions

Figure 1. Estimation of the exponential growth rate, doubling time of epidemics and detection rate of infected individuals in 8 European countries and the US. Red and black symbols show the daily counts of new case confirmation and new death, respectively. Dots denote data used for parameter inference; whereas open circles denote data that are not used for parameter inference. We simulated the exponential model using sampled parameter combinations that are able to explain the data shown in dots (see the Uncertainty Quantification section in the Supplementary Material). The colored bands denote the area between the lower and upper bounds of simulated/predicted true daily infection incidence (blue), daily case confirmation (red) and daily death (grey) assuming no intervention efforts nor changes in surveillance intensity. Deviation of open circles from the corresponding bands thus indicates either changes in surveillance intensity or impacts of control measures.

Figure 2. Point estimates and estimated ranges of the exponential growth rate, r and the detection rate, θ, in each country. BE: Belgium; FR: France; GR: Germany; IT: Italy; NT: Netherlands; SP: Spain; SW: Switzerland; UK: United Kingdom; US: United States.

Figure 3. Strong control measures are needed to achieve measurable benefits, no matter the goal is mitigation ('flatten the curve') or containment. (A) Predicted time to epidemic peak for different growth rates. To delay epidemic peak beyond 18 months, a growth rate less than 0·027/day is needed. This threshold is denoted by dashed lines. Red area shows the ranges of growth rates estimated for the eight European countries, the US and Wuhan, China. **(B)** Final fraction of infected individuals after 18 months. A growth rate less than 0·023/day (doubling time more than 30 days) is needed to achieve the goal that less than 10% of individuals are infected (dashed lines). However, the benefit, i.e. fraction of uninfected individuals, increases exponentially when the growth rate is further reduced beyond the threshold. **(C)** Similar levels of efforts, measured as fractions of transmission reduction, are needed to achieve containment, i.e. reverting epidemic growth (dots), or mitigation, i.e. the final fraction of infected individuals is 1% (x) or 10% (open circle). We assumed epidemic growth rates of 0·19 (red) or 0·29/day (blue).

448 **Table 1. Description of parameter and their values.** See the supplementary material for
449 discussions of choice of parameter values.

Parameters	Description	Value	Ranges used in uncertainty analysis	References
r	Exponential growth rate	Estimated from data	$0 \cdot 1 - 0 \cdot 35$ /day	
I_0^*	I_0^* is the number of total infected individual at time 0 (Jan. 20)	Estimated from data	$0 \cdot 0001 - 10$ on a log scale	
β	Infectivity in the SEIR model	Calculated from r		See Supplementary material
$1/k$	The mean latent period, i.e. from infection to becoming infectious	3 days	2-5 days	6,25
$1/g$	The mean duration from infection to case confirmation	8 days	6-10 days	6,12
n	Shape parameter for the duration from symptom onset to death.	4	4-5	
$1/d$	Mean duration from onset of infectious period to death	$18 \cdot 5$ days	$16 \cdot 5 - 20 \cdot 5$ days	6,26
X	Infection fatality ratio	$0 \cdot 01$	$0 \cdot 004 - 0 \cdot 015$	14,15

450
451