

The unseen and pervasive threat of COVID-19 throughout the US

Emily Javan, Dr. Spencer J. Fox, Dr. Lauren Ancel Meyers

Corresponding author:

Lauren Ancel Meyers

The University of Texas at Austin

laurenmeyers@austin.utexas.edu

Abstract

For each US county, we calculated the probability of an ongoing COVID-19 epidemic that may not yet be apparent. Based on confirmed cases as of April 15, 2020, COVID-19 is likely spreading in 86% of counties containing 97% of US population. Proactive measures before two cases are confirmed are prudent.

Research Letter

The unprecedented threat of COVID-19 could kill hundreds of thousands to millions of Americans (1,2). Over the course of a few weeks, it has emerged in all 50 US states (3). Officials are weighing the potentially enormous economic and societal costs of strict social distancing measures against the *unseen* risks of COVID-19 hospitalizations and mortality in their communities. To support decision making, we have estimated the likelihood that each county in the US already has extensive community transmission (Figure).

To make these risk assessments, we applied a model that we developed for another *silent spreader*—Zika—which threatened to emerge in southern US states in 2016 (4). Using stochastic simulations, we determine the probability that a county already has an ongoing COVID-19 epidemic by the time the county reports just one, two, three or more cases. The levels of risk displayed in Figure 1 are based on reported cases as of April 15, 2020 (5) and account for under-reporting, the uncertainty in the transmission rate of COVID-19, the likelihood of super-spreading events—like those that fueled SARS outbreaks in 2003 and have been reported recently for COVID-19 (6)—and the impact of social distancing measures that have been enacted across the US in recent weeks (Appendix).

By the time a first case of COVID-19 is reported in the county, we estimate that there is over a 50% chance that a full-blown epidemic is underway. As of April 15, 2020, we found 86% of US counties with 97% of the national population have reported at least one case (Figure). Even counties that have not *yet* reported a confirmed case have an estimated 9% chance of sustained local transmission. If an epidemic is indeed underway, then our model predicts a delay of 7.5 (95% CI 3.9-16.3) weeks between the first reported case and the total number of COVID-19 exceeding 1,000; by the time the tenth case has been reported this lead time shrinks to 4.4 (95% CI 2.1-11.4) weeks (Appendix).

Given the speed and stealth with which COVID-19 spreads (7), the entire map will likely be fully red within a week or two. Proactive social distancing can block community transmission, avert catastrophic surges in hospitalizations, and save lives (8). For those still weighing if, when, and how aggressively to act, there is little time for delay.




Figure. Probability of ongoing COVID-19 epidemics in the 3142 counties of the United States as of April 15, 2020. The risk of an unseen epidemic in a county with only a single reported case is 50%. The risk in counties that have not yet reported cases is 9%.

References

1. Ferguson N, Laydon D, Nedjati Gilani G, Imai N, Ainslie K, Baguelin M, et al. Report 9: Impact of non-pharmaceutical interventions (NPIs) to reduce COVID19 mortality and healthcare demand. 2020 Mar 16.
2. IHME COVID-19 health service utilization forecasting team, Murray CJL. Forecasting COVID-19 impact on hospital bed-days, ICU-days, ventilator-days and deaths by US state in the next 4 months. medRxiv. 2020 Mar 30.
3. Fauver JR, Petrone ME, Hodcroft EB, Shioda K, Ehrlich HY, Watts AG, et al. Coast-to-coast spread of SARS-CoV-2 in the United States revealed by genomic epidemiology. medRxiv. 2020 Mar 26.
4. Castro LA, Fox SJ, Chen X, Liu K, Bellan SE, Dimitrov NB, et al. Assessing real-time Zika risk in the United States. BMC Infect Dis. 2017 May 4;17(1):284.
5. Smith M, Yourish K, Almukhtar S, Collins K, Ivory D, Harmon A. The New York Times Coronavirus (Covid-19) Data in the United States. 2020 [cited 2020 Apr 16]. Available from: <https://github.com/nytimes/covid-19-data>
6. Lloyd-Smith JO, Schreiber SJ, Kopp PE, Getz WM. Superspreading and the effect of individual variation on disease emergence. Nature. 2005 Nov 17;438(7066):355–9.
7. Li R, Pei S, Chen B, Song Y, Zhang T, Yang W, et al. Substantial undocumented infection facilitates the rapid dissemination of novel coronavirus (COVID-19). Science. 2020 Mar 16.
8. Cowling BJ, Ali ST, Ng TWY, Tsang TK, Li JCM, Fong MW, et al. Impact assessment of non-pharmaceutical interventions against COVID-19 and influenza in Hong Kong: an observational study. medRxiv. 2020 Mar 16.