Protein Classifier for Thyroid Nodules Learned from Rapidly Acquired Proteotypes

Yaoting Sun[1,2#], Sathiyamoorthy Selvarajan[3#], Zelin Zang[4#], Wei Liu[1,2,5#], Yi Zhu[1,2#], Hao Zhang[6#], Hao Chen[1,2], Xue Cai[1,2], Huanhuan Gao[1,2], Zhicheng Wu[1,2], Lirong Chen[7], Xiaodong Teng[8], Yongfu Zhao[9], Sangeeta Mantoo[3],Tony Kiat-Hon Lim[3], Bhuvaneswari Hariraman[10], Serene Yeow[11], Syed Muhammad Fahmy bin Syed Abdillah[3], Sze Sing Lee[11], Guan Ruan[1, 2], Qiushi Zhang[1, 2], Tiansheng Zhu[1, 2], Weibin Wang[12], Guangzhi Wang[9], Junhong Xiao[9], Yi He[13], Zhihong Wang[6], Wei Sun[6], Yuan Qin[6], Qi Xiao[1, 2], Xu Zheng[14], Linyan Wang[15], Xi Zheng[16], Kailun Xu[16], Yingkuan Shao[16], Kexin Liu[5], Shu Zheng[16], Ruedi Aebersold[17,18], Stan Z. Li[4*], Oi Lian Kon[11*], N. Gopalakrishna Iyer[10,11*], Tiannan Guo[1,2,19*]

[1]Key Laboratory of Structural Biology of Zhejiang Province, School of Life Sciences, Westlake University, 18 Shilongshan Road, Hangzhou 310024, Zhejiang Province, China;

[2]Institute of Basic Medical Sciences, Westlake Institute for Advanced Study, 18 Shilongshan Road, Hangzhou 310024, Zhejiang Province, China;

[3]Department of Pathology, Singapore General Hospital, Republic of Singapore;

[4]School of Engineer, Westlake University, 18 Shilongshan Road, Hangzhou 310024, Zhejiang Province, China;

[5]Department of Clinical Pharmacology, College of Pharmacy, Dalian Medical University, No.9 West Section Lvshun South Road, Dalian 116044, Liaoning Province, China;

[6]Department of Thyroid Surgery, the First Hospital of China Medical University, No.155 Nanjing Bei Street, Shenyang 110001, Liaoning Province, China;

[7]Department of Pathology, The Second Affiliated Hospital of College of Medicine, Zhejiang University, 88 Jiefang Road, Hangzhou 310009, Zhejiang Province, China;

[8]Department of Pathology, the First Affiliated Hospital, College of Medicine, Zhejiang University, 79 Qingchun Road, Hangzhou 310024, Zhejiang Province, China;

[9]Department of General Surgery, The Second Hospital of Dalian Medical University, 467 Zhongshan Road, Dalian 116027, Liaoning Province, China;

[10]Division of Surgical Oncology, National Cancer Centre Singapore, Republic of Singapore;

[11]Division of Medical Sciences, National Cancer Centre Singapore, Republic of Singapore;

[12]Department of Surgical Oncology, the First Affiliated Hospital, College of Medicine, Zhejiang University, 79 Qingchun Road, Hangzhou 310024, Zhejiang Province, China;

[13]Department of Urology, The Second Hospital of Dalian Medical University, 467 Zhongshan Road, Dalian 116027, Liaoning Province, China;

[14]Liaoning Laboratory of Cancer Genetics and Epigenetics and Department of Cell Biology, College of Basic Medical Sciences, Dalian Medical University, No.9 West Section Lvshun South Road, Dalian 116044, Liaoning Province, China;

[15]Department of Ophthalmology, The Second Affiliated Hospital, Zhejiang University School of Medicine, 88 Jiefang Road, Hangzhou 310009, Zhejiang Province, China;

[16]Cancer Institute (Key Laboratory of Cancer Prevention and Intervention, China National Ministry of Education, Key Laboratory of Molecular Biology in Medical Sciences, Zhejiang Province, China), The Second Affiliated Hospital, Zhejiang University School of Medicine, 88 Jiefang Road, Hangzhou 310009, Zhejiang Province, China;

[17]Department of Biology, Institute of Molecular Systems Biology, ETH Zurich, 8093 Zurich, Switzerland

44   [18]Faculty of Science, University of Zurich, 8093 Zurich, Switzerland

45   [19]Lead contact

46   [#]Co-first authors;

47   [*]Correspondence: Stan.ZQ.Li@westlake.edu.cn (S.Z.L.); kairos712@singnet.com.sg (O.L.K.);

48   gopaliyer@singhealth.com.sg (N.G.I.); guotiannan@westlake.edu.cn (T.G.)

SUMMARY

Up to 30% of thyroid nodules cannot be accurately classified as benign or malignant by

cytopathology. Diagnostic accuracy can be improved by nucleic acid-based testing, yet

a sizeable number of diagnostic thyroidectomies remains unavoidable. In order to

develop a protein classifier for thyroid nodules, we analyzed the quantitative proteomes

of 1,725 retrospective thyroid tissue samples from 578 patients using pressure-cycling

technology and data-independent acquisition mass spectrometry. With artificial neural

networks, a classifier of 14 proteins achieved over 93% accuracy in classifying

malignant thyroid nodules. This classifier was validated in retrospective samples of 271

patients (91% accuracy), and prospective samples of 62 patients (88% accuracy) from

four independent centers. These rapidly acquired proteotypes and artificial neural

networks supported the establishment of an effective protein classifier for classifying

thyroid nodules.

Keywords: Thyroid nodule; Proteomics; Data-independent acquisition; Artificial neural

networks; Pressure cycling technology; Mass spectrometry

INTRODUCTION

Advances in imaging technology and liberal screening practices have identified

thyroid nodules in up to 50% of the general population, but only a small minority (7-15%)

eventually prove to be malignant by histology, and even fewer among these are

clinically relevant (Burman and Wartofsky, 2015; Jameson, 2012). Beyond clinical

assessment and ultrasonography, fine needle aspiration (FNA) followed by

cytopathology is considered the most reliable pre-surgical technique for differentiating

benign from malignant thyroid tumors (Burman and Wartofsky, 2015; Faquin et al.,

2011). Yet up to one-third of thyroid nodules are deemed indeterminate by FNA-

cytopathology (Alexander et al., 2012), and surgery remains the only option for accurate

diagnosis. However, the majority of thyroid surgeries are diagnostic procedures

undertaken to exclude thyroid cancer, of which no more than 25% accomplish any

therapeutic purpose (Ahn et al., 2014). Patients whose thyroid glands are removed in

part or entirely often require daily and lifelong thyroxine-replacement therapy and

medical monitoring. Given that only 10% of resected glands prove to be malignant, the

current clinical approach results in substantial over-treatment with unwarranted surgical

risks for patients who could otherwise be treated conservatively (Vaccarella et al.,

2016).

Molecular tests adjunctive to FNA-cytopathology have focused on RNA expression

or DNA mutational profiling of aspirates obtained prior to surgery, using small quantities

of RNA or DNA that can be amplified (Eszlinger et al., 2017). The Afirma® gene
expression classifier (GEC) determines RNA abundances of 142 genes with high
sensitivity and negative predictive value (NPV) of up to 92% and 93%, respectively.
However, positive predictive value (PPV) and specificity for diagnosing malignancy are
only 47% and 52%, respectively (Alexander et al., 2012). The latest version of Afirma®
called GSC has achieved a sensitivity of 91%, but the specificity was only 68% (Patel et
al., 2018). The ThyroSeq (v3.0) test combines mutation and expression profiling for
better accuracy (Nikiforova et al., 2013), but has other limitations i.e. the need for fresh
tissue samples with un-degraded RNA, and test data processing by central 'black box'
analysis. A recent review of the GEC test showed sensitivities of 83-100% but low
overall specificity, ranging from 10 to 52% across multiple centers. This may reflect
variability in sample composition, issues with tissue quality and the fragility of nuclei
acid-based testing in general (Wang and Sosa, 2018). While nucleic acid-based
approaches continue to be refined, for example with successive iterations of ThyroSeq
panels, there is evident scope for alternative approaches to address this diagnostic
dilemma.
101
Until recently, proteomics-based analyses were limited to large tissue quantities and
fresh/snap frozen samples. Proteotyping hundreds of biopsy-level tissue samples from
clinical cohorts remains unfeasible with conventional methods. We developed a
pressure cycling technology (PCT) protocol for proteomic analysis of tissue biopsy
samples (Guo et al., 2015) which can be performed on minimal amounts (0.2–1 mg) of
fresh-frozen tissue samples (Shao et al., 2016; Shao et al., 2015). The method was
recently extended to generate high quality proteome data from formalin-fixed, paraffin-
embedded (FFPE) tissue samples (Zhu et al., 2019), and in this study on fine-needle
thyroid gland aspirates. Coupled with Sequential Window Acquisition of all Theoretical
fragment ions (SWATH) mass spectrometry (MS), a data-independent acquisition (DIA)-
MS method (Gillet et al., 2012), this technique now permits practical proteomic analysis
of biopsy size FFPE tissue samples at high sample throughput. In this study we apply
this technology to analyze a large number of tissue samples and show that the high
quality proteotype data generated a robust panel of protein markers which differentiates
benign from malignant thyroid disease with high 'rule in' and 'rule out' accuracy.
117

RESULTS

Study design and clinical characteristics

We applied PCT-DIA on a total of 931 nodules from 911 patients using tissue cores
(1 mm diameter with 0.5-1 mm thickness) punched out from regions of interest marked
on FFPE tissue blocks, or from fine-needle aspiration (FNA) biopsies. The samples
comprise (i) a discovery set from Singapore General Hospital (n = 579 nodules) where
histopathological diagnoses were confirmed on central review by a board-certified
pathologist; and (ii) independent test sets from four hospitals in China consisting of
retrospective test sets of FFPE samples (n = 288 nodules) and prospective test set of
FNA biopsies (n = 64 nodules) (Figure 1A). The discovery set comprised FFPE samples
from 40 normal thyroid tissues (N), 203 multinodular goiters (MNG), 137 follicular

adenomas (FA), 75 follicular carcinomas (FTC) and 124 papillary carcinomas (PTC)
(Table 1; Table S1). For subsequent analyses, these were divided into benign
(comprising N, MNG and FA) and malignant (comprising FTC and PTC) thyroid nodules.
For each nodule in the discovery set, three punches were obtained from the region of
interest as biological replicates. In total, we analyzed 1,725 samples randomly
distributed into 121 batches to minimize batch effects (Figure 1B) using 45 min DIA-MS,
and an additional 56 randomly selected samples as technical replicates. These
technical replicates distributed randomly in the first 84 batches for the training sample
set. Although higher proteomic depth could be obtained with a longer LC gradient, we
adopted a reasonably fast analysis time to minimize batch effects without substantial
compromise of proteome depth (Sun, 2020), thus facilitating effective downstream
machine learning to establish a robust classifier.
141

Global proteomic profiling of thyroid nodules

To analyze the DIA data, we built a specific thyroid-tissue spectral library from FFPE
tissues containing 52 DDA files (Table S2) using a 120-min liquid chromatography (LC)
gradient on a Q Exactive Hybrid Quadrupole-Orbitrap mass spectrometer. Relatively
long gradient was adopted for DDA analysis to maximize the isolation of peptide
precursors. We constructed a spectral library containing 33,802 peptides from 5,190
protein groups using our previously established computational pipeline (Zhu, 2020). The
number of proteins in this library is relatively smaller than that of other tissue types
analyzed using a similar workflow (Guo et al., 2015; Shao et al., 2019; Zhu, 2020; Zhu
et al., 2019) or other proteomic protocols (Jiang et al., 2019; Sinha et al., 2019; Zhang
et al., 2016) as thyroid tissue contains a high abundance protein, thyroglobulin, which
weakens the signal for the rest of the proteome (M et al., 2018). Using OpenSWATH
(v2.0) and our thyroid library, we analyzed 1,781 DIA maps (1,725 FFPE cores and 56
technical replicates). We identified and quantified 30,915 peptides from 3,708 high-
confidence proteotypic proteins (Table S3). Details on quality control and reproducibility
(Figure S1) are detailed in the respective supplementary sections.
From these, we computed the average intensities of 2,617 proteotypic proteins
which were quantified with high confidence for each thyroid nodule, as visualized in a
tissue-type supervised heatmap (Figure 2A). Malignant tissue samples expressed
higher number of proteins than benign samples, indicating more diverse proteome of
tumor cells. To check whether the thus acquired proteotypes classifies different tissue
types, we applied uniform manifold approximation and projection (UMAP) algorithm to
visualize the 579 proteotypes from five tissue types (Figure 2B). The plot shows that
PTC samples are well isolated from the rest, indicating PTC samples are vastly different
in terms of proteome expression. We then grouped N, MNG and FA as benign group,
while FTC and PTC as the malignant group (Figure 2C). The UMAP visualization shows
that malignant samples are well resolved from benign samples with some overlap. We
further narrowed our focus to the N and MNG groups and found that their proteotypes
share high degree of similarity (Figure 2D). Not surprisingly, FA exhibited significant
overlap with both benign and malignant subsets, particularly between FA and FTC
(Figure 2E), corroborating biological overlap between the two pathologies, while there
were sufficient features that could distinguish FTC from PTC (Figure 2F). FA and FTC

could not be separated neither (Figure 2E). Pair-wise comparison of each two groups
are shown in Figure S2. The above analyses show that the proteotype maps thus
measured reasonably reflected the clinical phenotype of these samples.
177

Feature selection and classifier development

To derive a protein-based signature to differentiate benign from malignant thyroid
tumors, we developed a customized feature selection process combined with artificial
neural network algorithms based on the discovery dataset of 500 samples (Figure 3A).
The rest 79 samples serve as internal validation set. Here we limited the number of
selected features to no more than 20 so that they may be practically measured by
targeted proteomics or antibodies in clinic. We paid special attention to maximize the
specificity while keeping the sensitivity above 90% since the major problem of thyroid
nodule evaluation is over-diagnosis. Briefly, protein features selected by a genetic
algorithm (Mitchell, 1998) combined with 5-fold cross validation led us to identify a
classifier-panel of 14 proteins (Table 2) to separate benign and malignant nodules.
Individual protein expression levels are shown in Figure 4. Ten of these proteins have
been previously reported in thyroid cancers, including annexin A1 (Ciregia et al., 2016),
galectin-3 (Bartolazzi et al., 2018), SH3 domain-binding glutamic acid-rich-like protein 3
(Martínez-Aguilar et al., 2016), alpha 2-HS glycoprotein (Farrokhi Yekta et al., 2018),
myosin heavy chain 9 (Wang et al., 2017), phosphatidylethanolamine-binding protein 1
(Kim et al., 2010), clusterin (Kashat et al., 2010), LDL receptor related protein 2 (He et
al., 2020), calreticulin (Schürch et al., 2019) and moesin (Smith et al., 2019). In addition,
this list also includes two proteins involved thyroid functions, including tubulin folding
cofactor A (Figliozzi et al., 2017) and histone cluster 1 H1 family member c (Brix et al.,
1998). No previous association with thyroid disease has been reported for the remaining
two (Thy-1 cell surface antigen and sialic acid acetylesterase).
We next compared seven different machine learning models for classification using
these 14 selected proteins. Receiver operating characteristics (ROC) plots showed the
model based on artificial neural networks described here achieved the highest area
under the curve (AUC) value of 0.95 and accuracy of 0.91 (Figure 3B). Using the 14
protein features in our established neural network model, each specimen was re-
classified into benign or malignant in the 500 samples in the discovery cohort. The
model comprised a 'feature extraction sub-model' which extracts and maps features
from protein data into a feature vector in latent space, and a 'classification sub-model'
which assigns a score (from 0 to 1) to the feature vector indicating the likelihood of
malignancy for each sample. The 'feature extraction sub-model' was trained using
contrastive loss function, while cross-entropy loss function was used to train the
classification sub-model. Details of the neural network model are described in
Supplementary Methods. We validated this model using the 79 internal validation
samples (Table S4; Figure S3A-S3B). ROC plot showed that our model achieved an
AUC value of 0.96 for the training dataset (n=500) used to derive this algorithm and
0.96 for the cross-validation dataset (n=79) (Figure 3C). t-SNE plot of the feature latent
space showed remarkable separation between malignant and benign tissue using the
14-protein panel.
218

Performance of the protein classifier

In order to validate this 14-protein model in an independent patient cohort, we first analyzed 288 pathologist-reviewed FFPE tissues (n = 271 patients) from three high-volume hospitals, comprising 144 benign and 144 malignant tissue samples. To ensure rigorous validation, the diagnoses were blinded during data acquisition and analyses. Each sample was analyzed using the PCT-DIA workflow in technical duplicates. Analysis of the resulting 576 DIA maps identified 3,527 high-confidence proteotypic proteins (Table S3). Individual ROC plots for samples from each of the three hospitals using the 14-protein model showed AUC of >0.91 for the retrospective FFPE samples (Figure 3D). Both scatter and t-SNE plots demonstrated distinct separation between benign and malignant thyroid tissues (Figure 3D; Table S4), although there were variations between individual sites (Figure S3C). The overall sensitivity and specificity were 90% and 91%, respectively, with negative- (NPV) and positive-predictive values (PPV) of 90% and 91%, respectively. Further details are provided in Table 3.

Given that the eventual objectives were to apply this analysis to pre-surgical FNA biopsies, we extended validation to a separate prospective cohort of 64 FNA samples (n=62 patients) obtained from a fourth independent clinical center. Even from these small amounts of FNA biopsy tissues, we were able to generate a high-quality protein matrix containing 3,310 proteotypic proteins using PCT-DIA technology (Table S3). The model achieved an AUC value of 0.89 (Figure 3E) and correctly identified 56 of 64 samples with sensitivity, specificity, PPV and NPV of 87%, 89%, 95% and 73%, respectively (Table 3). The predictive accuracy for all subtypes exceeded 91% except FA (89%) and FTC (83%) (Figure 3F), likely due to similarity of histology (and potential biological overlap) between these two pathologies.

Biological insights on thyroid tumor subtypes

We next asked whether the proteomic data could be used to reveal biological insights of specific subtypes of thyroid neoplasms. In recent years, differentiated thyroid cancers have been further sub-classified based on specific morphological features or their expected clinical course. Hürthle cell adenomas and carcinomas are deemed as distinct entities, with the latter demonstrating a higher propensity for metastasis (Ganly et al., 2018; Gopal et al., 2018). The proteomic data of this study showed these to be well-resolved from other neoplasms, even from the closely related FA and FTC subtypes (Figure 5A; Figure S4A-S4C). Hürthle cell tumors are known for their oncocytic morphology and increased glucose uptake in FDG-PET scans (Grani et al., 2018), and indeed, our data showed that 91 of 109 proteins substantially elevated (fold change > 2 and adjusted p-value < 0.01) compared to follicular neoplasms are mitochondrial proteins participating in multiple metabolic pathways including the TCA cycle and oxidative phosphorylation (Figure 5B). Our data therefore uncovered biochemical processes contributing to the elevated metabolism of Hürthle cell tumors. Compared to the other four complexes in the oxidative phosphorylation pathway, the most strongly upregulated proteins (7/16) were in complex V which catalyzes ATP synthesis and potentially enhances tumor growth.

Follicular-variant papillary thyroid cancers (fvPTC) is a subtype with mixed

morphology, and we therefore examined specific differences between FTC, classical
papillary cancers (cPTC) and fvPTC. There were no significant proteotypic differences
between FTC and fvPTC (Figure S4D). However, 45 proteins were differentially
regulated in fvPTC compared to cPTC (Figure S4D; proteins listed in Table S5). Our
proteotypic data showed that fvPTC overlapped with both FTC and cPTC, but
resembled FTC more closely, indicating that fvPTC is potentially an intermediate entity
between FTC and cPTC (Figure 5C). This is consistent with genomic classifiers which
suggest that FTC and fvPTC share common alterations including in the RAS pathway
(Agrawal et al., 2014). Compared to cPTC, 50 proteins upregulated in histologically
distinct FTC participate in multiple biological processes including endoplasmic reticulum
stress and the unfolded protein response (Figure 5D, S4E), both of which are implicated
in bypassing Ras-driven oncogenic senescence (Blazanin et al., 2017). The signaling
network involving key upregulated proteins in FTC is shown in Figure S4F. In contrast,
the 97 proteins upregulated in cPTC compared to FTC (Figure 5E) mapped to major
oncogenic (TP53, MYC), dendritic cell maturation pathways and inflammatory response,
suggesting that canonical nocogenic pathways and inflammation are involved in the
pathogenesis of cPTC, which has been associated with thyroid inflammation (Figure
5F). Consistent with this suggestion, three proteins involved in inflammation, ANXA1,
LGALS3 and SOD2, showed greatest fold-change among proteins that mapped to the
TP53- and MYC- related networks in cPTC (Figure 5G). Intercellular adhesion molecule
1 (ICAM-1) and signal transducer and activator of transcription 1 (STAT1) are over-
expressed in cPTC and are potential immune-modulating targets.
285

DISCUSSION

Molecular diagnostics for thyroid nodules have been limited to nucleic acid-based
testing thus far due to the feasibility of analyzing small clinical samples and the
increasing affordability of next-generation sequencing. Several nucleic acid-based tests
are commercially available through central-lab testing, their performance in clinic is
suboptimal in terms of specificity, especially in malignancies with low mutational burden
as rigorously examined by Wang et al (Wang and Sosa, 2018). Since proteins are more
stable than RNA in biopsy tissue samples (Shao et al., 2019), we posit that our protein
panel can be developed as future point-of-care diagnostic tests through widely available
techniques such as mass spectrometry and/or immunohistochemistry, as a complement
to the nucleic acid-based test. The FFPE-PCT-DIA methodology used here was able to
derive protein abundance data of 3,779 proteins in 931 samples, generating 2,421 DIA
proteome data sets, including replicates. The pipeline generated the first repository of
proteome data of various thyroid pathologies. This enabled artificial neural network
analysis to mine large proteomic datasets for protein biomarkers of thyroid cancers. A
panel of 14 proteins differentiated benign from malignant disease with diagnostic
accuracy over 90% with sensitivity and specificity over 91% for retrospective test sets
and accuracy of 89% for prospective FNA-derived test set. The fact that 12 out 14 of
these have been previously implicated in thyroid physiology or pathology provides
orthogonal validation for the inclusion of these proteins in our classifier. These metrics
from proteomics data exhibited high degree of both sensitivity and specificity as shown
in Table 3. Our method works for small tissue samples obtained from both FFPE tissues

and FNA biopsies, making it broadly applicable for standard clinical practice, bypassing
RNA assays due to the fragility of RNA integrity.

Despite the accuracy in distinguishing benign from malignant thyroid nodules, the
major limitation for most algorithms is distinguishing FA from FTC. Indeed, the
proteotype data presented here even suggests that this entity may represent a
continuum of disease, in which differences exist at the extremes of phenotypes, but with
significant overlap in-between. Alternatively, the overlapping benign categories may
represent precursor lesions diagnosed prior to overt capsular or vascular invasion, even
though pre-requisite conditions for invasion are already present in the tumors.

In conclusion, we present the first protein-based artificial neural networks classifier
for classifying thyroid nodules. Although this test has been retrospectively validated in
three clinical centers and prospectively validated in a fourth independent center, further
validation using FNA biopsy specimens from larger prospective cohorts are required.
This large-scale thyroid proteome profile of 931 thyroid nodules coupled with artificial
neural networks demonstrates the power of a protein-based disease classifier with rapid
potential to be translated into clinical practice. The thus established protein classifiers
may complement nucleic acid-based tests in multiple clinical applications. Expanding
this robust workflow to other carefully curated clinical cohorts may offer unprecedented
opportunities to gain fundamental insights into molecular pathogenesis of diseases and
address critical unmet clinical needs beyond thyroid cancer.

ACKNOWLEDGEMENTS

This work was supported by grants from National Natural Science Foundation of China
(81972492) and National Natural Science Fund for Young Scholars (21904107),
Zhejiang Provincial Natural Science Foundation for Distinguished Young Scholars
(LR19C050001), Hangzhou Agriculture and Society Advancement Program
(20190101A04). Further support for this project was obtained from the National Cancer
Centre Research Fund (Peter Fu Program) and National Medical Research Council
Clinician-Scientist Award (NMRC/CSAINV/011/2016), both in Singapore. The
prospective study was supported by National Natural Science Foundation of
China (81902726), China Postdoctoral Science Foundation
(NO. 2018M641739), Natural Science Foundation of Liaoning Province (20180530090).

AUTHOR CONTRIBUTIONS

T.G., N.G.I., O.L.K., S.Z.L., Y.Zhu and Y.S. designed the project. SS performed central
pathology review of the discovery set. H. Z. collected the samples for prospective test
set. L.C., X.T., Y.Zhao, G.W., J.X., Z.W., W.S., Y.Q., Y.H., L.W. and S.Z. procured and
annotated the Chinese thyroid tissue samples, while SS, SM, TKHL, SY, SMFSA, SSL,
BH procured and annotated the Singapore tissue samples. Y.S., W.L., X.C., X.Z., Q.X.,
H.G., X.Z. and K. X. performed the experiments. Y.S., W.L., W.W., H.C., T.Z. and Q.Z.
conducted proteomic data analysis. Z.Z. and S.Z.L. performed the learning analysis.
Y.S., N.G.I., O.L.K. and T.G. wrote the manuscript with inputs from all co-authors. T.G.,

N.G.I., O.L.K. and S.Z.L. supervised the project.

DECLARATION OF INTERESTS

The research group of T.G. is supported by Pressure Biosciences Inc, which provides sample preparation instrumentation. Authors declare no competing interests.

DATA AVAILABILITY

All data are available in the manuscript or the supplementary material. MS raw data were deposited in iProX (IPX000444000). All the data will be publicly released upon publication.

REFERENCES

Agrawal, N., Akbani, R., Aksoy, B., Ally, A., Arachchi, H., Asa, S., Auman, J., Balasundaram, M., Balu, S., Baylin, S., *et al.* (2014). Integrated genomic characterization of papillary thyroid carcinoma. Cell *159*, 676-690.

Ahn, H.S., Kim, H.J., and Welch, H.G. (2014). Korea's Thyroid-Cancer "Epidemic" — Screening and Overdiagnosis. N Engl J Med *371*, 1765-1767.

Alexander, E.K., Kennedy, G.C., Baloch, Z.W., Cibas, E.S., Chudova, D., Diggans, J., Friedman, L., Kloos, R.T., LiVolsi, V.A., Mandel, S.J., *et al.* (2012). Preoperative diagnosis of benign thyroid nodules with indeterminate cytology. N Engl J Med *367*, 705-715.

Bartolazzi, A., Sciacchitano, S., and D'Alessandria, C. (2018). Galectin-3: The Impact on the Clinical Management of Patients with Thyroid Nodules and Future Perspectives. International journal of molecular sciences *19*.

Becht, E., McInnes, L., Healy, J., Dutertre, C., Kwok, I., Ng, L., Ginhoux, F., and Newell, E. (2018). Dimensionality reduction for visualizing single-cell data using UMAP. Nature biotechnology.

Blazanin, N., Son, J., Craig-Lucas, A.B., John, C.L., Breech, K.J., Podolsky, M.A., and Glick, A.B. (2017). ER stress and distinct outputs of the IRE1α RNase control proliferation and senescence in response to oncogenic Ras. Proc Natl Acad Sci USA *114*, 9900-9905.

Brix, K., Summa, W., Lottspeich, F., and Herzog, V. (1998). Extracellularly occurring histone H1 mediates the binding of thyroglobulin to the cell surface of mouse macrophages. J Clin Invest *102*, 283-293.

Burman, K.D., and Wartofsky, L. (2015). CLINICAL PRACTICE. Thyroid Nodules. N Engl J Med *373*, 2347-2356.

Ciregia, F., Giusti, L., Molinaro, A., Niccolai, F., Mazzoni, M.R., Rago, T., Tonacchera, M., Vitti, P., Giannaccini, G., and Lucacchini, A. (2016). Proteomic analysis of fine-needle aspiration in differential diagnosis of thyroid nodules. Translational research : the journal of laboratory and clinical medicine *176*, 81-94.

Eszlinger, M., Lau, L., Ghaznavi, S., Symonds, C., Chandarana, S.P., Khalil, M., and Paschke, R. (2017). Molecular profiling of thyroid nodule fine-needle aspiration cytology. Nat Rev Endocrinol *13*, 415-424.

Faquin, W.C., Bongiovanni, M., and Sadow, P.M. (2011). Update in thyroid fine needle aspiration. Endocr Pathol *22*, 178-183.

Farrokhi Yekta, R., Arefi Oskouie, A., Rezaei Tavirani, M., Mohajeri-Tehrani, M.R., and Soroush, A.R. (2018). Decreased apolipoprotein A4 and increased complement component 3 as potential markers for papillary thyroid carcinoma: A proteomic study. Int J Biol Markers *33*, 455-462.

Figliozzi, R.W., Chen, F., and Hsia, S.V. (2017). New insights on thyroid hormone mediated regulation of herpesvirus infections. Cell Biosci *7*, 13.

Ganly, I., Makarov, V., Deraje, S., Dong, Y., Reznik, E., Seshan, V., Nanjangud, G., Eng, S., Bose, P., Kuo, F., *et al.*

(2018). Integrated Genomic Analysis of Hurthle Cell Cancer Reveals Oncogenic Drivers, Recurrent
Mitochondrial Mutations, and Unique Chromosomal Landscapes. Cancer Cell *34*, 256-270 e255.
Gillet, L.C., Navarro, P., Tate, S., Rost, H., Selevsek, N., Reiter, L., Bonner, R., and Aebersold, R. (2012). Targeted
data extraction of the MS/MS spectra generated by data-independent acquisition: a new concept for consistent
and accurate proteome analysis. Mol Cell Proteomics *11*, O111.016717.
Gopal, R.K., Kubler, K., Calvo, S.E., Polak, P., Livitz, D., Rosebrock, D., Sadow, P.M., Campbell, B., Donovan, S.E.,
Amin, S., *et al.* (2018). Widespread Chromosomal Losses and Mitochondrial DNA Alterations as Genetic Drivers
in Hurthle Cell Carcinoma. Cancer Cell *34*, 242-255 e245.
Grani, G., Lamartina, L., Durante, C., Filetti, S., and Cooper, D.S. (2018). Follicular thyroid cancer and Hürthle cell
carcinoma: challenges in diagnosis, treatment, and clinical management. The Lancet Diabetes & Endocrinology
6, 500-514.
Guo, T., Kouvonen, P., Koh, C.C., Gillet, L.C., Wolski, W.E., Rost, H.L., Rosenberger, G., Collins, B.C., Blum, L.C.,
Gillessen, S., *et al.* (2015). Rapid mass spectrometric conversion of tissue biopsy samples into permanent
quantitative digital proteome maps. Nat Med *21*, 407-413.
He, Y., Cao, L., Wang, L., Liu, L., Huang, Y., and Gong, X. (2020). Metformin Inhibits Proliferation of Human
Thyroid Cancer TPC-1 Cells by Decreasing LRP2 to Suppress the JNK Pathway. Onco Targets Ther *13*, 45-50.
Jameson, J.L. (2012). Minimizing unnecessary surgery for thyroid nodules. N Engl J Med *367*, 765-767.
Jiang, Y., Sun, A., Zhao, Y., Ying, W., Sun, H., Yang, X., Xing, B., Sun, W., Ren, L., Hu, B., *et al.* (2019). Proteomics
identifies new therapeutic targets of early-stage hepatocellular carcinoma. Nature *567*, 257-261.
Kashat, L., So, A.K.C., Masui, O., Wang, X.S., Cao, J., Meng, X., Macmillan, C., Ailles, L.E., Siu, K.W.M., Ralhan, R.,
et al. (2010). Secretome-based identification and characterization of potential biomarkers in thyroid cancer.
Journal of proteome research *9*, 5757-5769.
Kim, H.-S., Kim, G.Y., Lim, S.-J., and Kim, Y.W. (2010). Raf-1 kinase inhibitory protein expression in thyroid
carcinomas. Endocrine pathology *21*, 253-257.
Kohavi, R. (1995). A Study of Cross-Validation and Bootstrap for Accuracy Estimation and Model Selection.
International Joint Conference of Artificial Intelligence.
M, G., A, W., M, P., Ł, M., M, C., K, J., D, L., R, J., A, G., and P, W. (2018). Proteome profiles of different types of
thyroid cancers. Molecular and cellular endocrinology *472*, 68-79.
Martínez-Aguilar, J., Clifton-Bligh, R., and Molloy, M.P. (2016). Proteomics of thyroid tumours provides new
insights into their molecular composition and changes associated with malignancy. Sci Rep *6*, 23660.
Mitchell, M. (1998). Elements of Generic Algorithms. An Introduction to Generic Algorithms, 158.
Nikiforova, M.N., Wald, A.I., Roy, S., Durso, M.B., and Nikiforov, Y.E. (2013). Targeted next-generation
sequencing panel (ThyroSeq) for detection of mutations in thyroid cancer. The Journal of clinical endocrinology
and metabolism *98*, E1852-1860.
Patel, K.N., Angell, T.E., Babiarz, J., Barth, N.M., Blevins, T., Duh, Q.Y., Ghossein, R.A., Harrell, R.M., Huang, J.,
Kennedy, G.C., *et al.* (2018). Performance of a Genomic Sequencing Classifier for the Preoperative Diagnosis of
Cytologically Indeterminate Thyroid Nodules. JAMA Surg *153*, 817-824.
Rost, H.L., Rosenberger, G., Navarro, P., Gillet, L., Miladinovic, S.M., Schubert, O.T., Wolski, W., Collins, B.C.,
Malmstrom, J., Malmstrom, L., *et al.* (2014). OpenSWATH enables automated, targeted analysis of data-
independent acquisition MS data. Nat Biotechnol *32*, 219-223.
Schubert, O.T., Gillet, L.C., Collins, B.C., Navarro, P., Rosenberger, G., Wolski, W.E., Lam, H., Amodei, D., Mallick,
P., MacLean, B., *et al.* (2015). Building high-quality assay libraries for targeted analysis of SWATH MS data. Nat
Protoc *10*, 426-441.
Schürch, C.M., Roelli, M.A., Forster, S., Wasmer, M.-H., Brühl, F., Maire, R.S., Di Pancrazio, S., Ruepp, M.-D.,
Giger, R., Perren, A., *et al.* (2019). Targeting CD47 in Anaplastic Thyroid Carcinoma Enhances Tumor
Phagocytosis by Macrophages and Is a Promising Therapeutic Strategy. Thyroid *29*, 979-992.

Shao, S., Guo, T., Gross, V., Lazarev, A., Koh, C.C., Gillessen, S., Joerger, M., Jochum, W., and Aebersold, R. (2016).
Reproducible Tissue Homogenization and Protein Extraction for Quantitative Proteomics Using MicroPestle-
Assisted Pressure-Cycling Technology. Journal of proteome research *15*, 1821-1829.
Shao, S., Guo, T., Koh, C.C., Gillessen, S., Joerger, M., Jochum, W., and Aebersold, R. (2015). Minimal sample
requirement for highly multiplexed protein quantification in cell lines and tissues by PCT-SWATH mass
spectrometry. Proteomics *15*, 3711-3721.
Shao, W., Guo, T., Toussaint, N.C., Xue, P., Wagner, U., Li, L., Charmpi, K., Zhu, Y., Wu, J., Buljan, M., *et al.* (2019).
Comparative analysis of mRNA and protein degradation in prostate tissues indicates high stability of proteins.
Nat Commun *10*, 2524.
Sinha, A., Huang, V., Livingstone, J., Wang, J., Fox, N.S., Kurganovs, N., Ignatchenko, V., Fritsch, K., Donmez, N.,
Heisler, L.E., *et al.* (2019). The Proteogenomic Landscape of Curable Prostate Cancer. Cancer Cell *35*, 414-427
e416.
Smith, A., Galli, M., Piga, I., Denti, V., Stella, M., Chinello, C., Fusco, N., Leni, D., Manzoni, M., Roversi, G., *et al.*
(2019). Molecular signatures of medullary thyroid carcinoma by matrix-assisted laser desorption/ionisation
mass spectrometry imaging. J Proteomics *191*, 114-123.
Steward, D.L., Carty, S.E., Sippel, R.S., Yang, S.P., Sosa, J.A., Sipos, J.A., Figge, J.J., Mandel, S., Haugen, B.R.,
Burman, K.D., *et al.* (2019). Performance of a Multigene Genomic Classifier in Thyroid Nodules With
Indeterminate Cytology: A Prospective Blinded Multicenter Study. JAMA Oncol *5*, 204-212.
Sun, R., et al, Guo, T. (2020). Accelerated Protein Biomarker Discovery from FFPE Tissue Samples Using Single-
Shot, Short Gradient Microflow SWATH MS. Journal of Proteome Research.
Vaccarella, S., Franceschi, S., Bray, F., Wild, C.P., Plummer, M., and Dal Maso, L. (2016). Worldwide Thyroid-
Cancer Epidemic? The Increasing Impact of Overdiagnosis. N Engl J Med *375*, 614-617.
Wang, T.S., and Sosa, J.A. (2018). Thyroid surgery for differentiated thyroid cancer - recent advances and future
directions. Nat Rev Endocrinol *14*, 670-683.
Wang, Y., He, H., Li, W., Phay, J., Shen, R., Yu, L., Hancioglu, B., and de la Chapelle, A. (2017). MYH9 binds to
lncRNA gene PTCSC2 and regulates FOXE1 in the 9q22 thyroid cancer risk locus. Proc Natl Acad Sci USA *114*,
474-479.
Zhang, H., Liu, T., Zhang, Z., Payne, S.H., Zhang, B., McDermott, J.E., Zhou, J.Y., Petyuk, V.A., Chen, L., Ray, D.,
et al. (2016). Integrated Proteogenomic Characterization of Human High-Grade Serous Ovarian Cancer. Cell
166, 755-765.
Zhu, T., et al, Guo, T. (2020). DPHL: A pan-human protein mass spectrometry library for robust biomarker
discovery using Data-Independent Acquisition and Parallel Reaction Monitoring. Genomics, Proteomic and
Bioinformatics, https://www.biorxiv.org/content/10.1101/2020.1102.1103.931329v931321.
Zhu, Y., Weiss, T., Zhang, Q., Sun, R., Wang, B., Yi, X., Wu, Z., Gao, H., Cai, X., Ruan, G., *et al.* (2019). High-
throughput proteomic analysis of FFPE tissue samples facilitates tumor stratification. Mol Oncol.

Table 1. Clinico-pathologic characteristics of the study cohorts.

| | Discovery dataset | Independent test datasets | | All |
		Retrospective test datasets	Prospective test dataset	
Total no.				
Clinical centers	1	3	1	5
patients	578	271	62	911
nodules	579	288	64	931
FFPE punches	1725	288	0	2013
Fine needle aspiration biopsies	0	0	64	64
DIA files	1781	576	64	2421
Histopathology diagnosis				
Normal thyroid tissues	40 (6.9%)	16 (5.6%)	0 (0.0%)	56 (6.0%)
Multinodular goiter	203 (35.1%)	44 (15.3%)	13 (20.3%)	260 (27.9)
Follicular adenoma[b]	137 (23.7%)	84 (29.2%)	5 (7.8%)	226 (24.3%)
Follicular thyroid carcinoma[b]	75 (13.0%)	52 (18.1%)	0 (0.0%)	127 (13.6%)
Papillary thyroid carcinoma	124 (21.4%)	92 (31.9%)	46 (71.9%)	262 (28.1%)
Diagnosis age				
Mean	52.8	47.5	43.6	50.6
Median	54	49	44	52
Range	13-85	15-82	24-74	13-85
< 18 y	7 (1.2%)	1 (0.4%)	0 (0.0%)	8 (0.9%)
18-55 y	288 (49.8%)	173 (63.8%)	48 (77.4%)	509 (55.9%)
≥55 y	283 (49.0%)	97 (35.8%)	14 (22.6%)	394 (43.2%)
Gender				
Female	433 (74.9%)	191 (66.3%)	54 (87.1%)	678 (74.4%)
Male	145 (25.1%)	80 (27.7%)	8 (12.9%)	233 (25.6%)
Nodule size[a] on ultrasonography				
Mean	3.3	2.9	2.5	3.1
Median	3.0	2.6	2.2	2.9
Range	0.2-13.0	0.2-8.9	0.6-5.5	0.2-13.0
< 1 cm	15 (2.8%)	37 (14.0%)	2 (3.1%)	54 (6.3%)
1~4 cm	342 (64.5%)	143 (54.0%)	53 (82.8%)	538 (62.6%)
≥4 cm	173 (32.6%)	85 (32.1%)	9 (14.1%)	267 (31.1%)

473 [a]Nodule size information of nine patients in discouvery dataset and seven patients in test dataset were not
474 record.
475 [b]Hurthle cell adenoma and carcinoma were allocated into follicular thyroid adenoma and carcinoma,
476 respectively.
477
478

Table 2. Fourteen proteins selected by genetic algorithm, functional pathways and previously known associated thyroid physiology or pathology.

Uniprot ID	Symbol	Entrez Gene Name	Thyroid cancer related	Thyroid function related
O75347	TBCA	tubulin folding cofactor A	-	Yes
P02765	AHSG	alpha 2-HS glycoprotein	Yes	-
P04083	ANXA1	annexin A1	Yes	Yes
P04216	THY1	Thy-1 cell surface antigen	-	-
P10909	CLU	clusterin	Yes	-
P16403	HIST1H1C	histone cluster 1 H1 family member c	-	Yes
P17931	LGALS3	galectin 3	Yes	-
P26038	MSN	moesin	Yes	-
P27797	CALR	calreticulin	Yes	Yes
P30086	PEBP1	phosphatidylethanolamine binding protein 1	Yes	Yes
P35579	MYH9	myosin heavy chain 9	Yes	-
P98164	LRP2	LDL receptor related protein 2	Yes	-
Q9H299	SH3BGRL3	SH3 domain binding glutamate rich protein like 3	Yes	Yes
Q9HAT2	SIAE	sialic acid acetylesterase	-	-

483   **Table 3. Performance of the protein classifier.**

Center	Retrospective test sets				Prospective test set	All
	ZY	ZE	DL	All retrospective study	FNA	
Sample type	FFPE	FFPE	FFPE	FFPE	Biopsy	FFPE+Biopsy
Malignant nodules No.	35	61	48	144	46	190
Benign nodules No.	19	79	46	144	18	162
Prevalence[a]	64.81 (53.62 - 74.59)	43.57 (36.84 - 50.54)	51.06 (42.65 - 59.42)	50.00 (45.16 - 54.84)	71.88 (61.86 - 80.11)	53.98 (49.58 - 58.31)
Predict M/M[b]	31	58	41	130	40	170
Predict B/B[c]	17	71	43	131	16	147
Sensitivity	88.57 (76.80 - 94.78)	95.08 (88.29 - 98.02)	85.42 (75.12 - 91.91)	90.28 (85.43 - 93.63)	86.96 (76.67 - 93.12)	89.47 (85.23 - 92.61)
Specificity	89.47 (72.59 - 96.46)	89.87 (82.88 - 94.21)	93.48 (84.73 - 97.37)	90.97 (86.23 - 94.19)	88.89 (71.30 - 96.26)	90.74 (86.28 - 93.85)
PPV	93.94 (86.86 - 97.76)	87.88 (82.57 - 91.70)	93.18 (88.11 - 96.67)	90.91 (87.79 - 93.39)	95.24 (88.81 - 98.12)	91.89 (89.01 - 93.87)
NPV	80.95 (71.33 - 88.61)	95.95 (91.91 - 97.77)	86.00 (79.27 - 91.03)	90.34 (87.01 - 92.79)	72.73 (63.51 - 81.45)	88.02 (84.92 - 90.63)
Accuracy	88.89 (79.89 - 94.16)	92.14 (87.54 - 95.14)	89.36 (82.96 - 93.54)	90.62 (87.40 - 93.09)	87.50 (79.12 - 92.82)	90.06 (87.11 - 92.39)
Youden's index	0.78	0.85	0.74	0.81	0.76	0.80

485   Each value was calculated to 95% Wilson confidence intervals.

486   [a]The ratio of carcinoma in total nodules.

487   [b]The number of benign nodules identified as benign.

488   [c]The number of malignant nodules identified as malignant.

FIGURE LEGEND

Figure 1. **Schematic view of the study and batch design.** (A) The project design and workflow of the FFPE PCT-DIA pipeline. (B-C) Batch design (B) The discovery group of 579 thyroid nodules from 578 patients consisted of 40 normal thyroid, 203 multinodular goiter, 137 follicular thyroid adenoma, 75 follicular thyroid carcinoma and 124 papillary thyroid carcinomas with unblinded diagnoses. Each nodule was represented by three cores as biological replicates. 1,781 thyroid FFPE punches and 56 technical replicates were randomly allocated into 121 discovery batches in order to minimize batch effect for this large-scale sample preparation. (C) The independent test group contained two parts, one is retrospective test group and the other is prospective test group. Retrospective test group included 288 thyroid nodules of blinded diagnoses from 271 patients. Each nodule was analyzed in technical duplicates, without biological replicates. A total of 288 FFPE cores and 288 corresponding technical duplicates were divided into 44 batches for analysis. Prospective test group contained 64 fine needle aspiration biopsies of thyroid nodules from 64 patients which were divided into 5 batches. Each batch consisted of 15 thyroid samples, one mouse liver sample and one pooled thyroid sample.

Figure 2. Global thyroid proteome profile. (A) Heatmap showing global protein expression profiles of the five main histological types of thyroid tissues. 579 thyroid tissues from 578 patients are in columns and 2,617 proteins (missing value <90%) in rows. The color bar indicates the intensity of the proteins. (B-F) UMAP plots showing global snapshots comparing the indicated types thyroid tissues using 2,617 proteins for (B) all subtypes; (C) benign versus malignant; (D) only benign; (E) FA versus FTC; and (F) only malignant tissue types.

Figure 3. Classifier development, performance testing and validation in independent blinded datasets. (A) Schematic of principal classifier model (details in Supplementary Methods). (B) ROC plots of seven different machine learning model under 14 selected features. (C) t-SNE plot showing the separation between benign malignant groups in the discovery set using 14 protein features with latent space; and ROC plots of the training and validation sample subgroups of the discovery set. (D-E) t-SNE and ROC plots of the performance for (D) retrospective test sets; and (E) prospective test set (total cohort and individual hospital sites as indicated). (F) Overall performance metrics of prediction of the neural network model for five specific histopathological types per site. Graduated colors in the shaded bar indicate accuracy levels. Numbers in the boxes indicate the number of correctly identified samples/total sample number.

Figure 4. Protein expression plots for 14 selected protein features in the five types of thyroid tissues in the discovery cohort. Y-axis shows Log2 values of protein expression intensity and x axis indicates tissue type. P-value was calculated by one-way ANOVA. Further protein details are listed alongside each plot.

Figure 5. Biological insights of thyroid tumor subtypes based on proteotypic data.
(A) UMAP plot for 109 proteins distinguishing Hürthle cell from other follicular
neoplasms. (B) Network map showing expression of key mitochondrial proteins
implicated in Hürthle cell neoplasms. (C) UMAP plot for 175 proteins distinguishing FTC
from cPTC, with fvPTC as an intermediate phenotype. (D) Heatmap showing
differentially expressed proteins in FTC compared with fvPTC and cPTC, with pathways
as indicated. X-axis of the vertical bar plot indicates -log2(p-value) based on right-tailed
Fisher's exact test from the IPA database. (E) Volcano plot showing 50 up-regulated
proteins in FTC and 97 up-regulated proteins in cPTC with two-fold-change cutoff and
adjusted p-value threshold less than 0.01. (F) Proteins in the cPTC network participate
in immune pathways and oncogenesis (MYC and TP53 pathways). (G) Box plots
showing three proteins specifically overexpressed in cPTC compared to other
histological subtypes.

MATERIALS AND METHODS

Patients and tissue samples

Tissue cores (1 mm diameter, approximate weight 0.6 mg, including was) were punched from blocks of formalin-fixed paraffin embedded (FFPE) thyroid tissues obtained from four clinical centers in Singapore and China spanning 2011-2019, with ethics approval of each hospital.

The discovery sample set of 579 thyroid nodules from 578 patients comprised follicular adenomas (FA, 137 cases), multinodular goiters (MNG, 203 cases), papillary thyroid carcinoma (PTC, 124 cases) and follicular thyroid carcinoma (FTC, 75 cases) from the Singapore General Hospital. Hematoxylin and eosin-stained slides from tissue blocks of each patient were reviewed by an experienced histopathologist who marked out the disease region for tissue coring. Normal thyroid tissues (N, 40 cases) were taken from cases of laryngectomy or pharyngo-laryngo-esophagectomy in which the thyroid gland was surgically removed incidental to radical surgery. These patients had no history of thyroid disease, prior chemotherapy or radiation. Three tissue cores were made for each case as biological replicates.

For multi-center blinded test sets, we firstly analyzed a total of 288 FFPE tissue cores from 271 cases composed of 44 MNG, 84 FA, 52 FTC and 92 PTC cases in retrospective test sets. Sixteen cores were of adjacent normal thyroid tissue (N). A single punch was made from each case. Furthermore, we tested 64 fine needle aspiration biopsies containing 13 MNG, 5 FA and 46 PTC cases in prospective test set. Histological diagnoses of these cores were blinded during the entire validation workflow of sample processing, mass spectrometry analysis and predictive data analysis.

Cases of microcarcinoma, extensive thyroiditis and/or inflammation were excluded from the discovery sample set.

Batch design

To minimize batch effects among different lots of analyzed samples, 1,725 FFPE cores from 579 thyroid nodules and 56 peptides extracted from cores selected as technical replicates from the discovery sample set were randomly distributed into 121 batches. Each batch contained 15 thyroid samples, one mouse liver sample as quality control (QC) for PCT and one thyroid pooled sample containing all five types of thyroid tissues for mass spectrometry (MS). In this discovery phase analysis, the histopathology diagnosis of each tissue core was known from which models of data segregation were established.

In the test phase analysis, 288 FFPE cores were analyzed in technical duplicates for a total of 576 samples in 44 batches for retrospective test sets and 64 fine needle biopsies in 5 batches for prospective test set (Figure S1).

Dewaxing, rehydration and hydrolysis of FFPE tissues

For each case in the discovery sample set, three biological replicates of FFPE
tissue cores were processed. Sample weights were recorded before dewaxing in
heptane (Sigma) and successively in 100% ethanol (Sigma), 90% ethanol, 75% ethanol
at room temperature. Formic acid (0.1%) (Sigma) was added next for achieve C-O
hydrolysis of protein methylol products and then washed with 100 mM Tris-HCl (pH10,
Sigma) to establish conditions for base hydrolysis at 95℃. The sample was then snap
cooled to 4℃.
595

Tissue lysis, protein extraction and protein digestion

Dewaxed samples were lysed in 6M urea (Sigma) and 2M thiourea (Sigma) using
pressure cycling technology (PCT) programmed for 90 cycles of 25 s at 45,000 p.s.i.
and 10 s at ambient pressure at 30℃. After lysis, 10 mM Tris(2-carboxyethyl) phosphine
hydrochloride and 40 mM iodoacetamide were simultaneously added to the solution and
incubated in the dark with gentle vortexing for 30 min, after which lysC (Hualishi
Scientific) was added at a ratio of 40:1 (protein to lysC). PCT-assisted lysC digestion
was performed with the following setting: 45 cycles of 50 s at 20,000 p.s.i. and 10 s at
ambient pressure at 30℃. Final tryptic digestion was performed at a ratio of 40:1
(protein to trypsin) by PCT with the following setting: 90 cycles of 50 s at 20,000 p.s.i.
and 10 s at ambient pressure at 30℃. Peptides were desalted before LC-MS analysis.
607

DIA library construction

DIA (data-independent acquisition) library was built as described previously (Guo et
al., 2015; Schubert et al., 2015). To build the spectral library for analyzing DIA files from
thyroid tissue samples, we collected tissue samples from the all five types of patient
groups into a single pool. The tissue samples were either fractionated into six fractions
using strong cation exchange (SCX) or processed with PCT-assisted lysis and in-
solution digestion, or PCT-assisted lysis and PCT-assisted digestion. Finally, peptides
were desalted on C18 columns.
Desalted peptides were separated by Ultimate 3000 nanoLC-MS/MS system
(Dionex LC-Packing, Amsterdam, The Netherlands) equipped with 15 cm* 75 μm ID
fused silica column custom packed with 1.9 μm 100 Å C18 AQUA. Peptides were
separated on a 120 min (148 min inject-to-inject) LC gradient at 300 nL/min in a 3-25%
linear gradient (buffer A: 2% acetonitrile,0.1% formic acid; buffer B: 98% acetonitrile,
0.1% formic acid). Eluted peptides were ionized at a potential of +2.0 kV into Q Exactive
HF mass spectrometer (Thermo Fisher, Bremen, Germany). The full MS was measured
at resolution 60,000 (at m/z 200) in an Orbitrap using an AGC target value of 3E6
charges. The top 20 peptide signals (charge-states +2 and higher) were submitted to
fragmentation in HCD cell (higher-energy collision, 27% normalized collision energy),
and then transferred to Orbitrap for MS/MS analysis. MS/MS spectra were acquired at
resolution 30,000 (at m/z 200) in Orbitrap using an AGC target value of 1E5 charges, a
maxIT of 80 ms and the dynamic exclusion time was 30 s.
In total, we acquired 52 DDA files including 20 SCX fraction files, 18 in-solution

digestion files and 14 PCT-assisted digestion files on a QE-HF mass spectrometer in DDA mode (Table S2). All DDA files were centroided and converted to mzXML using msConvert from ProteoWizard with parameters '—mzXML --filter "peakPicking true [1,2]"'. We analyzed DDA files using pFind version 3.1.3 with SwissProt fasta files (20,269 protein sequences), and identified 47,229 transition groups, 33,802 peptides, 5,190 protein groups and 4,048 proteotypic proteins.

DIA-MS data analysis

Peptides were separated by using Ultimate 3000 or nanoLC-MS/MS system (DIONEX UltiMate 3000 RSLCnano System, Thermo Fisher Scientific™, San Jose, USA) equipped with 15cm*75 µm ID fused silica column custom packed with 1.9 µm 120 Å C18 aqua. Peptides were separated on a 45 min (68 min inject-to-inject) LC gradient at 300 nL/min in a 3-25% linear gradient (buffer A: 2% acetonitrile,0.1% formic acid; buffer B: 98% acetonitrile, 0.1% formic acid). Peptides eluted from analytical columns were ionized at a potential +2.0 kV into Q Exactive HF mass spectrometer (Q Exactive Hybrid Quadrupole-Orbitrap, Thermo Fisher Scientific™, San Jose, USA). A full MS scan was acquired analyzing 390-1010 *m/z* at resolution 60,000 (at *m/z* 200) in the Orbitrap using an AGC target value of 3E6 charges and maximum injection time of 100 ms. After the full MS scan, 24 MS/MS scans were acquired, each with a 30,000 resolution (at m/z 200), AGC target value of 1E6 charges, normalized collision energy of 27%, with the default charge state set to 2, maximum injection time set to auto. The cycle of 24 MS/MS scans (center of isolation window) with three kinds of wide isolation window was as follows (m/z): 410, 430, 450, 470, 490, 510, 530, 550, 570, 590, 610, 630, 650, 670, 690, 710, 730, 770, 790, 820, 860, 910, 970. The entire cycle of MS and MS/MS scan acquisition took approximately 3 seconds and was repeated throughout the LC/MS analysis. DIA raw files were analyzed using OpenSWATH v2.0 (Rost et al., 2014).

Data quality control

We first assessed data quality by analyzing samples. The QC samples were mouse liver samples (PCT-QC) and pooled thyroid samples (DIA-QC) in each batch. Additional QC samples were analyzed as technical replicates for MS. Biological replicates were also analyzed to determine the extent of heterogeneity of thyroid diseases. Reproducibility of spiked-in mouse liver samples and thyroid pooled samples showed that PCT and MS instruments were stable during data acquisition (Figure S2A and B), with median coefficient of variance (CV) less than 0.03. MS data of 56 randomly selected paired thyroid samples in the discovery cohort had a median Spearman correlation coefficient of 0.9 (Figure S2C). CV for proteins in biological replicates was 0.2, indicating minimal tissue heterogeneity in the biology of thyroid disease (Figure S2D). Spearman correlations of four pooled samples and mouse liver samples are higher than 0.9 in prospective test set (Figure S2E and F)

Development of neural network classifier

A neural network classifier is developed to classify any given proteome data sample (of 14 features) into one of two classes, benign (B) or malignant (M), so as to achieve the best accuracy. This comprises 4 stages:

    (1) Protein data preprocessing;

    (2) Protein feature selection using a genetic algorithm (GA);

    (3) Learning a neural network (DNN)-based classifier;

    (4) Using the trained DNN for sample classification.

The following details the algorithms for the 4 stages.

Stage 1: Data preprocessing

Four datasets from 5 cohorts, denoted SG, ZY, ZE, DL, FNA, were used for the development of the DNN model. There, 500 random samples from the SG cohort were used to train the model, 79 from the same cohort were used for internal validation, and data from the ZY, ZE, DL and FNA cohorts were used for external validation. Note that the external validation datasets are different those of training and internal validation. A data sample in these datasets consists of 3,708 proteins.

The preprocessing consists of two steps: (1) missing value imputation and (2) normalization. Missing values are inevitably a feature of protein intensity data. Considering most missing values occur when the protein content is below a detection threshold, the imputation is done by filling in all the missing values with $0.8*D_{min}$, where D_{min} is the minimum of all feature values in the training and validation sets and for this work, $D_{min} = 13$.

Thereafter, for each feature D after the imputation step, the mean μ and variance σ of that feature are estimated from the training and validation sets, and the feature of every training sample is normalized according to the following

$$D^n = \frac{D - \mu}{\sigma} \qquad (a)$$

where D^n is the normalized feature. The obtained μ and σ are applied to the corresponding protein features in the testing phase afterwards.

Python's pandas library is used to complete data preprocessing.

Stage 2: Features selection

Feature selection is needed for two reasons: (1) most of the 3,708 proteins are useless or may be conflicting for the classification and they should be removed from consideration in DNN-based classifier learning; (2) it is desired to minimize the number of proteins in clinical applications. It is done in two steps (Figure S5A).

The first step is feature screening. Out of the initial 3,708 protein features, 521 were selected from differentially expressed protein of benign and malignant samples in the SG dataset and from published literatures related to thyroid or thyroid cancer. Among the 521, 64 did not appear in our datasets and are excluded, with 477 candidates remaining. Further, if such a protein has a deletion rate greater than 45%, then it is removed, with 275 proteins left. Even further, a pair of protein are removed if the absolute value of the Pearson correlation between them is less than 0.1, yielding the 243 candidate proteins.

As the second step, a combinatorial optimization is performed to select a best
combination of 14 proteins from the 243. While no algorithms can guarantee a globally
optimal solution with efficiency, here a genetic algorithm (GA)(Mitchell, 1998) was used
to find the best 14 proteins.
In GA, evolutionary operation (crossover, mutation, and selection operations) are
used to generate new protein feature combinations from existing protein features
combinations. At every iteration, the GA eliminates the low fitness combinations, and
generates new combinations based on the remaining high fitness combinations.
A fitness value is calculated for each candidate combination solution (gene chain).
For gene chain C, it is defined as

724
$$F^C = \frac{1}{5}\sum_{k=1}^{5} A_k^C \qquad\qquad (b)$$

where A_k^C is the accuracy of the k-th cross-validation(Kohavi, 1995), which is
computed from the difference between the output of the classifier and the true label.
Python's deap library is used to complete features selection.
Stage 3: Model training
The DNN classifier is a nonlinear function which takes a vector of 14 selected
protein features as the input and produces a class label of either 1 (for malignant) or 0
(for benign) as the output. This consists of the following 3 steps: (1) model structure
design; (2) loss function design; and (3) model training.
A multi-layer perceptron (MLP) structure is chosen for the DNN, shown in Figure
S5B, The MLP model is divided into feature extraction sub-model and classification sub-
model, trained in sequence as will be described below. The feature extraction sub-
model extracts effective feature vectors (V_p), and the classification sub-model completes
classification diagnosis based on the classification information.
Once the DNN structure is defined, the parameterized nonlinear function is trained
to achieve the best objective value as measured by a loss function. The loss function of
the MLP is defined as
741
$$L = \alpha_1 L_c + \alpha_2 L_e + \alpha_3 L_r \qquad\qquad (c)$$
742   where L_c is a contrastive loss (Becht et al., 2018) for training the feature extraction
743   sub-model, L_e is a cross-entropy loss for training the classification sub-model, L_r is an
744   L2 regularization loss for reducing overfitting, and α_1, α_2, $\alpha_3 > 0$ are the weights.
745   The contrastive loss L_c is defined under the Siamese Network (SN) structure. The
746   contrastive loss L_c is calculated as

747
$$L_c(p,q) = \begin{cases} \|V_p - V_q\|^2 & if\ \text{p, q}\ in\ the\ same\ class \\ max(M - \|V_p - V_q\|^2, 0)^2 & others \end{cases} \qquad (d)$$

748   where feature vectors V_p, V_q extracted from protein samples X_p and X_q, $\|V_p - V_q\|$ is
749   the distance, and M is the margin (M=1 in this work),. The contrastive loss encourages
750   reduction of within-class feature scatter and increase of between-class feature scatter.
751   The cross-entropy loss L_e is calculated as

$$L_e = -\left[\beta_1 Y \log\hat{Y} + \beta_2(1-Y)\log\left(1-\hat{Y}\right)\right] \qquad (e)$$

where Y is the real label of the patient, $\hat{Y}$ is the classification score predicted by classification sub-model, and β_1, β_2 are penalty parameters ($\beta_1 = 0.725, \beta_2 = 1.275$ in this work). The L2 regularizer is defined as the 2-norm of MLP weight W as follows

$$L_r = \|W\|_2^2 = \sum_{i=1}^{N} w_i^2 \qquad (f)$$

where N is the number of the layer.

The MLP training is done with the help of the training dataset (SG), in which the class label is provided for each sample as the supervisor to guide the training process. The MLP training process consists of two parts. The first part trains the feature extraction sub-model for 100 epochs, with $\alpha_1 = 1$, $\alpha_2 = 0$, $\alpha_3 = 0.04$ and learning rate 1×10^{-5}. The second part uses the cross-entropy loss training the classification sub-model for 900 epochs, with $\alpha_1 = 0$, $\alpha_2 = 1$, $\alpha_3 = 0.029$ and learning rate 5×10^{-4}, while the feature extraction sub-model parameters are not frozen for faster convergence. The obtained accuracy for the training set is 93%, and that for the validation set is 91%.

Python's pytorch library is used to complete model training.

Stage 4: Classification

The trained MLP is used as the classifier for diagnosis of unknown samples. Given the 14 features, the MLP outputs a classification score $\hat{Y}$ between 0 and 1 and the class prediction P is calculated as follows

$$P = \begin{cases} 1 & if\ \hat{Y} > 0.5 \\ 0 & if\ \hat{Y} < 0.5 \end{cases} \qquad (g)$$

where P=0 means the tissue is predicted to be benign, and P=1 malignant.

Statistical analysis

Statistical analysis was performed using R software (version 3.5.1) with pheatmap, UMAP and R package plot functions. CV was calculated as the ratio of the standard deviation to the mean. The prevalence for each cohort was based on the ratio of malignant to total tissues.   Sensitivity, specificity, PPV and NPV values were calculated following the established methodology and each value was calculated with 95% Wilson confidence intervals (Steward et al., 2019). Biological insights were analyzed by Ingenuity Pathway Analysis (IPA version 49309495). P values were calculated by one-way ANOVA in the expression of 14 protein features.

Figure S1. Data quality evaluation. (A) CV of identified protein numbers for 116 and 36 pooled thyroid samples in the discovery set and retrospective test sets, respectively. (B) CV of identified protein numbers for 112 and 18 mouse liver samples in discovery and retrospective test sets, respectively. (C) Spearman correlation of paired technical replicates from 56 randomly selected thyroid samples in the discovery group and 288 in the retrospective test sets. (D) CV for the number of proteins in biological replicates of the discovery set. (E) Spearman correlation of four pooled samples in prospective test set. (F) Spearman correlation of four mouse liver samples in prospective test set.

Figure S2. Uniform manifold approximation and projection (UMAP) analysis of five subtypes of thyroid tissues. 2,617 proteins for which missing value was less than 90% were used in data analysis. (A) All tissue types, showing FA distributed across benign (N and MNG) and malignant (FTC and PTC) tissues, (B) FA, (C) FTC, (D) PTC vs N, respectively, and (E) FA, (F) FTC and (G) PTC vs MNG, respectively. Normal tissue is generally well separated from all other lesional tissue, while MNG shows some overlap with FA, FTC and PTC.

Figure S3. Cross-validation of the classifier on discovery set and performance on test sets. (A) Scatter diagram showing the predicted malignancy scores for discovery (training and validation), and tests sets (retrospective and prospective test sets), and the probability for each sample to be malignant. X axis indicates the probability of malignancy. Y axis represents the number of thyroid tissues in different sets. (B) t-SNE plots showing specific tissue types (benign and malignant) based on the 14 protein features in the training set, validation set, retrospective and prospective test sets, labeled by each of the 5 subtypes. (C) t-SNE plots based on the 14 proteins for each of the four clinical sites for the different test sets.

Figure S4. Biological insights into Hürthle cell tumors, follicular (FTC), classical papillary (cPTC) and follicular-variant PTC (fvPTC). (A) Heatmap showing proteotype expression of thyroid tissue samples highlighting differentially expressed proteins in the Hürthle cell neoplasm marked in red frames. The subtype label of heatmap were based on reviewed slides by experienced pathologist. (B) Graph showing cellular component analysis of the 109 over-expressed proteins in Hürthle cell neoplasms. Mitochondrial proteins are the most dominant group. (C) Graph showing enriched pathways based on 109 over-expressed proteins of Hürthle cell tumors by IPA analysis. Y-axis shows -log2(p-value) based on right-tailed Fisher's exact test (IPA) based on the IPA database. (D) Volcano plots showing differentially expressed proteins between fvPTC vs FTC and cPTC vs fvPTC. Protein intensities used were from the average intensity of three biological replicates. We compared pair-wise groups, and highlighted proteins that were significantly different with two-fold-change cutoff and adjusted p-value threshold less than 0.01. (E) Pathway enrichment of 50 upregulated proteins in FTC compared with cPTC as analyzed by IPA. Y axis indicates -log2(p-value) based on right-tailed Fisher's exact test based on the IPA database. (F) Network map showing overexpressed proteins in FTC.

Figure S5. Structures of models. (A) Flow diagram of genetic algorithm for protein

features selection, i is the index of iteration, g is the index of gene chains. N_g is the

population size (number of the gene chains), N_i is the maximum number of iterations.

$C_{i,g}^b$ is the binary codes of i-th iteration g-th gene chian. $F^{C_{i,g}^b}$ is the fitness function of

$C_{i,g}^b$. (B) DNN's structure diagram with optimal parameters. p is the index of the

patients, X_p is the protein feature inputting to the feature extraction sub-model, V_p is

the output of the extraction sub-model, $\hat{Y}_p$ is the output of the classification sub-model

and the classification score predict by DNN, N is the number of neurons in the layer.

Figure 1. Schematic view of the study and batch design.

Figure 2. Global thyroid proteome profile.

Figure 3. Classifier development, performance testing and validation in
independent blinded datasets.

Figure 4. Classifier development, performance testing and validation in
independent blinded datasets.

849
850

851   **Figure 5. Biological insights of thyroid tumor subtypes based on proteotypic data.**

854　**Figure S1. Data quality evaluation.**

857 **Figure S2. Uniform manifold approximation and projection (UMAP) analysis of**
858 **five subtypes of thyroid tissues.**

Figure S3. Cross-validation of the classifier on discovery set and performance on test sets.

A

B

C

Figure S4. Biological insights into Hürthle cell tumors, follicular (FTC), classical papillary (cPTC) and follicular-variant PTC (fvPTC).

869   **Figure S5. Structures of artificial neural network models.**

(b) DNN's structure diagram with optimal parameters

871