

1 **Beginning section**

2 **Title page:**

3 Full title:

4 AN EPIDEMIOLOGICAL MODEL TO AID DECISION-MAKING

5 FOR COVID-19 CONTROL IN SRI LANKA

6

7 Short title:

8 AN EPIDEMIOLOGICAL MODEL FOR COVID-19 DECISION-MAKING

9

10 Authors:

11 Dileepa Senajith Ediriweera¹, Nilanthi Renuka de Silva², Neelika Gathsaurie Malavige³,

12 Hithanadura Janaka de Silva⁴

13

14 Affiliations:

15 ¹ Centre for Health Informatics, Biostatistics and Epidemiology, Faculty of Medicine,

16 University of Kelaniya, Ragama, Sri Lanka.

17 ² Department of Parasitology, Faculty of Medicine, University of Kelaniya, Ragama, Sri

18 Lanka

19 ³ Department of Microbiology, Faculty of Medical Sciences, University of Sri

20 Jayewardenepura, Nugegoda, Sri Lanka

21 ⁴ Department of Medicine, Faculty of Medicine, University of Kelaniya, Ragama, Sri Lanka

22

23

24 Corresponding author:

25 Dileepa Ediriweera email: dileepa@kln.ac.lk

26

27 **Abstract**

28 Background: Sri Lanka diagnosed its first local case of COVID-19 on 11 March 2020. The
29 government acted swiftly to contain transmission, with extensive public health measures. At
30 the end of 30 days, Sri Lanka had 197 cases, 54 recovered and 7 deaths; a staged relaxing of
31 the lockdown is now underway. This paper proposes a theoretical basis for estimating the
32 limits within which transmission should be constrained in order to ensure that the case load
33 remains within the capacity of the health system.

34

35 Methods: We used Susceptible, Infected, Recovered model to estimate the ICU bed
36 requirement at different levels of R_0 values after lockout. We developed a web application
37 that enables visualization of cases and ICU bed requirements with time, with adjustable
38 parameters that include: population exposed; proportion asymptomatic; number of active and
39 recovered cases; infectious period; R_0 or doubling time; proportion critically ill; available
40 ICU beds and duration of ICU stay.

41

42 Results: The three-day moving average of the caseload suggested two waves of transmission
43 from Day 0 to 17 ($R_0=3.32$, 95% CI 1.85 - 5.41) and from Day 18 - 30 ($R=1.25$, 95% CI: 0.93
44 - 1.63). We estimate that if there are 156 active cases with 91 recovered at the time of
45 lockout, and R increases to 1.5 (doubling time 19 days), under the standard parameters for Sri
46 Lanka, the ICU bed capacity of 300 is likely to be saturated by about 100 days, signalled by
47 18 new infections (95% CI 15 - 22) on Day 14 after lockout.

48

49 Conclusion: Our model suggests that to ensure that the case load remains within the available
50 capacity of the health system after lockout, transmission should not exceed $R=1.5$. This

- 51 model and the web-based application may be useful in other low- and middle-income
- 52 countries which have similar constraints on health resources.

53 **Introduction**

54 COVID-19 is caused by a new coronavirus (SARS CoV-2) that emerged in China in
55 December 2019. Although it causes an asymptomatic or mild infection in most instances, it
56 can cause severe respiratory illness or even death. Transmission is mainly via droplets
57 released into the air when an infected person coughs or sneezes. Aerosol and fomite
58 transmission of SARS-CoV-2 is also possible (1) (2). There is no vaccine at present, nor is
59 there any antiviral agent of proven efficacy. Thus traditional measures that control the spread
60 of infectious diseases such as quarantine, contact tracing, isolation of positives and contacts
61 as well as social distancing and hand-washing are of vital importance.

62

63 The basic reproduction number (R_0) is a central concept in infectious disease epidemiology,
64 representing the average number of new infections generated by an infectious person in a
65 completely susceptible population. For COVID-19, R_0 has been estimated by the WHO to be
66 1.4 – 2.5. Others have placed it higher, at a median of 2.79 with an interquartile (IQR) range
67 of 1.16 (3). For comparison, seasonal flu has a reported median R_0 of 1.28 (IQR, 1.19–1.37),
68 while measles has an R_0 of 12–18 (4).

69 **Situation in Sri Lanka**

70 The 1st case of COVID-19 was diagnosed in Sri Lanka on 27 January 2020, in a tourist from
71 China. The 2nd case was detected nearly 6 weeks later, on 11 March, in a tour guide who
72 probably contracted the infection from Italian tourists. Since then, the spread of infection has
73 been relatively slow, and mostly confined to returnees from countries with high transmission,
74 and their contacts. However, it must be noted that in four of the 190 cases diagnosed in the
75 30 days from 11 March to 10 April 2020, it was not possible to identify the source of
76 infection. It took nearly a week for the caseload to double from 50 (on 19 March) to 100 on

77 (25 March). It had not yet doubled again as of 11th April, when the count was 197 cases, with
78 54 recovered and 7 deaths (5). The epidemic has not yet reached the stage of full-blown
79 community transmission, and almost all cases still occur in clusters where the chain of
80 transmission can be traced.

81

82 The government of Sri Lanka acted swiftly to contain transmission, with very stringent public
83 health measures and social distancing: complete island-wide lockdown, contact tracing and
84 isolation, and quarantine of all inbound passengers were all adopted almost simultaneously.

85 The airport has been closed for inbound passengers since 19 March. The national policy with
86 regard to testing was that all symptomatic individuals clinically suspected of infection with
87 SARS-CoV-2, should be tested in one of seven designated laboratories, using PCR as a
88 diagnostic tool. All positive individuals (regardless of severity of illness) are managed in one
89 of three state hospitals, designated for management of COVID-19. These hospitals are also
90 equipped with intensive care units and ventilators for management of the critically ill.

91 However, the control measures have exacted a very heavy social and economic cost, and the
92 state is now about to implement a phased relaxation of preventive measures. For economic
93 and social reasons, the government will be forced to re-open Sri Lanka's borders in the near
94 future, while the pandemic is still going on elsewhere.

95 **Potential impact of COVID-19**

96 It has been suggested that most people infected with SAR-CoV-2 show no symptoms but are
97 still able to infect others. Blanket testing of an isolated village of about 3000 individuals in
98 northern Italy found that 50 – 75% of infected individuals were asymptomatic (6). Analysis
99 of the outbreak in China found that 81% of symptomatic individuals had mild illness,
100 whereas 14% developed severe illness (i.e., dyspnea, respiratory frequency $\geq 30/\text{min}$, blood

101 oxygen saturation $\leq 93\%$, partial pressure of arterial oxygen to fraction of inspired oxygen
102 ratio < 300 , and/or lung infiltrates $> 50\%$ within 24 to 48 h) and another 5% became critically
103 ill with respiratory failure, septic shock, and/or multiple organ dysfunction or failure (7). It is
104 the provision of effective care for this last group of patients, who may require ventilation for
105 2 – 3 weeks, that is the crucial limiting factor in any health system.

106

107 The global numbers as of 10 April were 1,617,204 cases, 364,686 recovered, and 97,039
108 deaths, which suggests a case fatality rate of 5.5% (8). Of the first 140 patients treated for
109 COVID-19 at the Infectious Disease Hospital in Sri Lanka, where the majority of patients
110 have been managed, nine (6.4%) have required intensive care; a similar proportion to that
111 reported from Wuhan. Sri Lanka's case fatality rate has been 3.7% (7/197) as of 11 April
112 2020.

113

114 If the spread of infection is not controlled, the R_0 of SARS-CoV-2 is such that it sweeps
115 swiftly through the susceptible population, resulting in a large number of very ill persons
116 within a short period of time, thus overloading the health system and causing it to collapse.
117 However, it is clearly possible to slow down transmission, as has been demonstrated in Sri
118 Lanka. The availability of beds and ventilators in hospital intensive care units (ICU), to care
119 for critically ill patients, is the major constraining factor that has been observed in all
120 countries with large epidemics. Sri Lanka will need to closely monitor and control the rate of
121 spread of infection so that the requirement for ICU beds and ventilators remains within the
122 available capacity.

123

124 This paper proposes a theoretical basis for estimating the limit within which the reproduction
125 number should be constrained, in order to ensure that the infection spreads slowly, and the
126 COVID-19 case load remains within the capacity of Sri Lanka's health system.

127

128 **Middle section**

129 **Materials and Methods**

130 We used publicly available data for the analysis. The 3-day moving average of cases
131 diagnosed each day during the period 11 March to 15 April were plotted (see Figure 1).
132 These numbers are based on a policy of screening all symptomatic individuals clinically
133 suspected of infection with SARS-CoV-2, using PCR as a diagnostic tool, as recorded in the
134 daily situation reports released by the Epidemiology Unit of the Ministry of Health. It should
135 be noted that an exception to this policy was made on 31 March, when screening was
136 extended to contacts, and 10 of the 21 cases reported on 1 April were asymptomatic positives.
137 Using the maximum likelihood method in the R0 package in R programming language (9),
138 we calculated R over the first 35 days.

139

140 We used the Susceptible, Infected, Recovered (SIR) model to explore the number of new
141 infections and estimated ICU bed requirements at different levels of R_0 values after lockout.
142 These R_0 values were selected to represent the range within which transmission may be
143 constrained, and assuming that it will increase after lockout (Table 1).

144

145

146 Table 1: R_0 values and doubling time of infections

R_0	Doubling time of active infections
1.3	32 days
1.4	24 days
1.5	19 days
1.6	16 days
1.7	14 days
1.8	12 days

147

148

149 We developed a web-based interactive application using an R Shiny package (available
150 through this link: bit.ly/COVID19_ICU) that enables visualization of cases and ICU bed
151 requirements with time under different values of R , with the following adjustable parameters
152 that include the total population exposed; the proportion of asymptomatic individuals among
153 those infected; the number of active cases; the number of recovered cases; the infectious
154 period in days; R_0 or doubling time in days; the percentage who are expected to become
155 critically ill; the available number of ICU beds; the average duration of ICU stay in days; and
156 uncertainty of projection.

157

158 We calculated the scenarios that emerge at different values of R , in terms of active infections
159 and ICU requirements subsequent to lockout on 20 April 2020, under the following
160 assumptions:


- 161 1. the entire population of Sri Lanka (22 million) is susceptible to infection,
- 162 2. there are 156 active cases, and 91 recovered (as recorded by the Epidemiology Unit of
- 163 the Ministry of Health on 19 April),
- 164 3. 50% of all infections are asymptomatic or pre-symptomatic and therefore
- 165 undiagnosed,
- 166 4. the average infectious period is 14 days,
- 167 5. 5% of symptomatic patients will require ICU care,
- 168 6. the average duration of ICU stay is 2 weeks,
- 169 7. maximum critical care capacity = 300 ICU beds and ventilators

170 At present, the state hospitals in Sri Lanka have a total of about 670 functional ICU beds with
171 ventilators. While retaining capacity for management of patients with other illnesses, we
172 assumed that up to 300 of these ICU beds may be made available for management of
173 COVID-19 patients at the peak of the epidemic.

174


175 Results

176 The three-day moving average of daily new cases over the first month (Figure 1) is
177 suggestive of two waves of transmission, and so we calculated R separately for these two
178 periods. The first wave, from Day 0 to Day 17 was largely due to infections among foreign
179 returnees ($R_0 = 3.32$ [95%CI: 1.85 - 5.41]). The second wave was largely due to local
180 transmission among their contacts ($R_2 = 1.25$ [95%CI: 0.93 - 1.63]).



181
182 Fig 1. Three-day moving average of new cases, 11th March to 15th April. Red dashed line
183 indicate day 17.

184
185 Figure 2 shows the possible course of the epidemic if transmission remained at the initial
186 level seen during the first wave of transmission ($R=3.32$). This model suggests that the
187 epidemic would have peaked in about 3 months, with more than 5,000,000 affected
188 individuals at the peak of the epidemic.




189

190

Fig 2. Natural progression of COVID-19 epidemic when $R=3.32$.

191

192 Figure 3 shows the interface of the web-based application. This web-based application plots
193 the expected epidemic curve under the user input parameters and provides projections on
194 expected new infections on day 7, 14, 21 and 30, new infections by day 7, 14, 21 and 30, day
195 of the peak epidemic, infected patients at peak, critically ill patients at peak, required ICU
196 bed days at peak and the day of the ICU saturation under each scenario.



197

198

Fig 3. Interface of the web-based application (URL: bit.ly/COVID19_ICU)


199

200

201 Figure 4 shows how the spread of infection could progress at each of the selected levels of

202 R_0 . It can be seen that as the value of R_0 decreases, the curve becomes flatter: the peak arrives

203 progressively later, and affects a smaller number of individuals at any one time.



204


205

Fig 4. The epidemic curve over time at selected values of R

206

207 Figure 5 shows how saturation of ICU bed capacity (300 beds) could be delayed, as the value
208 of R becomes lower. The curves suggest that while saturation of ICU bed capacity would not
209 occur until about 6 months have elapsed at the lowest value of R selected (R=1.3), this would
210 happen in about 2 months if R=1.8, the highest value selected.

211



212

213

214 Fig 5. Saturation of ICU bed capacity: changes with time at selected values of R

215

216 We then estimate active infections predicted on Days 7, 14, 21 and 30 after lockout and the
217 day of ICU saturation (300 beds) at different values of R (see Table 2), and the predicted new
218 infections over this same period (see Table 3). This suggests that a R value of 1.5 or above
219 would result in saturation of ICU capacity within about 3 months of lockout, and this would
220 be likely if the number of active infections reaches 515 (95% CI 414 - 615) and
221 approximately 18 (95% CI 14 - 22) new infections on Day 14 after lockout. Based on our
222 assumption that 50% of infections are asymptomatic, this means the number of active
223 symptomatic cases on Day 14 after lockout would have increased to about 255 and the
224 number of new symptomatic cases would be about 9.

225 Table 2. Predicted active infections and ICU bed saturation at selected values of R after
 226 lockout

R value (Day of the expected peak)	Active infections after 7 days (95% CI)	Active infections after 14 days (95% CI)	Active infections after 21 days (95% CI)	Active infections after 30 days (95% CI)	ICU saturation on day (95% CI)
1.3 (Day 426)	363 (291-434)	421 (339-504)	489 (394-585)	594 (477-710)	171 (163-181)
1.4 (Day 337)	381 (306-456)	466 (374-557)	569 (457-680)	736 (591-880)	128 (122-136)
1.5 (Day 280)	401 (322-479)	515 (414-615)	661 (531-790)	911 (733-1090)	103 (98-109)
1.6 (Day 240)	421 (339-504)	569 (457-680)	768 (617-918)	1129 (908-1351)	86 (81-91)
1.7 (Day 210)	443 (356-530)	629 (505-752)	892 (717-1067)	1399 (1125-1674)	74 (70-78)
1.8 (Day 187)	466 (374-557)	695 (559-831)	1037 (833-1240)	1734 (1394-2074)	64 (61-68)

227

228

229 Table 3. Expected new infections on day 7, 14, 21 and 30 at selected values of R

R value (Day of the expected peak)	New infections on day 7 (95% CI)	New infections on day 14 (95% CI)	New infections on day 21 (95% CI)	New infections on day 30 (95% CI)
1.3 (Day 426)	8 (6-9)	9 (7-11)	10 (8-12)	13 (10-15)
1.4 (Day 337)	11 (9-13)	13 (10-16)	16 (13-19)	21 (17-25)
1.5 (Day 280)	14 (11-17)	18 (14-22)	23 (19-28)	32 (26-38)
1.6 (Day 240)	18 (14-21)	24 (19-29)	32 (26-38)	47 (38-57)
1.7 (Day 210)	22 (17-26)	31 (25-37)	44 (35-52)	68 (55-82)
1.8 (Day 187)	26 (21-31)	39 (31-46)	58 (46-69)	96 (78-115)

230

231

232 Discussion

233 Our findings suggest that the multiple control measures adopted in Sri Lanka during March
234 2020, which includes prompt contact tracing and isolation, border closure and complete
235 lockdown, have enabled reduction in transmission from an initial level ($R=3.0$) that would
236 have almost certainly overwhelmed Sri Lanka's health system within a month, peaking in
237 about 3 months, with well over 5 million active infections at that point.

238 The simple SIR model we developed enables visualization of how different levels of control
239 would affect the speed at which ICU capacity in our country reaches saturation and the
240 number of cases that would signal the likelihood of this occurring in 2-3 months. Our
241 projections suggest that transmission should be controlled so that R does not exceed 1.5 for
242 any prolonged length of time, in order to avoid overloading the ICU capacity. The model can
243 also be used to envisage the impact of varying levels of control in different areas within Sri
244 Lanka, such as in the 6 districts in Sri Lanka categorized as having a high risk of transmission
245 compared to the other 19 districts which have a lower risk. This could inform healthcare
246 decision making at a more local level.

247 It may be argued that the SIR model is not applicable in the Sri Lankan context, because there
248 is, as yet, no full-blown community transmission. However, it is likely that the COVID-19
249 epidemic in Sri Lanka will move into this phase, as has happened in many other countries
250 over the past three months, and the SIR model is widely accepted as a means of
251 conceptualizing the spread of an infectious disease through a population over time (10).

252 The validity of the projections derived from our model depend a great deal on the accuracy of
253 the assumed parameters, such as the proportion of asymptomatic individuals, the average
254 period of infectiousness, the proportion of symptomatic individuals who require ICU care, the
255 duration of ICU stay, etc. The estimates presented here are based on data reported from other
256 countries where the epidemic is more advanced, and may not necessarily be appropriate in

257 the Sri Lankan context. However, the availability of the app enables the user to change the
258 parameters as required as more data becomes available.

259 Other web-based applications have been developed, such as the Epidemic Calculator
260 available at <https://gabgoh.github.io/COVID/index.html>. This application uses a SEIR
261 (Susceptible, Exposed, Infectious, Removed) model, and although it does not enable
262 calculation of the saturation of ICU bed capacity, the results produced by our model in terms
263 of active infections, susceptible individuals and recovered patients are on par with the
264 Epidemic Calculator under no intervention scenario. We chose not to use a SEIR model
265 because the data available at this stage in Sri Lanka was insufficient to estimate all the
266 parameters necessary for such a model.

267 Another web-based app developed at the London School of Hygiene & Tropical Medicine is
268 available at https://cmmid-lshtm.shinyapps.io/hospital_bed_occupancy_projections/, to
269 estimate projected hospital bed occupancy in the UK. However, this app can be used to
270 forecast COVID-19 bed requirements in a given location for only up to 21 days (e.g. a
271 healthcare facility, a county, a state) and our estimates were similar to this app.

272 We believe that the model and web based app, which we developed primarily for use in Sri
273 Lanka, may also be appropriate for use in other low and middle income countries that have
274 similar constraints for ICU care of COVID-19 patients, but are unable to enforce stringent
275 lockdown measures for a prolonged period of time due to social and economic reasons.

276

277 **Ending section**

278 **Acknowledgements:**

279 We thank Prof Deirdre Hollingsworth, Prof Don Bundy, Prof Rajitha Wickremasinghe and
280 Dr Sudath Samaraweera for helpful guidance and comments on the draft manuscript and Dr
281 Prasad Ranatunga for helpful comments on the web-based application.

282

283 **References**

- 284 1. COVID-19 Basics [Internet]. Johns Hopkins University & Medicine. 2020 [cited 2020
285 Apr 1]. Available from: <https://coronavirus.jhu.edu/#covid-19-basics>
- 286 2. van Doremalen N, Bushmaker T, Morris DH, Holbrook MG, Gamble A, Williamson
287 BN, et al. Aerosol and Surface Stability of SARS-CoV-2 as Compared with SARS-
288 CoV-1. *N Engl J Med*. 2020;
- 289 3. Liu Y, Gayle AA, Wilder-Smith A, Rocklöv J. The reproductive number of COVID-19
290 is higher compared to SARS coronavirus. *Journal of travel medicine*. 2020.
- 291 4. Lake MA. What we know so far: COVID-19 current clinical knowledge and research.
292 *Clinical medicine (London, England)*. 2020.
- 293 5. Epidemiology Unit Ministry of Health Sri Lanka [Internet]. Epidemiology Unit
294 Ministry of Health Sri Lanka. [cited 2020 Apr 15]. Available from:
295 <http://www.epid.gov.lk/web/index.php?lang=en>
- 296 6. Day M. Covid-19: identifying and isolating asymptomatic people helped eliminate
297 virus in Italian village. *BMJ*. 2020;
- 298 7. Wu Z, McGoogan JM. Characteristics of and Important Lessons From the Coronavirus
299 Disease 2019 (COVID-19) Outbreak in China. *JAMA*. 2020;

- 300 8. COVID-19 Dashboard [Internet]. Johns Hopkins University & Medicine, Coronavirus
301 Resource Centre. 2020 [cited 2020 Apr 6]. Available from:
302 <https://coronavirus.jhu.edu/map.html>
- 303 9. Obadia T, Haneef R, Boëlle PY. The R0 package: A toolbox to estimate reproduction
304 numbers for epidemic outbreaks. *BMC Med Inform Decis Mak*. 2012;
- 305 10. White PJ. *Mathematical Models in Infectious Disease Epidemiology*. In: *Infectious*
306 *Diseases* [Internet]. 4th ed. Elsevier; 2017. p. 49-53.e1. Available from:
307 <https://linkinghub.elsevier.com/retrieve/pii/B9780702062858000058>
308
309

310

311 **Conflict of Interest Statement**

312 All authors have completed the ICMJE uniform disclosure form at
313 www.icmje.org/coi_disclosure.pdf and declare: no support from any organization for the
314 submitted work; no financial relationships with any organizations that might have an interest
315 in the submitted work in the previous three years; no other relationships or activities that
316 could appear to have influenced the submitted work.

317

318

319