

A vulnerability-based approach to human-mobility reduction for countering COVID-19 transmission in London while considering local air quality

Manu Sasidharan^{1*}, Ajit Singh^{2*}, Mehran Eskandari Torbaghan^{3*}, Ajith Kumar Parlikad¹

¹Department of Engineering, University of Cambridge, CB2 1PZ, United Kingdom

5 ²School of Geography, Earth and Environmental Sciences, University of Birmingham, Edgbaston, Birmingham, B15 2TT, United Kingdom.

³Department of Civil Engineering, School of Engineering, University of Birmingham, Edgbaston, Birmingham, B15 2TT, United Kingdom.

10 *Correspondence to: mp979@cam.ac.uk (M.S.); a.singh.2@bham.ac.uk (A.S.); m.eskandaritorbaghan@bham.ac.uk (M.E.T.)

Abstract:

15 An ecologic analysis was conducted to explore the correlation between air pollution, and COVID-19 cases and fatality rates in London. The analysis demonstrated a strong correlation ($R^2 > 0.7$) between increment in air pollution and an increase in the risk of COVID-19 transmission within London boroughs. Particularly, strong correlations ($R^2 > 0.72$) between the risk of COVID-19 fatality and NO₂ and PM_{2.5} pollution concentrations were also found. Although this study assumed the same level of air pollution across a particular London borough, it demonstrates the possibility to employ air pollution as an indicator to rapidly identify the city's vulnerable regions. Such an approach can inform the decisions to suspend or reduce the operation of different public transport modes within a city. The methodology and learnings from the study can thus aid in public transport's response to COVID-19 outbreak by adopting different levels of human-mobility reduction strategies based on the vulnerability of a given region.

20 **One Sentence Summary:** This study introduces air pollution levels as an indicator for a region's vulnerability to COVID-19 and suggests human-mobility reduction measures.

25 Introduction:

The current outbreak of novel coronavirus COVID-19 or severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), has resulted in the World Health Organization (WHO) declaring it as a global pandemic [1]. Reported first within the city of Wuhan, Hubei Province of China in December 2019, the COVID-19 exhibits high human-to-human transmissibility and has spread rapidly across the world [2]. The human-to-human transmission of COVID-19 can occur from individuals in the incubation stage or showing symptoms, and also from asymptomatic individuals who remain contagious [3]. The COVID-19 has been reported to transmit via the inhalation of exhaled respiratory droplets [4] that remain airborne for up to 3 hours [5]. The long-term exposure to air pollution has been reported to increase the risk of experiencing severe COVID-19 outcomes [6]. The extent to which COVID-19 induces respiratory stress in infected individuals may also be influenced by underlying respiratory conditions [7] like acute respiratory inflammation, asthma and cardiorespiratory diseases [8]. The simultaneous exposure to air pollutants such as particulate matter (PM) and Nitrogen dioxide (NO₂) alongside COVID-19 virus is expected to exacerbate the level of COVID-19 infection and risk of fatality [9, 10]. Moreover, the adsorption of the COVID-19 virus on PM could also contribute to long-range transmission of the virus [4]. For example, the

2003 severe acute respiratory syndrome coronavirus 1 (SARS-CoV-1) infected patients who lived in moderate air pollution levels were reported to be 84% more likely to die than those in regions with lower air pollution [11]. The aerosol and surface stability of the COVID-19 or SARS-CoV-2 is reported to be similar to that of SARS-CoV-1 [5].

Given the limited understanding of the epidemiology of COVID-19, social-distancing and human-mobility reduction measures can contribute greatly to tailoring public health interventions [12]. Consequently, countries across the world have enforced lockdowns and other coordinated efforts to reduce human-mobility [13, 14, 15, 16]. The UK's national framework for responding to a pandemic states that public transport should continue to operate normally during a pandemic, but users should adopt good hygiene measures, and stagger journeys where possible [17]. Within the UK, London has recorded the highest COVID-19 related fatalities (i.e. 30.2% of UK's deaths as of 31 March 2020) [18]. On 18 March 2020, further to the UK government's advice, Transport for London (TfL) closed 40 out of 270 London Underground (LU) stations that do not serve as interchanges with other lines and announced a reduced service across its network [19]. This is also because 30% of TfL's drivers, station staff, controllers and maintenance teams were not able to come to work, including those self-isolating or ill with COVID-19 [20]. The UK's current human-mobility reduction response reflects the need to maintain business continuity, near normal functioning of society and enable critical workers to make essential journeys [17, 21]. However, a statistically significant association exists between human-mobility through public transport and transmissions of acute respiratory infections (ARI) [21, 22]. It was found that using public transport in the UK during a pandemic outbreak has an approximately six-fold increased risk of contracting an ARI [21]. Moreover, boroughs with access to LU interchange stations have historically higher pandemic case rates [22], as users are exposed to higher number of individuals in comparison to through stations. The methodology and results from this study can be employed to rapidly identify local authorities/regions that are highly vulnerable to COVID-19 and accordingly inform human-mobility reduction measures across the city's public transport network.

Results

As the COVID-19 is an evolving pandemic, the available data at the time of writing this paper on 31 March 2020 on COVID-19 morbidity and mortality for different boroughs in London was collected from [23, 18]. The fatality rate across each London boroughs was estimated by dividing the number of reported deaths by the number of reported positive COVID-19 cases. The average air pollution data associated with NO₂ and PM_{2.5} concentration was collected for London boroughs from [24]. NO₂ data was available for 15 boroughs namely Barking and Dagenham, Bexley, Wandsworth, City of London, Croydon, Greenwich, Havering, Hillingdon, Kensington and Chelsea, Lewisham, Reading, Redbridge, Sutton, Tower Hamlets and Westminster. While, the PM_{2.5} data was available only for 8 boroughs (Barking and Dagenham, Wandsworth, City of London, Croydon, Hillingdon, Kensington and Chelsea, Lewisham). An ecologic analysis was conducted to explore the correlation between short-term air pollution and COVID-19 cases and fatality rates. A linear regression model was fitted to the data for the boroughs with more than 100 reported cases and 10 deaths related to COVID-19 as of 31 March 2020. A strong correlation between short-term NO₂ and PM_{2.5} pollution concentrations and COVID-19 cases with R² values of 0.82 ($COVID-19\ cases = -29.345 + 10.306 * NO_2\ concentration$) and 0.72 ($COVID-19\ cases = -215.63 + 40.997 * PM_{2.5}\ level$) were observed respectively (see Figure 1). In particular, COVID-19 fatality rate increased with increase in short-term air pollution, where a significant correlation between COVID-19 fatality and NO₂ and PM_{2.5} pollution concentrations with R² of 0.90 (*fatality*

$rate = 1.864 + 0.5787 * NO_2 \text{ level}$) and 0.67 ($fatality \ rate = -7.733 + 2.3399 * PM_{2.5} \text{ level}$) were found (see Figure 2).

As per TfL's guidance to LU users [25], 40 LU stations across its network were closed as part of the UK Government's response to COVID-19. The median $PM_{2.5}$ levels recorded by [26] for 27 of these 40 closed stations range from $0-50 \mu\text{g m}^{-3}$ (5 stations), $50-100 \mu\text{g m}^{-3}$ (9 stations), $100-200 \mu\text{g m}^{-3}$ (5 stations), $200-300 \mu\text{g m}^{-3}$ (6 stations) and greater than $300 \mu\text{g m}^{-3}$ (2 stations) (see Table S1). Of the 230 operating stations, the median $PM_{2.5}$ levels recorded for 219 stations range from $0-50 \mu\text{g m}^{-3}$ (56 stations), $50-100 \mu\text{g m}^{-3}$ (15 stations), $100-200 \mu\text{g m}^{-3}$ (15 stations), $200-300 \mu\text{g m}^{-3}$ (18 stations) and greater than $300 \mu\text{g m}^{-3}$ (7 stations) [26] (see Table S1). This suggests that approximately 40% of the stations in operation during the current COVID-19 outbreak in London are up to 26 times more polluted than the ambient background locations and the roadside environment (that have median $PM_{2.5}$ level of $14 \mu\text{g m}^{-3}$ [26]). Moreover, the average NO_2 concentrations within the LU network was reported to be $51 \mu\text{g m}^{-3}$ [27]; which is 27.5% higher than the NO_2 limit values for the protection of human health [28].

Discussion

Various studies have reported an association between air pollution levels and excess morbidity and mortality from respiratory diseases [29, 30]. Both long-term and short-term exposure to air pollution has been associated with a variety of respiratory conditions [11] with children and elderly people being at most risk [31]. Our analysis shows that short-term exposure to air pollution (both NO_2 and $PM_{2.5}$) is significantly correlated with an increased risk of contracting and dying from COVID-19. It was reported by [6] that long-term exposure to fine particulate matter ($PM_{2.5}$) increases the risk of COVID-19 deaths. Biologically, either long-term or short-term exposure to air pollutants such as $PM_{2.5}$ and NO_2 can compromise lung function and therefore increasing the risk of death from COVID-19 [7]. Also, the median level of airborne $PM_{2.5}$ in LU during summer is often several times higher than other transport environments such as cycling ($35 \mu\text{g m}^{-3}$), bus ($30.9 \mu\text{g m}^{-3}$), cars ($23.7 \mu\text{g m}^{-3}$) [32, 26]. The greatest $PM_{2.5}$ concentrations across the LU network was reported to be on the Victoria Line (around 16 times higher than roadside environment), followed by Northern, Bakerloo and Piccadilly line [26]. The routine cleaning and maintenance of LU ranges from litter removal to preventing safety incidents rather than on reducing PM concentrations. Our study is in agreement with [22] who reported that there are higher pandemic case rates for London boroughs with access to interchange stations, as individuals would interact with more people in comparison to through stations.

An analysis by [33] of the UK's National Health Service (NHS) records have reported that 20% of England's population are at risk of mortality from COVID-19 due to underlying conditions and age. We support the UK government's existing COVID-19 guidance [34] to exercise good hygiene and to avoid unnecessary travel. While considering the evidence that COVID-19 can be transmitted from an asymptomatic individual [3], we do not support the current countermeasure of suspending LU operations on the stations that do not serve as interchanges. This is because of (i) the statistically significant risk of contracting ARI's on UK's public transport and higher pandemic case rates within London boroughs that have comparatively easier access to interchange stations [21, 22] and, (ii) relatively higher air pollution levels in LU stations than ambient background locations or road side environment. Since isolating towns or even cities is not yet part of the UK government's action plan [34], we recommend a vulnerability-based assessment of different boroughs in London and to accordingly suspend or reduce operations on LU stations within highly vulnerable regions. For instance, London Borough of Kensington and Chelsea is seen to be highly

vulnerable to COVID-19 fatality from our analysis (see Figure 2a). Table S1 shows that, all the through stations and 3 out of 4 interchange stations (South Kensington, Sloane square, Earl's court, Notting Hill gate) in this borough are currently operational. Such a vulnerability-based assessment might aid decision-makers in selecting appropriate human-mobility reduction measures to COVID-19 in London's different local authorities/boroughs (such as apportion of transport staff across railway stations, arranging dedicated shuttling services for key workers, scheduling bus operations etc.) while adhering to the UK's national framework for response to pandemic outbreaks [17].

Given that the immunity to the 2003 SARS-CoV-1 was reported to be relatively short-lived (around 2 years) [35], achieving herd immunity for such diseases would be unlikely without overwhelming the healthcare system [16]. Moving forward, human-mobility reduction measures provides the greatest benefit to COVID-19 mitigation [15, 14] as prevention is potentially cost-effective than cure [22] or death. One of the most controversial debate in pandemic countermeasures is the potential benefit of human-mobility reduction and social-distancing attained by closure of public transport systems [21]. From a public policy perspective, there is a need to achieve a trade-off between the potential public health benefits of closing public transport during a pandemic thereby delaying the community spread, against the socio-economic impacts of curtailing/reducing human mobility. Determining the vulnerability of regions/locations to COVID-19 might be helpful in achieving such trade-offs. To this end, this study demonstrates that the air pollution levels can serve as one of the indicators to assess a region's vulnerability to COVID-19. It has to be noted that the number of positive COVID-19 cases considered within this study are only those reported at the hospitals and does not include the growing number of people who are self-isolating due to COVID-19. While the individual risk of contracting and dying from COVID-19 is dependent on various factors (including age, underlying conditions, availability of health care, population density etc.), these results are informative for both scientists and decision-makers in their efforts to reduce the transmission and impact of the ongoing COVID-19 outbreak through appropriate human-mobility reduction strategies.

References and Notes:

- [1] World Health Organization, "WHO Director-General's opening remarks at the media briefing on COVID-19 - 11 March 2020," 11 March 2020. [Online]. Available: <https://www.who.int/dg/speeches/detail/who-director-general-s-opening-remarks-at-the-media-briefing-on-covid-19---11-march-2020>. [Accessed 31 March 2020].
- [2] L. Qun, G. Xuhua, W. Peng, W. Xiaoye, Z. Lei, T. Yeqing, R. Ruiqi, L. Kathy S.M, L. Eric, W. Jessica Y., X. Xuesen, X. Nijuan, W. Yang, L. Chao, C. Qi, L. Dan, L. Tian, Z. Jing, L. Man, T. Wenxiao and C. Chuding, "Early Transmission Dynamics in Wuhan, China, of Novel Coronavirus-Infected Pneumonia," *The New England Journal of Medicine*, vol. 382, pp. 1199-1207, 2020.
- [3] Y. Bai, L. Yao, T. Wei, F. Tian, D. Y. Jin, L. Chen and M. Wang, "Presumed asymptomatic carrier transmission of COVID-19.," *JAMA*, 10.1001/jama.2020.2565, 2020.
- [4] Q. Guangbo, L. Xiangdong, H. Ligang and J. Guibin, "An Imperative Need for Research on the Role of Environmental Factors in Transmission of Novel Coronavirus (COVID-19)," *Environmental Science & Technology*, vol. acs.et.0c01102, 23 March 2020.
- [5] v. D. Neeltje, B. Trenton and H. M. Dylan, "Aerosol and Surface Stability of SARS-CoV-2 as Compared with SARS-CoV-1," *The New England Journal of Medicine*, DOI: 10.1056/NEJMc2004973, 2020.
- [6] W. Xiao, N. Rachel C, S. Benjamin M, B. Danielle and D. Francesca, "Exposure to air pollution and COVID-19 mortality in the United States" doi.org/10.1101/2020.04.05.20054502, 2020.

- [7] L. Wei, T. Zhao-Wu, W. Lei, Y. Ming-Li, L. Kui, Z. Ling, W. Shuang, D. Yan, L. Jing, L. Hui-Guo, Y. Ming and H. Yi, "Analysis of factors associated with disease outcomes in hospitalized patients with 2019 novel coronavirus disease," *Chinese Medical Journal*, no. 10.1097/CM9.0000000000000775, 2020.
- [8] Centers for Disease Control and Prevention, "People Who Are at Higher Risk for Severe Illness," 2020. [Online]. Available: https://www.cdc.gov/coronavirus/2019-ncov/need-extra-precautions/people-at-higher-risk.html?CDC_AA_refVal=https%3A%2F%2Fwww.cdc.gov%2Fcoronavirus%2F2019-ncov%2Fspecific-groups%2Fhigh-risk-complications.html. [Accessed 31 March 2020].
- [9] Transport & Environment, "Air pollution probably increasing coronavirus death rates - experts," 19 March 2020. [Online]. Available: <https://www.transportenvironment.org/news/air-pollution-probably-increasing-coronavirus-death-rates-experts>. [Accessed 31 March 2020].
- [10] European Public Health Alliance, "Coronavirus threat greater for polluted cities," 16 March 2020. [Online]. Available: <https://epha.org/coronavirus-threat-greater-for-polluted-cities/>. [Accessed 31 March 2020].
- [11] C. Yan, Z. Zuo-Feng, F. John, Z. Jinkou, W. Hua, Y. Shun-Zhang and D. Roger, "Air pollution and case fatality of SARS in the People's Republic of China: an ecologic study," *Environmental Health: A Global Access Science Source*, vol. 2, no. 15, pp. 1-5, 2003.
- [12] L. Shengjie, B. Isaac, R. Nick, W. Alexander, L. Xin, Y. Weizhong, Y. Hongjie, K. Kamran and T. Andrew J, "Assessing spread risk of Wuhan novel coronavirus within and beyond China, January-April 2020: a travel network-based modelling study," *Health Sciences*, 10.1101/2020.02.04.20020479, 2020.
- [13] European Commission, "Coronavirus response: Travel and transportation," 26 March 2020. [Online]. Available: https://ec.europa.eu/info/live-work-travel-eu/health/coronavirus-response/travel-and-transportation_en. [Accessed 31 March 2020].
- [14] A. Roy M, H. H, K. D and H. T Deirdre, "How will country-based mitigation measures influence the course of the COVID-19 epidemic?," *The Lancet*, vol. 395, no. 10028, pp. 931-934, 2020.
- [15] C. Matteo, T. D. Jessica, A. Marco, G. Corrado, L. Maria, M. Stefano, P. Ana Pastore y, M. Kunpeng, R. Luca, S. Kaiyuan, V. Cécile, X. Xinyue, Y. Hongjie, H. M. Elizabeth and M. L. Ira, "The effect of travel restrictions on the spread of the 2019 novel coronavirus (COVID-19) outbreak," *Science*, 10.1126/science.aba9757, 2020.
- [16] D. B. Edward, R. Daniele and M. Yves, "Modeling the COVID-19 outbreaks and the effectiveness of the containment measures adopted across countries," *medRxiv*, doi.org/10.1101/2020.04.02.20046375, 2020.
- [17] Department of Health, "Pandemic Flu – A national framework for responding to an influenza pandemic," Department of Health, London, 2007.
- [18] National Health Services, "COVID-19 Daily Deaths," 2020. [Online]. Available: <https://www.england.nhs.uk/statistics/statistical-work-areas/covid-19-daily-deaths/>. [Accessed 31 March 2020].
- [19] Transport for London, "Planned services to support London's critical workers," 18 March 2020. [Online]. Available: <https://tfl.gov.uk/info-for/media/press-releases/2020/march/planned-services-to-support-london-s-critical-workers>. [Accessed 31 March 2020].
- [20] Transport for London, "London's Transport Commissioner Mike Brown on our response to coronavirus," 26 March 2020. [Online]. Available: <https://tfl.gov.uk/info-for/media/press-releases/2020/march/london-s-transport-commissioner-mike-brown-on-our-response-to-coronavirus>. [Accessed 31 March 2020].
- [21] T. Joy, M. Puja, G. Jack, H. Ahmed, E. Joanne, K. Susan, P. Christopher, A. Shahid, H. Andrew and V.-T. Jonathan Nguyen, "Is public transport a risk factor for acute respiratory infection?," *BMC Infectious Diseases*, vol. 11, no. 16, 2011.
- [22] G. Lara and J. Anders, "Analysing the link between public transport use and airborne transmission: mobility and contagion in the London underground," *Environmental Health*, vol. 17, no. 84, pp. 1-11, 2018.
- [23] "Public Health England," March 2020. [Online]. Available: <https://www.gov.uk/government/publications/covid-19-track-coronavirus-cases>. [Accessed 31 March 2020].
- [24] King's College London, "Data feeds for London air quality," 2020. [Online]. Available: <http://www.londonair.org.uk/LondonAir/API/>. [Accessed 31 March 2020].
- [25] Transport for London, "Coronavirus (COVID-19)," March 2020. [Online]. Available: <https://tfl.gov.uk/campaign/coronavirus-covid->. [Accessed 31 March 2020].

- [26] J. Smith, B. Barratt, G. Fuller, F. Kelly, M. Loxham, E. Nicolosi, M. Priestman, A. Tremper and D. Green, "PM2.5 on the London Underground," *Environmental International*, vol. 134, 2020.
- [27] S. James David, M. Christina, K. Nutthida, B. Ben M, W. Heather A, T. Jonathon G, A. Hugh Ross, K. Frank J and B. Sean D, "London Hybrid Exposure Model: Improving Human Exposure Estimates to NO2 and PM2.5 in an Urban Setting," *Environmental Science & Technology*, vol. 50, pp. 11760-11768, 2016.
- [28] European Environment Agency, "Nitrogen dioxide - Annual limit values for the protection of human health," 22 August 2014. [Online]. Available: <https://www.eea.europa.eu/data-and-maps/figures/nitrogen-dioxide-annual-limit-values-for-the-protection-of-human-health>. [Accessed 31 March 2020].
- [29] G. Adamkiewicz, S. Ebel, M. Syring, J. Slater, F. E. Speizer, J. Schwartz, H. Suh and D. R. Gold, "Association between air pollution exposure and exhaled nitric oxide in an elderly population," *Environmental Exposure*, vol. 59, pp. 204-209, 2004.
- [30] D. W. Dockery, "Epidemiologic evidence of cardiovascular effects of particulate air pollution," *Environmental Health Perspective*, vol. 109, no. 4, pp. 483-486, 2001.
- [31] Department for Environment, Food & Rural Affairs, "Air Quality: a briefing for directors of public health," DEFRA, London, 2017.
- [32] M. Vania, M. Teresa, M. Maria Cruz, A. Fulvio, d. M. Eladio, C. Marta and Q. Xavier, "Exposure to airborne particulate matter in the subway system," *Science of the Total Environment*, vol. 511, pp. 711-722, 2015.
- [33] B. Amitava, P. Laura, H. Steve, G.-I. Arturo, T. Ana, S. Laura, N. Mahdad, P. Deenan, P. Christina, W. Wai Keong, L. Claudia, W. Bryan, D. Spiros and H. Harry, "Estimating excess 1- year mortality from COVID-19 according to underlying conditions and age in England: a rapid analysis using NHS health records in 3.8 million adults," *medRxiv*, doi.org/10.1101/2020.03.22.20040287, 2020.
- [34] Department of Health & Social Care, "Coronavirus: action plan," 2020.
- [35] W. Li-Ping, W. Nai-Chang, C. Yi-Hua, T. Xiang-Yi, N. Dan-Yu, Z. Li-Yuan, Z. Lei, L. Tao, W. Lin-Fa and L. Guo-Dong, "Duration of antibody responses after severe acute respiratory syndrome," *Emerging infectious diseases*, vol. 13, no. 10, pp. 1562-1564, 2007.

Acknowledgments: We would like to thank Anne Baker from the Office for National Statistics for very kindly providing her input to the COVID-19 related fatality data. We appreciate the work of J.D. Smith, B.M. Barratt, G.W. Fuller and team that captured the levels of PM2.5 exposure in London Underground. **Funding:** MS's time on this research at the University of Cambridge was funded by the Engineering and Physical Science Research Council (EPSRC) through the grant EP/N021614/1 (CSIC Innovation and Knowledge Centre Phase 2) and Innovate UK through the grant 920035 (Centre for Smart Infrastructure and Construction). MET's time on this research at the University of Birmingham was funded by EPSRC through the grant EP/N010523/1 (Balancing the Impact of City Infrastructure Engineering on Natural Systems using Robots). **Author contributions:** The authors confirm contribution to the paper as follows: study conception and design: M Sasidharan, A. Singh, M. Eskandari Torbaghan, A. K. Parlikad; analysis and interpretation of results: M Sasidharan, A. Singh; draft manuscript preparation: M Sasidharan, A. Singh, M. Eskandari Torbaghan. All authors reviewed the results and approved the final version of the manuscript. **Competing interests:** Authors declare no competing interests; and **Data and materials availability:** The data collected for this analysis are all publicly accessible and is available in the main text or the supplementary materials

Supplementary Materials:

Materials and Methods

Figures S1-S2

Table S1




Fig. 1. Relationship between **a)** NO₂ and **b)** PM_{2.5} pollution concentrations and reported COVID-19 cases at London boroughs using data during March 2020.

5




Fig. 2. Relationship between **a)** NO₂ and **b)** PM_{2.5} pollution concentrations and COVID-19 fatality rate for each London borough. The fatality rate was calculating by dividing the number of reported deaths by the number of reported positive COVID-19 cases (associated data was available only for the London boroughs plotted in the figures).

10