Patient-derived mutations impact pathogenicity of SARS-CoV-2

Hangping Yao[1]†, Xiangyun Lu[1]†, Qiong Chen[2]†, Kaijin Xu[1], Yu Chen[1], Linfang Cheng[1],

Fumin Liu[1], Zhigang Wu[1], Haibo Wu[1], Changzhong Jin[1], Min Zheng[1]*, Nanping Wu[1]*,

Chao Jiang[2,3]*, Lanjuan Li[1]*

[1]State Key Laboratory for Diagnosis and Treatment of Infectious Diseases, National

Clinical Research Center for Infectious Diseases, First Affiliated Hospital, Zhejiang

University School of Medicine, Hangzhou, China.

[2]Life Sciences Institute, Zhejiang University, Hangzhou, China.

[3]Zhejiang Provincial Key Laboratory of Pancreatic Disease, First Affiliated Hospital,

Zhejiang University School of Medicine, Hangzhou, China.

†Equal contributions.

*Corresponding authors: M.Z. (minzheng@zju.edu.cn), N.W. (flwnp2013@163.com),

C.J. (jiang_chao@zju.edu.cn), and L.L. (ljli@zju.edu.cn).

Summary (130 words)

The sudden outbreak of the severe acute respiratory syndrome–coronavirus

(SARS-CoV-2) has spread globally with more than 1,300,000 patients diagnosed and a

death toll of 70,000. Current genomic survey data suggest that single nucleotide variants

(SNVs) are abundant. However, no mutation has been directly linked with functional

changes in viral pathogenicity. We report functional characterizations of 11

patient-derived viral isolates. We observed diverse mutations in these viral isolates,

including 6 different mutations in the spike glycoprotein (S protein), and 2 of which are

different SNVs that led to the same missense mutation. Importantly, these viral isolates

show significant variation in cytopathic effects and viral load, up to 270-fold differences,

when infecting Vero-E6 cells. Therefore, we provide direct evidence that the

SARS-CoV-2 has acquired mutations capable of substantially changing its pathogenicity.

Introduction

Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2; previously referred to as

2019-nCoV), associated with the ongoing outbreak of atypical pneumonia, has already

caused a global pandemic, despite China's extensive systematic effort to contain the

outbreak. As of April 7, 2020, SARS-CoV-2 has infected more than 1.3 million people

around the world with a death toll of 70,000. The numbers are still increasing rapidly.

The estimate of the incubation period of SARS-CoV-2 (mean, 5.1 days; range, 4.5 to 5.8

days) (Lauer et al., 2020) is in line with those of other known human coronaviruses, such

as SARS (mean, 5 days; range, 2 to 14 days) (Varia et al., 2003) and MERS (mean, 5 to 7

days; range, 2 to 14 days) (Virlogeux et al., 2016). The reproductive number of

SARS-CoV-2 is likely to be from 1.4 to 6.5, with a mean of 3.3 (Liu et al., 2020), which

is slightly higher than SARS, i.e., 2-5 (Bauch et al., 2005; Lipsitch et al., 2003) and

MERS, i.e., 2.7-3.9 (Lin et al., 2018). More than half of patients with SARS-CoV-2

showed no signs of fever before hospitalization (Guan et al., 2020). Strikingly,

Coronavirus Diease-2019 (COVID-19) can be transmitted by asymptomatic patients, who

show no fever, gastrointestinal or respiratory symptoms, and have normal chest computed

tomography (Bai et al., 2020; Hu et al., 2020), making it much more challenging to

prevent the spread of COVID-19. Moreover, SARS-CoV-2 can remain viable and

infectious in aerosols for multiple hours and up to 7 days on surfaces (van Doremalen et

al., 2020). Although multiple *in vitro* studies or clinical trials on inhibitors or drugs were

carried out, no effective cures or vaccines have been found so far (Cao et al., 2020;

Hoffmann et al., 2020; Wang et al., 2020). The World Health Organization (WHO)

declared COVID-19 a Public Health Emergency of International Concern on 30 January

2020, and raised the threat of the SARS-CoV-2 pandemic to the "very high" level on

February 28, 2020.

SARS-CoV-2 is the seventh member of enveloped RNA beta-coronavirus

(Sarbecovirus subgenus) (Zhu et al., 2020); SARS-CoV-2, SARS-CoV and MERS-CoV

can lead to devastating diseases, while HKU1, NL63, OC43 and 229E are related with

mild symptoms (Corman et al., 2018). So far, no recombination events were detected (Yu,

2020), although this could be at least partially due to the fact that most viral isolates were

sequenced with short-reads platform. The transmembrane spike (S) glycoprotein mediates

viral entry into host cells through homotrimers protruding from the viral surface. The S

protein includes two domains: S1 for binding to the host cell receptor and S2 for fusion of

the viral and cellular membranes, respectively (Tortorici and Veesler, 2019). Both

SARS-CoV-2 and SARS-CoV use the angiotensin converting enzyme 2 (ACE2) to enter

target cells (Walls et al., 2020). ACE2 is expressed in human nasal epithelial cells and

lung, spermatogonia, leydig, sertoli, gastric, duodenal, and rectal epithelial cells (Wang

and Xu, 2020; Xiao et al., 2020; Zhao et al., 2020). The receptor binding domain (RBD)

in the S protein is the most variable genomic part in the betacoronavirus group (Wu et al.,

2020; Zhou et al., 2020), and some sites of S protein might be subjected to positive

selection (Lv et al., 2020). Despite the abundant variability of SARS-CoV-2, one key

question remains as to whether these mutations have any real functional impact on the

pathogenicity of SARS-CoV-2. This is crucial in our understanding of the viral infectious

mechanisms and dictates the strategy of drug and vaccine development in preparation for

the next stage of the pandemic.

To address this, we characterized 11 SARS-CoV-2 viral isolates from patients

admitted to Zhejiang University-affiliated hospitals in Hangzhou, China, situated 757

KMs to the east of Wuhan (see Materials and Methods). Super-deep sequencing of the 11

viral isolates on the Novaseq 6000 platform identified 1-5 mutations in the coding

sequences among the viral isolates. Mixed viral populations (representing quasi-species)

were also observed. We infected Vero-E6 cells with 11 viral isolates and quantitatively

assessed their viral load at 1, 2, 4, 8, 24, and 48 hours post infection (P.I.) and their viral

cytopathic effects (CPE) at 48 and 72 hours P.I.. Our results show that the observed

mutations can have a direct impact on the viral load and CPE when infecting Vero-E6

cells, as much as 270-fold differences between the extremities. This finding suggests that

the observed mutations in our study, and possibly in the viral isolates collected around

the world, can significantly impact the pathogenicity of SARS-CoV-2.

Results

Summary of the epidemiological history of the patients

The samples of the 11 patients involved in this study were collected during the early

phase of the COVID-19 breakout in China, dates ranging from 1/22/2020 to 2/4/2020

(Table 1). 10 of the 11 patients had clear connections with Wuhan city, where the

SARS-CoV-2 was originally identified. 5 of the 11 people either worked in or traveled to

Wuhan before they were diagnosed, and another 5 had close contact with people who

lived in Wuhan, and the remaining person had contact with people who were COVID-19

victims. Notably, patients ZJU-4, -5, -9 attended the same business conference where a

few colleagues from Wuhan were present. These patients therefore constitute 1st and 2nd

generations of the viral victims based on their epidemiological history. The 11 patients

include 8 males and 3 females, with ages ranging from 4 months to 71 years old. There

are no clear criteria in selecting these patients other than the fact that they were all

admitted into Zhejiang University-affiliated hospitals in Hangzhou. All except one of the

patients had moderate or worse symptoms. 3 patients had co-morbidity conditions and

one patient needed ICU treatment. Luckily, all of the patients have recovered as of the

time of writing this article.

ID	Sex	Age	Sample	Sample collection date	Virus isolating date	Epide-miology	Viral gen	Onset sypmtoms	Severity	Blood presure	ICU	Onset date	Admission	Discharge
ZJU_1	M	30-40	Sputum	1.25	1.29	Contact with people from Wuhan	2	Fever	Moderate	0	0	1.23	1.25	2.23
ZJU_2	M	30-40	Sputum	1.26	1.29	Lived in Wuhan	1	Fever	Severe	0	0	1.23	1.24	2.23
ZJU_3	M	30-40	Sputum	1.25	1.30	Traveled to Wuhan	2	Fever	Severe	0	0	1.18	1.19	2.12
ZJU_4	M	30-40	Sputum	1.24	1.28	Conference with colleagues from Wuhan	2	Fever, fatigue	Moderate	0	0	1.17	1.21	2.9
ZJU_5	F	20-30	Sputum	1.22	1.26	Conference with colleagues from Wuhan	2	Fever	Severe	0	0	1.21	1.22	2.9
ZJU_6	M	70-80	Sputum	2.2	2.6	Contact with people who had COVID19 Wuhan.	2 or 3	Fever, fatigue	Severe	1	0	1.22	1.26	2.19
ZJU_7	F	0-1	Nasopharyngeal swab	2.3	2.8	Contact with people from Wuhan	2	Fever	Mild	0	0	1.29	1.29	2.20
ZJU_8	M	50-60	Sputum	1.26	1.30	Lived in Wuhan	1	Fever, fatigue	Critical	1	1	1.17	1.22	2.19
ZJU_9	M	30-40	Stool	1.28	2.7	Conference with colleagues from Wuhan	2	Fever	Severe	0	0	1.18	1.21	2.5
ZJU_10	F	30-40	Stool	2.3	2.8	Lived in Wuhan	1	Fever	Severe	0	0	1.19	1.27	2.12
ZJU_11	M	60-70	Stool	2.4	2.9	Lived in Wuhan	1	Coughing	Severe	1	0	1.19	1.26	3.15

Table 1. A summary of the epidemiological information of the 11 patients involved in

this study. The "Viral gen" (viral generation) was inferred based on their exposure

history.

Ultra-deep sequencing reveals diverse mutations in the patient-derived viral isolates

To assess the mutational spectrum of these 11 viral isolates, ultra-deep sequencing of the

isolated viral genomic RNA was performed on the Illumina Novaseq 6000 platform,

generating on average 245 million post-cleaning reads/67.16 Gb per sample (Table S1;

average coverage exceeding 2,000,000 X). This extraordinary depth is partially due to the

small genome of the SARS-CoV-2, which enables us to identify mutations with high

confidence. Moreover, in cases where the viral populations are not homogenous, the

depth could help us to characterize alleles with very low frequency.

In total, 33 mutations were identified (including 10 mutations observed in

mixed-populations), and 19 of these mutations were novel, according to the comparison

with 1111 genomic sequences available at GISAID on 3/24/2020 (Fig. 1, S1, and S2).

Specifically, G11083T and G26144T were found in ZJU-1, and both of these mutations

are known as founding mutations for a large group of viruses (Capobianchi et al., 2020).

C8782T and T28144C were found in two of our viral isolates, ZJU-2 and ZJU-8, and

these two are known as the founding mutations for another large group of viral isolates

(Capobianchi et al., 2020). Interestingly, mutation T22303G was found in five viral

isolates (ZJU-2, -5, -9, -10, and -11) and ZJU-5 and ZJU-9 were exposed to the same

potential source of infection during a business conference (Table 1). Previously, only one

viral isolate identified in Australia had the T22303G mutation. Strikingly, the viral isolate

from patient ZJU-4, who attended the same conference as ZJU-5 and ZJU-9, has a novel

mutation, A22301C, which causes the same missense mutation at the protein level

(S247R in the S protein) as T22303G, mutating the 1st instead of the 3rd position in the

respective codon. Observations of these two single nucleotide variants can only be

coincidental, albeit very unexpected. Finally, the ZJU-11 has 4 mutations in the ORF7b

gene, 3 of which are consecutive and introduce 2 mutations at the protein level.

Di-nucleotide and Tri-nucleotide mutations are of course rarer than SNV, but not

exponentially so, according to previous mutational accumulation studies in prokaryotes

(Lynch, 2007).

It is important to note that while the sequence data deposited in GISAID are very

helpful in tracking inter-personal variation of the virus, we still do not know much about

intra-personal viral evolutionary dynamics. For example, in ZJU-4 and ZJU-10, alleles of

two separate sites have very similar frequency distributions, indicating that these two

sites are probably linked, representing at least two haplotypes within the viral populations.

And as revealed by this study, 6 of the identified mutations would have been ignored if

using the consensus sequences for analyses. Taken together, despite only 11

patient-derived isolates being analyzed in this study, we observed abundant mutational

diversity, including several founding mutations for different major clusters of viruses

now circulating globally. This diverse mutational spectrum is consistent with their

relatively early sampling time and relative proximity to Wuhan city, where the first viral

strain was identified. The full mutational diversity of the virus in Wuhan city in the early

days is still unknown to this day, due to limited sampling (Lu et al., 2020; Zhou et al.,

2020).

Fig. 1. A summary of the mutations identified in each of the 11 viral isolates. Each ORF

of the viral genome was denoted based on the annotations of NC_045512.2 as provided

by NCBI. If a mutation was observed in the context of a mixed population, the respective

percentages of the top two alleles are provided. Changes at the amino acid level are

provided if applicable. Blue color indicates novel mutation at the time of writing the

article.

Phylogenetic analysis of the patient-derived viral isolates reveals their diverse

evolutionary history

To understand the phylogenetic context of 11 viral isolates with respect to the corpus of

available SARS-CoV-2 sequencing data, we acquired 725 high quality and high coverage

SARS-CoV-2 genomes from GISAID (downloaded on 3/21/2020), including the Yunnan

RaTG13 viral strain and the Guangdong pangolin viral strain as the outgroup. We aligned

the 736 genomic sequences with MAFFT (see Materials and Methods) and trimmed the

full-length alignment with trimAL (see Materials and Methods) to remove any spurious

parts of the alignment. We used iqtree (see Materials and Methods) to construct a

1000-times bootstrapped maximum-likelihood phylogenic tree of the 736 viral sequences

based on 835 parsimony informative sites (Fig. 2A). The resulting phylogenetic tree is

largely consistent with the phylogenetic analysis being updated on GISAID (Fig. S3). We

want to emphasize that due to the rapidly developing COVID-19 situation, tens or even

hundreds of new sequences are being uploaded to the GISAID every day. As a result,

new observations may be generated when more data are available.

Fig. 2. Characterizations of the patient-derived SARS-CoV-2 isolates. (A) Phylogenetic

analyses of the 11 viral isolates in the context of 725 SARS-CoV-2 sequences

downloaded from GISAID. The 1000-times bootstrapped maximum likelihood tree was

constructed to demonstrate the phylogenetic context of the 11 viral isolates. Major and

minor clusters were color-coded and denoted as shown in the "colored ranges" inset box.

All ZJU- samples were color-codded as green. The width of a branch indicates bootstrap

supporting level. (B) Fluorescent labeling of the viral S protein indicates that isolated

SAR-CoV-2 viral particles (Green) bind to the peripherals of the Vero-E6 cells (DNA

stained as Blue) prior to entry. Scale bars, 50 μm. (C) A representative TEM picture of

the isolated SAR-CoV-2 viral particles, arrows indicate the iconic "crown" consisted of S

proteins (Spike). Scale bar, 100 nm.

We observed quite a few sets of founding mutations. Specifically, we note the

following three biggest clusters in our phylogenetic analysis: 1. Three nucleotide

mutations C241T (silent), C14408T (silent), and A23403G (D614G in S) found a group

of 231 viral sequences (Fig. 2A; S-D614G cluster), most of which were isolated in

Europe; 2. Two nucleotide mutations C8782T (silent) and T28144C (L84S in ORF8),

found a group of 208 viral sequences (Fig. 2A; ORF8-L84S cluster), which is not

monophyletic in our analysis (Fig. 2A and S3). However, a distinct monophyletic

subclade of 92 sequences within the ORF8-L84S cluster can be observed, mainly

composed of viral sequences isolated from Seattle, USA (Fig. 2A;

ORF8-L84S-USA-WA-clade); 3. Two nucleotide mutations, G11083T (L3606F in

ORF1a), and G26144T (G251V in ORF3a) found a group of 34 viral sequences, most of

which were from Netherlands and England. Several smaller monophyletic clusters,

defined by different sets of founding mutations (bootstrap supporting value >95), can be

observed. For examples: 1. the G1937A (V378I in ORF1a) mutation founds a cluster of

31 viral sequences; 2. the G1440A and G2891A mutations, resulting in G392D and

A876T mutations in the ORF1a gene, founds a cluster of 12 viral sequences, mostly from

Germany or Netherlands; 3. the C15325T and C29303T mutations, resulting in P344S

mutation in the N gene, founds a small cluster of 8 sequences, all of which are from

China or Japan. When integrating the characterized 11 viral isolates into the phylogenetic

analysis, they are dispersed across the entire phylogenetic space. ZJU-1 clusters with the

ORF1a-L3606F & ORF3a-G251V groups, as it has both of the two defining mutations

(Fig. 2A). ZJU-2 and ZJU-8, on the other hand, cluster with the ORF8-L84S cluster

because they both have the two founding mutations (Fig. 2A). ZJU-9 and ZJU-11 cluster

with an Australian isolate because of the aforementioned T22303G mutation. The rest of

the group either have few mutations or novel mutations that do not cluster with any

known sizable groups, reflecting the extensive diversity within our 11 samples. Taken

together, some monophyletic clusters of viruses do show obvious geographic patterns

(Europe and WA-USA especially), but this could be due to the founding effect of

respective mutations that happened early during the initial phase of the pandemic.

Patient-derived SARS-CoV-2 isolates show significant variation in viral copy

number and cytopathic effects when infecting Vero-E6 cells

There is much speculations and many theories behind the observed mutations in

sequenced viral isolates. Theoretically, one usually needs not often invoke selection

arguments in explaining the origin of these mutations, as the human to human infection

process is a series of repeated naturally-occurring bottlenecking events, in which the

seeding viral population can be as little as hundreds of viral copies (Forni et al., 2017).

Therefore, a significant portion of the genetic diversity or even population-specific

fixations could be due to this process, where selection plays a small role (Renzette et al.,

2017). We conducted Tajima's test of neutrality using the constructed alignment of viral

sequences and Taijima's D is -2.8874 with a nucleotide diversity (π) of 0.000641 (p <

0.05 according to simulations performed in (Tajima, 1989), indicating that the

SARS-CoV-2 genome has an excess of low-frequency alleles due to recent population

expansions, consistent with the repeated bottlenecking events during viral infections.

However, certain mutations do provide selection advantages or disadvantages under

specific circumstances, as shown by the discovery that adaptive mutations are highly

enriched in the interface between the S protein and the human ACE2 receptor (Ou et al.,

2020).

To examine the mutational impact of the patient-derived SARS-CoV-2 isolates, we

conducted *in vitro* infectivity assay. We chose *in vitro* assay because COVID-19 patients

show a wide variety of clinical symptoms ranging from asymptotic to death, and

epidemiological research have shown that the clinical outcomes are heavily influenced by

individual's age, complications, and other potential unknown parameters (Guan et al.,

2020). We first examined whether the viral isolates could successfully bind to Vero-E6

cells as expected (Fig. 2B), and visually identified the viral particles with the iconic

"crown" formed by S proteins (Fig. 2C and S4A). We then infect the Vero-E6 cells with

all 11 patient-derived viral isolates and harvest the cells at 1, 2, 4, 8 (in quadruplicates),

24, and 48 (in duplicates) hours P.I. (see Materials and Methods); we included the

supernatant because cell death releases viral particles. DIC micrographs of the cells at 48

hours and 72 hours P.I. were also taken to assess the CPE. We used specific real-time

reverse transcriptase–polymerase chain reaction (RT-PCR) targeting ORF1a, E, and N

genes to detect the presence of SARS-CoV-2 (see Materials and Methods). Cycle

threshold values, C_t, were used to quantify the viral load, with lower values indicating

higher viral load. Because the results based on the three genes are highly consistent (R >

0.99, p < 2.2e-16), we will only discuss the results of the ORF1a gene (Fig. 3A). We

failed to detect any significant signals from our negative controls, hence we simply

assigned a C_t value of 40 for all them.

Briefly, the C_t values of samples remained mostly flat with small fluctuations for all

of the viral isolates at 1, 2, and 4 hours P.I. (Fig. 3A and B). During these early hours,

viral particles are binding to gain access into the cells, and replications would rarely

occur (Schneider et al., 2012). At 8 hours P.I., we observed significant decreases in Ct

value (increases in viral load) for ZJU-6, ZJU-7, ZJU-9, ZJU-10, and ZJU-11. At 24

hours P.I., we observed significant decreases in C_t values for all of the viral isolates

except for ZJU-2 and ZJU-7, although some of the viral isolates, namely ZJU-10 and

ZJU-11, decreased much faster than the others (Fig. 3A and B). At 48 hours P.I., we

observed small decreases for all viral isolates except for ZJU-10 and ZJU-11, both of

which had presumably already plateaued at 24 hours P.I. (Fig. 3A and B). Notably, at 24

hours P.I., ZJU-2 and ZJU-8, members of the ORF-8-L84S cluster (majority of USA

WA-Seattle isolates are in this group), showed considerably lower viral loads (Fig. 4A).

On the other hand, ZJU-1, which clusters with the S-D614G clade (mostly found in

Europe), has a viral load 19 times ($2^{4.25}$) higher than ZJU-2 and ZJU-8 (Fig. 4A). In

addition, a near 270-fold difference ($2^{8.09}$) in viral load was observed between ZJU-10

and ZJU-2 at 24 hours P.I. (Fig. 4A). These differences became statistically significant at

48 hours P.I., and are reproducible when analyzing data on gene E and N (Fig. S4B and C;

Fig. S5A). Therefore, different viral isolates, which are defined by different mutations in

their genomes, exhibit a significant variation of viral load when infecting Vero-E6 cells.

We next examined whether a higher viral load leads to more cell death (Fig. 4B).

When examining these cell lines under a microscope at 48 hours and 72 hours P.I., the

CPE, or the cell death rate, are highly consistent with the viral load data (Fig. 4C and S5B,

C, and D; at 48 hours P.I., C_t vs CPE, R = -0.72, p = 0.015), indicating that a higher viral

load leads to a higher cell death ratio. Note that the C_t numbers are negatively correlated

with the CPE because a lower C_t number means a higher viral load.

Fig. 3. The infectivity assay reveals temporal variation in the viral load of the

patient-derived SARS-CoV-2 isolates when infecting Vero-E6 cells. (A) Time-series

plots of the C_t values (corresponding to the multiplicative inverse of viral load) of the

SAR-CoV-2 ORF1a gene over the course of infectivity assay. Each viral isolate plus the

negative control "C" was color-coded accordingly. P-values were calculated using the

ANOVA method to compare the means of all 11 viral isolates at each time point,

excluding the negative control "C". (B) Time-series plots of the C_t values of the

SAR-CoV-2 ORF1a gene for each of the 11 patient-derived viral isolates. P-values were

calculated between consecutive time points using the t-test and adjusted p-values are

shown.

Fig. 4. The changes in CPE and viral load are highly correlated. (A) Significant variations in viral load were observed at each time point. Mean C_t values of selected viral isolates are displayed and color-coded, respectively. P-values were calculated using the ANOVA method to compare the means of all 11 viral isolates at each time point. Pair-wise p-values were calculated between isolates using the t-test and adjusted p-values

are shown. (B) Cytopathic effects were visible under the microscope 48 hours P.I., white

arrows indicate representative cells undergoing lysis. (C) Cytopathic effects (CPE) were

highly correlated with viral load (CT) in viral infectivity assay. Pearson correlations were

calculated and p-values were adjusted accordingly; only correlations with adjusted

p-value < 0.05 are shown. Note that the CT values are negatively correlated with CPE

values because CT values represent the inverse number of viral loads.

Discussion

The quickly-developing COVID-19 pandemic has already infected millions of victims

and caused 70,000 deaths globally. While many ongoing research projects are attempting

to track the evolutionary origin of the virus, find the mechanisms of infection, and

produce vaccines or drugs against the virus, we sought to establish the

genotype-phenotype link behind the abundant diversity being observed as a result of

global sequencing efforts (GISAID). Due to the extremely wide variety of clinical

symptoms shown in the patients, establishing a genotype-phenotype link in patients

would be very difficult. The *in vitro* cell line provides an ideal system to examine the

mutational impact of different isolates of viruses, when all other confounding factors are

removed. Although the Vero-E6 cell line was not derived from human, the ACE2 protein

of the Vero-E6 cell line is highly similar to that of Human (Fig. S6) and we provided

direct evidence that the SARS-CoV-2 can infect the cell line (Fig. 2B).

Several findings stand out in our study: 1. A diverse collection of mutations was

identified in the 11 viral isolates, including two sets of founding mutations for two major

clusters of viruses currently infecting the world population. In addition, 19 of the 31

identified mutations are novel, despite the relatively early sampling dates, indicating that

the true diversity of the viral strains is still largely underappreciated; 2. remarkably, the

T22303G and A22301C mutations result in the same S247R mutation in the S protein

(Fig. 1 and S1), mapping to the existing structure revealed that this residue is located in a

flexible loop region within the N-terminal domain of the S1 subunit of S protein,

although the exact position of S247 could not be determined (Fig. S7, red arch). While

the N-terminal domain is not directly involved with binding to ACE2 (Walls et al., 2020)

we note that this domain is positioned right next to the C-terminal domain, which binds to

ACE2. Interestingly, the T22303G mutation was observed in 5 viral isolates, albeit in

different proportions, indicating that this specific mutation was already present in the

early days of pandemic, and probably in a significant number of people of Wuhan,

despite the fact that it is still largely missing from current GISAID collection. This could

be due to the founding effect of mutations, in which case the T22303G mutation was not

transmitted out of the China during the early days; 3. The tri-nucleotide mutation in

ZJU-11 is unexpected; we note that this specific viral isolate is quite potent in our viral

load and CPE assay, and its patient remained positive for an astounding period of 45 days

and was only recently discharged from the hospital (Table 1). Investigating the functional

impact of this tri-nucleotide mutation would be highly interesting. We note that in the

current database, another trinucleotide mutation (G28881A, G2882A and G28883C) has

been identified, which also results in two missense mutations at the protein level (Fig. S8).

It leads to a cluster of more than 300 viral strains as of the time of writing this article, and

its mutational impact on the viral pathogenicity would be worth investigating. Finally, in

contrary to the recent report that a viable viral isolate could not be obtained from stool

samples, three of our viral isolated were extracted from stool samples, indicating that the

SARS-CoV-2 is capable of replicating in stool samples (Woelfel et al., 2020).

In short, our study provides direct evidence that mutations currently occurring in the

SARS-CoV-2 genome have the functional potential to impact the viral pathogenicity.

Therefore, viral surveillance should be also performed at the cellular level when possible,

in addition to the accumulating genomic sequencing data. Furthermore, characterizations

of all founding mutations in the major geo-based clusters of viruses could be very useful

in helping determining if there are actionable pathogenicity differences to aid the current

battle against the virus. Finally, similar to flu, drug and vaccine development, while

urgent, need to take the impact of these accumulating mutations, especially the founding

mutations, into account to avoid potential pitfalls.

Figure legends

Table 1. A summary of the epidemiological information of the 11 patients involved in

this study. The "Viral gen" (viral generation) was inferred based on their exposure

history.

Fig. 1. A summary of the mutations identified in each of the 11 viral isolates. Each ORF

of the viral genome was denoted based on the annotations of NC_045512.2 as provided

by NCBI. If a mutation was observed in the context of a mixed population, the respective

percentages of the top two alleles are provided. Changes at the amino acid level are

provided if applicable. Blue color indicates novel mutation at the time of writing the

article.

Fig. 2. Characterizations of the patient-derived SARS-CoV-2 isolates. (A) Phylogenetic

analyses of the 11 viral isolates in the context of 725 SARS-CoV-2 sequences

downloaded from GISAID. The 1000-times bootstrapped maximum likelihood tree was

constructed to demonstrate the phylogenetic context of the 11 viral isolates. Major and

minor clusters were color-coded and denoted as shown in the "colored ranges" inset box.

All ZJU- samples were color-codded as green. The width of a branch indicates bootstrap

supporting level. (B) Fluorescent labeling of the viral S protein indicates that isolated

SAR-CoV-2 viral particles (Green) bind to the peripherals of the Vero-E6 cells (DNA

stained as Blue) prior to entry. Scale bars, 50 μm. (C) A representative TEM picture of

the isolated SAR-CoV-2 viral particles, arrows indicate the iconic "crown" consisted of S

proteins (Spike). Scale bar, 100 nm.

Fig. 3. The infectivity assay reveals temporal variation in the viral load of the

patient-derived SARS-CoV-2 isolates when infecting Vero-E6 cells. (A) Time-series

plots of the C_t values (corresponding to the multiplicative inverse of viral load) of the

SAR-CoV-2 ORF1a gene over the course of infectivity assay. Each viral isolate plus the

negative control "C" was color-coded accordingly. P-values were calculated using the

ANOVA method to compare the means of all 11 viral isolates at each time point,

excluding the negative control "C". (B) Time-series plots of the C_t values of the

SAR-CoV-2 ORF1a gene for each of the 11 patient-derived viral isolates. P-values were

calculated between consecutive time points using the t-test and adjusted p-values are

shown.

Fig. 4. The changes in CPE and viral load are highly correlated. (A) Significant

variations in viral load were observed at each time point. Mean C_t values of selected viral

isolates are displayed and color-coded, respectively. P-values were calculated using the

ANOVA method to compare the means of all 11 viral isolates at each time point.

Pair-wise p-values were calculated between isolates using the t-test and adjusted p-values

are shown. (B) Cytopathic effects were visible under the microscope 48 hours P.I., white

arrows indicate representative cells undergoing lysis. (C) Cytopathic effects (CPE) were

highly correlated with viral load (CT) in viral infectivity assay. Pearson correlations were

calculated and p-values were adjusted accordingly; only correlations with adjusted

p-value < 0.05 are shown. Note that the CT values are negatively correlated with CPE

values because CT values represent the inverse number of viral loads.

STAR Methods

Experimental Model and Subject Details

The study was approved by the Clinical Research Ethics Committee of The First Affiliated Hospital, School of Medicine, Zhejiang University (Approval notice 2020-29) for emerging infectious diseases. Patients with confirmed COVID-19 were admitted in the First Affiliated Hospital from Jan 19 to Mar 5, 2020. The First Affiliated Hospital, located in Hangzhou, Zhejiang Province, China, is one of the major provincial hospitals designated to receive patients with COVID-19 infection across the Zhejiang Province; therefore, patients with severe symptoms outside of Hangzhou were also admitted. Starting Jan 10, 2020, all patients presenting to the hospital's fever clinic were screened by clinical staff for COVID-19 infection utilizing criteria for suspected cases as defined by the National Health Commission of China's clinical diagnosis and management guideline for COVID-19 (China National Health Committee, 2020). Briefly, patients were screened based on their clinical symptoms and their risk of epidemiological exposure, including past travel to Hubei Province or close contact with people who had visited Hubei Province during the COVID-19 outbreak. As the pandemic continued to spread, the probability of transmission outside of Hubei Province increased. The epidemiological exposure to Hubei Province was not a prerequisite for suspected cases.

419   All suspected cases were determined by laboratory tests and based on positive results of

420   qRT-PCR assay for COVID-19. Patients were excluded if two qRT-PCR tests 24 hours

421   apart both suggested negative results. Patients' clinical samples which PCR test C_t value

422   less than 28 were collected to isolate SARS-Cov-2.

Method Details

Sample collection, Viral isolation, cell infection, and electron microscopy

All samples, sources including sputum, nasopharyngeal swab, and stool, were collected

from patients with COVID-19 with consent from all patients. All collected samples were

sent to BSL-3 lab for viral isolation within 4 hours. The sputum, stool, and

nasopharyngeal swab samples were pre-processed by first mixing with appropriate

volume (Sputum, 5-10 volumes; Stool, 2 ml/100 mg; Nasopharyngeal swab, 1 volume) of

MEM medium with 2% FBS, Amphotericin B (100 ng/ml), Penicillin G (200 units/ml),

Streptomycin (200 µg/ml), and TPCK-trypsin (4 µg/ml). The supernatant was collected

after centrifugation at 3000 rpm at room temperature. Before infecting Vero-E6 cells, all

collected supernatant was filtered using a 0.45 µm filter to remove cell debris etc.

For viral infection and isolation, 3 ml of filtered supernatant was added to Vero-E6

cells in a T25 culture flask. After incubation at 35°C for 2h to allow binding, the

inoculum was removed and replaced with fresh culture medium. The cells were incubated

at 35°C and observed daily to evaluate cytopathic effects (CPE). The supernatant was

tested for SARS-CoV-2 by qRT-PCR (see below for qRT-PCR protocol). Once the

qRT-PCR test shows positive (typically after 4-5 days of incubation), the viral particles

were collected from culture supernatant by ultra-speed centrifugation (100,000x g for 2

hours) for downstream sequencing, infectivity assay, and were observed under 200 kV

Tecnai G2 electron microscope.

Immunofluorescence staining

Vero-E6 Cells were infected by SARS-CoV-2 for 24 hours, and then fixed in 80%

acetone (chilled at -20°C) at room temperature for 10 min. The cells were washed three

times with ice-cold PBS, blocked with 1% BSA for 30 min, and incubated with

SARS-CoV-2 Spike rabbit monoclonal antibody (dilution ratio 1:200) at room

temperature for 1 hour. The cells were again washed three times in ice-cold PBS, and

then stained with the Alexa Fluor488®-conjugated Goat Anti-rabbit IgG secondary

antibody (Abcam, Cat No. ab150077) for 1 hour at room temperature in the dark. The

cells were washed three times and then incubated in 0.5 μg/mL DAPI (nuclear DNA stain)

for 5 min. Immunofluorescence was detected and picture were taken using the IX81

Olympus microscope equipped with a fluorescence apparatus.

Viral infectivity assay

Vero-E6 cells were grown in a 24-well plate and infected with different SARS-CoV-2

isolates in duplicates at MOI of 0.5. The inoculum was removed at 1 hours P.I. for the

1-hour timepoint group and at 2 hours P.I. for other timepoint groups. After incubation,

the cultures were rinsed with PBS for three times and replenished with 1mL fresh culture

medium. Then, the cultures were subjected to freezing immediately at -80°C for the 1-

and 2-hours samples, or continued to grow for the other groups (4, 8, 24 and 48 hours)

until harvest. Finally, all frozen samples from each timepoint were thawed together and

the viral nucleic acid abundance was measured with SARS-CoV-2 qRT-PCR Kits,

targeting ORF1a, E, and N genes (Liferiver Biotech, Shanghai). Results from the first

two time points reflect the capacity of viral attachment or entry into the target cells, while

results from the latter four time points represent the viral replication dynamics.

Cytopathic effect (CPE) evaluation

Vero-E6 cell monolayers were grown and infected by different patient-derived

SARS-CoV-2 isolates as described in the viral infectivity assay. At 24, 48, and 72 hours

P.I., virus induced cytopathic effects were observed with a digital microscope (Bio-Rad)

and pictures were taken. No obvious CPE was observed at 24 hours P.I.. Pictures taken at

48 and 72 hours P.I. were evaluated first by expert opinions and then quantitated by cell

death ratio.

Sequencing library construction

The total RNA in each deactivated viral sample was extracted using a viral RNA mini kit (Qiagen, Germany). The sequencing library was constructed using the Total RNA-Seq Kit (Kapa, Switzerland) and deep-sequenced on the illumina Novaseq 6000 platform (2 x 151 bases; Illumina Inc., San Diego, CA) by BGI genomics.

Quantification and Statistical Analysis

Statistical Analyses and visualization

The majority of statistical analyses and visualizations were done in Rstudio and R (at the time of writing, 1.0143 for Rstudio and 3.4.0 for R), with necessary aid from customized python scripts (2.7.4) and shell scripts (Linux). The primary R packages are mostly maintained by the Bioconductor project (https://www.bioconductor.org/, along with all their dependencies). The essential ones used are ggplot2 (2.2.1), reshape2 (1.4.3), RColorBrewer (1.1-2), scales (0.5.0), corrplot (0.84), Hmisc (4.1-1), ggrepel (0.7.0), cluster (2.0.6), factoextra (1.0.5), plyr (1.8.4), dplyr (0.7.4), psych (1.7.8), devtools (1.13.4), ggpubr (0.1.6), tidyverse (1.2.1), gridExtra (2.3), ggsci (2.8), ggbeeswarm (0.6.0), ggpmisc (0.2.16), colorspace (1.3-2).

In general, parametric statistical tests (t-test, Anova, and Pearson correlation) were

used when the data distribution conforms to normality distribution (such as qPCR

measurements), and non-parametric statistical tests (Wilcoxon test, Kruskal-Wallis, and

Spearman correlation) were used when datasets do not conform to the normality

assumption. We adjust the p values using the Benjamini & Hochberg (BH) method

(Benjamini and Yekutieli, 2001) to control for False Discovery Rate (FDR), when

multiple comparisons are concerned, including p value matrix constructed when

calculating correlations matrix among different features or samples.

The 3D structure of the S protein was visualized and downloaded from

https://www.rcsb.org/3d-view/6VSB/1.

Sequence data processing, de novo assembling, and mutation identifications

Sequencing data was generated from Novaseq 6000 and first filtered of low quality and

high barcode contamination by Soapnuke and then mapped to 43 complete genome

references of 2019-nCoV (SARS-CoV-2) by BWA-MEM (Li and Durbin, 2009).

References of SARS-CoV-2 were downloading from NCBI on date February 28th, 2020.

Further, mapping reads that longer than 100nt were extracted for de novo assembly by

SPAdes (Bankevich et al., 2012) (v3.1.3) using an iterative short-read genome assembly

module for pair-end reads. K-values were selected automatically at 33nt, 55nt and 77nt

for these samples. After assembling, contigs was blasted to nt database (20190301) to

confirm their origins, and only contigs belonging to coronavirus were retained for base

correction. Next, filtering reads of each sample were mapped back to retained assembled

contigs and bam-readcount was applied (--min-mapping-quality=5, other parameter was

set default) to calculate the base frequency of every post of each assemble contigs.

Meanwhile, Haplotypecaller of gatk was applied to call snp/indel based on the assembled

contigs with reads quality higher than 20. Finally, bam files were inspected in igv

manually to verify each mutation based on the number of reads mapped, the balance

between reads mapped to plus and minus strands of the reference genome, and the

relative positions of the mutations on these reads.

Phylogenetic analysis

We acquired 725 high quality and high coverage SARS-CoV-2 genomes from GISAID

(downloaded on 3/21/2020), including the Yunnan RaTG13 viral strain and the

Guangdong pangolin viral strain as the outgroup. We aligned the 736 genomic sequences

with MAFFT (Katoh and Standley, 2013) with options --thread 16 --globalpair

--maxiterate 1000 and trimmed the full-length alignment with trimAL (Capella-Gutiérrez

et al., 2009) using the -automated1 option to remove any spurious parts of the alignment,

which could introduce noise to the phylogenetic analysis process. We used iqtree

(Nguyen et al., 2015) with options -bb 1000 -alrt 1000 -nt 64 -asr to construct a

1000-times bootstrapped maximum-likelihood phylogenic tree of the 736 viral sequences

based on 835 parsimony informative sites. The resulting phylogenetic tree was imported

in iTOL and visualized (Letunic and Bork, 2011). We conducted Tajima's test of

neutrality based on the constructed alignment of viral sequences using MEGA 7 (Kumar

et al., 2016).

Data and Software Availability

541   The full-genome sequences of the 11 viral isolates have been deposited to the GISAID

542   collection with the following IDs: EPI_ISL_415709, EPI_ISL_416042, EPI_ISL_416044,

543   EPI_ISL_416046, EPI_ISL_415711, EPI_ISL_416047, EPI_ISL_416425,

544   EPI_ISL_416473, EPI_ISL_416474, EPI_ISL_418990, and EPI_ISL_418991.

Supplementary Figure Legends

Table S1. A summary of the sequencing statistics of the 11 viral isolates involved in the study, Related to Figure 1. Note that ZJU_10 and ZJU_11 were sequenced in a different batch.

Fig. S1. A summary of the nucleotide mutations that lead to the S247R mutations observed in the 11 patient-derived isolates, Related to Figure 1. Note that some of the mutations are in the form of minor alleles. Images were produced by IGV.

Fig. S2. A summary of additional mutations in the S gene and the tri-nucleotide mutation, Related to Figure 1. Note that some of the mutations are in the form of minor alleles. Images were produced by IGV.

Fig. S3. Phylogenetic analyses produced from GISAID using time (top) or number of mutations (bottom) as the branch length, Related to Figure 2. Note that all three major clusters described in the study are labeled accordingly. The major distinction is that the ORF8-L84S clade is not monophyletic in our more computationally intensive and bootstrapping-supported approach.

Fig. S4. The characterizations of the 11 viral isolates, Related to Figure 2 and Figure 4.

(A) A representative TEM picture of the isolated SAR-CoV-2 viral particles, arrows

indicate the iconic "crown" consisted of S proteins (Spike). (B) Time-series plots of the

C_t values (corresponding to the multiplicative inverse of viral load) of the SAR-CoV-2 E

gene (top) and N gene (bottom) over the course of infectivity assay. Each viral isolate

plus the negative control "C" was color-coded accordingly. P-values were calculated

using the ANOVA method to compare the means of all 11 viral isolates at each time

point, excluding the negative control "C". (C) Time-series plots of the C_t values of the

SAR-CoV-2 E gene (left) and N gene (right) for each of the 11 patient-derived viral

isolates. P-values were calculated between consecutive time points using the t-test and

adjusted p-values are shown.

Fig. S5. Significant variations were observed in viral load and viral CPE among the 11

patient-derived isolates, Related to Figure 4. (A) Significant variations in viral load can

be observed based on E gene (left) and N gene (right). (B) CPE at 48h and 72h P.I. as

evaluated by an expert's opinions. (C) CPE at 48h and 72h P.I. evaluated by quantitively

calculating the cell death ratio (ratio.dead) for 1-3 images per viral isolate. The results

from (B) and (C) are highly correlated (R > 0.89, p < 0.001). (D) Representative images

used for CPE evaluation, arrows indicate cells facing immediate death. Scale bars, 100

μm.

Fig. S6. The alignment of ACE2 protein sequences from human (Homo), Chimpanzee,

and green monkey (from which the Vero-E6 cell line was derived), Related to Figure 2.

Note that overall the ACE2 proteins are highly similar to each other. The alignment and

image were produced by Jalview.

Fig. S7. The 3D structure of the S-protein with the S247R overlay, Related to Figure 1.

The top (A), bottom (B), side (C), and close-up view (D) were provided. Note that the

actual position of S247 was not determined in the original structure, hence a small red arc

was in place to represent to the potential flexible loop conformation for (C) and (D). Also

note that the protein complex is trimeric, but only one of the three mutations was labeled.

The 3D structure of the S protein was visualized and downloaded from

https://www.rcsb.org/3d-view/6VSB/1.

Fig. S8. The trinucleotide mutation (G28881A, G28882A, and G28883C) was identified

in the GISAID dataset and is the founding mutation for a large cluster of viral isolates

within the S-D614G group (European clade), Related to Figure 1.

Author Contributions

M.Z., C.J., N.W., and L.L conceived and supervised the study. H.Y. and X.L. performed

all the experiments, with help from K.X., Y.C., L.C., F.L., Z.W., H.W., and C. Jin. C.J.,

M.Z., and Q.C. performed all the data analyses. C.J., Q.C., H.Y., and M.Z. drafted and

revised the manuscript with input from all authors.

Acknowledgments

We gratefully acknowledge Drs. X. Zhu, L. Xiang, J. Jensen, and M. Lynch for their

helpful discussions. We thank Dr. V. Billing for her help on improving the manuscript.

Funding

This work was supported by funds from Major Project of Zhejiang Provincial Science

and Technology Department #2020C03123, National Science and Technology Major

Project for the Control and Prevention of Major Infectious Diseases in China

Declaration of Interests

None

References

Bai, Y., Yao, L., Wei, T., Tian, F., Jin, D.Y., Chen, L., and Wang, M. (2020). Presumed Asymptomatic

Carrier Transmission of COVID-19. JAMA - J. Am. Med. Assoc.

Bankevich, A., Nurk, S., Antipov, D., Gurevich, A.A., Dvorkin, M., Kulikov, A.S., Lesin, V.M., Nikolenko,

S.I., Pham, S., Prjibelski, A.D., et al. (2012). SPAdes: A new genome assembly algorithm and its

applications to single-cell sequencing. J. Comput. Biol. *19*, 455–477.

Bauch, C.T., Lloyd-Smith, J.O., Coffee, M.P., and Galvani, A.P. (2005). Dynamically modeling SARS and

other newly emerging respiratory illnesses: Past, present, and future. Epidemiology *16*, 791–801.

Benjamini, Y., and Yekutieli, D. (2001). The control of the false discovery rate in multiple testing under

dependency. Ann. Stat. *29*, 1165–1188.

Cao, B., Wang, Y., Wen, D., Liu, W., Wang, J., Fan, G., Ruan, L., Song, B., Cai, Y., Wei, M., et al. (2020). A

Trial of Lopinavir-Ritonavir in Adults Hospitalized with Severe Covid-19. N. Engl. J. Med. 1–13.

Capella-Gutiérrez, S., Silla-Martínez, J.M., and Gabaldón, T. (2009). trimAl: A tool for automated

alignment trimming in large-scale phylogenetic analyses. Bioinformatics.

Capobianchi, M.R., Rueca, M., Messina, F., Giombini, E., Carletti, F., Colavita, F., Castilletti, C., Lalle, E.,

Bordi, L., Vairo, F., et al. (2020). Molecular characterization of SARS-CoV-2 from the first case of

COVID-19 in Italy. Clin Microbiol Infect *0*.

China National Health Committee (2020). COVID-19 clinical diagnosis and management guideline issued

by National Health Commission of China, the 5th edition.

Corman, V.M., Muth, D., Niemeyer, D., and Drosten, C. (2018). Hosts and Sources of Endemic Human

Coronaviruses. In Advances in Virus Research, pp. 163–188.

van Doremalen, N., Bushmaker, T., Morris, D.H., Holbrook, M.G., Gamble, A., Williamson, B.N., Tamin,

A., Harcourt, J.L., Thornburg, N.J., Gerber, S.I., et al. (2020). Aerosol and Surface Stability of

SARS-CoV-2 as Compared with SARS-CoV-1. N. Engl. J. Med.

Forni, D., Cagliani, R., Clerici, M., and Sironi, M. (2017). Molecular Evolution of Human Coronavirus

Genomes. Trends Microbiol. *25*, 35–48.

Guan, W., Ni, Z., Hu, Y., Liang, W., Ou, C., He, J., Liu, L., Shan, H., Lei, C., Hui, D.S., et al. (2020).

Clinical characteristics of 2019 novel coronavirus infection in China. N. Engl. J. Med.

Hoffmann, M., Kleine-Weber, H., Schroeder, S., Krüger, N., Herrler, T., Erichsen, S., Schiergens, T.S.,

Herrler, G., Wu, N.-H., Nitsche, A., et al. (2020). SARS-CoV-2 Cell Entry Depends on ACE2 and

TMPRSS2 and Is Blocked by a Clinically Proven Protease Inhibitor. Cell 1–10.

Hu, Z., Song, C., Xu, C., Jin, G., Chen, Y., Xu, X., Ma, H., Chen, W., Lin, Y., Zheng, Y., et al. (2020).

Clinical characteristics of 24 asymptomatic infections with COVID-19 screened among close contacts in

Nanjing, China. Sci. China Life Sci.

Katoh, K., and Standley, D.M. (2013). MAFFT Multiple Sequence Alignment Software Version 7:

Improvements in Performance and Usability. Mol. Biol. Evol. *30*, 772–780.

Kumar, S., Stecher, G., and Tamura, K. (2016). MEGA7: Molecular Evolutionary Genetics Analysis

Version 7.0 for Bigger Datasets. Mol. Biol. Evol.

Lauer, S.A., Grantz, K.H., Bi, Q., Jones, F.K., Zheng, Q., Meredith, H.R., Azman, A.S., Reich, N.G., and

Lessler, J. (2020). The Incubation Period of Coronavirus Disease 2019 (COVID-19) From Publicly

Reported Confirmed Cases: Estimation and Application. Ann. Intern. Med.

Letunic, I., and Bork, P. (2011). Interactive Tree of Life v2: Online annotation and display of phylogenetic

trees made easy. Nucleic Acids Res. *39*, W475–W478.

Li, H., and Durbin, R. (2009). Fast and accurate short read alignment with Burrows-Wheeler transform.

Bioinformatics *25*, 1754–1760.

Lin, Q., Chiu, A.P.Y., Zhao, S., and He, D. (2018). Modeling the spread of Middle East respiratory

syndrome coronavirus in Saudi Arabia. Stat. Methods Med. Res. *27*, 1968–1978.

Lipsitch, M., Cohen, T., Cooper, B., Robins, J.M., Ma, S., James, L., Gopalakrishna, G., Chew, S.K., Tan,

C.C., Samore, M.H., et al. (2003). Transmission dynamics and control of severe acute respiratory syndrome.

Science (80-.). *300*, 1966–1970.

Liu, Y., Gayle, A.A., Wilder-Smith, A., and Rocklöv, J. (2020). The reproductive number of COVID-19 is

higher compared to SARS coronavirus. J. Travel Med.

Lu, R., Zhao, X., Li, J., Niu, P., Yang, B., Wu, H., Wang, W., Song, H., Huang, B., Zhu, N., et al. (2020).

Genomic characterisation and epidemiology of 2019 novel coronavirus: implications for virus origins and

receptor binding. Lancet *395*, 565–574.

Lv, L., Li, G., Chen, J., Liang, X., and Li, Y. (2020). Comparative genomic analysis revealed specific

mutation pattern between human coronavirus SARS-CoV-2 and Bat-SARSr-CoV RaTG13. BioRxiv

2020.02.27.969006.

Lynch, M. (2007). The Origins of Genome Architecture.

Nguyen, L.T., Schmidt, H.A., Von Haeseler, A., and Minh, B.Q. (2015). IQ-TREE: A fast and effective

stochastic algorithm for estimating maximum-likelihood phylogenies. Mol. Biol. Evol.

Ou, J., Zhou, Z., Zhang, J., Lan, W., Zhao, S., Wu, J., Seto, D., Zhang, G., and Zhang, Q. (2020). RBD

mutations from circulating SARS-CoV-2 strains enhance the structure stability and infectivity of the spike

protein. BioRxiv 2020.03.15.991844.

Renzette, N., Pfeifer, S.P., Matuszewski, S., Kowalik, T.F., and Jensen, J.D. (2017). On the Analysis of

Intrahost and Interhost Viral Populations: Human Cytomegalovirus as a Case Study of Pitfalls and

Expectations. J. Virol. *91*, e01976-16.

Schneider, M., Ackermann, K., Stuart, M., Wex, C., Protzer, U., Schätzl, H.M., and Gilch, S. (2012). Severe

Acute Respiratory Syndrome Coronavirus Replication Is Severely Impaired by MG132 due to

Proteasome-Independent Inhibition of M-Calpain. J. Virol. *86*, 10112–10122.

Tajima, F. (1989). Statistical method for testing the neutral mutation hypothesis by DNA polymorphism.

Tortorici, M.A., and Veesler, D. (2019). Structural insights into coronavirus entry. In Advances in Virus

Research, pp. 93–116.

Varia, M., Wilson, S., Sarwal, S., McGeer, A., Gournis, E., Galanis, E., and Henry, B. (2003). Investigation

of a nosocomial outbreak of severe acute respiratory syndrome (SARS) in Toronto, Canada. Cmaj *169*,

285–292.

Virlogeux, V., Fang, V.J., Park, M., Wu, J.T., and Cowling, B.J. (2016). Comparison of incubation period

distribution of human infections with MERS-CoV in South Korea and Saudi Arabia. Sci. Rep. *6*.

Walls, A.C., Park, Y.-J., Tortorici, M.A., Wall, A., McGuire, A.T., and Veesler, D. (2020). Structure,

Function, and Antigenicity of the SARS-CoV-2 Spike Glycoprotein. Cell.

Wang, Z., and Xu, X. (2020). scRNA-seq profiling of human testes reveals the presence of ACE2 receptor,

a target for SARS-CoV-2 infection, in spermatogonia, Leydig and Sertoli cells. Viruses 1–16.

Wang, C., Li, W., Drabek, D., Okba, N.M.A., Haperen, R. van, Osterhaus, A.D.M.E., Kuppeveld, F.J.M.

van, Haagmans, B.L., Grosveld, F., and Bosch, B.-J. (2020). A human monoclonal 1 antibody blocking

SARS-CoV-2 infection. BioRxiv 2020.03.11.987958.

Woelfel, R., Corman, V.M., Guggemos, W., Seilmaier, M., Zange, S., Mueller, M.A., Niemeyer, D., Vollmar,

P., Rothe, C., Hoelscher, M., et al. (2020). Clinical presentation and virological assessment of hospitalized

cases of coronavirus disease 2019 in a travel-associated transmission cluster. MedRxiv

2020.03.05.20030502.

Wu, F., Zhao, S., Yu, B., Chen, Y.M., Wang, W., Song, Z.G., Hu, Y., Tao, Z.W., Tian, J.H., Pei, Y.Y., et al.

(2020). A new coronavirus associated with human respiratory disease in China. Nature *579*, 265–269.

Xiao, F., Tang, M., Zheng, X., Liu, Y., Li, X., and Shan, H. (2020). Evidence for gastrointestinal infection

of SARS-CoV-2. Gastroenterology.

Yu, W.-B. (2020). Decoding evolution and transmissions of novel pneumonia coronavirus (SARS-CoV-2)

using the whole genomic data Comparative analyses of the chloroplast genome in carnivorous plants View

project. ChinaXriv.

Zhao, Y., Zhao, Z., Wang, Y., Zhou, Y., Ma, Y., and Zuo, W. (2020). Single-cell RNA expression profiling

of ACE2, the putative receptor of Wuhan 2019-nCov. BioRxiv 2020.01.26.919985.

Zhou, P., Yang, X. Lou, Wang, X.G., Hu, B., Zhang, L., Zhang, W., Si, H.R., Zhu, Y., Li, B., Huang, C.L.,

et al. (2020). A pneumonia outbreak associated with a new coronavirus of probable bat origin. Nature *579*,

270–273.

Zhu, N., Zhang, D., Wang, W., Li, X., Yang, B., Song, J., Zhao, X., Huang, B., Shi, W., Lu, R., et al. (2020).

A novel coronavirus from patients with pneumonia in China, 2019. N. Engl. J. Med. *382*, 727–733.