Multinational modeling of SARS-CoV-2 spreading dynamics: Insights on the heterogeneity of COVID-19 transmission and its potential healthcare burden

Erick R. Martinez-Loran[1], J. Jesús Naveja[2], Omar Y. Bello-Chavolla[3], Flavio F. Contreras-Torres[4]

[1]Department of NanoEngineering, University of California, San Diego, 9500 Gilman Drive, La Jolla, CA 92093, USA
[2]Instituto de Química, Universidad Nacional Autónoma de México, Circuito Exterior S/N C.U. 04510, Ciudad de México, Mexico
[3]Instituto Nacional de Geriatría, Anillo Perif. 2767, San Jerónimo Lídice, La Magdalena Contreras, 10200, Ciudad de México, Mexico
[4]Escuela de Ingeniería y Ciencias, Tecnologico de Monterrey, Monterrey, Nuevo León 64849, Mexico

April 14, 2020

ABSTRACT

Background: Despite efforts aimed at forecasting the course of the pandemic extrapolating from reports of the early onset in Wuhan, China, heterogeneity in the transmission seems to be an important characteristic of SARS-CoV-2 between countries. The study of the progress of the disease with a single set of transmission parameters is complicated by the application of different levels of containment measures at different stages of the pandemic.

Methods: We studied the variation in transmission of COVID-19 globally and individually for 18 countries. We developed a piecewise fitting methodology to model the progress of the COVID-19 epidemic for a set of countries with a number of confirmed infections in excess of 1000 cases. Our study uses the susceptible-exposed-infected-removed (SEIR) model to assess the contagious contact rate (β) of SARS-CoV-2 at different stages of the pandemic as a metric of the effectiveness of policies aimed at reducing the spread of COVID-19. We extrapolate the results of our fittings to evaluate the effect of a reduction of β on the level of burden laid on local health care infrastructure considering different levels of intervention and to estimate the maximum number of cases and potential peak date to inform future public health interventions.

Results: According to the current trend, we project that the peak in the number of infections worldwide will take place after the third quarter of 2020. For 12 out of the 17 countries analyzed, the projections suggest that the number of severe infections will surpass the healthcare capacity. Additionally, it is observed that 4 out of the 17 analyzed will undergo a significant delay in the peak of the infection for a 90% reduction of the transmission rate, and thus extending the course of the epidemic further than the 365 day simulation window. COVID19 transmission dynamics are

highly heterogeneous and influenced by both, the rate of spread and local response of the public health measures and to a lesser extent, to the compliance of the population.

Funding: No funding received.

KEYWORDS: COVID-19; SEIR model; SARS-CoV-2; Public health; Compartmental models;

1.   INTRODUCTION

The SARS-CoV-2 virus responsible for the coronavirus disease 2019 (COVID-19) emerged in Wuhan, China at the end of 2019. Early situation reports from the World Health Organization (WHO) database indicated that most of the regions with more than a thousand cases of SARS-CoV-2 infection exhibited an exponential progression in the number of cases after the first three weeks of the first confirmed case.[1] The first reports of the epidemic suggested an alarming death rate of ~3%, which has increased with the time up to more than 5% [2]. As of April 2020, the COVID-19 pandemic has globally reached more than 1,350,000 confirmed cases and 80,000 deaths [3]. However, the lethality rate and the number of confirmed cases notably differs between countries,[2] indicating distinct effects on the disease severity and transmission rate. Recent reports from Chinese COVID-19 cases have provided valuable information at the early stages of the epidemic and thus a general guidance to control the propagation within China has been considered in other countries.[4,5] The control measures aimed at reducing the contact rate, such as social distancing policies and isolation of infected populations, have proven to be effective in managing the epidemic. Nevertheless, difficulties in the implementation of strict measures have led to a fast increase in the number of cases.[6,7] For instance, Europe and the US have surpassed the number of fatalities in China despite the strict policies on social distancing that were adopted weeks ago. Due to the heterogeneity of the spread,

it is expected that the healthcare systems of most economies will face a considerable burden in the coming months because of the COVID-19 outbreak. [8]

In the present circumstance of the COVID-19 spread, understanding of the dynamics of this epidemic is necessary to forecast and control the transmission of the disease, as well as to evaluate the effectiveness of potential interventions in modifying its degree of infectiousness.[5] In particular, the first report for the transmissibility of COVID-19 estimated a basic reproduction number (R) ranging from 2.24 to 3.58 based on the time series of two other well-known coronavirus diseases (MERS and SARS). [9] However, the main goal of several studies was focused on the prediction of the trend of the epidemic as it expanded across mainland China, limiting the comparability of disease courses worldwide. Despite considerable efforts to model COVID-19 epidemic courses, the local dynamics of transmission is currently not well described for other countries.

The susceptible-exposed-infected-removed (SEIR) compartmental model has been recently used to model the progression of the epidemic in China.[4] Models of higher complexity to extrapolate information from early stages of the outbreak may not necessarily be more reliable in making predictions due to the larger number of model parameters to be estimated.[10] The SEIR model has been successfully used to model many other infectious diseases,[5,9,11–13] and can offer an adequate tradeoff between accuracy and complexity (e.g. between compartmental models of less complexity like the susceptible-infected-removed (SIR) model[14] and models of larger number of compartmental interactions, like the one described by Arenas *et al.* 2020[15]). This makes the SEIR model suitable to estimate COVID-19 rates by fitting the time series of the infected population. While it is common practice to fit only the infected compartment, the

removed population should also be considered when fitting, because its evolution determines the extent to which a decline in the number of infections will occur throughout the course of the epidemic. In this work, we use the SEIR model without vital kinetics (i.e., assuming a constant population size) to simultaneously fit the progression of active infections and removed populations for selected countries. We present a piecewise methodology that allows us to compare the progression of the epidemic at different stages and make comparisons of the infectiousness across countries. We use the extracted epidemic parameters to make relevant projections on the dynamics of the disease in forthcoming months. Information about the local dynamics of the epidemic can be useful to anticipate the impact of public policy interventions, including social distancing as well as to project healthcare supply needs in the short-term.

2.   METHODS

2.1   Data sources

The time series of the COVID-19 were extracted from official reports provided by the Johns Hopkins Coronavirus Resource Center (hereafter JHCRC), starting from January 21st to April 12th, 2020.[16,17] In our multinational modeling of SARS-CoV-2 spreading, we selected 18 countries corresponding to a ~10% of the total number of countries reported in JHCRC. The particular choice of countries was made based on the consistency of the data with the SEIR model ($R^2 > 0.8$), and taking into account a number of confirmed cases larger than one per 100k, with a number of total deaths exceeding 200. Country-level data on population size was extracted from the latest report by the Department of Economic and Social Affairs of the United Nations.[18] The lethality rates were used to assess the status of the COVID-19 outbreak for all the analyzed countries;

we computed the current lethality rate (in %) as the quotient of deaths by COVID-19 divided by confirmed cases in last reported values as obtained from JHCRC. To evaluate the burden that the pandemic lays on the country-level healthcare system, we used the number of hospital beds and acute care beds based on data extracted from the World Bank Database. [19]

2.2   The SEIR model

The total population N is considered to be constant and is partitioned into four disjoint groups, namely: susceptible,$S(t)$; exposed, $E(t)$; infected, $I(t)$, and recovered, $R(t)$, which all are functions of time. The SEIR model considers that individuals who have had contact with an infected person are not themselves infectious instantaneously. Instead susceptible individuals $S(t)$interacting with the infectious population $I(t)$ move to an intermediate compartment $E(t)$ where they develop infectious symptoms at a rate σ[20,21]. The fractions of populations follow the restriction $[S(t) + E(t) + I(t) + R(t)] / N = 1$. A set of coupled ordinary differential equations (ODEs) describing the spreading of the disease and the evolution of the four partitions of N define the SEIR model as follows:

$$S(t) + E(t) + R(t) + I(t) = N(t),$$

Equivalently, dividing all terms by $N(t)$:

$$s(t) + e(t) + i(t) + r(t) = 1,$$

And the system of ODEs is given by:

$$\frac{ds}{dt} = -\beta \, s(t) \, i(t)$$

$$\frac{de}{dt} = \beta \, s(t) \, i(t) - \sigma E$$

$$\frac{di}{dt} = \sigma E - \gamma \, i(t)$$

$$\frac{dr}{dt} = \gamma\, i(t)$$

where, the coefficients to be determined $\beta \geq 0$, $\gamma \geq 0$ and $\sigma \geq 0$ are the effective infectious contact rate of spread, the removal (including deaths and recovered patients) rate, and the incubation rate, respectively. A local epidemic scenario occurs when the number of infected patients increases, i.e., $\left(\frac{di}{dt} > 0\right)$. The basic reproductive number commonly gives the number of secondary infections generated by the first infected individual, $R = \beta/\gamma$.

We solve the coupled system for the SEIR model using SciPy's implementation of the explicit Runge-Kutta method of order 8 (DOP853);[22] this enables us to estimate the transmission rate (β), recovery rate (γ), and the initially exposed population ($E_0 \equiv E(t = 0)$) by fitting the progression of $R(t)$ and $I(t)$. We use the trust-reflective non-linear least-squares method to minimize the residuals between the SEIR model and the data as obtained from JHCRC time series repository. The values $R(t)$ and $I(t)$ were taken as linear combinations of the number of confirmed cases per day $C(t)$, confirmed recoveries per day $R_r(t)$ and confirmed deaths per day $R_d(t)$ with $R(t) = R_r(t) + R_d(t)$ and $I(t) = C(t) - R(t)$. The initial value of susceptible cases taken as $S(0) = N - E(0) - I(0) - R(0)$ with $E(0)$ a result from the fitting. $R(0) = 0$ from the JHCRC for all countries except for China, where 48 recovered cases were identified. For purposes of following the progression at different stages of the epidemic, the values of $I_n = I(t = t_n)$, $R_n = R(t = t_n)$ and $E_n = E(t = t_n)$ can be used as inputs of the SEIR model to determine the evolution of the system.

2.3   SEIR parameters tuning

We calculate the transmission rate (β) by fitting the data in three different stages for the current COVID-19 spread. The parameter β is hard to estimate accurately because the transmission rate can be altered by public policies. Hence, its value can range several orders of magnitude, mostly depending on the government response to contain the outbreak locally. For the Wuhan case, early reports [23] estimated a value of 0.064 at the initial stage of the spread but it further leaps to 0.322 after clinical support was adopted in the modelling. On the other hand, the recovery and latency rates (γ and σ, respectively) are not directed by population controls; according to empirical data it was shown that these rates could range in the order of days (from 2 to 14 days) [24,25]. Because of this, we impose restrictions on the possible values that these fitted parameters can take for predictions. We set the bounds for the fitted parameters as follows: $1/\beta \in [0,10]$, $1/\gamma \in [1\ day, 1\ year]$, $E(t = 0) \in [0, N]$ years, and $1/\sigma = 5.1$ days which is a robust estimation of the average incubation time.[26]

2.4   Piecewise SEIR fitting and projection

To assess COVID-19 dynamics at different epidemic stages, we fitted $I(t)$ and $R(t)$ piecewise and optimized the interval ranges until the value of R^2 was higher than 0.8. To compare changes in epidemic curves where at least two inflection points were identified, we composed a ratio of transmission rates ($Q_n = \beta_{n+1} / \beta_n$), for $n = 1,2,3$ being the interval number (i.e. stages). This estimate can reflect the changes in the COVID-19 epidemic as induced by public health policies. Functional prediction bands $pb(t) = f(x) \pm \delta(x)$ were obtained using $\delta(x) =$(variance)×(inverse of the t-student cumulative distribution).

2.5   Model projections on healthcare burden

A fundamental question for decision makers in countries where the pandemic is developing consists of whether their healthcare systems will be coped with the increasing demands posed by SARS-CoV-2 [27–30]. To add this practical perspective into our analysis, we included: a) the number of hospital beds per capita as reported in official sources, and b) split the number of confirmed COVID-19 cases according to the reported proportions to which 80% of the infected will likely develop mild symptoms only (not requiring hospitalization), 15% would display severe disease (requiring to stay at the hospital) and 5% would reach a critical state (requiring admission to the Intensive Care Unit) [31]. Moreover, we have performed three kinds of projections: first, following the best fit of the SEIR model at the latest stage of the time series, and other two other projections assuming a 50% and 90% decrease in the spreading of the disease, as assessed by the β parameter. As a metric of the level of burden that the epidemic will lay on the health care systems, we define the health care stress as the ratio between the number of infection cases that require hospitalization at the peak of the disease and the number of available beds as reported by the World Bank database.

3.   RESULTS

3.1   Confirmed cases and lethality rates in different countries

The current officially reported lethality for COVID-19 is approximately 3.4% [1]. However, the data fitted indicates a global lethality rate of about 6.17%; this number considers only confirmed cases and could be an overestimation over reporting biases for symptomatic and high-risk cases. A cluster of European countries (UK, Italy, Spain and France) whose calculated lethality rate range from 10.3 to 12.7% and their number of

confirmed cases exceeds 100 cases per 100k is visibly the most affected region (**Figure 1**). Germany, Austria, Switzerland and the US reported >100 cases/100k and their lethality rate similar or lower to than China (4.021%). South Korea, Singapore, Australia and Canada also cluster together, being countries with a lower lethality (< 3%) and confirmed cases lower than 100 per 100k. Brazil, Ecuador and Iran have higher lethality rates than China and their number of confirmed cases is currently still increasing (>10 per 100k). Finally, Japan and Mexico show a number of confirmed cases lower than 10 per 100k; however, the lethality rate for Mexico (6.4%) is higher than that of China. In this study, Mexico shows the lowest number of confirmed cases per 100k habitants. Although this relationship among lethality rate and confirmed cases could be an indication that the pandemic has different dynamics of transmissibility when comparing countries to each other, it must be kept in mind that those countries have responded heterogeneously to COVID-19, both in testing and in the enforcement of public policies.

3.2 Differences in SEIR model fitting across countries

The number of active infections and removed individuals shows an apparent change in both transmission and removal rates, leading to a crossover in February 28, with a transitory decline in the number of active infections, followed by a second crosspoint around March 16, when the number of active infections takes over the number of recoveries (**Figure 2**). These features can be regarded because of a lag between the pandemic's onset across different regions of the globe, which also introduces an additional heterogeneity due to local reporting and containment policies, and demonstrates that SEIR modelling of individual countries is required to address the heterogeneity of the COVID-19 pandemic.

The first crossing point between the infected and removed curves, where the number of recovered cases surpasses active infectious cases is currently not observed for most of the countries studied, with a clear exception for China, South Korea and Austria (see **Supplementary Material**). This is because (1) several countries are still at the early stages of an exponential spread, and (2) an apparent lag between the application of the social distancing measures and the effective change in the rate of the expansion of the epidemic. Because of the inherently high effectiveness of the transmission of SARS-CoV-2, it is expected that the pandemic will extend for months. According to the current trend, we project that the peak in the number of infections worldwide will take place after the third quarter of 2020 (**Figure 2C**). When assessing the projection of COVID-19 cases worldwide, we simulated the same scenarios over different ranges of β (see **Table 1**), indicating the potential of reduced transmission by the initiation of public policy measures. As observed in **Figure 2D**, an earlier peak of infection and a reduced burden on healthcare systems can be achieved by targeting drastic reductions on transmission using social distancing measures.

So far, European countries still lead in the number of confirmed cases and COVID-19 deaths per capita, followed by the United States (**Figure 1, Table 1**). Most countries, with the exception of Japan and Mexico, have managed to slow down the spreading of the epidemic, as revealed by lower β_3 values, as well as $Q_{1,2}$ values<1 (**Figure 3**). In the case of Mexico, the modelling might be limited due to the implementation of a sentinel surveillance model, which samples only a few of all SARS-CoV-2 infections; therefore, the effect of public policy measures might not be accurately estimated using our model [32]. The sharpest declines in spreading rates were observed for Austria, China and South

Korea. Notably, Austria was an early responder of the epidemic, enforcing more stringent restrictions before its neighbors, and as a result it is reportedly now preparing to lift lockdown.[33] Removal rates are also increasing in many countries, which goes along with increases in the estimations of lethality rates, probably it might also be related with a shortening in recovery times or to an increase in recovery rates as the pandemic progresses.

Two distinct values of Q are defined per country to reflect changes in β for the three fitted intervals (**Figure 3**). Values of Q above the dashed line indicate an increase of β from interval n to interval $n + 1$, whereas values below that threshold indicate a reduction in the rate of spread of the disease. Q_1 gives a metric of how contagious contact rate is effectively changing as more reports on new cases become available at the early stages of contagion for each country. With the exception of Switzerland, Austria, Iran China and Japan all studied countries show $Q_1 > 1$. According to available data, the later result indicates that cases at the early onset of the disease are following a consistent reporting trend over different stages of the pandemic. Overall, most of the studied countries show $Q_2 > 1$ with the exception of Japan and Mexico, which recently reported a surge in new cases.

3.3 Model projections on the healthcare burden and effects of containment measures

Assuming that the treatment and prevention of the disease is not affected by other means that are not accounted by the model, then the projections suggest a very strong decrease of the spreading of the disease necessary in most countries to avoid surpassing the healthcare system capacities. Our results also suggest that more stringent measures are still required in most countries; only a few exceptions were observed for Australia,

Austria, China, Iran and South Korea, which could either continue the same or even relax the containment measures to some controlled extent (**Figure 4**). As expected, a monotonic decrease in the number of cases at the peak of the infection I_{peak} is observed with a decrease in the value of β. For 12 out of the 18 countries analyzed, the current trend indicates that probably the number of severe infections will surpass the healthcare capacity (**Figure 4C**). According to our projections, Germany, South Korea, Spain and Switzerland would avoid a saturation of the health case capacity with just a 10% reduction in their β_{latest}. We observe that for four out of the 18 countries analyzed, a 50% reduction on the β_{latest} is enough to avoid the saturation of the available health care infrastructure. We also see that a reduction of 90% in β_{latest} will be a failsafe intervention for all of the analyzed countries. For this level of intervention, it is expected that 11 out of 18 countries (not including Australia, Austria, China and South Korea, which seem to have already passed their infection peaks) reach their peak before the third quarter of 2020. Additionally, it is observed that 5 out of the 18 analyzed will see a significant delay in the peak of the infection for a 90% reduction of β_{latest} extending the course of the epidemic further than the 365 day simulation window, namely, Brazil, Ecuador, Japan, the United Kingdom and the United States.

4.   DISCUSSION

Multinational SEIR modeling of the COVID-19 pandemic demonstrates a significant heterogeneity of the disease across different countries; this suggests that there is no consistent public policy, which will be universally applicable, and that the course of the pandemic will likely be extended given that peak infection rates will occur at different times across countries. Our results suggest that the measures that most countries have

imposed so far have not been sufficient at the time they were enforced to contain the spread. In some scenarios, the likelihood of significantly reducing the number of cases and mortality rates would require significant additional measures to reinforce social distancing. Countries that acted earlier were benefited from the social lockdown; those countries facing a more challenging COVID-19 epidemic should enforce the most stringent measures to avoid overwhelming the capacities of their healthcare systems. Should the situation progress as it has so far, our model projects that the global peak of confirmed cases will be reached by after the third quarter of 2020, though significant fluctuations of this timing are expected among countries. Our piecewise analysis also shows that the rate of spread and impact of COVID-19 has slowed down in most countries, and this fact will likely modify predictions with regard to the size and temporality of the peaks in coming weeks.

Our approach to estimate different stages of the COVID-19 pandemic would likely be better at reflecting changes in both government response and human mobility throughout the course of the epidemic.[34] Notably, our proposed Q_T metric evidences changes in transmission rates and elucidates country-specific changes in the rate of spread after implementation of public policy measures to contain the virus. The social distancing policy has been implemented heterogeneously across countries, most likely to reduce the socio-economic impact on vulnerable populations.[35] Preliminary data from China demonstrates that public health interventions including traffic restriction, social distancing, home quarantine, and universal symptom survey was temporally associated with reduced effective reproduction number of SARS-CoV-2:[36] this is confirmed in our data, whereby China showed significant reductions in β values indicating containment of

SARS-CoV-2 spread. Austria and South Korea are additional examples of public health measures, which have effectively curbed the total number of cases and reduced the impact of the COVID-19 epidemic.[37,38] Additional modelling efforts are required to retrospectively assess the effect of social distancing in curbing the pandemic; nevertheless, our results show that most countries in which social distancing has been implemented show a decrease in the rate of contact transmission. Moreover, the latter likely indicates that those countries with early phases of spread will also experience long-term benefits with more strict policies. Notably, our analyses show that stringent decreases in contact transmission of SARS-CoV-2 could lower the healthcare stress, and decrease the number of total cases. However, depending on the local progress of the pandemic, even a stringent reduction in β could have the effect of either shortening the length of the disease, or prolong the time to reach the peak number of infections. The relevance of decreasing the contagious contact rate stems from the fact that the lethality could be exacerbated once the number of severe cases exceeds the healthcare capacity. [39] With respect to mobility, the citywide quarantine at Wuhan since January 23, 2020, had negligible effect on the epidemic trajectories for the rest of the country.[4] A 50% reduction in inter-city mobility leads to a negligible effect on epidemic dynamics.[4] Therefore, we concluded that accounting for inter-city and inter-country mobility in modeling countries where the epidemic is already present is not of main importance and could be mostly neglected. Given the heterogeneity of the COVID-19 pandemic, our projections should be helpful in understanding the effect of reducing β under different circumstances.

Unlike most SEIR models reported for COVID-19, we fit the data of both confirmed and removed cases, which provides higher confidence in our estimations. However, some

limitations to our analysis can significantly influence the model predictions. For instance, our analysis does not consider the fact that the number of confirmed cases could be underestimated. To address this issue, some countries such as South Korea, Germany and to some extent the US have enacted massive testing which has shown benefits in reporting the spread of the disease,[40] whilst others like Mexico have enacted a sentinel surveillance because of effectiveness in tracking the outbreak in other respiratory virus infections.[41] It is noteworthy that using only the official reported numbers then our model lessens the effect of methodological biases as introduced by different reporting approaches. However, an underestimation in the number of active infections arising from the use of potentially under-sampled data[42] could lead to offsets in our forecasted peak date and in the expected number of peak infections. A second limitation of our approach is that, in contrast with Bayesian models, no prior distribution for constraining the parameter values to a credible interval is used when fitting the models; this might decrease the precision of the estimations in countries with limited data due to a lag in the pandemic's spread.

Because the parameters model were unconstrained to some extent (see Methods), the basic reproductive number obtained for some countries do not precisely agree with other reports for the early onset (see the Supplementary Information). We emphasize the interpretation on the variations in the parameter values between countries and mainly across different stages as defined based on the time intervals that best fitted the SEIR model (i.e., those used in the piecewise methodology).

5.   CONCLUSIONS

Ongoing data collection and epidemiological analysis are essential parts of assessing the impacts of mitigation strategies against SARS-CoV-2 pandemic. In this study, a multinational SEIR modeling considering three different epidemiological stages was used to compare the COVID-19 spreading dynamics. This allowed us to identify the effectiveness of the containment measures across 18 countries and determine the potential burden to the healthcare system, under the assumption of different levels of reduction of the contagious contact rate. Our estimations for the number of active infections place the global peak after the third quarter of 2020; however, several countries may experience a sharper decline within the coming weeks. We found that the peak number of infections monotonically decreases with the decrease in the contagious contact rate. Nevertheless, there is a mixed outcome in the duration of the epidemic since for some countries, the implementation of even more stringent containment measures would lead to a prolonged development of the COVID-19 epidemic. For all studied countries a 90% reduction in the contagious contact rate is required to avoid an excessive burden on the health care capacity. COVID-19 exhibits a heterogeneous spread for the countries studied, a fact that can be related to variations in both timing and efficacy with which the containment measures were implemented in each country. Our results provide informative guidelines to avoid healthcare overcapacity and weigh the long-term effects of more lenient policies.

References

1   Sohrabi C, Alsafi Z, O'Neill N, *et al.* World Health Organization declares global emergency: A review of the 2019 novel coronavirus (COVID-19). *Int J Surg* 2020; **76**: 71–6.

2   Baud D, Qi X, Nielsen-Saines K, Musso D, Pomar L, Favre G. Real estimates of mortality following COVID-19 infection. *Lancet Infect Dis* 2020; published online

March 12. DOI:10.1016/S1473-3099(20)30195-X.

World Health Organization. Coronavirus disease 2019 (COVID-19) Situation Report – 76. World Health Organization, 2020 https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200405-sitrep-76-covid-19.pdf?sfvrsn=6ecf0977_4 (accessed April 5, 2020).

Wu JT, Leung K, Leung GM. Nowcasting and forecasting the potential domestic and international spread of the 2019-nCoV outbreak originating in Wuhan, China: a modelling study. *Lancet* 2020; **395**: 689–97.

Lin Q, Zhao S, Gao D, *et al.* A conceptual model for the outbreak of Coronavirus disease 2019 (COVID-19) in Wuhan, China with individual reaction and governmental action. *Int J Infect Dis* 2020; published online March 4. DOI:10.1016/j.ijid.2020.02.058.

Anderson RM, Heesterbeek H, Klinkenberg D, Hollingsworth TD. How will country-based mitigation measures influence the course of the COVID-19 epidemic? *Lancet* 2020; **395**: 931–4.

Remuzzi A, Remuzzi G. COVID-19 and Italy: what next? *Lancet* 2020; published online March 13. DOI:10.1016/S0140-6736(20)30627-9.

Tanne JH, Hayasaki E, Zastrow M, Pulla P, Smith P, Rada AG. Covid-19: how doctors and healthcare systems are tackling coronavirus worldwide. *BMJ* 2020; **368**: m1090.

Zhao S, Lin Q, Ran J, *et al.* Preliminary estimation of the basic reproduction number of novel coronavirus (2019-nCoV) in China, from 2019 to 2020: A data-driven analysis in the early phase of the outbreak. *Int J Infect Dis* 2020; **92**: 214–7.

Roda WC, Varughese MB, Han D, Li MY. Why is it difficult to accurately predict the COVID-19 epidemic? *Infect Dis Model* 2020; **5**: 271–81.

Read JM, Bridgen JR, Cummings DA, Ho A, Jewell CP. Novel coronavirus 2019-nCoV: early estimation of epidemiological parameters and epidemic predictions. *medRxiv* 2020; published online Jan 24. DOI:10.1101/2020.01.23.20018549.

Liu T, Hu J, Kang M, *et al.* Transmission Dynamics of 2019 Novel Coronavirus (2019-nCoV). *SSRN Journal* 2020. DOI:10.2139/ssrn.3526307.

Rachah A, Torres DFM. Mathematical modelling, simulation, and optimal control of the 2014 ebola outbreak in west africa. *Discrete Dyn Nat Soc* 2015; **2015**: 1–9.

Kermack WO, McKendrick AG. A Contribution to the Mathematical Theory of Epidemics. *Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences* 1927; **115**: 700–21.

Arenas A, Cota W, Gomez-Gardenes J, *et al.* A mathematical model for the

spatiotemporal epidemic spreading of COVID19. *medRxiv* 2020; published online March 23. DOI:10.1101/2020.03.21.20040022.

GitHub - CSSEGISandData/COVID-19: Novel Coronavirus (COVID-19) Cases, provided by JHU CSSE. https://github.com/CSSEGISandData/COVID-19 (accessed April 7, 2020).

Dong E, Du H, Gardner L. An interactive web-based dashboard to track COVID-19 in real time. *Lancet Infect Dis* 2020; published online Feb 19. DOI:10.1016/S1473-3099(20)30120-1.

United Nations Population Division | Department of Economic and Social Affairs. https://www.un.org/en/development/desa/population/index.asp (accessed April 7, 2020).

Hospital beds (per 1,000 people) | Data. https://data.worldbank.org/indicator/sh.med.beds.zs (accessed April 7, 2020).

Hethcote HW. The Mathematics of Infectious Diseases. *SIAM Rev* 2000; **42**: 599–653.

Hethcote HW. THE BASIC EPIDEMIOLOGY MODELS: MODELS, EXPRESSIONS FOR r_0 , PARAMETER ESTIMATION, AND APPLICATIONS. In: Mathematical understanding of infectious disease dynamics. WORLD SCIENTIFIC, 2008: 1–61.

Hairer E, Nørsett SP, Wanner G. Solving ordinary differential equations I Nonstiff Problems. Berlin, Heidelberg: Springer Berlin Heidelberg, 1993 DOI:10.1007/978-3-540-78862-1.

Zhao S, Chen H. Modeling the epidemic dynamics and control of COVID-19 outbreak in China. *Quant Biol* 2020; 1–9.

Symptoms of Coronavirus | CDC. https://www.cdc.gov/coronavirus/2019-ncov/symptoms-testing/symptoms.html?CDC_AA_refVal=https%3A%2F%2Fwww.cdc.gov%2Fcoronavirus%2F2019-ncov%2Fabout%2Fsymptoms.html (accessed April 7, 2020).

World Health Organization. Novel Coronavirus(2019-nCoV)Situation Report - 7. World Health Organization, 2020 https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200127-sitrep-7-2019--ncov.pdf (accessed April 10, 2020).

Lauer SA, Grantz KH, Bi Q, *et al.* The Incubation Period of Coronavirus Disease 2019 (COVID-19) From Publicly Reported Confirmed Cases: Estimation and Application. *Ann Intern Med* 2020; published online March 10. DOI:10.7326/M20-0504.

Ji Y, Ma Z, Peppelenbosch MP, Pan Q. Potential association between COVID-19

mortality and health-care resource availability. *Lancet Glob Health* 2020; **8**: e480.

Legido-Quigley H, Asgari N, Teo YY, *et al.* Are high-performing health systems resilient against the COVID-19 epidemic? *Lancet* 2020; **395**: 848–50.

Willan J, King AJ, Jeffery K, Bienz N. Challenges for NHS hospitals during covid-19 epidemic. *BMJ* 2020; **368**: m1117.

Legido-Quigley H, Mateos-García JT, Campos VR, Gea-Sánchez M, Muntaner C, McKee M. The resilience of the Spanish health system against the COVID-19 pandemic. *Lancet Public Health* 2020; published online March 18. DOI:10.1016/S2468-2667(20)30060-8.

World Health Organization. Coronavirus disease 2019 (COVID-19)Situation Report – 46. World Health Organization, 2020 https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200306-sitrep-46-covid-19.pdf?sfvrsn=96b04adf_2 (accessed April 6, 2020).

WHO | Sentinel Surveillance. https://www.who.int/immunization/monitoring_surveillance/burden/vpd/surveillance_type/sentinel/en/ (accessed April 14, 2020).

Mahbubani R. Austria gets ready to lift coronavirus lockdown as early as next week. Business Insider. 2020; published online April 9.

Kraemer MUG, Yang C-H, Gutierrez B, *et al.* The effect of human mobility and control measures on the COVID-19 epidemic in China. *Science* 2020; published online March 25. DOI:10.1126/science.abb4218.

Lewnard JA, Lo NC. Scientific and ethical basis for social-distancing interventions against COVID-19. *Lancet Infect Dis* 2020; published online March 23. DOI:10.1016/S1473-3099(20)30190-0.

Pan A, Liu L, Wang C, *et al.* Association of Public Health Interventions with the Epidemiology of the COVID-19 Outbreak in Wuhan, China. *JAMA* 2020; published online April 10. DOI:10.1001/jama.2020.6130.

Park SW, Sun K, Viboud C, Grenfell BT, Dushoff J. Potential roles of social distancing in mitigating the spread of coronavirus disease 2019 (COVID-19) in South Korea. *medRxiv* 2020; published online March 30. DOI:10.1101/2020.03.27.20045815.

Bendtsen Cano O, Cano Morales S, Bendtsen C. COVID-19 Modelling: the Effects of Social Distancing. *medRxiv* 2020; published online March 30. DOI:10.1101/2020.03.29.20046870.

Real-time data show virus hit to global economic activity | Financial Times. https://www.ft.com/content/d184fa0a-6904-11ea-800d-da70cff6e4d3 (accessed April

14, 2020).

40 In one Italian town, we showed mass testing could eradicate the coronavirus | Andrea Crisanti and Antonio Cassone | Opinion | The Guardian. https://www.theguardian.com/commentisfree/2020/mar/20/eradicated-coronavirus-mass-testing-covid-19-italy-vo (accessed April 14, 2020).

41 Cortes-Alcala R, Dos Santos G, DeAntonio R, *et al.* The burden of influenza A and B in Mexico from the year 2010 to 2013: An observational, retrospective, database study, on records from the Directorate General of Epidemiology database. *Hum Vaccin Immunother* 2018; **14**: 1890–8.

42 Lachmann A. Correcting under-reported COVID-19 case numbers. *medRxiv* 2020; published online March 18. DOI:10.1101/2020.03.14.20036178.

Author Contributions

Source code

The Python code for this work is available at https://github.com/erickmartinez/covid19

Funding Sources

No funding received.

ORCID

Erick Martinez-Loran: 0000-0003-3841-4981
J. Jesús Naveja: 0000-0001-8640-6690
Omar Y. Bello-Chavolla: 0000-0003-3093-937X
Flavio F. Contreras-Torres: 0000-0003-2375-131X

TABLES

Table 1. COVID-19 fitting results to the SEIR model using piecewise analysis for each evaluated country.

Country	First stage				Second stage				Third stage				Qt_1	Qt_2
	β1	γ1	E1	R^2	β2	γ2	E2	R^2	β3	γ3	E3	R^2		
United Kingdom	0.253 (0.173-0.370)	0.013 (0.004-0.045)	0 (0-N)	0.910	0.546 (0.545-0.547)	0.014 (0.014-0.014)	35 (35-35)	0.994	0.167 (0.157-0.177)	0.016 (0.014-0.018)	12290 (11112-13595)	0.995	2.157	0.305
France	0.026 (0.015-0.046)	0.022 (0.018-0.027)	6 (5-8)	0.824	0.579 (0.578-0.58)	0.005 (0.005-0.005)	171 (171-172)	0.991	0.172 (0.158-0.188)	0.040 (0.036-0.043)	13586 (11445-16126)	0.977	22.066	0.298
United States	0.026 (0.019-0.037)	0.016 (0.013-0.019)	8 (7-9)	0.924	0.686 (0.686-0.686)	0.005 (0.005-0.005)	73 (73-73)	0.996	0.133 (0.124-0.143)	0.012 (0.01-0.014)	134660 (124437-145725)	0.996	26.057	0.194
Iran	2.084 (1.867-2.325)	0.115 (0.099-0.133)	27 (18-42)	0.994	0.166 (0.165-0.166)	0.055 (0.055-.055)	4920 (4908-4933)	0.987	0.012 (0.000-2.456)	0.115 (0.108-0.124)	15315 (9980-23502)	0.976	0.079	0.073
Brazil	0.660 (0.586-0.744)	0.003 (0.000-0.095)	2 (1-3)	0.990	0.912 (0.905-0.919)	0.006 (0.005-0.007)	55 (50-59)	0.996	0.183 (0.172-0194)	0.008 (0.006-0.011)	2452 (2143-2806)	0.995	1.381	0.201
South Korea	0.1 (0.072-0.138)	0.015 (0.009-0.026)	10 (8-13)	0.898	0.238 (0.238-0.238)	0.003 (0.003-0.003)	1888 (1885-1892)	0.979	0.001 (0.015-0.044)	0.044 (0.042-0.045)	1669 (1244-2240)	0.963	2.388	0.019
Ecuador	0.042 (0.023-0.079)	0.003 (0-0.017)	9 (8-11)	0.990	0.451 (0.449-0.453)	0.006 (0.006-0.006)	526 (524-528)	0.972	0.161 (0.138-0.187)	0.013 (0.009-0.019)	758 (492-1168)	0.979	10.679	0.356
Germany	0.029 (0.018-0.046)	0.003 (0.001-0.01)	12 (11-13)	0.983	0.446 (0.446-0.446)	0.003 (0.003-0.003)	400 (400-400)	0.991	0.075 (0.063-0.091)	0.050 (0.047-0.054)	43680 (33319-449790)	0.963	15.51	0.169
Spain	0.119 (0.095-0.15)	0.025 (0.017-0.037)	0 (0-1)	0.920	0.524 (0.524-0.524)	0.029 (0.029-0.029)	399 (399-399)	0.988	0.072 (0.067-0.077)	0.055 (0.054-0.056)	49072 (46722-51540)	0.996	4.396	0.137
Mexico	1.579E-16 (1.58E-16-1.579E-16)	0.034 (0.019-0.062)	7 (6-8)	0.883	0.280 (0.279-0.280)	0.007 (0.007-0.007)	134 (134-135)	0.993	0.321 (0.180-0.571)	0.084 (0.061-0.115)	245 (4-15573)	0.801	1.77E 15	1.146
Australia	0.055 (0.039-0.077)	0.018 (0.014-0.023)	7 (6-8)	0.973	0.420 (0.419-0.420)	0.011 (0.011-0.011)	25 (25-25)	0.995	0.006 (0.001-0.023)	0.021 (0.019-0.023)	3229 (2978-3501)	0.993	7.677	0.014

Austria	0.746 (0.719-0.774)	0.005 (0.002-0.009)	7 (6-8)	0.999	0.329 (0.328-0.329)	0.003 (0.003-0.003)	1156 (1154-1158)	0.996	1.31E-10 (0-N)	0.043 (0.040-0.046)	6534 (5863-7281)	0.974	0.441	3.67E-10
Canada	0.031 (0.023-0.042)	0.014 (0.011-0.017)	5(4-5)	0.937	0.521 (0.518-0.524)	0.005 (0.004-0.005)	0 (0-3.15E7)	0.986	0.143 (0.129-0.159)	0.039 (0.036-0.043)	5875 (5213-6622)	0.986	16.753	0.274
Singapore	0.122 (0.101-0.147)	0.036 (0.03-0.043)	19 (16-23)	0.912	0.193 (0.031-1.207)	0.003 (0-15734.204)	31 (6-149)	0.975	0.158 (0.151-0.166)	0.024 (0.023-0.026)	103 (88-122)	0.992	1.584	0.817
Switzerland	0.633 (0.545-0.736)	0.004 (0-0.081)	43 (34-56)	0.985	0.349 (0.346-0.352)	0.005 (0.004-0.005)	1597 (1575-1620)	0.995	0.031 (0.020-0.047)	0.050 (0.047-0.054)	9546 (8239-11060)	0.966	0.551	0.089
Japan	0.166 (0.121-0.227)	0.007 (0.002-0.031)	7 (5-10)	0.936	0.14 (0.139-0.141)	0.019 (0.019-0.019)	92 (91-92)	0.994	0.180 (0.176-0.184)	0.016 (0.014-0.018)	0 (0-N)	0.994	0.843	1.289
Global	0.866 (0.768-0.976)	0.017 (0.002-0.121)	0 (0-N)	0.907	0.003 (0.000-0.024)	0.040 (0.038-0.041)	25092 (21822-28852)	0.979	0.124 (0.117-0.132)	0.028 (0.026-0.030)	328676 (301318-358517)	0.992	0.004	36.462
Italy	0.011 (0.004-0.032)	0.003 (0.001-0.008)	1 (1-1)	0.982	0.415 (0.415-0.415)	0.035 (0.035-0.035)	947(947-947)	0.995	0.064 (0.060-0.068)	0.027 (0.026-0.028)	42634 (40231-45182)	0.992	37.203	0.155
China	0.469 (0.425-0.518)	0.013 (0.008-0.022)	3441 (2902-4079)	0.993	1E-14 (0.000-*)	0.033 (0.033-0.034)	38804 (38196-39421)	0.984	0 (0.005-0.084)	0.082 (0.083-0.083)	3995 (3785-4215)	0.999	0.000	0.000

Table 2. Effect of diminishing beta in peak date and peak cases for the evaluated countries using the SEIR model as well as piecewise model fit. Cases and total beds are standardized per 100k habitants.

Country	Recovery rate	Lethality rate	Peak date	Peak cases*	Peak date (0.90β)	Peak cases (0.90β)*	Peak date (0.50β)	Peak cases (0.50β)	Peak date (0.10β)	Peak cases (0.10β)*	Beds*
United Kingdom	0.73	12.47	20/07/2020	43802739	29/07/2020	40591113	05/10/2020	30800646	12/04/2021	48689	280
France	20.54	10.78	12/07/2020	54085747	25/07/2020	21231960	28/10/2020	9975739	12/04/2020	71927	650
United States	5.94	3.96	04/08/2020	266212175	11/08/2020	209272719	25/10/2020	161850887	12/04/2021	560983	290
Iran	61.23	6.24	05/04/2020	34887	12/04/2020	27779	12/04/2020	27779	12/04/2020	27779	150
Brazil	0.73	5.66	04/08/2020	174282221	13/08/2020	161299611	23/10/2020	140743932	13/04/2021	374534	220
South Korea	70.09	2.03	12/04/2021	17687	12/04/2020	9620	12/04/2020	9620	12/04/2020	9620	1150
Ecuador	6.71	4.46	25/07/2020	14141657	04/08/2020	11473186	14/10/2020	9115332	12/04/2021	21909	150
Germany	47.16	2.36	07/10/2020	69493925	10/04/2021	2495398	20/04/2020	80197	16/04/2020	75211	830
Spain	37.39	10.31	01/11/2020	9110718	12/04/2021	474852	18/04/2020	83053	16/04/2020	78857	300
Mexico	42.00	6.47	30/06/2020	108231568	24/07/2020	30834404	10/11/2020	12300244	12/04/2020	4930	150
Australia	28.59	0.95	26/03/2020	2625	23/04/2020	4448	22/04/2020	4398	22/04/2020	4353	380
Austria	50.10	2.50	07/04/2020	7552	17/04/2020	9711	17/04/2020	9711	17/04/2020	9711	760
Canada	29.31	2.93	07/08/2020	31076420	04/09/2020	10395885	04/02/2021	3825390	17/04/2020	13008	270
Singapore	22.11	0.31	31/07/2020	5228560	11/08/2020	2698684	09/11/2020	1701218	12/04/2020	2148	240
Switzerland	49.97	4.35	26/03/2021	4664165	18/04/2020	17772	16/04/2020	17369	16/04/2020	17085	470
Japan	11.29	1.60	07/08/2020	74401596	18/08/2020	78164736	08/11/2020	60829798	12/04/2021	11610	1340
global	22.83	6.17	04/10/2020	2025863750	05/10/2020	2740579323	19/03/2021	1317196421	20/04/2020	666098	8390
Italy	21.87	12.72	31/01/2021	6545277	10/01/2021	9330226	13/04/2021	291549	22/04/2020	61993	340
China	93.77	4.02	03/03/2020	61142	12/04/2020	19955	12/04/2020	19955.3371	12/04/2020	19955	420

Figure 1. Number of confirmed COVID-19 cases per 100k habitants versus the lethality rate (in %) for different countries. The lethality rate was calculated as the quotient of deaths by COVID-19 divided by the confirmed cases according to the report published on April 12, 2020, from JHCRC.

Figure 2. Fit of the SEIR model to worldwide data. In all cases, the x-axis is the period of time in format mm/dd. Upper left: total infections, recoveries, deaths and infected cases as per official reports from JHCRC; upper right: model fit and confidence intervals for both infected and removed (shadow intervals indicate the piecewise fitting for obtaining the effective infectious contact rate of spread for each period of time); bottom left: projections by the best-fit model, bottom right: solid lines represent splitting the projected curve into severity categories, whereas dashed lines represent the projection with 90% reduction of current spreading rate. Figure 2a, shows the global picture of the progress of COVID19 epidemic as reported by JHCRC. Indicated in solid orange circles is the total number of confirmed cases per day $C(t)$, the number of confirmed recoveries $R_r(t)$ and deaths $R_d(t)$ are indicated in open green squares and open red triangles respectively. The active number of infections at time t estimated as $I(t) = C(t) - R_r(t) - R_d(r)$ is indicated by open blue circles. Figure 2b shows the the selection of intervals used to fit the mode globally: (a) from the date of the first detected case to the point 1 indicated on the infections curve, (b) on the range indicated between the arrows labeled as 2 on the infections curve and (c) from the point labeled as 3 on the infections curve to April 12th 2020.

Figure 3. Comparison of the fitted values for the SEIR model for different countries. From top to bottom (a) total confirmed cases, recoveries and deaths per of SEIR, in orange, green and red respectively. (b) Fitted β(infectious rate) for each of the different intervals (c) Fitted removal rate (γ) for each of the different intervals (e) Ratio $Q_n = \beta_{n+1}/\beta_n$ between the different intervals, indicated as a measure of the effectiveness of the containment measures. For panels (b) through (d) missing values indicate that only two intervals were identified in the corresponding dataset. The scale of the vertical axis in (d) was adjusted to a maximum of 100. Nonetheless, Mexico overpasses this limit drastically, which could be attributed to a low early detection rate due to the implementation of a sentinel model for epidemiological surveillance (see Table 1).

Figure 4. Projected infections for different countries for different values of the contagious contact rate β related to the effectiveness of the containment: cyan: current value of β estimated by our model, blue: 0.9β, purple 0.5β and magenta 0.1β. (a) Shows the estimated time until the local peak of the epidemic has been reached. The dashed line indicates the maximum simulation time. (b) Shows the total number of cases at the local peak of the epidemic. (c) Shows the health care stress, defined as the number of cases at the peak requiring hospitalization (20% of the total number of cases) divided by the number of available hospital beds as reported in the World Bank database.