Fatigue in perinatal stroke is associated with resting-state functional connectivity

Wrightson, J.G [1,2], Carlson, H.L [1,3,4] Condliffe, E.G [1,2,3,4] and Kirton, A [1,2,3,4].

1. Hotchkiss Brain Institute, University of Calgary, Calgary, AB, Canada
2. Department of Clinical Neurosciences, Cumming School of Medicine, University of Calgary, Calgary, AB, Canada
3. Alberta Children's Hospital Research Institute, Calgary, AB Canada
4. Department of Pediatrics, Cumming School of Medicine, University of Calgary, Calgary, AB, Canada

Corresponding Author: Dr. Adam Kirton
Email: Adam.Kirton@Albertahealthservices.ca
Phone: +1.403.955.7816

Abstract

Fatigue is prevalent in youth with perinatal stroke, but the causes are unclear. Predictive coding models of adult post-stroke fatigue suggest that fatigue may arise from dysfunction in predictive processing networks. To date, the association between fatigue and neural network connectivity in youth with perinatal stroke has not been examined. The present study examined the association between fatigue and the functional connectivity of predictive processing neural networks, measured using resting-state functional magnetic resonance imaging, in individuals with perinatal stroke. Participants who reported experiencing fatigue had weaker functional connectivity between the non-lesioned middle frontal and supramarginal gyri and between the non-lesioned intracalcarine cortex and the lesioned paracingulate cortex. In contrast, participants reporting fatigue had stronger functional connectivity between the lesioned inferior temporal gyrus and non-lesioned insula. These results suggest that fatigue in youth with hemiparetic cerebral palsy caused by perinatal stroke is associated with the functional connectivity of hubs previously associated with predictive processing and fatigue. These results suggest potential cortical and behavioral targets for the treatment of fatigue in individuals with perinatal stroke.

Keywords: Cerebral palsy, Resting-state fMRI, Predictive coding, Active inference

Introduction

Perinatal stroke is the leading cause of hemiparetic cerebral palsy and results in lifelong physical disability [1,2]. Fatigue is common in individuals with hemiparetic cerebral palsy, including those with perinatal stroke [3–6]. Fatigue significantly reduces health-related quality of life in individuals with cerebral palsy [7–9], and reducing fatigue is caregivers' highest priority [10]. However, the mechanisms which give rise to fatigue are not well understood, and consequently, there are few evidence-based treatments.

Predictive coding models of fatigue suggest that insult to the central nervous system causes an incongruity between the efferent sensory predictions encoded by the brain's generative model of the body and the afferent sensory prediction errors that arise from an action [11,12]. It has been proposed that dysfunction in the predictive processing networks gives rise to fatigue [11]. Predictive processing theoretical frameworks suggest that the brain contains a generative model of the body-state [13]. Within these frameworks, fatigue arises from an ongoing and unresolvable discrepancy

between the brain's prediction of the body-state and the afferent sensory information signalling the actual-body state (the prediction) [11,12,14]. These prediction errors signal a failure of the brain to predict the body state, i.e. an inability to maintain homeostasis. This ongoing discrepancy between predicted and actual body state is consciously perceived as the feeling of fatigue, which acts as a signal for allostatic behavioral adjustment: rest. Two predictive coding accounts of fatigue have been proposed; the first suggests that fatigue arises from a failure to attenuate from proprioceptive or exteroceptive sensory prediction errors, specifically during motor actions [15]. An opposing account posits that fatigue arises through the metacognitive awareness of an ongoing discrepancy between predicted and actual interoceptive information, processed in an interoceptive predictive processing network [12]. These two accounts are often presented as disparate, assigning fatigue to either interoceptive or exteroceptive and proprioceptive predictive processing (see Kuppuswamy 2021; Manjaly et al. 2019 for reviews). However, these systems likely interact to predict proprioceptive and interoceptive signals in the pursuit of goal-states [16] and thus are both likely to contribute to the perception of fatigue [17].

Unlike in adult stroke, where pre-stroke conditions and post-stroke responses are frequently assumed to contribute to the subsequent development of fatigue [18,19], fatigue in perinatal stroke is likely to be a consequence of the abnormal development of the brain [6]. The interoceptive predictive processing account identifies several regions in the brain that may compose a 'fatigue network,' including the anterior insula, the anterior cingulate cortex and frontal/prefrontal areas [12]. The exteroceptive and proprioceptive predictive coding account of fatigue predicts that in addition to the insula, networks comprised of sensorimotor, visual, and attentional hubs also contribute to fatigue [15]. There is emerging evidence from resting-state functional magnetic resonance imaging studies that connectivity between these areas is associated with fatigue perception [20–22]. Proprioception is impaired in individuals with perinatal stroke and is shown to be related to connectivity of sensory networks [23,24], supporting that the connectivity in these networks may contribute to fatigue. Following the recent suggestion that fatigue likely arises from *both* interoceptive and proprioceptive predictive processing [17], many of the hubs whose connectivity is associated with fatigue have also been suggested to be hubs within a global predictive processing network [25–27] At present, the relationship between fatigue in individuals with perinatal stroke and the functional connectivity of these networks has not been studied.

The aim of the present study was to examine the associations between fatigue and functional brain connectivity in individuals with perinatal stroke and compare this to the functional

connectivity of typically developed youth. We have previously reported that corticospinal excitability was associated with fatigue severity in individuals with perinatal stroke [6]. Thus, we hypothesized that the functional connectivity of the networks involved in proprioceptive and exteroceptive predictive processing would be associated with fatigue severity.

Results

Seventy-five participants were identified through the Alberta Perinatal Stroke Project database. Of these, eleven were ineligible due to incomplete data, and four were removed due to excessive head motion in the scanner. The data for 60 stroke participants and 59 TDC were included in the analysis. Participant demographics and outcomes are summarized in Table 1. Lesion overlay maps for AIS participants are shown in Figure 1. The median age for the full sample was 11 years, range 7-19 years, and 35% were female. Forty-five percent (27/60) of the participants with perinatal stroke reported experiencing problems with fatigue (PEDSQL-CP fatigue score ≤68.75) and were assigned to the "Fatigued" group [6] and the remaining 33 participants were assigned to the "Not Fatigued" group. There was no difference in age between the stroke groups. TDC were significantly older than the "Not Fatigued" group (see Table 1). There were no other differences in demographics between groups. (p>0.05).

Table 1 Here please

Figure 1 Here please

Fisher-transformed correlation coefficients for ROI-to-ROI connections, age, PSOM score and stroke type (AIS or PVI) were entered into the model. The model selected three FC values: FC between the non-lesioned supramarginal gyrus and middle frontal gyrus, the non-lesioned insula and lesioned inferior temporal gyrus, and the lesioned paracingulate cortex and non-lesioned intracalcarine cortex. These connections are shown in Figure 2.

Figure 2. Here please

Associations between Fatigue and FC

While associations between fatigue and FC were strong (as identified previously), the directionality of such associations differed by ROI-ROI pair. In participants with perinatal stroke, fatigue (PEDSQL score) was positively associated with FC between the non-lesioned middle frontal gyrus and supramarginal gyrus (β=19.6, SE=6.4, p=0.002, Figure 3A) and between the lesioned paracingulate gyrus and the non-lesioned intracalcarine cortex (β=31.6, SE=12.6, p=0.015, Figure 3C). In contrast, fatigue was negatively associated with FC between the non-lesioned insula and the lesioned inferior temporal gyrus (β=−42.3, SE=7.7, p<0.001, Figure 3B). Age, stroke type or sensorimotor function (PSOM) were not identified as predictors.

Figure 3. Here please

Differences in functional connectivity between groups

Results from the Welch's ANOVA for the Fisher-transformed correlations selected by the elastic net regression for fatigued, not fatigued and TDC participants are shown in Table 2 and Figure 4. Welch's ANOVA revealed a difference between groups for all three FC ROI pairs. FC between the non-lesioned middle frontal gyrus and supramarginal gyrus was stronger in not fatigued participants compared to fatigued participants ($p_{(fdr)}$=0.015, d=0.67), and TDC ($p_{(fdr)}$=*0.002*, d=1.88), whilst it was stronger in fatigued participants than in TDC ($p_{(fdr)}$=0.002, d=1.14). The negative FC between the non-lesioned insula and lesioned temporal gyrus was stronger in not fatigued participants compared to fatigued participants ($p_{(fdr)}$=0.002, d=0.88) and TDC ($p_{(fdr)}$=0.002, d=1.12) but not different between fatigued participants and TDC ($p_{(fdr)}$=0.273, d=0.35). The negative FC between the non-lesioned intracalcarine cortex and the lesioned paracingulate cortex was stronger in fatigued participants compared to not fatigued ($p_{(fdr)}$=0.045, d=0.56), and TDC participants ($p_{(fdr)}$=0.002, d=0.94), and not different between not fatigued and TDC participants ($p_{(fdr)}$=0.149, d=0.30).

Table 2 here please.
Figure 4. Here please

Discussion

The aim of the present study was to examine potential associations between fatigue and resting-state functional connectivity in the brains of individuals with perinatal stroke. Compared to participants who were not fatigued, those who reported problems with fatigue had weaker connectivity between the non-lesioned middle frontal gyrus and posterior supramarginal gyrus, and between the non-lesioned intracalcarine cortex and lesioned paracingulate gyrus. In contrast, functional connectivity between the non-lesioned anterior insula cortex and the lesioned inferior temporal gyrus was stronger in individuals experiencing fatigue. This study is the first to explicitly test the relationship between functional brain connectivity and perceived fatigue in young people with perinatal stroke. These results suggest that fatigue is associated with the functional connectivity of the interoceptive and salience prediction networks in individuals with perinatal stroke.

Here, we provide evidence showing that fatigue in perinatal stroke is associated with altered connectivity between hubs in previously identified global predictive coding networks [25–27]. These include hubs frequently linked to proprioceptive and interoceptive processing and allostatic control [28–30]. Within predictive coding frameworks, fatigue is suggested to arise from an ongoing and unresolvable discrepancy between the brain's prediction of the body-state and the afferent sensory information signalling the actual-body state, the prediction errors [11,12,14]. Unresolvable prediction errors signal a failure for the brain to predict the body state, i.e. an inability to maintain allostatic control, consciously perceived as the feeling of fatigue [12]. We found that individuals experiencing more fatigue had weaker FC between the non-lesioned supramarginal gyrus and middle frontal gyrus compared to those who did not experience fatigue. Recent evidence suggests that these areas are structurally connected via the superior longitudinal fasciculus [31]. Perhaps surprisingly, the weaker connectivity seen in fatigued participants reflected the pattern in typically developed participants, where there was virtually no functional connectivity between these areas. In adults and children with stroke, the supramarginal gyrus and middle frontal gyrus have been related to successful proprioceptive processing [32–34], while individuals with perinatal stroke may have impaired proprioceptive function [23,24]. Within the predictive processing framework, both areas are associated with both prediction formation and error [26,27]. When comparing participants who did not experience fatigue with TDC, we speculate that stronger functional connectivity between these areas seen in not fatigued participants represents possible compensatory neural plastic

alterations related to impaired proprioceptive prediction and prediction error processing.
Kuppuswamy [11] suggests that fatigue arises from dysfunction in the systems involved in
processing proprioceptive prediction errors, suggesting that the weaker functional connectivity, or
an inefficient compensation mechanism, between these areas may be related to the increased
perceptions of fatigue.

A similar pattern can be observed in the functional connectivity between lesioned inferior temporal
gyrus and non-lesioned insula, wherein not fatigued participants had stronger (negative)
functional connectivity than fatigued participants, who more closely resembled typically developed
participants. The insula, cingulate and inferior temporal gyrus are suggested to be hubs within a
neural network that supports interoceptive processing and allostasis [30]. An alternative predictive
processing account of fatigue suggests that fatigue arises from continued and unresolvable
interoceptive, not proprioceptive, prediction errors which signal failures in allostatic control [12].
Again, when interpreted through the predictive processing theoretical framework, the stronger
connectivity seen in not fatigued participants suggests adaptive plasticity supporting interoceptive
predictive processing. The present data indicate the involvement of both proprioceptive and
interoceptive predictive processing in the development of fatigue in individuals with perinatal
stroke. Although the proprioceptive and interoceptive predictive processing accounts of fatigue
are often contrasted [14,15], it is more likely that fatigue can arise from ongoing and unreconcilable
proprioceptive and interoceptive prediction errors, which both signal challenges to allostatic
control [17]. Since fatigue is believed to be the affective response signalling ongoing unresolvable
prediction errors and thus failure to exert allostatic control, these results support the predictive
processing accounts of fatigue [14,15]. At present, there is considerable uncertainty about how
predictive processing networks function and much more evidence of the strength and direction of
relationships is required before this suggestion can be genuinely considered.

Somewhat surprisingly, and in contrast to our hypothesis, we did not find any associations
between fatigue and the connectivity between primary or secondary sensorimotor areas. This is
in contrast to an earlier analysis of this data, where we used a theory-driven approach and placed
seeds within these areas [35], as well as our previous study where we used transcranial magnetic
stimulation to probe the association between fatigue and corticomotor excitability [6] and previous
evidence of altered FC within the sensorimotor network associated with motor function [36–38].
However, this is not to say that functional connectivity within sensorimotor areas is not associated
with fatigue, indeed we have shown that it is [29]. Instead, what the current data show is that of

more relevance to the perception of fatigue in individuals with perinatal stroke is the connectivity
of hubs within predictive processing networks.
70
This study has some limitations which should be acknowledged. We have interpreted our results
using the predictive coding theories of the brain and affect. While our data support these models,
we did not measure the activity of this network during an action [39], nor did we have a measure of
predictive coding during performance [40]. These are the necessary next steps in examining fatigue
in youth with perinatal stroke. Although we applied post-hoc corrections for head movement, fMRI
remains a challenging technique in a pediatric population, and we were required to remove
several participants from our sample. The replication of these effects in adults with perinatal stroke
may overcome this limitation and provide a method to replicate and extend these results. Our
sample was composed of participants capable of completing an MRI, which is necessarily the
less disabled and higher functioning portion of the perinatal stroke population.

Conclusion

We found that fatigue in individuals with perinatal stroke was associated with altered functional
connectivity of a global predictive processing network. These findings provide interesting avenues
for both future research and therapeutic interventions. The effects of emerging therapies, such as
novel training paradigms and non-invasive brain stimulation, on both neural functional connectivity
and fatigue are warranted.
87

Method and Materials

Participants

Participants were recruited via the Alberta Perinatal Stroke Project [41] and the Healthy Infants and
Children Clinical Research Program (www.hiccupkids.ca) research cohorts at the Alberta
Children's Hospital. Inclusion criteria for cases were age 6-19 years; term birth; magnetic
resonance imaging (MRI)-confirmed unilateral perinatal arterial ischemic stroke (AIS) or
periventricular venous infarction (PVI), symptomatic hemiparetic cerebral palsy including
Pediatric Stroke Outcome Measure >0.5; Manual Ability Classification Scale levels I-IV. Inclusion
criteria for typically developed controls (TDC) were right-handed, ages 6-19 years, and no
neurological conditions, medications, or MRI contraindications. Participants and parents provided

written informed assent/consent. Experiments were carried out with the approval of the Research

Ethics Board at the University of Calgary and in accordance with the declaration of Helsinki [42].

Fatigue

Self-reported fatigue was assessed in participants with perinatal stroke using the fatigue subscale

of the Pediatric Quality of Life Inventory 3.0 cerebral palsy module (PEDSQL-CP), a validated

measure of quality of life in children with CP [43]. The fatigue subscale consists of four statements:

"I feel tired," "I feel physically weak (not strong)," "I rest a lot," and "I don't have enough energy to

do things that I like to do." Individuals were asked to give a rating of 0-4 for each statement if they

never (0), almost never (1), sometimes (2), often (3), or almost always (4) experienced a problem

with the four areas of fatigue highlighted in the statements. Parents assisted children under eight

years with the rating using the PEDSQL 3.0 CP module parent report for young children. The

PEDSQL scores for fatigue were reverse scored and linearly transformed to a 0 to 100 scale [43].

The average score across all categories in each subscale was calculated. Lower scores in each

subscale represent higher levels of fatigue. Participants were classified as experiencing fatigue if

they scored $\leq$68.75, indicating a problem with at least two or more of the categories of fatigue, or

if they scored 4 ("almost always" a problem) for one of the categories [5].

MRI collection

MRI sequences were acquired according to a standardized perinatal stroke neuroplasticity

protocol at the Alberta Children's Hospital Diagnostic Imaging Suite using a 3.0 Tesla GE

MR750w MRI scanner (GE Healthcare, Waukesha, WI) with an MR Instruments 32-channel head

coil. High-resolution T1-weighted anatomical images were acquired in the axial plane [166

contiguous slices; voxel size = 1.0 mm isotropic; repetition time (TR) = 8.5 ms; echo time (TE) =

3.2 ms]. Resting-state fMRI acquisition used 150 T2*-weighted whole-brain echo-planar volumes

(EPI; 36 interleaved contiguous slices; voxel size = 3.6 mm isotropic; TR/TE = 2000/30 ms).

Participants were told to fixate on a centrally presented cross. Images were reoriented such that

the stroke was located in the right hemisphere for all patients so that lesioned hemispheres of

cases corresponded to non-dominant hemispheres in controls.

Functional connectivity analyses

Resting-state functional connectivity analyses were performed using the SPM12 (Statistical

Parametric Mapping, Wellcome Trust, UCL, UK) Functional Connectivity Toolbox (CONN) [44] in

Matlab version r2018a (Mathworks, Natick, MA, USA). Slice timing correction, realignment, and
co-registration were performed, and head motion parameters were estimated. Co-registered
images were segmented using subject-specific tissue probability maps. Images were normalized
into Montreal Neurological Institute (MNI) space via direct non-linear transformations using the
standard 152-average template. Direct normalization results in structural and functional volumes
being separately normalized to MNI space. Images were then smoothed with a 6 mm^3 full-width
at half-maximum Gaussian kernel. Head motion exceeding 0.9 mm of translational head
movement and global intensity outliers (z-score>5) were identified using the Artifact Repair
Toolbox [45]. Time courses of blood oxygenation level-dependent (BOLD) responses were
extracted for the GM, CSF and WM. The CSF, WM and lesion time courses were regressed from
the general linear regression model (GLM). Outlier volumes and head motion were additionally
de-weighted in the GLM.
142
143   In a preliminary analysis of this data [35], we performed both ROI-to-ROI and seed to voxel analysis
144   in eight predetermined ROI's within the sensorimotor network. However, based on feedback to
145   this preprinted analysis, and subsequent experimental and theoretical advances in the study of
146   fatigue [15,21,22], we decided to adopt a data-driven analysis plan to reduce possible bias in seed
147   selection.
148
Individuals with perinatal stroke often have widespread morphological and functional differences
in brain structure and development ([46] for a review). Because of these differences, we could not
perform component identification whole-brain connectivity analyses (e.g., independent
component analysis, principal component analysis). Instead, we used a machine learning ROI-
to-ROI approach. All ROIs defined using the default Harvard-Oxford Atlas within CONN were
included in the analysis, except those in the cerebellum and vermis. To reduce the number of FC,
the two ROIs for the cerebellum were selected from the CONN ICA analysis of the Human
Connectome Project dataset [44], resulting in 108 ROIs entered into the analysis. BOLD signal time
series were calculated using the average of all voxels within the ROI. Fisher-transformed
correlation coefficients were interpreted as quantifications of functional connectivity (FC) strength
between areas. FC between each of the ROIs were identified in CONN and extracted for data
analysis using elastic net regression. FC values between ROIs identified as associated with
fatigue in children with perinatal stroke using the elastic-net regression model (described below)
were subsequently quantified in TDC to compare functional connectivity between participant
groups.

Sensorimotor function

We included a measure of sensorimotor function to quantify whether function mediated the relationship between fatigue and the connectivity of the sensorimotor prediction network. Sensorimotor function was assessed via subscales of the Pediatric Stroke Outcome Measure (PSOM). The score for the more affected hand was used. The PSOM is a reliable and valid measure of neurological function following stroke in children [47] and the sensorimotor subscale evaluations include motor, visual, hearing, and somatosensory function [48].

Data and statistical analysis

Statistical analyses were performed with the R statistical software package [49]. Analysis of variance (ANOVA), chi-square and Mann-Whitney U tests were used to compare the demographic data. To identify FC between ROIs that was associated with fatigue, stroke participants' Fisher-transformed coefficients (N = 5778 ROI-to-ROI connections), demeaned age, PSOM score and stroke type (AIS or PVI infarction) were entered into an elastic net regularized regression model, with fatigue score as the outcome variable. Ten-fold cross-validated elastic net regression was performed using the 'glmnet' package [50]. Elastic net regression is a regularized regression technique, used for variable selection in high dimension low N studies, which uses a penalty term to shrink regression coefficients of multiple predictors, setting some to 0 and leaving only the most relevant predictors in the model. A subsequent robust linear regression model containing predictors identified in the model was used to determine model goodness-of-fit and estimate coefficients. Connectivity identified in the elastic net model was compared between groups (Fatigued, Not fatigued, TDC) using Welch's analysis of variance (ANOVA). Significant main effects were followed up using *post hoc* Welch's t-tests. Shapiro-Wilk's test of normality was used to assess the distribution of the residuals. Cohen's *d* and its associated 95% confidence were used as an estimate of the effect sizes for differences in connectivity between groups. The threshold to reject the null hypothesis was set at $p<0.05$. Where necessary, the false-discovery rate was adjusted to control for multiple comparisons [51].

Funding Statement:

This work was supported by funding from the Canadian Institutes of Health Research (CIHR).

Data availability statement

Raw data is available upon reasonable request from the corresponding author. Code used for
analysis is availble from the Open Science Framework (https://osf.io/u4jvh).

Author Contributions:

Author contributions included conception and study design (JGW, HC, EGC, AK) data collection
and acquisition (HC, AK) statistical analysis (JGW), interpretation of results (JGW, HC, EGC,
AK), drafting the manuscript work or revising it critically for important intellectual content (JGW,
HC, EGC, AK)) and approval of final version to be published and agreement to be accountable
for the integrity and accuracy of all aspects of the work (All authors).
206

207   Compliance with Ethical Standards

208   Experiments were carried out with the approval of the Research Ethics Board at the University
209   of Calgary and in accordance with the declaration of Helsinki [42].
210

212   Conflicts of Interest
213   The authors have no conflicts of interest to declare

214 Supplementary material
215
The following is a description of the results from the first iteration of this study, where the analysis
of the association between fatigue core and ROI-ROI FC was constrained to the FC between
sensorimotor areas "The model identified two ROI-to-ROI connections as significant predictors of
fatigue status. FC between the non-lesioned S1 and lesioned SMA (β= −21.7, SE = 9.9, p = 0.032)
and the lesioned and non-lesioned thalamus (β= 15.5, SE = 7.5, p = 0.032) were significant
predictors of fatigue score (R^2 = 0.15, $F_{(2,57)}$ = 4.9, p = 0.017, Figure 2).

References

1. Sreenan, C., Bhargava, R. & Robertson, C. M. T. Cerebral infarction in the term newborn:

Clinical presentation and long-term outcome. *J. Pediatr.* **137**, 351–355 (2000).

2. Wu, Y. W. *et al.* Perinatal stroke in children with motor impairment: A population-based

study. **114**, 612–619 (2004).

3. Maher, C. *et al.* Fatigue is a major issue for children and adolescents with physical

disabilities. *Dev. Med. Child Neurol.* **57**, 742–747 (2015).

4. Russchen, H. A. *et al.* Focus on fatigue amongst young adults with spastic cerebral palsy. *J.*

NeuroEngineering Rehabil. **11**, 161 (2014).

5. Brunton, L. K. *et al.* More Than Just Adolescence: Differences in Fatigue Between Youth

With Cerebral Palsy and Typically Developing Peers. *Ann. Rehabil. Med.* **45**, 197–203

(2021).

6. Wrightson, J. G., Zewdie, E., Kuo, H.-C., Millet, G. Y. & Kirton, A. Fatigue in children with

perinatal stroke: clinical and neurophysiological associations. *Dev. Med. Child Neurol.* **62**,

234–240 (2020).

7. Brunton, L. K. Descriptive Report of the Impact of Fatigue and Current Management

Strategies in Cerebral Palsy. *Pediatr. Phys. Ther.* **30**, 135–141 (2018).

8. Brunton, L. K. & Bartlett, D. J. The bodily experience of cerebral palsy: a journey to self-

awareness. *Disabil. Rehabil.* **35**, 1981–1990 (2013).

9. Hilberink, S. R. *et al.* Health issues in young adults with cerebral palsy: towards a life-span

perspective. *J. Rehabil. Med.* **39**, 605–611 (2007).

10. Boyer, E. R., Palmer, M., Walt, K., Georgiadis, A. G. & Stout, J. L. Validation of the Gait

Outcomes Assessment List questionnaire and caregiver priorities for individuals with

cerebral palsy. *Dev. Med. Child Neurol.* **64**, 379–386 (2022).

11. Kuppuswamy, A. The fatigue conundrum. *Brain* **140**, 2240–2245 (2017).

12. Stephan, K. E. *et al.* Allostatic Self-efficacy: A Metacognitive Theory of Dyshomeostasis-

Induced Fatigue and Depression. *Front. Hum. Neurosci.* **10**, 550 (2016).

13. Friston, K. Hierarchical Models in the Brain. *PLOS Comput. Biol.* **4**, e1000211 (2008).

14. Manjaly, Z.-M. *et al.* Pathophysiological and cognitive mechanisms of fatigue in multiple

sclerosis. *J Neurol Neurosurg Psychiatry* jnnp-2018-320050 (2019) doi:10.1136/jnnp-2018-

320050.

15. Kuppuswamy, A. The Neurobiology of Pathological Fatigue: New Models, New Questions.

The Neuroscientist 1073858420985447 (2021) doi:10.1177/1073858420985447.

16. Pezzulo, G., Rigoli, F. & Friston, K. J. Hierarchical Active Inference: A Theory of Motivated

Control. *Trends Cogn. Sci.* **22**, 294–306 (2018).

17. Greenhouse-Tucknott, A. *et al.* Toward the unity of pathological and exertional fatigue: A

predictive processing model. *Cogn. Affect. Behav. Neurosci.* (2021) doi:10.3758/s13415-

021-00958-x.

18. Doncker, W. D., Dantzer, R., Ormstad, H. & Kuppuswamy, A. Mechanisms of poststroke

fatigue. *J Neurol Neurosurg Psychiatry* jnnp-2017-316007 (2017) doi:10.1136/jnnp-2017-

316007.

19. Wu, S., Mead, G., Macleod, M. & Chalder, T. Model of Understanding Fatigue After Stroke.

Stroke **46**, 893–898 (2015).

20. Baran, T. M., Zhang, Z., Anderson, A. J., McDermott, K. & Lin, F. Brain structural

connectomes indicate shared neural circuitry involved in subjective experience of cognitive

and physical fatigue in older adults. *Brain Imaging Behav.* **14**, 2488–2499 (2020).

21. Ondobaka, S., De Doncker, W., Ward, N. & Kuppuswamy, A. Neural effective connectivity

explains subjective fatigue in stroke. *Brain* (2021) doi:10.1093/brain/awab287.

22. Wylie, G. R., Yao, B., Genova, H. M., Chen, M. H. & DeLuca, J. Using functional

connectivity changes associated with cognitive fatigue to delineate a fatigue network. *Sci.*

Rep. **10**, 21927 (2020).

23. Kuczynski, A. M. *et al.* Sensory tractography and robot-quantified proprioception in

hemiparetic children with perinatal stroke. *Hum. Brain Mapp.* **38**, 2424–2440 (2017).

24. Kuczynski, A. M., Dukelow, S. P., Semrau, J. A. & Kirton, A. Robotic Quantification of

Position Sense in Children With Perinatal Stroke. *Neurorehabil. Neural Repair* **30**, 762–772

(2016).

25. D'Astolfo, L. & Rief, W. Learning about Expectation Violation from Prediction Error

Paradigms – A Meta-Analysis on Brain Processes Following a Prediction Error. *Front.*

Psychol. **8**, 1253 (2017).

26. Ficco, L. *et al.* Disentangling predictive processing in the brain: a meta-analytic study in

favour of a predictive network. *Sci. Rep.* **11**, 16258 (2021).

27. Siman-Tov, T. *et al.* Is there a prediction network? Meta-analytic evidence for a cortical-

subcortical network likely subserving prediction. *Neurosci. Biobehav. Rev.* **105**, 262–275

(2019).

28. Feldman Barrett, L. The theory of constructed emotion: an active inference account of

interoception and categorization. *Soc. Cogn. Affect. Neurosci.* **12**, 1–23 (2017).

29. Garfinkel, S. N. & Critchley, H. D. Interoception, emotion and brain: new insights link internal

physiology to social behaviour. Commentary on:: 'Anterior insular cortex mediates bodily

sensibility and social anxiety' by Terasawa et al. (2012). *Soc. Cogn. Affect. Neurosci.* **8**,

231–234 (2013).

30. Kleckner, I. R. *et al.* Evidence for a Large-Scale System Supporting Allostasis and

Interoception in Humand. *Nat Hum Behav* 1–37 (2017) doi:10.1038/s41562-017-0069.

31. Briggs, R. G. *et al.* Anatomy and White Matter Connections of the Middle Frontal Gyrus.

World Neurosurg. **150**, e520–e529 (2021).

32. Findlater, S. E. *et al.* Lesion locations associated with persistent proprioceptive impairment

in the upper limbs after stroke. *NeuroImage Clin.* **20**, 955–971 (2018).

33. Giudice, C., Rogers, E. E., Johnson, B. C., Glass, H. C. & Shapiro, K. A. Neuroanatomical

correlates of sensory deficits in children with neonatal arterial ischemic stroke. *Dev. Med.*

Child Neurol. **61**, 667–671 (2019).

34. Iandolo, R. *et al.* Neural correlates of lower limbs proprioception: An fMRI study of foot

position matching. *Hum. Brain Mapp.* **39**, 1929–1944 (2018).

35. Wrightson, J. G., Condliffe, E. G., Carlson, H. L. & Kirton, A. Fatigue in perinatal stroke is

associated with the functional connectivity of the sensorimotor network. *medRxiv* (2020).

36. Woodward, K. E. *et al.* Sensory-motor network functional connectivity in children with

unilateral cerebral palsy secondary to perinatal stroke. *NeuroImage Clin.* **21**, 101670 (2019).

37. Saunders, J., Carlson, H. L., Cortese, F., Goodyear, B. G. & Kirton, A. Imaging functional

motor connectivity in hemiparetic children with perinatal stroke. *Hum. Brain Mapp.* **40**,

1632–1642 (2019).

38. Carlson, H. L. *et al.* Structural and functional connectivity of motor circuits after perinatal

stroke: A machine learning study. *NeuroImage Clin.* **28**, 102508 (2020).

39. Turner, I. F., Ward, N. S. & Kuppuswamy, A. Fatigue, effort perception and central activation

failure in chronic stroke survivors: a TMS and fMRI investigation. *bioRxiv* 821959 (2019)

doi:10.1101/821959.

40. van der Schaaf, M. E. *et al.* Fatigue Is Associated With Altered Monitoring and Preparation

of Physical Effort in Patients With Chronic Fatigue Syndrome. *Biol. Psychiatry Cogn.*

Neurosci. Neuroimaging **3**, 392–404 (2018).

41. Cole, L. *et al.* Clinical characteristics, risk factors, and outcomes associated with neonatal

hemorrhagic stroke: A population-based case-control study. *JAMA Pediatr.* (2017)

doi:10.1001/jamapediatrics.2016.4151.

42. World Medical Association. World Medical Association Declaration of Helsinki: ethical

principles for medical research involving human subjects. *JAMA* **310**, 2191–2194 (2013).

43. Varni, J. W. *et al.* The PedsQL in pediatric cerebral palsy: reliability, validity, and sensitivity

of the Generic Core Scales and Cerebral Palsy Module. *Dev. Med. Child Neurol.* **48**, 442

(2006).

44. Whitfield-Gabrieli, S. & Nieto-Castanon, A. Conn: a functional connectivity toolbox for

correlated and anticorrelated brain networks. *Brain Connect.* **2**, 125–141 (2012).

45. Mazaika, P., Hoeft, F., Glover, G. & Reiss, A. Artifact Repair for fMRI Data from High Motion

Clinical Subjects. *Hum. Brain Mapp. San Franc. CA USA* (2007).

46. Kirton, A. *et al.* Perinatal stroke: mapping and modulating developmental plasticity. *Nat.*

Rev. Neurol. **17**, 415–432 (2021).

47. Kitchen, L. *et al.* The pediatric stroke outcome measure: a validation and reliability study.

Stroke **43**, 1602–1608 (2012).

48. deVeber, G. A., MacGregor, D., Curtis, R. & Mayank, S. Neurologic Outcome in Survivors of

Childhood Arterial Ischemic Stroke and Sinovenous Thrombosis. *J. Child Neurol.* **15**, 316–

324 (2000).

49. R Development Core Team, R. *R: A language and environment for statistical computing.* (R

foundation for statistical computing Vienna, Austria, 2011).

50. Friedman, J., Hastie, T. & Tibshirani, R. Regularization Paths for Generalized Linear Models

via Coordinate Descent. *J. Stat. Softw.* **33**, 1–22 (2010).

51. Benjamini, Y., Krieger, A. M. & Yekutieli, D. Adaptive linear step-up procedures that control

the false discovery rate. *Biometrika* **93**, 491–507 (2006).

343
344

Table 1. Demographics characteristics and fatigue scores

	Stroke (N = 60)	Fatigued (N = 27)	Not fatigued (N = 33)	TDC (N=59)
Sex (Female: Male)	21:39	11:16	10:23	29:30
Age (years, mean$\pm$SD)	11.5$\pm$2.8	11.5$\pm$2.5	11.5$\pm$3.1	14.2+3.0
Stroke side (Left: Right)	37:23	19:13	18:10	NA
Stroke type (Arterial: Periventricular Venous Infarction)	29:31	15:12	17:16	NA
Fatigue (PEDSQL, median [IQR])	68.8 [32.8]	50 [18.8]	81.3 [25.0]	NA

Table 2. Means + SD for Fisher-transformed correlations for the ROI-to-ROI analysis

	Fatigued (N = 27)	Not Fatigued (N = 33)	TDC (N = 59)	Group differences (Welch's ANOVA)
Non-lesioned (left) middle frontal gyrus – supramarginal gyrus (posterior)	0.27$\pm$0.30	0.50$\pm$0.34	-0.03$\pm$0.23	$F_{(2,51)}$=32.6, p<0.001
Non-lesioned (left) insula – lesioned (right) inferior temporal gyrus	0.02$\pm$0.21	-0.15$\pm$0.18	0.10$\pm$0.25	$F_{(2,65)}$=16.2, p<0.001
Non-lesioned intracalcarine cortex – lesioned paracingulate cortex	-0.17$\pm$0.15	-0.06$\pm$0.24	0.02$\pm$0.21	$F_{(2,65)}$=10.8, p<0.001

In TDC; lesioned = right, non=lesioned = left, TDC; typically developed control participants

346
347 Fig. 1. Lesion overlay maps for participants with arterial (AIS) strokes, shown are heat maps
348 corresponding to the number of patients that have a lesion in that area overlaid on axial slices
349 from a standard template in MNI space (MNI152). White shading indicates most lesion overlap
350 and dark red least overlap. Images are presented in neurological convention (i.e., right
351 hemisphere is on the right side).
352

353
354
355 Figure 2. FC identified by the 10-fold cross-validated elastic net regression. Red lines illustrate a
356 positive correlation, where a lower PEDSQL fatigue score (indicating more severe fatigue) was
357 associated with weaker FC. The lesioned hemisphere is displayed on the left of this figure.
358 MFG; middle frontal gyrus, SMG; supramarginal gyrus, ICC; intracalcarine cortex, PC;
359 paracingulate gyrus, INS; insula cortex, ITG; Inferior temporal gyrus.

362
363 Associations between FC identified by the elastic net regression and fatigue (PEDSQL fatigue score). A lower fatigue score
364 represents more severe fatigue. MFG; middle frontal gyrus, SMG; supramarginal gyrus, ICC; intracalcarine cortex, PC; paracingulate
365 gyrus, ITG; Inferior temporal gyrus.
366

Figure 3.

367
368 Figure 4. Fisher-transformed correlation coefficients for the connectivity between the A) non-lesioned primary middle frontal gyrus
369 (MFG) and supramarginal gyrus (SMG), B) lesioned inferior temporal gyrus (ITG) and non-lesioned insula and C) between the non-
370 lesioned intracalcarine cortex (ICC) and lesioned paracingulate cortex (PC). The black circle and error bars represent the mean±95%
371 confidence interval. TDC; Typically developed controls.* represents a different (p<0.05) FC compared to the not fatigued group.
372