Investigating spatial variability in COVID-19 pandemic

severity across 19 geographic areas, Spain, 2020

Authors:

Sushma Dahal[a]*, Kenji Mizumoto [b,c]*, Richard Rothenberg[a], Gerardo Chowell[a]

*Joint first authors

[a] Department of Population Health Sciences, School of Public Health, Georgia State

University, Atlanta, Georgia, USA

[b] Graduate School of Advanced Integrated Studies in Human Survivability, Kyoto University

Yoshida-Nakaadachi-cho, Sakyo-ku, Kyoto, Japan

[c] Hakubi Center for Advanced Research, Kyoto University, Yoshidahonmachi, Sakyo-ku,

Kyoto, Japan

Article type:

Major Article

Address for correspondence:

K Mizumoto, Graduate School of Advanced Integrated Studies in Human Survivability,

Kyoto University Yoshida-Nakaadachi-cho, Sakyo-ku, Kyoto, Japan, Tel: +81 75 762 2001

Fax: +81 75 762 2277, E-mail: mizumoto.kenji.5a@kyoto-u.ac.jp

Word count:

Abstract: 199   (Max 200)

Main: 3182

Abstract

Introduction: Spain has been disproportionately affected by the COVID-19 pandemic,

ranking fifth in the world in terms of both total cases and total deaths due to COVID-19 as of

May 20, 2020. Here we derived estimates of pandemic severity and assessed its relationship

with socio-demographic and healthcare factors.

Methods: We retrieved the daily cumulative numbers of laboratory-confirmed COVID-19

cases and deaths in Spain from February 20, 2020 to May 20, 2020. We used statistical

methods to estimate the time-delay adjusted case fatality risk (aCFR) for 17 autonomous

communities and 2 autonomous cities of Spain. We then assessed how transmission and

sociodemographic variables were associated with the aCFR across areas using multivariate

regression analysis.

Results: We estimated the highest aCFR for Madrid (25.9%) and the average aCFR in Spain

(18.2%). Our multivariate regression analysis revealed three statistically significant predictor

variables: population size, population density, and the unemployment rate.

Conclusions:

The estimated aCFR for 10 autonomous communities/cities in Spain are significantly higher

than those previously estimated for other geographic regions including China and Korea. Our

results suggest that public health interventions focused on densely populated areas and low

socioeconomic groups can ameliorate the mortality burden of the COVID-19 pandemic in

Spain.

Key words: COVID-19, Spain, time-delay adjusted CFR, 2020

Introduction

Since the emergence of COVID-19 in Wuhan, China in December 2019, the novel coronavirus (SARS-CoV-2) continues to spread throughout the world, straining and overloading healthcare systems and causing substantial morbidity and mortality burden during a short time period. As of May 20 2020, 4,789,205 confirmed cases including 318,789 deaths attributed to COVID-19 have been reported from 215 countries/territories/areas [1]. Thus far, the US has reported the highest number of cases (30.8%) and deaths (28.0%) globally. Other countries that follow the US in terms of COVID-19 death toll include the United Kingdom (11.1%), Italy (10.1%), France (8.8%) and Spain (8.7%) [1].

The case fatality risk is a useful metric to assess pandemic severity, which is typically estimated as the proportion of deaths among the total number of cases attributed to the disease [2, 3]. However, during the course of outbreak of an infectious disease such as COVID-19, real-time estimates of CFR need to be derived carefully since it is prone to ascertainment bias and right censoring [4, 5]. In particular, the disease spectrum for COVID-19 ranges from asymptomatic and mild infections to severe cases that require hospitalization and specialized supportive care. This may lead to overestimation of the CFR among ascertained cases. On the other hand, there is a delay from illness onset to death for severe cases [6], which could lead to an underestimation of the CFR [5, 7]. Therefore, statistical methods that help mitigate inherent

biases in estimates of the CFR should be employed to accurately plan for medical resources

such as ICU units and ventilators, which are essential resources to save the lives of critically ill

patients [8-10].

Several studies have reported CFR estimates for COVID-19 [11-13]. Overall, these estimates

have varied substantially across geographic regions even within the same country. For

example, a recent study estimated the time-delay-adjusted CFR at 12.2% for the ground zero of

the COVID-19 pandemic: the city of Wuhan [6], whereas for the most affected region in Italy

(Lombardia), the delay-adjusted CFR reached 24.7% [14]. The drivers behind the geographical

variations in the severity of the COVID-19 pandemic are yet to be investigated but could

provide critical information to mitigate the morbidity and mortality impact of this and future

pandemics [15].

In this study we aim to estimate the severity of COVID-19 pandemic across 19 geographic

areas in Spain and aim to explain how these estimates varied geographically as a function of

underlying factors. For the real-time estimation of severity, we adjust for right censoring using

established methods [16, 17] and report the estimates of the time-delay adjusted CFR of

COVID-19 for 17 autonomous communities (Comunidades Autonomas, CCAA), and 2

autonomous cities of Spain as well as for the entire Spain. We then assessed the association

between different transmission and socio-demographic factors and the estimated CFRs across

areas using multivariate regression analyses.

Methods

Study setting: Spain is situated on the Iberian Peninsula and is divided into 17 autonomous

communities (CCAA) and 2 African autonomous cities [18]. Ceuta and Melilla are the 2

African autonomous cities whereas the 17 CCAA include: Andalucía, Aragon, Asturias,

Balears, Canarias, Cantabria, Castilla-La Mancha, Castilla y Leon, Catalunya, C. Valenciana,

Extremadura, Galicia, Madrid, Murcia, Navarra, Pais Vasco, La Rioja [18].

Initial cases of COVID-19 in Spain: The first case of COVID-19 in Spain was confirmed on 31st

January, 2020 in La Gomera, Canary Islands in a person who was in contact with an infected

person while in Germany [19]. By February 27, there were total 12 cases which increased to 45

cases in March 1 and then continued to rise rapidly throughout the country [20]. The incidence

trajectory continued to rise for about 4 weeks before a gradual decline. The daily reported

number of new cases has not exceeded 1000 for 10 consecutive days since May 11, 2020 [21]

Data Sources:

The Ministry of Health of Spain releases daily reports on COVID-19 cases and deaths [21].

From these reports we retrieved the daily cumulative numbers of reported

laboratory-confirmed COVID-19 cases and deaths from February 20, 2020 to May 20, 2020.

We then stratified the data into 20 groups that included 17 CCAA, 2 African autonomous cities

and for the entire Spain.

For each CCAA we obtained data on demographic, socio-demographic, and healthcare factors

from the Statistics National Institute (Instituto Nacional de Estadística) [22] and the annual

report of the national health system [23]. The demographic factors we included were: total

population size, population density, proportion of population size by region, proportion of the

population older than 60 years, proportion of the population older than 70 years, and life

expectancy at birth. We included four socio-demographic factors: poverty risk rate, proportion

of population of low social class, unemployment rate, and infant mortality rate. To assess the

effect of healthcare factors, we obtained the data on percentage of total consolidated

expenditure on hospital and specialized services, and public health expenditure per GDP by

CCAA from the annual report issued by the national health system, 2018 [23]. Finally, we also

included three factors related to transmission dynamics: The number of days from the date the

first case was reported to the day the cumulative number cases surpassed one hundred cases,

the cumulative morbidity (total number of reported cases) and the cumulative morbidity rate

calculated by the cumulative number cases divided by the local population size. Table S1

summarizes the range and median of above variables for 17 CCAA regions in Spain.

Additionally, we obtained the shapefiles of the autonomous communities of Spain from the

national geographic information system of Spain [24].

Time-delay adjusted CFR estimation

The crude CFR is defined as the number of cumulative deaths divided by the number of

cumulative cases at a specific point in time. For the estimation of CFR in real time during the

course of outbreak, we employed the delay from hospitalization to death, h_s, which is assumed

to be given by $h_s = H(s) - H(s-1)$ for s>0 where H(s) is a cumulative density function of the

delay from hospitalization to death and follows a gamma distribution with mean 10.1 days and

SD 5.4 days, obtained from the previously published paper [6]. Let $\pi_{a,ti}$ be the time-delay

adjusted case fatality risk on reported day t_i in area a, the likelihood function of the estimate $\pi_{a,ti}$

is

$$L(\pi_{a,t_i}; D_{a,t}, c_{a,t})$$

$$= \prod_{t_i} \binom{\sum_{t=1}^{t_i} c_{a,t}}{D_{a,t_i}} \left(\pi_{a,t_i} \frac{\sum_{t=2}^{t_i} \sum_{s=1}^{t-1} c_{a,t-s} h_s}{\sum_{t=1}^{t_i} c_{a,t}} \right)^{D_{a,t_i}} \left(1 \right.$$

$$\left. - \pi_{a,t_i} \frac{\sum_{t=2}^{t_i} \sum_{s=1}^{t-1} c_{a,t-s} h_s}{\sum_{t=1}^{t_i} c_{a,t}} \right)^{\sum_{t=1}^{t_i} c_{a,t} - D_{a,t_i}}$$

where $c_{a,t}$ represents the number of new cases with reported day t in area a, and $D_{a,ti}$ is the

cumulative number of deaths until reported day t_i in area *a* [16, 17]. Second parenthesis

corresponds to the estimated crude case fatality risk at day t_i in area a. Among the cumulative

cases with reported day t in area a, $D_{a,ti}$ have died and the remainder have survived the

infection. The contribution of those who have died with biased death risk is shown in the

middle parenthetical term and the contribution of survivors is presented in the right

parenthetical term. We assume that $D_{a,ti}$ is the result of the binomial sampling process with

probability $\pi_{a,ti}$.

The priors specified in our Bayesian model were non-informative priors. We estimated model

parameters using a Monte Carlo Markov Chain (MCMC) method in a Bayesian framework.

Posterior distributions of the model parameters were estimated by sampling from the three

Markov chains. For each chain, we drew 100,000 samples from the posterior distribution after

a burn-in of 20,000 iterations. Convergence of MCMC chains were evaluated using the

potential scale reduction statistic [25, 26]. Estimates and 95% credibility intervals for these

estimates are based on the posterior probability distribution of each parameter and based on the

samples drawn from the posterior distributions.

Multivariate regression analysis

After quantifying the extent of spatial auto-correlation of CCAA-level time-delay-adjusted

CFR median estimates, with the use of Moran's I statistic with a Delauney triangulation

neighbours spatial mixing matrix [27, 28], we explored the association between

time-delay-adjusted CFR and predictor variables to identify simplified models with significant

factors linked to the variation in CFR estimates across geographic areas in Spain.

Multicollinearity tests were examined using variance inflation factors (VIF). Predictors

exceeding VIF value of 4.0 were excluded one by one from the multivariate analyses. By fitting

all combinations of models given a set of predictors, we derived best minimal models and

indicated in Table 2. By taking 1000 draws from the posterior estimates of time-delay-adjusted

CFR and running the linear regression 1000 times, the combination of variables with the largest

count was selected as the final model. With this model, pooled estimates accounting for both

within-model uncertainty and between models uncertainty were derived.

All statistical analyses were conducted in R version 3.6.1 (R Foundation for Statistical

Computing, Vienna, Austria).

Results:

As of May 20, a total of 233,037 cases and 27,940 deaths due to COVID-19 have been reported

in Spain. Moreover, the Madrid region has reported the highest number of cases at 67,049

(28.8%) and deaths at 8,931 (32.0%) followed by Catalunya with 55,888 cases (24.0%) and

6,021 deaths (21.5%).

Figure 1 displays the observed and posterior estimates of crude case fatality risk in Spain, 17

CCAAs and 2 autonomous cities (A through T). Day 1 corresponds to March 1st in 2020. Black

dots show the crude case fatality risks, and light and dark indicate 95% and 50% credible

intervals for posterior estimates, respectively. Our model-based crude CFR fitted the observed

data well in all the regions except Aragon where the model did not fit well for first 2 weeks.

There was a rapid rise in crude CFR in the Aragon, Asturias, Castilla-La Mancha, C.

Valenciana, Extremadura, and in Madrid.

Figure 2 illustrates observed and model based posterior estimates of time-delay-adjusted CFR

in the 20 areas. Black dots show crude case fatality risks, and light and dark indicate 95% and

50% credible intervals for posterior estimates, respectively. In most of the regions and at the

national level, we saw a greater difference between the time delay adjusted CFR and crude

CFR in the first 4 weeks of the epidemic followed by a slowly declining difference in the later

stage of the epidemic. The graph of the time-delay adjusted CFR varies considerably for

different areas. For instance, as the epidemic progressed, the adjusted CFR increased slightly in

CN and NC. In contrast, it showed a rapidly increasing trend for first three weeks and then

gradually declined in CL. For AR, it was at a high level for the first two weeks after which it

started to decline. For the most affected MD region, the time-delay adjusted CFR fluctuated for

first 25 days followed by a decline, a leveling off, and then moved upwards.

A summary of the time delay adjusted case fatality risk, range of median estimates and crude

CFR of COVID-19 across different areas of Spain are presented in Table 1. Our model based

posterior estimates of time-delay adjusted CFR are higher than the observed crude CFR. The

Madrid autonomous community had the highest time delay adjusted CFR of 25.9% [95%

credible interval: 25.3-26.3%] followed by Extremadura (EX) (21.7%) [95%CrI: 20.0-23.4%],

Aragon (AR) (21.3%) [95%CrI: 19.9-22.5%], and Asturias(AS) (20.3%) (95%CrI:

18.3-22.4%). The national estimate for Spain was 18.2% (95%CrI: 18.0-18.4%]. Of the 19

autonomous areas of Spain, 13 had the time-delay adjusted CFR greater than 10% (Table 1,

figure 3).

Our analysis of the statistical geospatial heterogeneities in the time-delay adjusted using

Moran's I statistic, which represents the extent of the degree of spatial autocorrelation, was not

statistically significant (Moran's Index = 0.08, P = 0.15). After inspecting VIF, the proportion

of population size by region, proportion of seniors aged 60 and above, life expectancy at birth,

public health expenditure per GDP, proportion of low social class, poverty risk rate,

cumulative morbidity, and the cumulative morbidity rate were excluded from the multivariate

analysis.

Autonomous communities with larger population size (p value= 0.10 (95%CrI: 0.05 – 0.20)),

higher population density (p value= 0.08 (95%CrI:0.05 – 0.12)) and with a higher

unemployment rate (p value=0.07 (95%CrI: 0.04 –0.14)) experienced higher time-delay

adjusted CFRs. These three significant factors explained 16% (95%CrI: 7 –24)) (adjusted

R-squared) of the variation of the pandemic severity across CCAAs (Table□2). The

correlation matrix of predictors with correlation coefficient (r) is presented in Figure S3.   We

found that the population size is strongly correlated (r >= 0.7 or r <= -0.7) with the proportion

of population size by region (r =1.00) and the cumulative morbidity while population density is

strongly correlated with the proportion of low social class (r = -0.77) and the cumulative

morbidity (r = 0.70). Meanwhile the unemployment rate is strongly correlated with the

poverty risk rate (r=0.91) and the life expectancy at birth (r=-0.76).

The range of variables included in the linear regression analysis are provided in supplementary

Table S1.

Discussion

In this paper, we have estimated the time delay adjusted case fatality risk of COVID-19 for 19

autonomous areas/cities of Spain. Our estimate of time-delay adjusted CFR in Spain was at

18.2%, but it varied widely across the 19 Spanish areas, with some areas exhibiting higher CFR

values such as in Madrid (25.9%), and Extremadura (21.7%) while other areas such as Melilla

(2.9%), Ceuta (4.6%), and Galicia (6.9%) experiencing relatively lower CFR values. Of the 19

areas, 13 experienced an adjusted CFR greater than 10%. We also observed a significant

positive association of the time-delay adjusted CFR with population size, population density,

and unemployment rate across 17 Spanish areas.   Our findings suggest the need for additional

control efforts as well as generating real time regional severity estimates particularly for areas

with high population density and unemployment rate, which have been hit hard by the

COVID-19 pandemic.

The adjusted CFR estimates in Spain (18.2%) is higher than the estimates for Wuhan

(12.2%)[6], and Korea (1.4%) [29] and Italy (17.4%) [14]. All of these studies employed same

methods and hence are comparable. When we compare the estimates for the most affected

areas across different countries, the risk in Madrid in Spain (25.9%) is higher than that

estimated for Wuhan in China (12.2%) [6], Daegu in Korea (2.4%) [29] and Lombardia in Italy

(24.7%) [14]. This difference across countries and regions may be partly explained by

differences in population age structure, density, other socio-demographic factors, and the scale

of the pandemic in different areas. The median age in Spain (44.9 years) [30] is comparable to

that of Italy (45.4 years) but higher than that for China (36.7 years) [31]. Indeed, the elderly

population is at risk of severe outcomes from COVID-19 [32-34], which could partly explain

the higher severity observed for Italy and Spain. Other factors that could have played a role in

these differences may be related to differences in testing strategies. For example, in Korea

extensive testing and rigorous contact tracing strategy were implemented [35] while testing

prioritized more severe cases in Italy [34] and Spain [36].

Our results from the multivariate analysis found a significant positive association of adjusted

CFR with population size and population density, emphasizing the importance of social

distancing. We also found a significant positive association between adjusted CFR and

unemployment rate across areas in Spain. Unemployment rate is not only an indicator of

performance of labor market but also an indicator of socioeconomic status for an area, and this

is supported by the strong positive correlation with poverty risk and negative correlation with

the life expectancy at birth. Unemployment rate can lead to decreased investment in health,

education, nutrition, and good housing condition [37]. In any pandemic situation like

COVID-19, the poorer tend to exhibit the highest morbidity and mortality rates. For instance,

lower socioeconomic groups were also disproportionately affected by the 1918 influenza

pandemic [38, 39]. Data from the US and the UK have shown marked ethnic and racial

disparities in COVID-19 death rates [40]. Moreover, preliminary COVID-19 mortality data

from the US also indicates a 2-fold age-adjusted death rate among Hispanic/Latino and

1.9-fold among Black/African American compared to Whites [41]. However, we do not know

whether the association between unemployment rate and adjusted CFR is inflated by a limited

access to COVID-19 testing among low socio-economic groups.

In our study we saw considerable variations in CFR trend across areas. For instance, as the

epidemic progressed, the adjusted CFR showed a slightly upward trend in CN and NC, a rapid

upward trend followed by the slow decline in CL, and a downward trend in AR. Likewise, for

CM the graph showed an upward trend followed by a relative decline and then again an upward

trend before declining gradually. The CFR trend for the 19 autonomous areas can be helpful in

the planning and implementation of health care services and prevention measures separately

for each of them. The downward trend in CFR as seen in some of the areas in our study suggest

the improvement in epidemiologic surveillance leading to the increased capture of mild or

asymptomatic cases. A higher number of mild and asymptomatic cases also indicate an

increase in human-to-human transmission leading to a prolonged epidemic which can be

controlled through effective social distancing measures until an effective vaccine or treatment

becomes available [6].

The upward trend in CFR indicates that the temporal disease burden exceeded the capacity of

healthcare facilities and the surveillance system probably missed many cases during the early

phase of the epidemic [6], particularly due to a significant presence of mild and asymptomatic

cases. It has been found that about 18% of the COVID-19 infections in Diamond Princess

Cruise ship were asymptomatic [42]. The increasing trend in CFR could further be explained

the nosocomial transmission affecting the health care workers, inpatients and their visitors [6].

In China, of 44672 confirmed COVID-19 cases, 3.8% was among the health care personnel

[43]. Similarly, Wang et al. in their study suspected 41% of the patients to have

human-to-human hospital associated transmission of COVID-19 [44].

Our study has some limitations. The preferential ascertainment of severe cases bias in

COVID-19 may have spuriously increased our estimate of CFR [5], which is a frequent caveat

in this type of studies [45, 46]. Similarly, given the long infection-death time for COVID-19

which ranges between 2 to 8 weeks [30], our estimate may have been affected by delayed

reporting bias [5, 7]. Similarly, in our data, the date of report reflects the date of reporting and

not the date of onset of illness. Finally, we assumed infant mortality and poverty risk rate as a

proxy for areas with low socio-economic groups.

Conclusion

The risk of death due to COVID-19 in Spain was estimated at 18.2%, but estimates varied

substantially across 19 geographic areas. The CFR was as high as 25.9% in Madrid and as low

as 2.9% in Melilla. Of the 19 autonomous areas/cities, 13 had a time-delay-adjusted CFR

greater than 10% reflecting a disproportionate severity burden of COVID-19 in Spain.

Importantly, our estimate of CFR for the most affected Madrid region is higher than previous

estimates for the most affected areas within China, Korea, and Italy. Our findings suggest a

significant association between unemployment rate, population density, and population size

with an increased risk of death due to COVID-19. Further studies with patient level data on

mortality, and risk factors could provide a more detailed understanding of the factors shaping

the risk of death related to COVID-19.

Note

Financial support.

KM acknowledges support from the Japan Society for the Promotion of Science (JSPS)

KAKENHI Grant Number 20H03940, and from the Leading Initiative for Excellent Young

Researchers from the Ministry of Education, Culture, Sport, Science & Technology of Japan.

GC acknowledges support from NSF grant 1414374 as part of the joint NSF-NIH-USDA

Ecology and Evolution of Infectious Diseases program; UK Biotechnology and Biological

Sciences Research Council grant BB/M008894/1.

322
323

Potential conflicts of interest.

No conflict.

Additional files

Additional file 1: Table S1. The range of major socio-demographic, and healthcare variables

for 17 regions in Spain used in our analyses (linear regression).

Additional file 2: Figure S1. Daily number of reported COVID-9 cases by region in Spain. Day

1 corresponds to March 1st in 2020.

Additional file 3: Figure S2. Daily number of reported COVID-9 deaths by region in Spain.

Day 1 corresponds to March 1st in 2020.

Additional file 4: Figure S3. Correlation matrix of predictors.

References

1. WHO, Coronavirus disease 2019 (COVID-19)-Situation report-121. 20 May **2020**.

2. Deng, X., et al., Case fatality risk of the first pandemic wave of novel coronavirus
disease 2019 (COVID-19) in China. *Clinical Infectious Diseases*, 15 May **2020**.

3. Rothenberg, R., Death and The Virus. *Clinical Infectious Diseases*, 30 May **2020**.

4. Mizumoto, K., et al., Estimating the risk of Middle East respiratory syndrome

(MERS) death during the course of the outbreak in the Republic of Korea, 2015.

International Journal of Infectious Diseases, **2015**. 39: p. 7-9.

5. Lipsitch, M., et al., Potential biases in estimating absolute and relative case-fatality

risks during outbreaks. *PLoS neglected tropical diseases*, **2015**. 9(7).

345 6. Mizumoto, K. and G. Chowell, Estimating Risk for Death from 2019 Novel

346  Coronavirus Disease, China, January-February 2020. *Emerging infectious diseases*,

347  **2020**. 26(6).

348 7. Kucharski, A.J. and W.J. Edmunds, Case fatality rate for Ebola virus disease in west

349  Africa. *The Lancet*, **2014**. 384(9950): p. 1260.

350 8. Lipsitch, M., D.L. Swerdlow, and L. Finelli, Defining the epidemiology of

351  Covid-19—studies needed. *New England Journal of Medicine*, **2020**.

352 9. Reich, N.G., et al., Estimating absolute and relative case fatality ratios from infectious

353  disease surveillance data. *Biometrics*, **2012**. 68(2): p. 598-606.

354 10. Wu, J.T., et al., Estimating clinical severity of COVID-19 from the transmission

355  dynamics in Wuhan, China. *Nature Medicine*, **2020**.

356 11. Novel, C.P.E.R.E., The epidemiological characteristics of an outbreak of 2019 novel

357  coronavirus diseases (COVID-19) in China. Zhonghua liu xing bing xue za zhi=

358  Zhonghua liuxingbingxue zazhi, **2020**. 41(2): p. 145.

359 12. WHO, Report of the WHO-China Joint Mission on Coronavirus Disease 2019

360  (COVID-19). **2020**, February.

361 13. Rodriguez-Morales, A.J., et al., Clinical, laboratory and imaging features of

362  COVID-19: A systematic review and meta-analysis. *Travel Medicine and Infectious*

363  *Disease*, **2020**: p. 101623.

364   14.   Mizumoto, K., S. Dahal, and G. Chowell, Spatial variability in the risk of death from

365   COVID-19 in Italy, **2020**. March 2020.

366   15.   Chowell, G., et al., Spatial-temporal excess mortality patterns of the 1918–1919

367   influenza pandemic in Spain. *BMC infectious diseases*, **2014**. 14(1): p. 371.

368   16.   Nishiura, H., et al., Early epidemiological assessment of the virulence of emerging

369   infectious diseases: a case study of an influenza pandemic. *PLoS One*, **2009**. 4(8).

370   17.   Tsuzuki, S., et al., Dynamics of the pneumonic plague epidemic in Madagascar,

371   August to October 2017. *Eurosurveillance*, **2017**. 22(46).

372   18.   NSD. Spain-Administrative division 4/6/2020]; Available from:

373   https://nsd.no/european_election_database/country/spain/administrative_divisions.htm

374   l

375   19.   Chavez, N. First case of coronavirus confirmed in Spain. January 31, **2020**

376   3/31/2020].

377   20.   WHO, Coronavirus disease 2019 (COVID-19): Situation reports 38, and 41. 2020**.**

378   21.   Spain, M.o.H. Situation of COVID-19 in Spain **2020**; Available from:

379   https://covid19.isciii.es/

380   22.   INE. Resident population by date, sex and age. **2020**; Available from:

381   https://www.ine.es/dyngs/INEbase/es/operacion.htm?c=Estadistica_C&cid=12547361

382   76951&menu=ultiDatos&idp=1254735572981

23. Informe anual del sistema nacional de salud, 2018 MINISTERIO DE SANIDAD,

CONSUMO Y BIENESTAR SOCIAL.

24. Geographical reference information 04/13/2020]; Available from:

http://centrodedescargas.cnig.es/CentroDescargas/equipamiento.do?method=mostrarE

quipamiento

25. Gamerman, D. and H.F. Lopes, Markov chain Monte Carlo: stochastic simulation for

Bayesian inference. **2006**: CRC Press.

26. Gelman, A. and D.B. Rubin, Inference from iterative simulation using multiple

sequences. *Statistical science*, **1992**. 7(4): p. 457-472.

27. Chowell G, Towers S, Viboud C, Fuentes R, Sotomayor V, Simonsen L, et al. The

influence of climatic conditions on the transmission dynamics of the 2009 A/H1N1

influenza pandemic in Chile. BMC Infect Dis. **2012**.12:298. doi:

10.1186/1471-2334-12-298.

28. Bivand, S., E. J. Pebesma and V. Gomez-Rubio. **2008**. Applied Spatial Data Analysis

with R (Use R), Springer-Verlag.

29. Shim Eunha, et al., Estimating the risk of COVID-19 death during the course of the

outbreak in Korea, February- March, 2020 **2020**.

30. Worldometer, Spain population. **2020**.

31.   World Population Prospects 2019, Online Edition. Rev. 1. **2019**, United Nations,

Department of Economic and Social Affairs, Population Division.

32.   Organization, W.H., Report of the WHO-China Joint Mission on Coronavirus Disease

2019 (COVID-19). **2020**, February.

33.   Surveillances, V., The Epidemiological Characteristics of an Outbreak of 2019 Novel

Coronavirus Diseases (COVID-19)—China, 2020. *China CDC Weekly*, **2020**. 2(8): p.

113-122.

34.   Onder, G., G. Rezza, and S. Brusaferro, Case-Fatality Rate and Characteristics of

Patients Dying in Relation to COVID-19 in Italy. *JAMA*, **2020**.

35.   Colbourn, T., COVID-19: extending or relaxing distancing control measures. The

Lancet Public Health, **2020**.

36.   Linde, P., Spain's coronavirus crisis: Why the numbers are failing to show the full

picture, in El Pais. April 5, **2020**.

37.   Qing WA. The effects of unemployment rate on health status of Chinese people.

Iranian journal of public health. 2015 Jan;44(1):28.

38.   Dahal, S., et al., Natality decline and spatial variation in excess death rates during the

1918–1920 influenza pandemic in Arizona, United States. *American journal of*

epidemiology, **2018**. 187(12): p. 2577-2584.

39.   Mamelund, S.E., 1918 pandemic morbidity: The first wave hits the poor, the second

wave hits the rich. *Influenza and other respiratory viruses*, **2018**. 12(3): p. 307-313.

40.   Bibbins-Domingo K. This time must be different: disparities during the COVID-19

pandemic. Available from: https://www.acpjournals.org/doi/full/10.7326/M20-2247

41.   Age adjusted fatal lab confirmed COVID-19 cases per 100,000 by race/ethnicity

group. April 6, **2020**.41.

42.   Mizumoto, K., et al., Estimating the asymptomatic proportion of coronavirus disease

2019 (COVID-19) cases on board the Diamond Princess cruise ship, Yokohama,

Japan, 2020. *Eurosurveillance*, **2020**. 25(10).

43.   Wu, Z. and J.M. McGoogan, Characteristics of and important lessons from the

coronavirus disease 2019 (COVID-19) outbreak in China: summary of a report of 72

314 cases from the Chinese Center for Disease Control and Prevention. *Jama*, **2020**.

44.   Wang, D., et al., Clinical characteristics of 138 hospitalized patients with 2019 novel

coronavirus–infected pneumonia in Wuhan, China. *Jama*, **2020**.

45.   Yu, H., et al., Human infection with avian influenza A H7N9 virus: an assessment of

clinical severity. *The Lancet*, **2013**. 382(9887): p. 138-145.

46.   Cauchemez, S., et al., Middle East respiratory syndrome coronavirus: quantification

of the extent of the epidemic, surveillance biases, and transmissibility. *The Lancet*

infectious diseases, **2014**. 14(1): p. 50-56.

Table 1. Summary results of time-delay adjusted case fatality risk of COVID-19 in the 19 areas in Spain, 2020 (As of May 20, 2020)

Area	Median estimate of time-delayed adjusted case fatality risk as of May 20, 2020	Range of median estimates of time-delayed adjusted case fatality risk during the study period by geographic region	Crude CFR[‡]
Spain (National)	18.2% (95%CrI[§]: 18.0-18.4%)	12.4-26.8%	12.0% (95%CI[¶]:11.9-12.1%) 27940/233037
Andalucia(AN)	13.0% (95%CrI: 12.4-13.7%)	9.8-50.8%	11.0% (95%CI:10.4-11.5%) 1375/12547
Aragon(AR)	21.3% (95%CrI: 19.9-22.5%)	15.9-87.5%	15.2% (95%CrI: 14.2-16.1%) 848/5588
Asturias(AS)	20.3% (95%CrI: 18.3-22.4%)	3.6-51.7%	12.9% (95%CrI: 11.6-14.3%) 307/2374
Balears(IB)	15.1% (95%CrI: 13.2-17.1%)	9.4-51.8%	10.9% (95%CrI: 9.6-12.4%) 221/2024
Canarias(CN)	8.4% (95%CrI: 7.2-9.7%)	6.4-52.5%	6.7% (95%CrI: 5.7-7.8%) 155/2307
Cantabria(CB)	9.5% (95%CrI: 8.3-10.8%)	7.4-51.8%	9.2% (95%CrI: 8.0-10.4%) 209/2279
Castilla-La Mancha(CM)	18.7% (95%CrI: 18.0-19.3%)	15.4-51.3%	17.4% (95%CrI: 16.8-18.0%) 2919/16789
Castilla y Leon(CL)	11.9% (95%CrI: 11.4-12.4%)	9.7-51.2%	10.5% (95%CrI: 10.1-11.0%) 1960/18627
Catalunya(CT)	14.8% (95%CrI: 14.5-15.2%)	11.6-40.5%	10.8% (95%CrI: 10.5-11.0%) 6021/55888
Ceuta(CE)	4.6% (95%CrI:1.7-10.0%)	4.4-51.8%	3.4% (95%CrI: 0.9-8.4%) 4/119
C.Valenciana(VC)	19.0% (95%CrI: 18.0-19.9%)	8.0-32.3%	12.6% (95%CrI: 12.0-13.2%) 1383/10987
Extremadura(EX)	21.7% (95%CrI: 20.0-23.4%)	15.8-49.5%	16.6% (95%CrI: 15.3-18.0%) 505/3042
Galicia(GA)	6.9% (95%CrI: 6.4-7.5%)	5.1-51.4%	6.7% (95%CrI: 6.2-7.2%) 608/9077
Madrid(MD)	25.9% (95%CrI: 25.3-26.3%)	13.9-30.5%	13.3% (95%CrI:13.1-13.6%) 8931/67049
Melilla(ME)	2.9% (95%CrI: 0.7-7.4%)	2.4-50.8%	1.7% (95%CrI: 0.2-5.8%) 2/121
Murcia(MC)	9.9% (95%CrI: 8.3-11.5%)	3.4-53.7%	9.5% (95%CrI: 8.1-11.0%) 149/1570
Navarra(NC)	11.3% (95%CrI: 10.3-12.3%)	6.4-51.8%	9.7% (95%CrI: 8.9-10.6%) 506/5195
Pais Vasco(PV)	17.4% (95%CrI: 16.6-18.2%)	10.9-71.8%	11.0% (95%CrI: 10.5-11.6%) 1483/13421
La Rioja(RI)	11.8% (95%CrI:10.6-13.1%)	3.9-49.7%	8.8% (95%CrI: 7.9-9.7%) 354/4033

[§]CrI: 95% credibility intervals (CrI), ‡ Estimates calculated from exact binomial test, [¶]95%CI: 95% confidence interval

Table 2. Final multivariate regression model of time-delay-adjusted CFR as a function of

socio-demographic variables across autonomous communities of Spain

Predictor	Coefficient	Standard error	P value	Adjusted R-squared
Baseline	8.7 (95%CrI§: 7.2 – 10.2)	4.8 (95%CrI: 4.4 – 5.3)	0.09 (95%CrI: 0.05 – 0.17)	0.16 (95%CrI: 0.07 –0.24)
Population Size (in 1,000), 2019	-0.001 (95%CrI:-0.001– -0001)	0.0006 (95%CrI: 0.0006 – 0.0007)	0.10 (95%CrI: 0.05 – 0.20)	
Population density (per square km), 2017	0.016 (95%CrI:0.014 –0.018)	0.008 (95%CrI:0.008 –0.009)	0.08(95%CrI:0.05 –0.12)	
Proportion of population size by region, 2019*				
Proportion of seniors aged 70 and above, 2019 (%)				
Proportion of seniors aged 60 and above, 2019 (%)*				
Life expectancy at birth, 2016*				
Infant mortality rate, 2016				
Public health expenditure per GDP*				
Expenditure on hospitals and specialized services (% of total consolidated public spending), 2016				
Proportion of population of low social class, 2017*				
Poverty risk rate, 2017*				
Unemployment rate (among population age 16 years and above), 2019	0.64 (95%CrI:0.52 –0.77)	0.33 (95%CrI:0.30 – 0.36)	0.07 (95%CrI:0.04 –0.14)	
Cumulative morbidity*				
Growth rate (number of days reaching to 100 cases)				
Cumulative morbidity rate (%)*				

Cumulative morbidity: cumulative cases, cumulative morbidity rate: cumulative cases divided by the population size

Multicollinearity tests test were examined using variance inflation factors (VIF). Predictors exceeding VIF value of 4 were excluded one by one for multivariate analysis, and marked with

*. Results of multivariate analysis are shown in the table. Fitting all combination of models
given a set of predictors, best models were selected. Predictors in final models are highlighted
in bold font.

Figure legend

Fig 1: Temporal variation of risk of death caused by COVID-19 by area, Spain, March-May,

2020: crude case fatality risk (cCFR)

Observed and posterior estimates from the Bayesian model of crude case fatality risk in (A) Spain (National), (B)

Andalucia(AN),(C) Aragon(AR), (D) Asturias(AS),(E) Balears(IB),(F) Canarias(CN),(G) Cantabria(CB),(H)

Castilla-La Mancha(CM),(I) Castilla y Leon(CL), (J) Catalunya(CT),(K) Ceuta(CE),(L) C.Valenciana(VC),(M)

Extremadura(EX),(N) Galicia(GA),(O) Madrid(MD),(P) Melilla(ME),(Q) Murcia(MC),(R) Navarra(NC),(S)

Pais Vasco(PV), and (T) La Rioja(RI). Day 1 corresponds to March 1st in 2020. Black dots show crude case

fatality risk, and light and dark indicate 95% and 50% credible intervals for posterior estimates from the Bayesian

model, respectively.

Fig 2: Temporal variation of risk of death caused by COVID-19 by area, Spain, March-May,

2020: time-delay adjusted case fatality risk

Observed and posterior estimates from the Bayesian model of time-delay adjusted case fatality risk in (A) Spain

(National), (B) Andalucia(AN),(C) Aragon(AR), (D) Asturias(AS),(E) Balears(IB),(F) Canarias(CN),(G)

Cantabria(CB),(H) Castilla-La Mancha(CM),(I) Castilla y Leon(CL), (J) Catalunya(CT),(K) Ceuta(CE),(L)

C.Valenciana(VC),(M) Extremadura(EX),(N) Galicia(GA),(O) Madrid(MD),(P) Melilla(ME),(Q)

Murcia(MC),(R) Navarra(NC),(S) Pais Vasco(PV), and (T) La Rioja(RI). Day 1 corresponds to March 1[st] in 2020.

Black dots show crude case fatality risk, and light and dark indicate 95% and 50% credible intervals for posterior

estimates from the Bayesian model, respectively.

Figure 3. Latest estimates of time-delay adjusted risk of death caused by COVID-19 by area,

2020, Spain (As of May 20, 2020).

Distribution of time-delay adjusted case fatality risks derived from the latest estimates (May 20, 2020) are

presented. Top to bottom: (A)Spain (National), (B) Andalucia(AN), (C)Aragon(AR), (D) Asturias(AS),(E)

Balears(IB), (F) Canarias(CN), (G) Cantabria(CB), (H) Castilla-La Mancha(CM), (I) Castilla y Leon(CL), (J)

Catalunya(CT), (K) Ceuta(CE),(L) C.Valenciana(VC),(M) Extremadura(EX),(N) Galicia(GA),(O)

Madrid(MD),(P) Melilla(ME),(Q) Murcia(MC),(R) Navarra(NC),(S) Pais Vasco(PV), and (T) La Rioja(RI)

Figure 4. Geographical variability of COVID-19 time-delay adjusted CFR, unemployment rate,

population size, and population density across 17 autonomous communities, Spain (As of May 20,

2020).

Distribution of time-delay adjusted case fatality risks derived from the latest estimates (May 20, 2020) are

presented. Top to b (A) time –delay adjusted case fatality risk as at May 20 2020 (B) Unemployment rate among

population aged 16 years and above in 2019 (C) Population size (in thousands) in 2019 (D) Population density

(per square km), 2017.

Time delay adjusted CFR by region
The latest estimate (As of May 20, 2020)