**Environmentally targeting surveillance using forest data and health facility surveys to detect malaria**

**transmission foci**

**Authors:** Kimberly M Fornace*[1,2], Ralph A Reyes[3], Maria Lourdes M Macalinao[3], Alison Paolo N Bareng[3],

Jennifer S Luchavez[3], Julius Clemence R Hafalla[1], Fe Esperanza J Espino[3], Chris J Drakeley[1]

1.  Faculty of Infectious and Tropical Diseases, London School of Hygiene and Tropical Medicine,

Keppel Street, London WC1E 7HT, United Kingdom

2.  Centre for Climate Change and Planetary Health, London School of Hygiene and Tropical

Medicine, Keppel Street, London WC1E 7HT, United Kingdom

3.  Department of Parasitology, Research Institute for Tropical Medicine, 9002 Research Drive,

Alabang, Muntilupa, 1781 Metro Manila, Philippines

**\* Corresponding author:**

Kimberly M Fornace

Kimberly.Fornace@lshtm.ac.uk

London School of Hygiene and Tropical Medicine

Keppel Street

London, WC1E 7HT, United Kingdom

+44 (0) 20 7927 2010

**Abstract:**

As malaria programmes move towards elimination, malaria transmission becomes increasingly spatially heterogenous, necessitating novel surveillance approaches to detect infections. Within Southeast Asia, forested landscapes are associated with both increased malaria transmission and reduced healthcare access. Here, we conduct health facility-based surveys using tablet-based applications and remote sensing data to geolocate and characterise environments of participant residences. Using a spatially explicit Bayesian process-based modelling approach, we quantified the detection probabilities of different surveillance approaches and estimated the distribution of malaria infections. We show that health facility surveys substantially increase spatial coverage of surveillance systems, with a probability of detection over three-fold greater than routine passive case detection (3.34, 95%BCI: 1.03 – 8.27). Our results also demonstrate closed canopy forested areas have higher proportions of subpatent malaria infections only identified by molecular diagnostics. Applying these findings, we show risk-based surveillance approaches incorporating forest data provide a cost-effective and operationally feasible method of identifying malaria foci.

**Significance:**

Surveillance is a critical component of malaria control and elimination programmes, aiming to identify malaria infections in heterogenous environments to target interventions. However, despite extensive research on the impact of landscapes on malaria transmission, landcover data rarely informs the design of malaria surveillance systems. Here, we adapt an ecological modelling framework to demonstrate how local environmental factors influence the spatial distributions of malaria infections, diagnostic sensitivity and detection probabilities at health facilities in the Philippines. Using convenience

sampling of health facility attendees, we demonstrate how risk-based surveillance incorporating forest

data can increase the probability of detecting malaria foci and estimate the underlying distribution of

malaria infections. This provides an operationally feasible and cost-effective method to characterise

transmission while accounting for imperfect detection from biased data. Our findings highlight the role

of environment in determining both malaria infection and detection rates and demonstrate the utility of

risk-based surveillance approaches.

**Background:**

Malaria transmission is highly variable spatially, driven by the geographical distribution of human

populations, mosquito vectors and the environments in which these populations interact (1).

Surveillance systems aim to identify these high-risk locations and populations in order to effectively

plan, implement and evaluate control policies and measures. As control programmes reduce incidence

and transmission decreases, spatial heterogeneity becomes more pronounced, necessitating

increasingly higher resolution data to detect foci of infection (2). This becomes increasingly critical in

rapidly changing environments, where changing human populations and vector habitats may cause

significant shifts in transmission patterns (3).

For passive surveillance systems relying on reported malaria case data, understanding spatial

distributions of risk is further challenged by underreporting due to health-seeking behaviour or

asymptomatic infections present in the community. Increasing evidence suggests the proportion of

asymptomatic malaria infections not detectable by standard diagnostics increases in low transmission

settings, resulting in large numbers of infections not detected by passive surveillance systems (4, 5).

These asymptomatic infections are commonly reported in older age groups, with potentially different

risk factors and spatial distributions from clinical malaria cases (6, 7). Population-based community

surveys remain the gold standard for assessing spatial patterns of infection; however, these sampling

approaches are highly resource intensive and may require very large sample sizes in low transmission

settings. Alternatively, more operationally feasible surveys of easy access groups, such as health facility

attendees or school children, are used to increase probability of detecting infections within the

community (e.g. (8-11)). However, both approaches targeting clinical cases and easy access groups are

inherently biased and do not fully capture the spatial distribution of infections.

Within ecology, estimates of species distribution or abundance are similarly challenged by imperfect

detection and spatially biased observation processes (12). Occupancy models are widely used to

estimate the probability that a species occupies a geographic location within a specified time period

while accounting for possible non-detection (13). This method partitions observation processes

determining detection probability and biological processes determining occupancy probability, each

associated with potentially overlapping spatial and environmental covariates. This makes the simple

assumption that a species cannot be detected if it is not present; however, if present, the species may or

may not be detected during sampling. In addition to allowing estimation of true occupancy states as a

latent variable, this method allows quantification of uncertainty in the observation process under

different sampling scenarios (14-16).

Here, we combine occupancy models with health facility surveys and molecular diagnostics to

evaluate the spatial coverage of surveillance approaches and their ability to detect locations with recent

malaria transmission. By estimating a spatially explicit sensitivity of different diagnostic methods, we

illustrate how environmental data can be used to develop operationally feasible risk-based surveillance

systems to increase the probability of detecting areas of infection while rationalising scarce resources.

This provides an adaptable framework to integrate convenience sampling approaches into existing

disease surveillance systems to target control measures and characterise spatial and environmental

drivers of infection from opportunistically collected data.

Using rolling cross-sectional health facility-based surveys in which all attendees regardless of

patient status or symptoms were screened for malaria using routine and molecular diagnostics, with

residences geolocated in real-time using tablet-based applications, we apply this approach to describe

the spatial distribution of malaria infections in Rizal, Palawan, The Philippines (Figure 1 (11)). While the

Philippines has made substantial progress towards malaria elimination, with 50 provinces declared

malaria free, Palawan contributes over 95% of malaria cases nationally, including 2718 cases from the

municipality of Rizal in 2018 (17). Deforestation rates have increased markedly within Rizal; of the 24%

decrease in forest cover between 2000 – 2018, over 50% of deforestation occurred after 2015, largely

driven by agricultural expansion (18). Within this region, malaria risks are strongly associated with

proximity to forested areas due to both vector ecology and occupational risks (19). Described as

"frontier malaria," factors associated with increased malaria risks, such as proximity to forest edges and

recent deforestation, are also associated with reduced healthcare access, resulting in potentially

complex interactions between detection bias and infection risks (20).

Our results illustrate that enhanced health facility-based surveys increase the probability of

detecting locations with malaria infections by markedly increasing the spatial coverage of the

surveillance system in addition to simply increasing the numbers of individuals screened. By comparing

locations of infections detected by routine diagnostics (microscopy and/ or rapid diagnostic tests (RDTs))

with sub-patent infections only detectable by molecular methods, we show the sensitivity of routine

diagnostics decreases in highly forested areas, with many locations of malaria transmission only

detectable by molecular methods. We demonstrate how these findings can be used to develop

operationally feasible and cost-effective environmentally targeted risk-based surveillance methods and

prioritise locations with high probabilities of infection not captured by existing surveillance systems.

**Results:**

*Impact of survey method on detection probability*

129             We conducted monthly rolling cross-sectional surveys at 27 health facilities across Rizal,

Palawan (Figure S1). All consenting individuals, regardless of symptoms or patient status, were screened

for malaria using microscopy or RDTs and polymerase chain reaction (PCR) as described by (21). Using

this data, we initially compared two surveillance approaches: 1. Standard - passive case detection (PCD),

including febrile patients screened using routine diagnostics (RDTs or microscopy) as per current

national surveillance guidelines; 2. Enhanced – standard PCD plus screening all health facility attendees

with both routine and molecular methods.

137             Between June 2016 – June 2017, 5767 individuals were enrolled in this study, including 1914

(33.2%) febrile patients screened for malaria by existing passive surveillance systems (21). Of these, 801

individuals (13.9%) were positive for malaria by molecular methods and 498 were positive by RDT or

microscopy. We geolocated all residences in Rizal (n=7313), identifying individuals screened by PCD from

698 unique locations while enhanced surveillance screened individuals from 2201 locations (Table S2).

Malaria infections were detected at 352 household locations using enhanced surveillance and 117

locations by standard PCD.

As control measures are targeted based on reported locations with malaria infection and over

80% of infected locations had only one infected individual detected, we chose to model whether malaria

infection was present or absent in a specific location (occupancy) rather than incidence. Detection

probabilities, the probability of screening at least one individual from a location during the study period,

varied geographically, with travel time to the nearest health facility negatively associated with detection

probability by both PCD and enhanced surveillance methods (Table 1). Enhanced surveillance increased

detection probabilities over three-fold compared to standard PCD (mean probability 3.34, 95%BCI: 1.03

– 8.27) in addition to markedly increasing spatial coverage of surveillance, particularly in rural

populations living in forested areas (Figure 2A, 2B).

*Spatial distribution of infection*

Incorporating these detection probabilities into hierarchical occupancy models revealed a much

wider spatial distribution of malaria detected by enhanced surveillance compared to PCD alone,

identifying areas of infection not captured by existing surveillance systems (Figure 2C, 2D). We identified

a range of differing spatial and environmental risk factors for infections detected by different diagnostic

methods; however, all infections were associated with proximity to closed canopy forests in more rural

populations (Table 1). For joint models for each surveillance scenario, incorporating a shared spatial

random effect between detection and infection probability improved model performance, suggesting a

common spatial process driving healthcare access and disease risks (Table S3).

**Table 1.** Posterior estimates of fixed effects and spatial range for joint models of A. standard PCD and B.
enhanced surveillance
168
169 A.

| | Mean | SD | 95% BCI |
|---|---|---|---|
| **Probability of detection** | | | |
| Distance to roads | 0.226 | 0.125 | -0.020, 0.227 |
| Travel time to clinic | -0.317 | 0.120 | -0.561, -0.090 |
| Distance to forest | -0.112 | 0.053 | -0.217, -0.009 |
| Spatial range (km) | 8.140 | 2.500 | 4.319, 14.050 |
| **Probability of infection** | | | |
| Distance from roads | 0.094 | 0.101 | -0.109, 0.287 |
| Population density | -0.603 | 0.145 | -0.894, -0.322 |
| Precipitation of wetness month | 0.212 | 0.107 | -0.006, 0.420 |
| Distance from closed canopy forest | -0.222 | 0.156 | -0.537, 0.078 |
| Spatial range (km) | 1.752 | 1.096 | 0.493, 4.643 |
| Scaling parameter for shared spatial effect | 0.590 | 0.163 | 0.282, 0.921 |

170 ** All covariates mean-centred and scaled
171
172 B.

| | Mean | SD | 95% BCI |
|---|---|---|---|
| **Probability of detection** | | | |
| Population density | -0.533 | 0.085 | -0.701, -0.368 |
| Travel time to clinic | -0.511 | 0.107 | -0.729, -0.309 |
| Aspect | 0.094 | 0.039 | 0.017, 0.171 |
| Spatial range (km) | 15.804 | 6.130 | 7.212, 30.957 |
| **Probability of infection** | | | |
| Distance from roads | 0.285 | 0.071 | 0.145, 0.423 |
| Upslope area | 0.181 | 0.111 | -0.038, 0.399 |
| Topographic wetness index | -0.243 | 0.120 | -0.481, -0.011 |
| Temperature annual range | 0.236 | 0.098 | 0.043, 0.428 |
| Distance from closed canopy forest | -0.326 | 0.111 | -0.548, -0.112 |
| Spatial range (km) | 0.897 | 0.232 | 0.532, 1.438 |
| Scaling parameter for shared spatial effect | 1.216 | 0.177 | 0.887, 1.179 |

173 ** All covariates mean-centred and scaled
174
175

To explore the factors determining these differing distributions of infections, we estimated

probability of patent malaria detectable by RDT or microscopy in all malaria infections identified.

Malaria infected individuals were identified from 435 locations and over one third of infected individuals

(37.8%, 95% CI: 34.5-41.3%) could only be detected by molecular methods. Subpatent malaria was

substantially more common in forested areas with the odds of patent infections increasing 1.23 (95%BCI

1.03-1.47) with every kilometre distant from closed canopy forested areas (Table 2). Using data from all

residence locations identified within Rizal, we predicted a location-specific probability of patent malaria,

equivalent to the sensitivity of routine diagnostics (RDT and/or microscopy) (Figure S2).

**Table 2.** Posterior estimates of fixed effects for malaria positive households detected by routine
diagnostics (RDT and microscopy) compared to molecular methods

|  | Mean | SD | 95% BCI |
|---|---|---|---|
| Distance from closed canopy forest* | 0.225 | 0.112 | 0.036, 0.476 |
| Annual precipitation* | -0.318 | 0.121 | -0.620, -0.145 |
| Precipitation of wettest month* | 0.256 | 0.085 | 0.091, 0.422 |
| Population density$^2$* | 0.093 | 0.087 | -0.078, 0.263 |

187    * Mean-centred and scaled
188
189

*Evaluating surveillance systems*

As our primary aim was to identify all locations of malaria infection, we evaluated surveillance

approaches against estimates of true infection status using molecular diagnostics and all available

spatial data. Accounting for spatial bias of collection data, we estimated 11.4% (95%BCI: 4.6%-21.9%) of

all locations in Rizal had malaria infections during the sampling period. While no surveillance method

identified all areas of infection perfectly, enhanced surveillance using molecular methods identified

38.7% (95% BCI: 33.6-43.8%) of infected locations while PCD only identified 5.7% (95% BCI: 0.1 – 11.9%).

Including molecular diagnostics in PCD only slightly improved the probability of detecting infections

while conducting health facilities surveys using only routine diagnostics decreased the number of

infected locations identified (Figure 3A, Figure S4). We additionally identified 247 locations with a very

low (<0.05% probability) of detection by any health facility-based surveillance.

Based on distributions of malaria infections and probability of detection by routine diagnostics,

we additionally explored the use of environmentally-stratified risk-based surveillance approaches. We

defined high-risk areas based on distances from closed canopy forest (Figure S3). Evaluating costs of

each surveillance system relative to baseline costs of standard PCD (Methods, SI), we estimated the total

cost per location with malaria infection identified (Figure 3B). As the cost per infection detected was

most sensitive to inclusion of molecular diagnostics, we defined a risk-based surveillance approach using

health facility surveys in all areas and only applying molecular diagnostics to locations within 100m of

closed canopy forest areas (Table 3, Figure S5). This risk-based surveillance approach almost halved the

cost of detecting a location of infection compared to enhanced surveillance, from USD 603.10 (95% BCI:

530.02 – 690.82) to USD 370.00 (95% BCI: 313.18- 444.04) while detecting only slightly fewer locations

of infections.

**Table 3.** Surveillance methods assessed
215

|  | Survey method | | Diagnostic method | | Total cost (USD) |
|---|---|---|---|---|---|
|  | Passive case detection | Health facility surveys | Routine (RDT/ microscopy) | Molecular |  |
| 1: Standard PCD | X |  | X |  | - |
| 2: Enhanced surveillance | X | X | X | X | 193,547.70 |
| 3: PCD + molecular | X |  | X | X | 56,654.40 |
| 4: Health facility surveys + routine | X | X | X |  | 22,844.50 |
| 5: Risk-based surveys + diagnostics | X | Risk zone only | X | Risk zone only |  |

216
217

**Discussion**

The spatial distribution of malaria is driven by a complex interplay of environmental and social

factors influencing both disease transmission and identification by health systems. Statistical methods

enabling examination of processes driving infection and detection provide an invaluable tool to evaluate

the roles of environmental factors and develop targeted surveillance approaches. Here, we demonstrate

a convenience sampling approach using health facility surveys markedly increases spatial coverage of

surveillance systems; incorporating these surveys with satellite-derived remote sensing data allows

estimation of the underlying distribution of infection not captured by passive surveillance. We

additionally show higher proportions of subpatent malaria infections in forested areas, highlighting the

limited utility of routine diagnostics within these regions. Applying these findings, we develop a cost-

effective and operationally feasible risk-based surveillance approach using environmental data and

illustrate how landscape data can be incorporated into disease surveillance systems.

Despite extensive research identifying proximity to forests as risk factors for malaria infection in

Southeast Asia (e.g. (3, 22-27)), landscape data are not routinely used to design or inform surveillance

systems. Malaria control programmes typically conduct community-based active case detection in

response to reported malaria cases (2); however, we show this may miss a substantial proportion of

active malaria foci due to biases in health-seeking behaviour and increased prevalence of subpatent

malaria within higher transmission areas. Although mechanisms driving this relationship between forest

cover and subpatent malaria are not known, patent malaria infections are more common in children in

this area and settlements in closer proximity to forests may have different demographic compositions

(e.g. logging and plantation camps) (21). Previous studies have also suggested a role for immunity in

high transmission areas, with individuals repeatedly exposed to malaria commonly having lower parasite

densities (4). Despite lower parasite densities, these subpatent infections contribute to infectious

reservoirs and may have a critical role in sustaining transmission in elimination settings, highlighting the

importance of identifying and treating these individuals (28).

While screening entire populations may be prohibitively costly and intensive, the increasing

availability of satellite-based remote sensing data provides new opportunities to use environmental data

to target surveillance activities (29). Surveillance systems for malaria, as well as for other low incidence

or emerging diseases, are challenged by the need to identify relatively rare events with shifting spatial

patterns. Widely used in veterinary epidemiology, risk-based surveillance uses known risk factors to

focus intensive surveillance activities on the populations where rare events are most likely to occur (30).

Integrating remote-sensing derived environmental data into this approach provides an adaptable

framework which can be easily updated in changing landscapes. Further, quantifying the detection

probabilities associated with these surveillance approaches allows estimation of biases in data and

identification of populations not captured by opportunistic sampling which can be used to plan active

case detection. For this example, using health facility surveys, this enables evaluation of spatial patterns

of health seeking behaviour as well as populations outside health system coverage. Screening all

individuals attending health facilities, including those accompanying patients or for nonfebrile illnesses,

vastly increases the spatial coverage of surveillance. The improved performance of models including a

common spatial term for both infection and detection suggests processes determining healthcare

coverage also influence infection risks, demonstrating the multiple utilities of measuring health seeking

behaviour in this context.

Within this study, remote sensing data was used not only as covariates but also to define the

populations at risk. The application of machine learning approaches to identify building footprints from

very high resolution satellite imagery has produced increasingly accurate estimates of household

distributions and allows estimation of spatial distribution of the populations (31). Development of

tablet-based methods including population data and satellite imagery enabled near real-time

identification of the residences of health facility attendees in a rural population with no formal

addresses and limited internet connectivity (11). These tools and datasets can be further expanded to

create accessible interfaces for local health workers to use environmental and spatial data and

incorporate risk-based decision pathways on screening procedures and diagnostic tests based on

household locations and travel history. This approach can also be easily modified to include multiple

diseases with different underlying environmental risk factors and updated to include new environmental

data, such as near real time deforestation alerts (32). Despite advances in using meteorological data to

forecast vector-borne disease epidemics, landscape data is rarely used operationally and may provide

more actionable information within rapidly changing environments.

Despite the utility of these methods, there were several limitations to this study. As this study

was designed to identify spatial locations of malaria infections within the sampling year, we did not

explore temporal patterns of infection or health seeking behaviours. However, the modelling approach

used is easily extendable to incorporate dynamic state-space models of changes in infection over time

(12). If health facility surveys were collected over longer periods of time, this could additionally be

expanded to incorporate seasonally varying meteorological data to further refine risk stratifications.

Additionally, while populations at risk were defined using multiple datasets, this is likely to have limited

coverage of highly mobile indigenous populations not residing in permanent structures. Future work

could explore the utility of satellite imagery to identify these populations, such as through monitoring of

forest disturbance or modelling movement patterns.

Even with these limitations, this provides a novel and adaptable surveillance approach for

environmentally driven diseases and demonstrates the role of landscapes in driving malaria infection

and detection within this region. Incorporation of forest data enables identification of cost-effective risk-

based surveillance approaches which increase probabilities of detecting malaria infections and can be

applied to support elimination efforts. Additionally, the process-based ecological modelling method

used provides a flexible framework to quantify detection probabilities and estimate the true spatial

distribution of infection using biased data from convenience-based sampling approaches.

**Methods:**

*Survey approaches*

We conducted cross-sectional health facility-based surveys at 27 facilities, including at one

Regional Health Unit (RHU), 9 Barangay Health Stations (BHS) and 17 RDT centres based in community

health worker households, established to increase access to malaria diagnosis and treatment (Figure S1).

For one week every month between June 2016 – June 2017, all individuals attending the health facility,

regardless of symptoms or patient status, were asked to participate in this survey. For consenting

individuals, malaria status was assessed using either microscopy or rapid diagnostic tests (RDTs), with

finger-prick blood samples collected on Whatman 3MM filter paper for subsequent polymerase chain

reaction (PCR) (21). We classified malaria as positive for any *Plasmodium* species as 75% of identified

species were *P. falciparum* and all malaria species are transmitted by common vectors in this area.

Household locations were identified on offline maps and basic demographic information was recorded

using GeoODK on Android tablets (11).

*Spatial and environmental covariates*

To define the locations of all households within Rizal, we extracted information on household

structure locations from the Facebook High Resolution Settlement Layer, a 30m resolution satellite-

based remote sensing derived dataset on all inhabited structures (33). This dataset was combined with

all reported households identified by survey participants and geolocated households from the 2015

Philippine census (34). We resampled all datasets to 50m resolution and removed duplicate locations to

account for spatial resolution of datasets and inaccuracies of reported household locations. To estimate

detection probabilities, we classified households included by a surveillance scenario if one or more

individuals were sampled. Similarly, households were classified as infected if one or more individuals

were identified as infected by diagnostics used by the surveillance approach.

Plausible covariates used to model detection or infection probabilities were assembled (Table

S1). Handheld GPS devices (Garmin, USA) were used to record locations of all sampled clinics and roads

within the region. Travel time to the nearest sampled clinic was calculated as accumulated cost from

friction surfaces extracted from (35). Additional environmental and spatial covariates included

population density (36), Euclidean distance from roads and bioclimatic variables (37). Elevation and

topographic measures, including topographic wetness index (TWI), upslope area and aspect, were

calculated from the ASTER Global Digital Elevation Model (38). Forest cover was classified as over 50%

canopy cover based on (18) and Euclidean distance was calculated to the forest edge, recent

deforestation within the past year and historical deforestation within the previous five years. We

additionally included closed canopy forest, defined as canopy cover over 90% with a minimum area of

0.5ha (39).

*Occupancy modelling*

Covariates were extracted for all identified household locations and Pearson correlation analysis

was used to assess multicollinearity between variables. To select variables for inclusion, nonspatial

binomial generalised linear models were fit separately for detection probabilities and infection

probabilities for each surveillance scenario using backwards stepwise model selection with a five-point

threshold for improvement in deviance information criteria (DIC) to minimise overfitting (40). Residual

spatial autocorrelation was assessed using Moran's I and performance assessed by area under the

receiver operating curve (AUC). Weakly informative priors of Normal (0, 1/0.01) were used for all

intercepts and coefficients. All models were implemented in Integrated Nested Laplace (INLA), with

10,000 posterior samples generated from the approximated posterior distribution to include uncertainty

in these estimates (41). Final models had an AUC of 85% for both surveillance approaches.

We modelled the distribution of infections under each surveillance scenario separately using

occupancy models in which the probability of detecting an infection ($y_i$) in location $i$ is dependent on the

probability of detection ($p_i$) and presence of infection ($\omega_i$) (13), modelled as:

$$y_i \sim \begin{cases} 0, & \omega_i = 0 \\ \text{Bernoulli}(p_i), & \omega_i = 1 \end{cases}$$

355    Where the linear predictor determining the probability of detection is modelled as:

$$\text{logit}(p_i) = \alpha_0 + X_i\alpha + u_i$$

358    Where $\alpha_0$ represents the intercept, $X_i\alpha$ represents a vector of covariate effects and $u_i$ is the spatial

359    effect modelled as a Matern covariance function using the stochastic partial differential equations

360    approach to represent the spatial process by Gaussian Markov random fields as implemented in INLA

361    (41, 42). The process determining the true state of malaria presence $\omega$ is determined by the true

362    probability of infection $\psi$:

$$\omega_i \sim \text{Bernoulli}(\psi_i)$$

364    With the linear predictor for the Bernoulli model specified as:

$$\text{logit}(\psi_i) = \beta_0 + X_i\beta + \gamma_i + Zu_i$$

Where $\beta_0$ represents the intercept, $\boldsymbol{X_i\beta}$ represents a vector of covariate effects and $\gamma_i$ represents the

spatial effect, modelled as described above. As processes influencing probability of detection

(healthcare access) additionally may impact infection, we include a shared spatial component $Zu_i$ with

scaling parameter $Z$ (43).

*Patent malaria distribution*

To explore factors affecting the distribution of locations of patent malaria infections (detected

by RDT or microscopy) compared to all infections, we subset all malaria infected locations. For $J_j$ malaria

infected individuals identified in each location, the number of patent infections observed ($m_i$) is

modelled as:

$$m_i \sim \mathrm{Binomial}(J_i, s_i)$$

379 With the linear predictor determining the probability of patent infections ($s_i$) modelled as:

$$\mathrm{logit}(s_i) = \kappa_0 + \boldsymbol{X_i\kappa}$$

Where $\kappa_0$ represents the intercept and $\boldsymbol{X_i\kappa}$ represents a vector of covariate effects. Using data from all

locations included in the study site, we then predicted a location specific sensitivity of routine

diagnostics. Based on these results, we used environmental data to define an area with higher

probabilities of malaria infections only detectable by molecular diagnostics.

*Evaluation of surveillance systems*

We modelled the true probability of infection from the infection process model using data from

all available diagnostics. To compare surveillance methods with different survey and diagnostic

methods, we estimated the number of infected locations not detected as:

$$\sum_{i=1}^{i} \omega_i(1 - p_i) + \omega_i p_i(1 - s_i)$$

Where $p_i$ is the probability of detection using different survey methods and $s_i$ represents diagnostic

sensitivity, with PCR considered the gold standard. We additionally included risk-based surveillance

methods, using health facility surveys and molecular diagnostics in high risk areas defined by proximity

to closed canopy forest. All derived quantities were estimated using 10,000 posterior samples.

To evaluate the cost effectiveness of different surveillance approaches, we estimated the

additional costs to health systems of including different survey and diagnostic approaches (Table S4).

This excluded capital costs and costs already covered by existing health systems (e.g. RDT and

microscopy diagnostics for febrile patients). Health facility survey costs included additional payments to

personnel, training, equipment, RDT and microscopy for non-febrile participants, sample collection and

molecular diagnostics for all attendees and salary for a technician to support data management and

sample analysis. The costs of molecular diagnostics included DNA extraction and PCR reagents, assuming

all DNA extraction was completed using Chelex with 10% of samples verified using a commercial Qiagen

kit. To account for varying numbers of samples screened from each location, we estimated the mean

cost of molecular diagnostics per location as the total cost of molecular diagnostics divided by the total

number of locations included. We evaluated these against the estimated number of locations correctly

identified as a measure of cost effectiveness. All analysis was completed in R statistical programming

language (v 3.6), with maps visualised in R or ArcGIS (ESRI, Redlands, USA).

*Ethics approval*

This study was approached by the Institutional Review Board of the Research Institute for Tropical

Medicine, Department of Health Philippines (IRB:2016-04) and the Research Ethics Committee of the

London School of Hygiene and Tropical Medicine (11597). Written informed consent was obtained from

all participants or parents or guardians and assent obtained from children under 18.

1.    Lambin EF, Tran A, Vanwambeke SO, Linard C, & Soti V (2010) Pathogenic landscapes:
interactions between land, people, disease vectors, and their animal hosts. *International journal*
*of health geographics* 9:54.
2.    World Health Organisation (2018) Malaria surveillance, monitoring and evaluation: a reference
manual. (World Health Organization, Geneva).
3.    Yasuoka J & Levins R (2007) Impact of deforestation and agricultural development on
anopheline ecology and malaria epidemiology. *The American journal of tropical medicine and*
*hygiene* 76(3):450-460.
4.    Okell LC*, et al.* (2012) Factors determining the occurrence of submicroscopic malaria infections
and their relevance for control. *Nature communications* 3:1237.
5.    Okell LC, Ghani AC, Lyons E, & Drakeley CJ (2009) Submicroscopic infection in Plasmodium
falciparum-endemic populations: a systematic review and meta-analysis. *The Journal of*
*infectious diseases* 200(10):1509-1517.
6.    Zhou G, Afrane YA, Malla S, Githeko AK, & Yan G (2015) Active case surveillance, passive case
surveillance and asymptomatic malaria parasite screening illustrate different age distribution,
spatial clustering and seasonality in western Kenya. *Malaria journal* 14(1):41.
7.    Hsiang MS*, et al.* (2019) Active case-finding for malaria: A three-year national evaluation of
optimal approaches to detect infections and hotspots through reactive case detection in the low
transmission setting of Eswatini. *Clin Infect Dis*.
8.    Ashton RA*, et al.* (2015) Geostatistical modeling of malaria endemicity using serological
indicators of exposure collected through school surveys. *The American journal of tropical*
*medicine and hygiene* 93(1):168-177.
9.    Stevenson JC*, et al.* (2013) Reliability of school surveys in estimating geographic variation in
malaria transmission in the western Kenyan highlands. *PloS one* 8(10):e77641.
10.    Stresman GH*, et al.* (2014) High levels of asymptomatic and subpatent Plasmodium falciparum
parasite carriage at health facilities in an area of heterogeneous malaria transmission intensity
in the Kenyan highlands. *The American journal of tropical medicine and hygiene* 91(6):1101-
1108.
11.    Fornace KM*, et al.* (2018) Use of mobile technology-based participatory mapping approaches to
geolocate health facility attendees for disease surveillance in low resource settings.
*International journal of health geographics* 17(1):21.
12.    Bailey LL, MacKenzie DI, & Nichols JD (2013) Advances and applications of occupancy models.
*Methods in Ecology and Evolution* 5(12):1269-1279.
13.    Hobbs NT & Hooten MB (2015) *Bayesian Models: A Statistical Primer for Ecologists* (Princeton
University Press, New Jersey, USA).
14.    Banks-Leite C*, et al.* (2014) Assessing the utility of statistical adjustments for imperfect detection
in tropical conservation science. *J. Appl. Ecol.* 51(4):849-859.
15.    Nelli L, Ferguson HM, & Matthiopoulos J (2019) Achieving explanatory depth and spatial breadth
in infectious disease modelling: Integrating active and passive case surveillance. *Stat Methods*
*Med Res*:962280219856380.
16.    Salvador LCM, Deason M, Enright J, Bessell PR, & Kao RR (2018) Risk-based strategies for
surveillance of tuberculosis infection in cattle for low-risk areas in England and Scotland.
*Epidemiol Infect* 146(1):107-118.
17.    Epidemiology Bureau (2018) Malaria surveillance report. ed Health PDo (Manila).
18.    Hansen MC*, et al.* (2013) High-resolution global maps of 21st-century forest cover change.
*Science* 342(6160):850-853.
19.    DOH P, WHO, & UCSF (2014) Eliminating malaria: case-study 6, Progress towards subnational
elimination in the Philippines. (World Health Organisation, Geneva).

20.  de Castro MC, Monte-Mor RL, Sawyer DO, & Singer BH (2006) Malaria risk on the Amazon
frontier. *Proceedings of the National Academy of Sciences of the United States of America*
103(7):2452-2457.
21.  Reyes RA, *et al.* (submitted) Improved detection of malaria infection through enhanced health
facility surveys.
22.  Sanann N, *et al.* (2019) Forest work and its implications for malaria elimination: a qualitative
study. *Malaria journal* 18(1):376.
23.  Hasyim H, *et al.* (2018) Spatial modelling of malaria cases associated with environmental factors
in South Sumatra, Indonesia. *Malaria journal* 17(1):87.
24.  Guerra CA, Snow RW, & Hay SI (2006) A global assessment of closed forests, deforestation and
malaria risk. *Annals of tropical medicine and parasitology* 100(3):189-204.
25.  Fornace KM, *et al.* (2016) Association between Landscape Factors and Spatial Patterns of
Plasmodium knowlesi Infections in Sabah, Malaysia. *Emerging infectious diseases* 22(2):201-208.
26.  Durnez L, *et al.* (2013) Outdoor malaria transmission in forested villages of Cambodia. *Malaria
journal* 12:329.
27.  Cui L, *et al.* (2012) Malaria in the Greater Mekong Subregion: heterogeneity and complexity.
*Acta tropica* 121(3):227-239.
28.  Slater HC, *et al.* (2019) The temporal dynamics and infectiousness of subpatent Plasmodium
falciparum infections in relation to parasite density. *Nature communications* 10(1):1433.
29.  Hay SI, George DB, Moyes CL, & Brownstein JS (2013) Big data opportunities for global infectious
disease surveillance. *PLoS medicine* 10(4):e1001413.
30.  Stark KD, *et al.* (2006) Concepts for risk-based surveillance in the field of veterinary medicine
and veterinary public health: review of current approaches. *BMC Health Serv. Res.* 6:20.
31.  Tiecke TG, *et al.* (2017) Mapping the world population one building at a time. *ArXiv*
arXiv:1712.05839.
32.  Hansen MC, *et al.* (2016) Humid tropical forest disturbance alerts using Landsat data.
*Environmental Research Letters* 11(3).
33.  Facebook Connectivity Lab & Center for International Earth Science Information Network
(CIESIN) Columbia University (2016) High resolution settlement layer (HRSL).  (CIESIN, Columbia
University, New York).
34.  Philippine Statistics Authority (2016) Philippine Population Density (Based on the 2015 Census of
Population).
35.  Weiss DJ, *et al.* (2018) A global map of travel time to cities to assess inequalities in accessibility
in 2015. *Nature* 553(7688):333-336.
36.  Lloyd CT, Sorichetta A, & Tatem AJ (2017) High resolution global gridded data for use in
population studies. *Sci Data* 4:170001.
37.  Fick SE & Hijmans RJ (2017) Worldclim 2: New 1-km spatial resolution climate surfaces for global
land areas. *International Journal of Climatology*.
38.  Land Processes Distributed Active Archive Center (LP DAAC) (2015) Advanced Spaceborne
Thermal Emission and Reflection Radiometer Global Digital Elevation Model (ASTER GDEM)
Version 2.  (NASA EOSDIS Land Processes DAAC, USGS Earth Resources Observatoin and Science
(EROS) Center, Sioux Falls, South Dakota).
39.  ALOS-2/ALOS Science Project Earth Observation Research Center (EORC) (2017) Global PALSAR-
2/PALSAR/JERS-1 Mosaic and Forest/Non-Forest map.  (Japan Aerospace Exploration Agency
(JAXA)).
40.  Redding DW, Tiedt S, Lo Iacono G, Bett B, & Jones KE (2017) Spatial, seasonal and climatic
predictive models of Rift Valley fever disease across Africa. *Philosophical transactions of the
Royal Society of London. Series B, Biological sciences* 372(1725).

41. Rue H, Martino S, & Chopin N (2009) Approximate Bayesian inference for latent Gaussian
models by using integrated nested Laplace approximations. *Journal of the Royal Statistical*
*Society* 71(Part 2):319-392.
42. Lindgren F & Rue H (2015) Bayesian Spatial Modelling with R-INLA. *Journal of Statistical*
*Software* 63(19).
43. Martins TG, Simpson D, Lindgren F, & Rue H (2013) Bayesian computing with INLA: new
features. *Computational Statistics and Data Analysis* 67:68-83.

520
521
522 **Figure 1.** Study area and forest cover in 2017
523
524 **Figure 2.** Posterior probability of infection under different sampling scenarios adjusted for detection
525 probabilities: A. detection probability using routine passive case detection; B. detection probability using
526 health facility-based surveys; C. probability of infection estimated from passive case detection using
527 routine diagnostics; D. probability of infection estimated from active case detection using molecular
528 diagnostics
529
530 **Figure 3.** Evaluation of surveillance methods described in Table 3 by A. estimated numbers of locations
531 with malaria infections not detected, and B. estimated additional costs per location with malaria
532 infections detected (relative to standard PCD)
533