Epidemiological and Genomic Analysis of SARS-CoV-2 in Ten Patients from a Mid-sized

City outside of Hubei, China in the Early Phase of the COVID-19 Outbreak

Jinkun Chen[1§], Evann E. Hilt[2§], Huan Wu[3], Zhuojing Jiang[1], QinChao Zhang[1], JiLing Wang[1],

Yifang Wang[3], Fan Li[4], Ziqin Li[5], Jialiang Tang[1*], Shangxin Yang[2,5*]

[1]Shaoxing Center for Disease Control and Prevention, Shaoxing, Zhejiang, China; [2]Department

of Pathology and Laboratory Medicine, University of California Los Angeles, Los Angeles, CA,

USA; [3]IngeniGen XunMinKang Biotechnology Inc., Shaoxing, Zhejiang, China; [4]Three Coin

Analytics, Inc. Pleasanton, CA, USA; [5]Zhejiang-Californina International Nanosystems Institute,

Zhejiang University, Hangzhou, Zhejiang, China

§Authors contributed equally.

*Corresponding authors: Jialiang Tang: 992488904@qq.com; Shangxin Yang, PhD, D(ABMM):

shangxinyang@mednet.ucla.edu

RUNNING TITLE: Genomic Epidemiology of SARS-CoV-2 in Shaoxing

KEYWORDS: 2019-nCoV, COVID-19, Genotype, Metagenomic Sequencing, Mutation Rate,

SARS-CoV-2

ABSTRACT

A novel coronavirus known as severe acute respiratory syndrome coronavirus 2 (SARS-

CoV-2) is the cause of the ongoing COVID-19 pandemic. In this study, we performed a

comprehensive epidemiological and genomic analysis of SARS-CoV-2 genomes from ten

patients in Shaoxing (Zhejiang Province), a mid-sized city outside of the epicenter Hubei

province, China, during the early stage of the outbreak (late January to early February, 2020).

We obtained viral genomes with > 99% coverage and a mean depth of 296X demonstrating that

viral genomic analysis is feasible via metagenomics sequencing directly on nasopharyngeal

samples with SARS-CoV-2 Real-time PCR C_t values less than 28. We found that a cluster of 4

patients with travel history to Hubei shared the exact same virus with patients from Wuhan,

Taiwan, Belgium and Australia, highlighting how quickly this virus spread to the globe. The

virus from another cluster of two family members living together without travel history but with

a sick contact of a confirmed case from another city outside of Hubei accumulated significantly

more mutations (9 SNPs vs average 4 SNPs), suggesting a complex and dynamic nature of this

outbreak. Our findings add to the growing knowledge of the epidemiological and genomic

characteristics of SARS-CoV-2 and offers a glimpse into the early phase of this viral infection

outside of Hubei, China.

INTRODUCTION

Coronaviruses (CoVs) are a large family of single-stranded RNA viruses that can be isolated from a variety of animals including camels, rats, birds and bats (Cascella et al., 2020). These coronaviruses can cause a range of disease states in animals including respiratory, enteric, hepatic and neurological disease (Weiss and Leibowitz, 2011). Before late 2019, there were six known CoVs capable of infecting humans (Hu-CoVs). The first four Hu-CoVs that cause mild disease are HKU1, NL63, OC43 and 229E and are known to circulate in the human population (Corman et al., 2018). The other two Hu-CoVs, known as severe acute respiratory syndrome-CoV (SARS-CoV) and middle east respiratory syndrome-CoV (MERS-CoV), caused two previous epidemics in 2003 (Rota et al., 2003) and 2012 (Zaki et al., 2012) respectively. Both SARS-CoV and MERS-CoV were the results of recent spillover events from animals. These two epidemics highlighted how easy it is for spillover events in CoVs to occur and cause outbreaks in humans.

In December 2019, another spillover event occurred and a seventh Hu-CoV appeared known as severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), previously named 2019-nCoV (Wu et al., 2020). SARS-CoV-2 has been spreading rapidly across the world since it was first reported in Wuhan, Hubei province, China (Wang et al., 2020a;Wu et al., 2020). The advances and accessibility of sequencing technologies have allowed researchers all over the world to quickly sequence the genome of SARS-CoV-2 (Shu and McCauley, 2017;Zhou et al., 2020). Zhou *et al.* 2020 showed that SARS-CoV-2 shared 79.6% sequence identity to SARS-CoV and 96% sequence identity to a bat CoV further supporting the theory of another spillover event (Zhou et al., 2020).

Genomic analysis of SARS-CoV-2 genomes suggested that there were two major

genotypes in the early phase of the outbreak, known as L type and S type, based on almost

complete linkage between two SNPs (Tang et al., 2020). Tang *et al.* 2020 proposed that the L

type may be more aggressive in replication rates and spreads more quickly (Tang et al., 2020). A

recent reclassification of SARS-CoV-2 was proposed, and most viruses of the S type and L type

described in the Tang's study are now classified as Lineage A and B, respectively (Rambaut et

al., 2020). Here we present a comprehensive epidemiological and genomic analysis of SARS-

CoV-2 genomes from 10 patients in Shaoxing (Zhejiang Province), a mid-sized city about 500

miles away from Wuhan at the early stages of the outbreak.

MATERIALS AND METHODS

Study design and Ethics

Ten remnant nasopharyngeal swab samples collected between 1/27/2020 and 2/7/2020,

and tested positive by a SARS-CoV-2 real-time PCR assay with cycle threshold (C_t) values of

less than 28, were included in this study. The samples were de-identified except the associated

epidemiological data were retained. Since the patient identification was removed and the samples

used in this study were remnant and otherwise would be discarded, the Shaoxing Center for

Disease Control and Prevention had determined that the institutional review boards (IRB)

approval was waived for this project, and the informed consent form was not required.

SARS-CoV-2 PCR & RNA Sequencing

Total nucleic acid was extracted from the nasopharyngeal swabs using the Total Nucleic

Acid Extraction Kit (IngeniGen XMK Biotechnologies, Inc. Zhejiang, China). Real-time PCR

was performed by using the IngeniGen XMKbio 2019-nCoV (SARS-CoV-2) RNA Detection kit,

which targets the highly specific sequences in the *ORF1ab* and *N* genes of the virus, on the ABI

7500 system (ThermoFisher Scientific, Inc. MA, USA). The RNA libraries were constructed

using the Ingenigen XMKbio RNA-seq Library Prep Kit (IngeniGen XMK Biotechnologies, Inc.

Zhejiang, China). Briefly, DNase was used to remove residual human DNA and the RNA was

fragmented, followed by double-strand cDNA synthesis, end-repair, dA-tailing and adapter

ligation. Sequencing was performed by using the 2X75bp protocol on the Nextseq 550 system

(Illumina, Inc. CA, USA).

Data Analysis

Quality control and trimming of paired-end reads was performed using custom Python

scripts as follows: 1) trim 3' adapters; 2) trim reads at ambiguous bases; 3) filter reads shorter

than 40bp; 4) filter reads with average quality score < 20. Host-derived reads were removed by

alignment against the GRCh38.p13 genome reference using bowtie2 (v2.3.4.3) with default

parameters. The retained reads were then mapped to 163 published SARS-CoV-2 reference

genomes obtained from GISAID (https://www.gisaid.org/CoV2020/, accessed March 2, 2020) by

bowtie2 (v2.3.4.3) with default parameters. snippy (v4.5.0) was used for variant calling and core

SNP alignment against the Wuhan-Hu-1 reference, FastTree (v2.1.3) was used for tree

construction using default parameters, and Figtree (v1.4.4) was used to visualize the resulting

phylogenetic tree. Additional statistical analyses and visualizations were performed using custom

Python scripts with the pandas (v0.25.0) and matplotlib (v3.1.1) modules. The L/S type was

determined according to methods previously described (Tang et al., 2020). Briefly, C at position

8782 and T at position 28144 was determined to be A type, and T at position 8782 and C at

position 28144 was determined to be B type. The A/B lineage was determined by using pipeline

pangolin (https://github.com/hCoV-2019/pangolin) as described previously (Rambaut et al.,

2020).

RESULTS

<u>Epidemiology of Shaoxing Patients</u>

All ten patients presented with symptoms (fever and cough) consistent with COVID-19 in

late January and early February of 2020. The majority of patients were male (60%) and the

average age was 44 (**Table 1**). The patients can be categorized into two epidemiologic groups

with either a travel history to the Hubei province or sick contact with a confirmed case (**Table 1**).

There was one case where we were unable to obtain a travel or exposure history (Shaoxing-8).

Table 1. Epidemiological History of the 10 Shaoxing Patients

ID	Age Range	History of Travel or Sick Contact	Date of Symptom Onset	Date of Sample Collection
Shaoxing-01	30-39	Family members traveled together to Hubei province (1/15 – 1/24)	1/24/20	1/27/20
Shaoxing-02	70-79		1/29/20	1/30/20
Shaoxing-03	60-66		1/28/20	1/30/20
Shaoxing-04	50-59		1/29/20	1/31/20
Shaoxing-05	50-59	Traveled to Hubei (1/16 – 1/23)	1/29/20	1/31/20
Shaoxing-06	30-39	Resident of Wuhan; Traveled to Shaoxing on 1/17	1/29/20	1/31/20
Shaoxing-07	<10	Traveled to Hubei (1/11 – 1/24); Two family members were confirmed cases	1/30/20	1/30/20
Shaoxing-08	50-59	Unknown	1/31/20	2/7/20
Shaoxing-09	30-39	Family members living together no travel history; Contact with a confirmed case from Ningbo, Zhejiang on 1/27	2/2/20	2/5/20
Shaoxing-10	30-39		2/5/20	2/6/20

There are two apparent clusters in these ten patients. The first cluster involves four

patients who are relatives and traveled together to Hubei province for a wedding in late January.

The first patient in this cluster had symptom onset on their last day in Hubei province while the

other three patients had symptom onset 4-5 days after coming back to Shaoxing (**Table 1**). The

second cluster involves two patients who are family members that live together and did not travel

to Hubei province. One of the family members (Shaoxing-09) had a sick contact with a

confirmed case who visited her but lived in Ningbo, a more populated city in Zhejiang province

(**Table 1**).

Metagenomic Sequencing

The patients were confirmed to have SARS-CoV-2 infection by a commercial Real-time

PCR assay. The average C_t values for the 10 patient samples were 23.17 for *ORF1ab* and 24.54

for *N* (**Table 2**). Metagenomic sequencing was performed to recover the full viral genome. The

total number of sequence reads per samples ranged from 10.4 million to 27.5 million with an

average of 17.1 million. A small percentage of these reads mapped to SARS-CoV-2 RNA (**Table

2**). The range of sequence reads that mapped to SARS-CoV-2 RNA was 2,413 to 163,158 with

an average of 49,066. We observed a clear negative correlation between the C_t values of each

gene (*ORF1ab* and *N*) and the log value of SARS-CoV-2 RNA reads (**Figure 1**). However, the

linearity is not significant (R^2=0.6628, 0.5595 for *ORF1ab* and *N*, respectively), indicating that

the number of RNA reads measured by metagenomics sequencing are only semi-quantitative and

cannot be interpreted directly as viral loads.

Table 2. Summary of Sequencing Results of 10 Shaoxing Patient Samples.

ID	Ct Value (ORF1ab)	Ct Value (N)	Total Reads (PE 75)	2019-nCoV RNA (Raw Reads)	2019-nCoV RNA (Log Value)	Genome Coverage (%)	Mean Depth (X)
Shaoxing-01	21.57	23.62	17,158,277	40,057	4.60	99.4	219
Shaoxing-02	18.86	20.93	13,602,710	149,682	5.18	99.9	929
Shaoxing-03	20.09	22.25	24,769,343	163,158	5.21	99.8	1024
Shaoxing-04	24.02	24.68	21,509,477	15,424	4.19	100.0	81
Shaoxing-05	21.81	23.81	14,043,326	99,521	5.00	99.9	591
Shaoxing-06	25.88	27.08	18,909,299	3,535	3.55	99.9	18
Shaoxing-07	23.11	23.87	10,480,051	5,063	3.70	99.8	26
Shaoxing-08	23.34	24.53	11,506,909	2,413	3.38	99.9	12

Shaoxing-09	26.81	27.85	27,517,291	8,897	3.95	99.9	47
Shaoxing-10	26.24	26.8	11,071,595	2,911	3.46	99.7	15
Min	18.86	20.93	10,480,051	2,413	3.38	99.4	12
Max	26.81	27.85	27,517,291	163,158	5.21	100.0	1024
Mean	23.17	24.54	17,056,828	49,066	4.22	99.8	296

With a large variation in the SARS-CoV-2 RNA mapped reads, we were still able to

obtain excellent coverage and depth when each genome was mapped to the first SARS-CoV-2

genome, Wuhan-Hu-1 (Wu et al., 2020) (**Figure 2A**). The coverage for all genomes was above

99% and the mean depth for the genomes ranged from 12X to 1024X (**Table 2, Figure 2B**).

Genomes sequenced to a relatively low mean depth (12X to 47X) were still able to be genotyped

successfully (see Results below) but our results suggest that SARS-CoV-2 read counts of at least

15,000 yield sufficiently high depth to characterize even low prevalence or rare mutations.

<u>Estimation of Mutation Rate</u>

To determine the single nucleotide polymorphisms (SNPs) of SARS-CoV-2 in these 10

patients, we mapped each genome to the original Wuhan-Hu-1 reference which was collected on

December 31, 2019 (Wu et al., 2020). The genomes contained a fairly moderate number of SNPs

(mean of 4 SNPs, range 1-9) (**Table 3**), consistent with previous reports of relatively low

mutation rates (Wang et al., 2020b). The genomes with the largest number of SNPs came from

individuals who had contact with a confirmed case from Ningbo, Zhejiang and no travel history

to the Hubei province (**Table 3, Shaoxing-9 and 10**).

Table 3. Summary of Genomic Descriptions for the Shaoxing SARS-CoV-2 Genomes

ID	Lineage	Type	No. of SNP[a]	No. of Days[b]	Mutation Rate (#SNP/day)	Mutation Rate (#SNP/day/nt)	Mutation Rate (#SNP/yr/nt)
Shaoxing-01	A	S	2	27	0.07	2.48E-06	9.04E-04
Shaoxing-02	A	S	2	30	0.07	2.23E-06	8.14E-04
Shaoxing-03	A	S	2	30	0.07	2.23E-06	8.14E-04
Shaoxing-04	A	S	2	31	0.06	2.16E-06	7.87E-04

Shaoxing-05	B	L	2	31	0.06	2.16E-06	7.87E-04
Shaoxing-06	A	S	3	31	0.10	3.24E-06	1.18E-03
Shaoxing-07	B	L	5	30	0.17	5.57E-06	2.03E-03
Shaoxing-08	B	L	1	38	0.03	8.80E-07	3.21E-04
Shaoxing-09	A	S	9	36	0.25	8.36E-06	3.05E-03
Shaoxing-10	A	S	9	37	0.24	8.13E-06	2.97E-03
Min			1	27	0.03	8.80E-07	3.21E-04
Max			9	38	0.25	8.36E-06	3.05E-03
Mean			4	32	0.11	3.74E-06	1.37E-03

[a] SNP calculated by mapping each genome to the genome of Wuhan-Hu-1 (Wu et al., 2020)

[b] Number of days between the date that the sample was collected and the date the Wuhan-Hu-1 genome was
published (12/31/2019)

Using the SNP analysis, we calculated the various mutation rates using the number of
days between the date that the sample was collected and the date the Wuhan-Hu-1 sample was
collected. The mutation rate (SNP per day) ranged from 0.03 to 0.25 (**Table 3**). We used this
mutation rate to calculate the nucleotide substitution per site per day and the nucleotide
substitution per site per year. We saw an average mutation rate of 3.74×10^{-6} nucleotide
substitution per site per day and an average mutation rate of 1.37×10^{-3} nucleotide substitution per
site per year (**Table 3**).

We investigated each SNP to determine if there were any non-synonymous mutations in
genes important to the virus lifecycle (**Table 4**). No non-synonymous mutations were found in
the S gene, which encodes the spike protein that's critical for viral binding to human receptor
ACE2 (Zhou et al., 2020). Notably in the cluster of the two family members (Shaoxing-9 and -
10), the two viruses are closely related but not identical. Shaoxing-9 was infected first and then
transmitted to Shaoxing-10, whose virus gained a non-synonymous mutation C9962T in the
ORF1ab gene (**Table 4**). This could be explained by the sequential transmission, however, we
could not rule out a possibility of intra-host viral heterogeneity in the two patients.

Table 4. Summary of SNPs in the Ten SARS-CoV-2 Genomes

SNP#	Position	Gene	Reference nt	Shaoxing-01	Shaoxing-02	Shaoxing-03	Shaoxing-04	Shaoxing-05	Shaoxing-06	Shaoxing-07	Shaoxing-08	Shaoxing-09	Shaoxing-10
1	207	non-coding	C									T (non-coding)	
2	889	orf1ab	T							C (A)			
3	946	orf1ab	T									C (G)	C (G)
4	5099	orf1ab	T									A (S->T)	A (S->T)
5	7420	orf1ab	C									T (I)	T (I)
6	8344	orf1ab	C								T (D)		
7	8782	orf1ab	C	T (S)	T (S)	T (S)	T (S)		T (S)			T (S)	T (S)
8	9962	orf1ab	C										T (H->Y)
9	11430	orf1ab	A									G (Y->C)	G (Y->C)
10	11916	orf1ab	C							T (S->L)			
11	15324	orf1ab	C					T (N)					
12	21676	S	C									T (Y)	T (Y)
13	22081	S	G							A (Q)			
14	25672	ORF3a	C							A (L->I)			
15	28000	ORF8	C							T (P->L)			
16	28144	ORF8	T	C (L->S)	C (L->S)	C (L->S)	C (L->S)		C (L->S)			C (L->S)	C (L->S)
17	29095	N	C							T (F)			
18	29303	N	C					T (P->S)					
19	29625	ORF10	C									T (S->F)	T (S->F)

L Type (B Lineage)   S Type (A Lineage)   Non-Synonymous   Synonymous

SNP analysis was based on NC_045512.2 (Wuhan-Hu-1) as the reference genome (Wu et al., 2020). Variants are denoted as nucleotides versus the reference

base. Amino acid changes listed in parentheses with synonymous mutations listed as a single residue. The non-synonymous mutations are bolded.

<u>SARS-CoV-2 Genotype and Phylogenetic Characteristics</u>

Previous reports demonstrate that SARS-CoV-2 has two genotypes known as L type and

S type in the early phase of the outbreak (Tang et al., 2020). The majority of the Shaoxing

patients in this study have the S type (70%). We decided to compare our ten SARS-CoV-2

genomes to 163 other published SARS-CoV-2 genomes obtained from GISAID (Shu and

McCauley, 2017). Although the majority of the SARS-CoV-2 genomes obtained from GISAID

are the L type/B lineage (**Figure 3, red**), the 10 Shaoxing SARS-CoV-2 genomes (**Figure 3,**

green dots) are distributed throughout the these genomes with more of them classified in the S

type/A lineage (**Figure 3, blue**). Only two outliers (Shenzhen_SZTH-001_2020 and

South_Korea_KUMC01_2020) were classified as Lineage A but not assigned to either L or S

type (not shown in Figure 3). Due to a lack of the original sequencing data of these two outliers,

we are not able to investigate further regarding the discrepancy between the L/S type vs. A/B

lineage on them, as either a recombination event or a sequencing error is possible.

Interestingly, four of the Shaoxing SARS-CoV-2 genomes (Shaoxing -1 to -4) were

identical to six other GISAID SARS-CoV-2 genomes (**Figure 3, Cluster 1**). These six other

genomes were isolated from patients all over the world: two from Wuhan, two from Taiwan, one

from Belgium and one from Australia (**Figure 3, Cluster 1**). Shaoxing-6 is identical to five other

genomes isolated in Shenzhen, Guangdong Province in Southern China (**Figure 3, Cluster 2**).

Notably, in all ten Shaoxing patients, we found no virus with D614G Spike gene mutation, which

was shown to start spreading in Europe in early February, and rapidly become the dominant form

in the rest of the world out of China (Korber et al., 2020).

DISCUSSION

In this study, we sequenced the SARS-CoV-2 genome from ten patient samples from

Shaoxing, Zhejiang, China. Using metagenomic sequencing, we were able to obtain above 99%

coverage and an average depth of 296X for all 10 SARS-CoV-2 genomes. Although not

statistically significant, there does appear to be a clear negative correlation between the C_t values

of both gene targets and the log count of SARS-CoV-2 RNA sequence reads acquired by

metagenomics sequencing. This suggests that the log value of RNA sequence reads by

metagenomics sequencing may be used as a semi-quantitative measurement for SARS-CoV-2

viral loads.

The rapid spread of this virus is highlighted by the fact that four SARS-CoV-2 genomes

from Shaoxing individuals were identical to six other SARS-CoV-2 genomes from patients all

over the world. Our data support recent publications that the virus had spread rapidly around the

world especially in Europe before the United States (Deng et al., 2020;Gonzalez-Reiche et al.,

2020;Schuchat, 2020).

Overall, we did not see a large number of SNPs in these SARS-CoV-2 genomes. The

greatest number of SNPs seen was 9 and these two SARS-CoV-2 genomes were from individuals

with no travel history to Hubei province (**Table 3, Shaoxing-9 and 10**). Instead, Shaoxing-9 and

10 had contact with a confirmed case from Ningbo, another city outside of Hubei. We can use

these data to infer that the virus accumulated more mutations when it was spread to another city

outside of Hubei first before coming to Shaoxing, compared to the virus from people traveled to

Shaoxing directly from Hubei.

We combined epidemiologic data with the SNP analysis to estimate the mutation rate of

the SARS-CoV-2 from these ten patients. We saw an average mutation rate of 1.37×10^{-3}

nucleotide substitution per site per year for SARS-CoV-2, which is consistent with other reports

on the mutation rate of SARS-CoV-2(Li et al., 2020;Wang et al., 2020b) and SARS-CoV-1 with

a reported mutation rate of 0.80-2.38 x10^{-3} nucleotide substitution per site per year (Zhao et al.,

2004). These data demonstrate that SARS-CoV-2 is similar in the mutation rate as other

coronaviruses.

The major limitation of this study is that we only had 10 samples analyzed due to the

requirement of sufficient SARS-CoV-2 RNA from a metagenomic sample. However, with the

development of a SARS-CoV-2 probe enrichment or multiplex PCR protocols, this type of viral

sequencing analysis may be applied to samples with lower viral loads, thereby enabling more

complete molecular epidemiological surveillance. In addition, the C_t value cut-off of 28

established in this study may not be directly applicable to other real-time PCR assays due to the

technical differences.

In summary, we demonstrated that a full viral genomic analysis is feasible via

metagenomics sequencing directly on nasopharyngeal samples, which allows retrospective

molecular surveillance on SARS-CoV-2 to understand the dynamics of the outbreak in the early

phase. The identical virus found in patients in Shaoxing, a mid-sized city outside of Hubei,

China, and patients in Europe and Australia was striking. Our analysis added to the growing

body of evidence that SARS-CoV-2 spread extremely quickly around the globe as early as

January. Although only ten patients were included in this study, we found both lineages (A & B)

/types (L & S) of viruses with numerous mutations (both synonymous and non-synonymous)

across the entire viral genome. Our study contributed to the understanding of the SARS-CoV-2

evolution in the early phase of the COVID-19 pandemic.

ACKNOWLEDGEMENTS

We would like to thank Yong-Zhen Zhang (Fudan University) and Eddie Holmes

(University of Sydney) for sharing the sequence of the first SARS-CoV-2 isolate in a very timely

manner. We would also like to thank Fanchao Meng, Bin Hu, Haihao Shou and Yuanyuan Cai

from Shaoxing IngeniGen XMK Biotechnologies, Inc. for their technical assistance. This

manuscript has been released as a pre-print at MedRxiv:

https://doi.org/10.1101/2020.04.16.20058560. (Chen et al., 2020)

DISCLOSURE

Authors Huan Wu and Yifang Wang were employed by the company Shaoxing

IngeniGen XMK Biotechnologies. Author Fan Li was the Chief Executive Officer of the

company Three Coin Analytics. The remaining authors declare that the research was conducted

in the absence of any commercial or financial relationships that could be construed as a potential

conflict of interest. The authors declare that this study received funding from Shaoxing

IngeniGen XMK Biotechnologies. The funder had the following involvement with this study:

providing sequencing data and preliminary bioinformatics analysis. All authors declare no

conflict of interest.

References
274
Cascella, M., Rajnik, M., Cuomo, A., Dulebohn, S.C., and Di Napoli, R. (2020). "Features,

Evaluation and Treatment Coronavirus (COVID-19)," in *StatPearls*. (Treasure Island

(FL): StatPearls Publishing

StatPearls Publishing LLC.).

Chen, J., Hilt, E., Wu, H., Jiang, Z., Zhang, Q., Wang, J., Wang, Y., Li, F., Li, Z., Tang, J., and

Yang, S. (2020). Epidemiological and Genomic Analysis of SARS-CoV-2 in Ten Patients

from a Mid-sized City outside of Hubei, China. *medRxiv*, 2020.2004.2016.20058560.

Corman, V.M., Muth, D., Niemeyer, D., and Drosten, C. (2018). Hosts and Sources of Endemic

Human Coronaviruses. *Adv Virus Res* 100, 163-188.

Deng, X., Gu, W., Federman, S., Du Plessis, L., Pybus, O., Faria, N., Wang, C., Yu, G., Pan, C.-

Y., Guevara, H., Sotomayor-Gonzalez, A., Zorn, K., Gopez, A., Servellita, V., Hsu, E.,

Miller, S., Bedford, T., Greninger, A., Roychoudhury, P., Famulare, M., Chu, H.Y.,

Shendure, J., Starita, L., Anderson, C., Gangavarapu, K., Zeller, M., Spencer, E.,

Andersen, K., Maccannell, D., Tong, S., Armstrong, G., Paden, C., Li, Y., Zhang, Y.,

Morrow, S., Willis, M., Matyas, B., Mase, S., Kasirye, O., Park, M., Chan, C., Yu, A.,

Chai, S., Villarino, E., Bonin, B., Wadford, D., and Chiu, C.Y. (2020). A Genomic

Survey of SARS-CoV-2 Reveals Multiple Introductions into Northern California without

a Predominant Lineage. *medRxiv*, 2020.2003.2027.20044925.

Gonzalez-Reiche, A.S., Hernandez, M.M., Sullivan, M., Ciferri, B., Alshammary, H., Obla, A.,

Fabre, S., Kleiner, G., Polanco, J., Khan, Z., Alburquerque, B., Van De Guchte, A.,

Dutta, J., Francoeur, N., Melo, B.S., Oussenko, I., Deikus, G., Soto, J., Sridhar, S.H.,

Wang, Y.-C., Twyman, K., Kasarskis, A., Altman, D.R., Smith, M., Sebra, R., Aberg, J.,

Krammer, F., Garcia-Sarstre, A., Luksza, M., Patel, G., Paniz-Mondolfi, A., Gitman, M.,

Sordillo, E.M., Simon, V., and Van Bakel, H. (2020). Introductions and early spread of

SARS-CoV-2 in the New York City area. *medRxiv*, 2020.2004.2008.20056929.

Korber, B., Fischer, W., Gnanakaran, S., Yoon, H., Theiler, J., Abfalterer, W., Foley, B., Giorgi,

E., Bhattacharya, T., Parker, M., Partridge, D., Evans, C., Freeman, T., De Silva, T.,

Labranche, C., and Montefiori, D. (2020). Spike mutation pipeline reveals the emergence

of a more transmissible form of SARS-CoV-2. *bioRxiv*, 2020.2004.2029.069054.

Li, X., Wang, W., Zhao, X., Zai, J., Zhao, Q., Li, Y., and Chaillon, A. (2020). Transmission

dynamics and evolutionary history of 2019-nCoV. *J Med Virol* 92**,** 501-511.

Rambaut, A., Holmes, E.C., Hill, V., O'toole, Á., Mccrone, J.T., Ruis, C., Du Plessis, L., and

Pybus, O.G. (2020). A dynamic nomenclature proposal for SARS-CoV-2 to assist

genomic epidemiology. *bioRxiv*, 2020.2004.2017.046086.

Rota, P.A., Oberste, M.S., Monroe, S.S., Nix, W.A., Campagnoli, R., Icenogle, J.P., Penaranda,

S., Bankamp, B., Maher, K., Chen, M.H., Tong, S., Tamin, A., Lowe, L., Frace, M.,

Derisi, J.L., Chen, Q., Wang, D., Erdman, D.D., Peret, T.C., Burns, C., Ksiazek, T.G.,

Rollin, P.E., Sanchez, A., Liffick, S., Holloway, B., Limor, J., Mccaustland, K., Olsen-

Rasmussen, M., Fouchier, R., Gunther, S., Osterhaus, A.D., Drosten, C., Pallansch, M.A.,

Anderson, L.J., and Bellini, W.J. (2003). Characterization of a novel coronavirus

associated with severe acute respiratory syndrome. *Science* 300**,** 1394-1399.

Schuchat, A. (2020). Public Health Response to the Initiation adn Spread of the Pandemic

COVID-19 in the United States. *MMWR Morbidity, Mortaility Weekly Report*, 551-556.

Shu, Y., and Mccauley, J. (2017). GISAID: Global initiative on sharing all influenza data - from

vision to reality. *Euro Surveill* 22.

Tang, X., Wu, C., Li, X., Song, Y., Yao, X., Wu, X., Duan, Y., Zhang, H., Wang, Y., Qian, Z.,

Cui, J., and Lu, J. (2020). On the origin and continuing evolution of SARS-CoV-2.

National Science Review.

Wang, C., Horby, P.W., Hayden, F.G., and Gao, G.F. (2020a). A novel coronavirus outbreak of

global health concern. *Lancet* 395**,** 470-473.

Wang, C., Liu, Z., Chen, Z., Huang, X., Xu, M., He, T., and Zhang, Z. (2020b). The

establishment of reference sequence for SARS-CoV-2 and variation analysis. *J Med*

Virol.

Weiss, S.R., and Leibowitz, J.L. (2011). Coronavirus pathogenesis. *Adv Virus Res* 81**,** 85-164.

Wu, F., Zhao, S., Yu, B., Chen, Y.M., Wang, W., Song, Z.G., Hu, Y., Tao, Z.W., Tian, J.H., Pei,

Y.Y., Yuan, M.L., Zhang, Y.L., Dai, F.H., Liu, Y., Wang, Q.M., Zheng, J.J., Xu, L.,

Holmes, E.C., and Zhang, Y.Z. (2020). A new coronavirus associated with human

respiratory disease in China. *Nature* 579**,** 265-269.

Zaki, A.M., Van Boheemen, S., Bestebroer, T.M., Osterhaus, A.D., and Fouchier, R.A. (2012).

Isolation of a novel coronavirus from a man with pneumonia in Saudi Arabia. *N Engl J*

Med 367**,** 1814-1820.

Zhao, Z., Li, H., Wu, X., Zhong, Y., Zhang, K., Zhang, Y.P., Boerwinkle, E., and Fu, Y.X.

(2004). Moderate mutation rate in the SARS coronavirus genome and its implications.

BMC Evol Biol 4**,** 21.

Zhou, P., Yang, X.L., Wang, X.G., Hu, B., Zhang, L., Zhang, W., Si, H.R., Zhu, Y., Li, B.,

Huang, C.L., Chen, H.D., Chen, J., Luo, Y., Guo, H., Jiang, R.D., Liu, M.Q., Chen, Y.,

Shen, X.R., Wang, X., Zheng, X.S., Zhao, K., Chen, Q.J., Deng, F., Liu, L.L., Yan, B.,

Zhan, F.X., Wang, Y.Y., Xiao, G.F., and Shi, Z.L. (2020). A pneumonia outbreak

associated with a new coronavirus of probable bat origin. *Nature* 579**,** 270-273.

348   **Figure 1 Correlation of C_t Values and SARS-CoV-2 RNA Reads.**

349   **(a)** *ORF1ab* **Gene Target.** The X-axis plots the log value of the SARS-CoV-2 RNA reads while

350   the Y-axis plots the C_t values for the *ORF1ab* gene for the ten Shaoxing patients. **(b)** *N* **Gene**

351   **Target.** The X-axis plots the log value of the SARS-CoV-2 RNA reads while the Y-axis plots

352   the C_t values for the *N* gene for the ten Shaoxing patients.

Figure 2. Coverage and Depth

(a) Coverage and Depth Map. The coverage and depth at each base are depicted by the dark red shading along the circle. (**b**) **Depth Ratio.** The X-axis plots the log value of the depth for each genome while the Y-axis plots the cumulative percentage of bases covered to the specified depth.

Figure 3 Phylogenetic Comparison of SARS-CoV-2 Genomes.

Phylogenetic comparison of 162 published genomes from GISAID (Shu and McCauley, 2017) and the ten Shaoxing genomes (green dots). Genomes are color-coded based on their type/lineage (blue=S type/A lineage, red=L type/B lineage).