Interim evaluation of emergency declaration in Japan in the first week of the COVID-19 outbreak

Junko Kurita[1], Tamie Sugawara[2], Yasushi Ohkusa[2]

1)Department of Nursig , Tokiwa University, Ibaraki,, Japan

2) National Institute of Infectious Diseases, Tokyo, Japan

Keywords: COVID-19, voluntary restriction to going out, emergency declaration, outbreak in Japan, SIR model,

Corresponding author: Junko Kurita, kuritaj@tokiwa.ac.jp

Abstract

Background:　Prime Minister of Japan Abe declared an emergency to control the

COVID-19 outbreak on April 7, 2020. He asked almost half of the population of Japan

to reduce their personal contacts by 70–80%.

Object: This study estimates the effectiveness of that emergency declaration.

Method: We applied a simple susceptible–infected–recovery model to data of patients

with symptoms in Japan for January 14 – April 13. We estimate the reproduction

number before the emergency declaration (R_0) as 0.083; its range was [0.067, 0.105].

Results: Results suggest that the emergency declaration reduced the reproduction

number dramatically. The finding implies that the outbreak will be contained in a few

weeks.

Introduction

To control the COVID-19 outbreak in Japan, the Prime Minister Abe declared an emergency on April 7 and applied countermeasures starting from April 8 [1]. The declared measures required residents of seven prefectures, collectively accounting for 44.5% of Japan's population, to restrict their trips outside the home voluntarily. Voluntary compliance means that the declared measures were not enforced as law. People were not arrested even if they do not comply with this government declaration. At the same time, the policy required the reduction of contact by 70–80%, based on the author's earlier research [2]. However, the share of the population cooperating with government requirements without law enforcement has remained uncertain. Therefore, this policy's effects must be evaluated as nearly and in as real-time a manner as possible. If the reproduction number after the emergency declaration (R_e) is less than one, then the government would be able to rescind the declaration earlier. If greater than one, and especially of greater than two, then a more powerful policy, including law enforcement, might be necessary. Therefore, we undertook preliminary evaluation of the effect as of April 14, at one week after the declaration was issued.

Method

The COVID-19 outbreak in Japan, which occurred from January 14, and its related countermeasures, are divisible into four periods. The national government required sports and entertainment events to be canceled in Japan for two weeks from February 26 through March 19 according to a government advisory. At the same time, it was advised that small business and private meetings be cancelled voluntarily. Moreover, from March 3, almost all schools were closed to control the spread of the infection until spring vacation began in the middle of March. The policy was designated as voluntary event cancellation (VECSC). Consequently, the COVID-19 outbreak can be divided into four distinct periods: before VCSEC through February 26, VCSEC through March 19, and after VCSEC through April 7, and the after emergency was declared. We defined the reproduction numbers for the respective periods as R_0, R_v, R_a, and R_e.

Onset dates were sometimes unreported. Therefore, we estimated the onset date. To do so, we inferred a distribution of the length from onset to reporting based on patients for whom onset dates were available. Then we applied this empirical distribution to patients for whom onset dates were not available. Letting $f(k)$ represent this empirical distribution and letting Nt denote the number of patients for whom onset dates were not available published at date t, then the number of patients for whom the onset date was

t-1. The number of patients for whom onset dates were not available was estimated as

$f(1)N_t$. Similarly, the number of patients with onset date *t*-2 and whose onset dates were

not available was estimated as $f(2)N_t$. Therefore, the total number of patients whose

onset date were not available, given an onset date of *s*, was estimated as $\Sigma_{k=1}f(k)N_{s+k}$

for a long time passed from *s*.

Moreover, the reporting delay for published data from MHLW might be

considerable. In other words, if *s+k* was larger than that in the current period *t*, then *s+k*

represents the future for period *t*: thus N_{s+k} was not observable. Such a reporting delay

causes an under-estimation bias in the number of patients. Therefore, we must adjust it

as $\Sigma_{k=1}^{t-s}f(k)N_{s+k} / \Sigma_{k=1}^{t-s}f(k)$. Similarly, patients for whom the onset dates were available

are expected to be affected by the reporting delay. Thus we have $M_{s|t} / \Sigma_{k=1}^{t-s}f(k)$, where

$M_{s|t}$ represents the reported number of patients for whom onset dates were within

period *s* extending until the current period *t*.

We applied a simple susceptible–infected–recovery (SIR) model [2–4] to the

epidemic curve for Japan, with 120 million populations. We assume an incubation

period calculated according to the empirical distribution.

Symptomatic and asymptomatic states continued for one week and then moved to a

recovery state with probability of one. We are not concerned about outcomes or

necessary medical resources in this model. Therefore, the state of death or

hospitalization was not incorporated into the model. Asymptomatic cases cannot be

observable unless complete laboratory-based surveillance is performed. One exceptional

study found the asymptomatic cases to be 3/23 among elderly people [5]. For these

analyses, the powers of infectivity among severe patients and mild patients were equal.

Moreover, we assumed that asymptomatic cases have the same power of infectivity as

symptomatic cases have [5]. The distribution of infectiousness in symptomatic and

asymptomatic cases was assumed to be 30% on the onset day, 20% the following day,

and 10% for the next five days [5].

We sought R_0, R_v, R_a, and R_e.to fit the data to minimize the sum of absolute values

of discrepancies among the bootstrapped epidemic curve and the fitted values. The

estimated distribution of three reproduction numbers was calculated using 10,000

iterations of bootstrapping from the empirical distribution of the data obtained for

symptomatic patients.

We estimated the results sequentially as follows: We first estimated R_0 as the best

fit to bootstrapped data in the pre-VECSC period. Then based on the obtained R_0 and

course of the outbreak in the pre-VECSC period, we estimated R_v as the best fit to

bootstrapped data for the VECSC period. Similarly, based on the obtained R_0 and R_v,

we estimated R_a as the best fit to bootstrapped data in the post-VECSC period. Finally,

based on the obtained R_0, R_v, and R_a, we estimated R_e as the best fit to a bootstrapped

data after the emergency was declared. For each step, reproduction numbers were

grid-searched in the interval of (0, 10) by 0.001 resolution.

Data source

Data used were the numbers of symptomatic patients during January 14 – April 13

reported by MLHW [6] as of April 14. During this period, 6084 cases with onset dates

were reported. We excluded imported cases and those who were presumed to have been

infected passengers and crew from the Diamond Princess: they were presumed not to be

community-acquired cases in Japan. Of those, onset dates were available for 3734 cases;

onset dates were not reported for other cases.

Ethical consideration

All information used for this study has been published. There is therefore no ethical

issue related to this study.

Results

Figure 1 depicts the empirical distribution among incubation periods of 91 cases for which exposure dates and onset dates were reported by MHLW. Its mode was six days. Its average was 6.6 days. Figure 2 depicts the empirical distribution of duration from onset to report. We adjusted the reporting delay based on this distribution.

The value of R_0 before VECSC was introduced was estimated as 2.184; its 95% range was [2.082, 2.251]. However, R_v during the VECSC period when measures were introduced was estimated as 1.028 [0.965, 1.191]. After VECSC, R_a was estimated as 3.237 [3.005, 3.441]. After the emergency declaration, R_e decreased dramatically to 0.083 [0.067, 0.105]. The null hypothesis that R_0, R_e, R_a, and R_e were all equal was rejected.

Discussion

We applied a simple SIR model to all cases, including asymptomatic cases that had not been incorporated into the model to date. An earlier study [7–10] estimated R_0 for COVID-19 as 2.24–3.58 in Wuhan. Our R_0 obtained before VECSC was similar. However, an earlier study [11] estimated R_0 in Japan as 0.6 as of the end of February, which was the period mainly before the VECSC was implemented. Such a low number might mislead policy efforts for countermeasures in Japan as it adheres to contact

tracing efforts to detect clusters. Outbreaks contained themselves. Emergency

declarations would be unnecessary if 0.6 is correct.

Results show a dramatic decrease in the reproduction number on the basis of the

emergency declaration. The outbreak will be contained in a few weeks if the success

continues at the same level. In other words, the 0.6 figure reported from an earlier study

was comparable with our estimation for the period after the emergency declaration,

even though no countermeasure was used before VECSC.

However, after VECSC, the reproduction number escalated again to become higher

than that before VECSC. Therefore, a similar phenomenon might occur several weeks

after countermeasures cease. One must be reminded that these results reflect only those

at one week after the emergency declaration. The reproduction number probably

changed thereafter.

Conclusion

Results demonstrated that the reproduction number declined dramatically after the

emergency declaration. The reproduction number must be monitored carefully and

continuously. We hope that the present study contributes to government

decision-making about cessation of the emergency declaration.

This study was based on the authors' opinions. The analyses and results do not reflect any stance or policy of our affiliations.

Acknowledgments

We acknowledge the great efforts of all staff at public health centers, medical institutions, and other facilities who are fighting the spread and destruction associated with COVID-19.

Reference

1. Japan Times. Japan state of emergency to cover Tokyo, Osaka and five other
    prefectures
    https://www.japantimes.co.jp/news/2020/04/06/national/japan-state-of-emergency-c
    ovid-19/#.XpPANv37RaQ [accessed on April 15,2020]

2. Sugishita Y, Kurita J, Sugawara T, OhkusaY. Forecast of the COVID-19 outbreak,
    collapse of medical facilities, and lockdown effects in Tokyo, Japan.
    https://medrxiv.org/cgi/content/short/2020.04.02.20051490v1[accessed on
    April 15,2020]

3. Sugishita Y, Kurita J, Sugawara T, OhkusaY. Preliminary evaluation of voluntary
    event cancellation as a countermeasure against the COVID-19 outbreak in Japan as
    of   11   March,   2020.
    https://www.medrxiv.org/content/10.1101/2020.03.12.20035220v1

4. Ohkusa Y, Sugawara T, Taniguchi K, Okabe N. Real-time estimation and
    prediction for pandemic A/H1N1(2009) in Japan. J Infect Chemother.
    2011;17:468-72.

5. Kimball A, Hatfield KM, Arons M, James A, Taylor J, Spicer K, Bardossy AC,
    Oakley LP, Tanwar S, Chisty Z, Bell JM, Methner M, Harney J, Jacobs JR, Carlson
    CM, McLaughlin HP, Stone N, Clark S, Brostrom-Smith C, Page LC, Kay M,
    Lewis J, Russell D, Hiatt B, Gant J, Duchin JS, Clark TA, Honein MA, Reddy SC,
    Jernigan JA; Public Health ? Seattle & King County; CDC COVID-19
    Investigation Team. Asymptomatic and Presymptomatic SARS-CoV-2 Infections in
    Residents of a Long-Term Care Skilled Nursing Facility - King County,

Washington, March 2020. Morb Mortal Wkly Rep. 2020 Apr 3;69(13):377-381. doi: 10.15585/mmwr.mm6913e1.

6. Japan Ministry of Health, Labour and Welfare. Press Releases. https://www.mhlw.go.jp/stf/newpage_10723.html (in Japanese) [accessed on April 15, 2020]

7. Lee PI, Hu YL, Chen PY, HuanYC, Hsueh PR.Lee PI, et al. Are children less susceptible to COVID-19 ? J Microbiol Immunol Infect. 2020 Feb 25:S1684-1182;30039-6.

8. Zhao S, Lin Q, Ran J, Musa SS, Yang G , Wang W, Lou Y, Gao D, Yang L, He D, Wang M. Preliminary Estimation of the Basic Reproduction Number of Novel Coronavirus (2019-nCoV) in China, From 2019 to 2020: A Data-Driven Analysis in the Early Phase of the Outbreak. Int J Infect Dis 2020[Online ahead of print]

9. Liu Y, Gayle AA, Wilder-Smith A, Rockly J. The reproductive number of COVID-19 is higher compared to SARS coronavirus. J Travel Med. 2020.DOI: 10.1093/jtm/taaa021

10. Lai C, Shih T, Ko W, Tang H, Hsueh P. Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) and Coronavirus disease-2019 (COVID-19): The Epidemic and the Challenges. Int J Antimicrob Agents. DOI:10.1016/j.ijantimicag.2020.105924

11. H Nishiura, H Oshitani, T Kobayashi, T Saito, T Sunagawa, Takaji Wakita, MHLW COVID-19 Response Team, Mi Suzuki. Closed environments facilitate secondary transmission of coronavirus disease 2019 (COVID-19). doi: https://doi.org/10.1101/2020.02.28.20029272

Figure 1: Empirical distribution of incubation period published by Ministry of Labour,

Health and Welfare, Japan

(%)

Notes: Bars indicates the number of patients by incubation period among 59 cases

whose exposed date and onset date were published by Ministry of Labour, Health and

Welfare, Japan.

Figure 2: Empirical distribution of duration from onset to report by Ministry of Labour,

Health and Welfare, Japan

(%)

Note: Bars indicates probability of duration from onset to report base on 657 patients

whose onset date was available in Japan. Data source was Ministry of Labour, Health

and Welfare, Japan.