Title: **The Enlightenment of mainland China's epidemic situation to the world**

through the intrinsic growth rule of infected and cured cases with COVID-19

Chuanliang Han[1△], Yimeng Liu[2,3,4△], Jiting Tang[2,3,4] , Yuyao Zhu[2,3,4] and Saini Yang[2,3,4*]

[1]State Key Laboratory of Cognitive Neuroscience and Learning & IDG/McGovern

Institute for Brain Research, Beijing Normal University, Beijing 100875, China

[2]Key Laboratory of Environmental Change and Natural Disaster, Ministry of Education,

Beijing Normal University, Beijing 100875, China

[3]State Key Laboratory of Earth Surface Processes and Resource Ecology, Beijing

Normal University, Beijing 100875, China

[4]Academy of Disaster Reduction and Emergency Management Ministry of Emergency

Management and Ministry of Education, Faculty of Geographical Science, Beijing

Normal University, Beijing 100875, China

[△]C.L.H and Y.M.L contributed equally to this work

*Correspondence authors:

Saini Yang,

Key Laboratory of Environmental Change and Natural Disaster, Ministry of Education

& State Key Laboratory of Earth Surface Processes and Resource Ecology & Academy

of Disaster Reduction and Emergency Management Ministry of Emergency

Management and Ministry of Education, Faculty of Geographical Science, Beijing

Normal University, Beijing 100875, China

Email: yangsaini@bnu.edu.cn

Conflict of Interest: The authors declare no competing conflicts of interest.

Acknowledgements: This study is sponsored by the National Key Research and

Development Program of China (2018YFC1508903), the National Natural Science

Foundation of China (41621061) and the support of International Center for

Collaborative Research on Disaster Risk Reduction (ICCR-DRR).

Abstract

The novel coronavirus disease (COVID-19) emerged at the end of 2019 has been

controlled in mainland China so far, while it is still spreading all over the world and

causing global panic. When the pandemic will end is an issue of great concern. A

logistic model depicting the growth rules of infected and cured cases in mainland China

may shed some light on this question. We extend this model to 31 countries outside

China experiencing serious COVID-2019 outbreaks. The model well explained data

from each country in our study (R-Square $\geq$ 0.95). For the infected case, the

semi-saturation period of these countries are ranging from 63 to 170 (Mar 3 to Jun 18).

For all countries in our study, the increasing rate (k) and semi-saturation period (t_0) of

the fitted curves are negatively correlated for both infected and cured cases. The

increasing rate of infected case is positively correlated with that of cured case, so is the

semi-saturation period. We used a simple linear model to bridge the relationship

between the increasing rules of infected and cured case. According to the connection of

infected and cured growth rules which is identified from cases in China, we predicted

that the semi-saturation period of the cured case outside China ranges from 82 to 196

(Mar 22 to July 8) for cured cases. More importantly, we found there exists strong

positive correlation between the semi-saturation period (t_0) of infected cases and the

timing of government's response, which could provide strong evidence for the

effectiveness of rapid epidemic prevention in various countries.

Introduction

In December 2019, a series of cases with unknown cause of pneumonia appeared in Wuhan, the capital of Hubei Province in mainland China (WHOₐ). Deep sequencing analysis later from lower respiratory samples confirmed a novel coronavirus, which was firstly named as 2019 novel coronavirus (2019-nCoV) on Jan 12, 2020 and may originate from bat (Zhou et al., 2020). The coronavirus (SARS-CoV-2) has been officially renamed as COVID-19 on February 12, 2020. Human to human transmission through COVID-19 has been confirmed not only in China (Wang et al., 2020; Li et al., 2020; Zhu et al., 2020; Hui et al., 2020), but in dozens of countries around world such as Republic of Korea (Choi et al., 2020; Ki et al., 2020; Shim et al., 2020), Italy (Livingston et al., 2020; Spina et al., 2020; Rosenbaum, 2020) and Iran (Tuite et al., 2020; Abdi, 2020; Zhuang et al., 2020). So far, the epidemic situation in mainland China has been effectively controlled since there are only sparse daily local new cases since Mar 19, 2020, and the cure rate in mainland China has risen up to 93.51% by April 13 (Chinese Center for Disease Control and Prevention, CCDC). However, the epidemic situation outside China is much more serious than before. As of Apr 13, there have been a total of 1788665 confirmed cases of COVID-19 globally, and the death and cured rate are 6.35% and 20.45 % respectively (Sina.com). When the inflection point will come for both infection and cure outside China remains unclear. The hope and lessons for COVID-19 control from China is necessary and worth to quantify (Azman & Luquero, 2020).

We applied a descriptive model that has been proved robust and stable (Han et al., 2020) to analyze the data on the COVID-19 global cases (infected and cure cases in mainland China, and infected cases countries outside China). We then obtained the parameter relationship between parameters of infected and cure cases in mainland China. Further, we used that relationship to map parameter space of infected case to the parameter space of cured cases for 31 countries globally. In the end, we tried to explore the relationship

between the model parameters and governmental control measures.

Method

Sources of Data:

The cumulative number of confirmed and cured COVID-19 cases in mainland China

was obtained from the National Health Commission of China, and the provincial Health

Commission of 30 provincial administrative regions (excluded Tibet, because the only

one confirmed infected COVID-19 case in Tibet has been cured on Feb. 12) in mainland

China (January 10 to March 19, 2020), which is publically available. All cases were

laboratory confirmed following the published standards made by the National Health

Commission of China (CCDC$_a$). The basic test procedure has been described in detail in

previous work (Zhou et al., 2020; Huang et al., 2020). This dataset was partially

analyzed in an initial work on this topic (Han et al., 2020).

We selected 31 countries outside China with serious coronavirus epidemic situation in

our analyses (January 10 to Apr 13, 2020), whose population counts for 39.4% of the

world's population (57.8% if includes China) and 81.6% of the world's infected

COVID-19 cases (86.2% if includes China) (WHO$_a$). The data of COVID-19 cases in

countries outside China were obtained from the situation reports on the official website

of World Health Organization (WHO$_a$), which is publicly available. The data used in this

study include the cumulative number of reported laboratory-confirmed COVID-19 cases.

Countries included in our study are Republic of Korea, Japan, Australia, Singapore,

which are in Western Pacific Region; Italy, Spain, France, Germany, Switzerland, the

United Kingdom, Netherlands, Sweden, Denmark, Austria, Belgium, Portugal, Czechia,

Finland, Ukraine, Slovakia, Bulgaria, Lithuania, which are in European Region; India in

South-East Asia Region, Islamic Republic of Iran, Lebanon, which are in Eastern

Mediterranean Region; United States of America, Canada, Brazil, Mexico, which are in

America; South Africa, Ethiopia, which are in Africa. All the laboratory-confirmed

cases are determined according to the WHO standard.

Governments of these countries all declared a state of national emergency, wartime, or a

blockade of their borders as a result of the COVID-19 outbreak. In this study, we

collected the time point of government declaration from mainstream authoritative media

of each country we selected.

Epidemic curve modeling

In this work, we used a sigmoid-formed descriptive model (Equation (1)), which is a

derivation from logistic form differential equations (Han et al., 2020).

$$N(t) = \frac{A}{1 + e^{-k(t - t_0)}} \quad (1)$$

where $N(t)$ is the general form of the cumulative numbers of infected or cured patients at

time t. A denotes the maximum number of infections or cures, k represents incidence

growth rate, t_0 is the inflection point of the sigmoid curve. This descriptive model has

been verified by not only the infected, death and cured cases of COVID-19, but also the

data of SARS in 2003 (Han et al., 2020). For infected case, there are three parameters (A,

k, t_0) in the model, and for cured case, we fixed A to the subtraction of the maximum

number of cumulative infections and death, which obeys the basic biological fact. It is

worth mentioning that in our model, t_0 is the inflection point in mathematical definition.

In this paper, we assumed that t_0 is the time of inflection point of epidemic situation in a

region.

We processed the data and modeling with custom scripts on MATLAB (the Math

Works). We adopted the nonlinear least square (NLS) algorithm for data fitting and

parameter estimation.

Results

With the cumulative number of confirmed and cured cases in 30 provinces of mainland China as well as in 31 countries and above-mentioned model, all data could be well explained (R^2>0.95) by the descriptive model (See Methods, Fig. 1A-C, Fig. 4A-C).

Relationship between infected and cured cases in mainland China

The epidemic situation of COVID-19 has been controlled in China (CCDC$_b$), with a cure rate of 93.51% (CCDC$_a$). Time series of infected and cured cases and their fitted curves of three example provinces (provinces of Anhui, Heilongjiang, Hebei) are shown in Fig. 1A-C. By observing the fitted curves of all provinces in mainland China (Supplementary Figure S1), we found the increasing rate (k) of infected case is larger than that of cured case and the semi-saturation period (t_0) of infected cases is shorter than that of cured cases.

As shown in Figure 2, the parameter space of k and t_0 are separated for infected (red circle) and cured cases (blue circle), and there is a significant negative correlation (for infected case, r=-0.46, p=0.0103; for cured case, r=-0.56, p=0.0012; Pearson correlation). As we compared these two parameters of infected and cured cases respectively, we found infection case has larger (t=6.9136, p<10^{-7}, right-tailed test) increasing rate (Fig. 2B) but smaller (t=-37.271, p<10^{-25}, left-tailed test) semi-saturation period (Fig. 2C) than that of cured cases. The increasing rate of infected and cured case is significant positively correlated (r=0.38, p=0.0359, Pearson Correlation), so is the semi-saturation period of infected and cured case (r=0.66, p=0.0001, Pearson Correlation). Each of them could be fitted to a linear model, with a slope and intercept of 0.36 (CI: [0.0268, 0.698]) and 0.1 (CI: [0.014, 0.191]), respectively (panel B), and that of panel C is 1.06 (CI: [0.59, 1.54]) and 15.36 (CI: [-0.85, 31.56]).

Intrinsic growth rules of COVID-19 infections outside China

With the dataset of infected case in mainland China, there shows time stability of this

164 model (Fig. 3). An example of Fujian province is shown in Figure 3A. It is clear that as

165 the increase of length of time-series data, the fitting curves converges better to the actual

166 number of cases (black dots).

167 We defined an index, the minimum time length (MTL), to measure how much data is

168 necessary to produce stable output. This index is the time length with which the change

169 of parameter value (k or t_0) is less than 5% in two consecutive days. With the data of 30

170 provinces in China, we found that the MTL for increasing rate (k) is 15 days (Fig. 3B)

171 and for semi-saturation period (t_0), is 26 days (Fig. 3C). This result provides some hint

172 on how much data is necessary to measure the epidemic situation.

174 After identifying the relationship between the increasing rate and the semi-saturation

175 period, we applied this relationship to the COVID-19 cases from 31 countries. We

176 applied the descriptive model to fit the data of the cumulative infected cases in each of

177 the 31 countries. Figure 4 illustrates three countries' time-series data of infection

178 (Canada, Germany and Iran, Fig. 4A-C) (See Supplementary Figure S2 for all 31

179 countries' plots). All data series of infected case could be well explained by our model

180 (R^2>0.95). The average change rate of these parameters is already lower than 5% (Fig 4

181 D-F) which indicates the parameters we estimated from the model is stable. We noticed

182 that the increasing rate of 31-country cases does not have a significant difference

183 (t=1.197, p=0.237 two-tailed test) with that of Chinese provinces, but the 31-country

184 cases have significantly longer (t=-17.4819, p<10^{-20}, right-tailed test) semi-saturation

185 period. The numbers indicated that the t_0 of infections in 31 countries is averaged at 98

186 (std: 20.5), and the date matched to which is around Apr 7 (ranging from Mar 3 to Jun

187 18).

189 **Prediction for growth rules of cured COVID-19 cases and its relation to the timing**

190 **of governmental control measures**

191 By observing the increasing rate (k) and semi-saturation period (t_0) of infected case in 31

countries, we found there was a weak negative correlation (r=-0.63, p=0.0001, Pearson

Correlation) (Fig. 5) between these two parameters, which shows similar tendency with

results of provinces in mainland China (Fig. 2B).To predict the increasing rate (k) and

inflection point (t_0) of the cured COVID-19 cases outside China, we mapped the

parameter space of infected cases into the parameter space of cured cases, based on the

parameter relationship between infected cases and cured cases obtained from China (Fig.

2BC). For the 31 countries, the increasing rate ranges from 0.08 to 0.45. The mean t_0 of

the fitted cured cases curves is 119.71 (std: 21.73), which means the mean

semi-saturation period (t_0) for these 31 countries will come around Apr. 29, 2020

(ranging from Mar 22 to July 8).

To explore the practical implication of these parameters, we took the timing of

governmental emergency control measures related to COVID-19 as a variable to analyze

its relationship with the key parameters (Fig. 6). We found the Semi-Saturation Period

(SSP) showed a significant positive correlation (r=0.73, p<0.0001, Spearman correlation)

with the timing of governmental control measures (Fig. 6 BE), and k showed a marginal

significant negative correlation (r=-0.3, p=0.0925, Spearman correlation), while the

increasing rate or the maximum of infected number do not show strong correlation

(r=0.02, p=0.91, Spearman correlation) with the control measures' timing (Fig. 6

ACDF). China released the earliest national control measures to prevent the spread of

COVID-19 (on Jan. 23) and its t_0 is the lowest (Figure 5, red circle). We also noticed

that the timing of governmental emergency control measure is significantly earlier than

SSP (Fig. 6B) (t=2.99, p=0.0043, two-tailed test) in time, which indicates that most of

the selected countries have taken measures at the early phase of the outbreak.

Discussion

We compared the relationship between growth characteristics of infected and cured case of 30 provinces in mainland China based on the data from Jan 20 to Mar 29, 2020, which are positively correlated (Fig. 2 BC). Further, we explained the data from 31 countries using the same model (See Methods). We predicted their increasing rate (k) and semi-saturation period (t_0) of cure case based on the knowledge of data from China. It is worth mentioning that there is a strong correlation between these parameters and the timing of governmental policy on COVID-19.

Comparison with previous work

Different from models like SIR or SEIR applied in COVID-19 (Li et al., 2020; Wang et al., 2020; Tian et al., 2020), our model is relatively simple, but robust. This work is an extension of our previous work (Han et al., 2020), we have verified that our model well explains the data in various geographic space of mainland China, as well as provided robust results with different lengths of time series (between 30 and 40 days). This model is able to capture the features of macroscopic dynamic of the epidemics worldwide ($R^2 > 0.95$) (Fig. 4) and we quantitatively investigated the minimum data length to produce stable output (Fig. 3). By now, the domestic infected cases in China is very limited, which is consistent with our prediction results. The cumulative number of infected and cured cases are almost saturated, and this means the spread of the virus has been effectively controlled, thus we believe that the parameters estimated in this work for Chinese provinces are reliable (Fig. 1A-C). The characteristics of two curves can depict the dynamic macroscopic features of the epidemic situation. Moreover, we extended this model to the data in each of these 31 countries (Fig 4 A-C), and found the sigmoid model could also fit the data very well ($R^2 > 0.95$) (Fig 4 A-C).

Relationship among model parameters and the prediction of the cured cases

We noticed that semi-saturation period of infected and cured cases in China showed a

positive correlation (p-value = 0.0001). This phenomenon obeys the common sense,

since the cure is always later than infection when a new virus comes. Although the

positive correlation between increasing rate of infected and cured cases is significant

(p-value = 0.0359), we noticed the correlation is much lower than that of semi-saturation

period. This result is reasonable, since the reasons cause the ascending increasing rate of

infected case are totally different from that of cured case. So that correlation may not be

that strong. As we compared the k and t_0 in any specific case no matter in mainland

China or worldwide region (Fig. 5), we found significant negative correlations. This is

consistent with our initial work (Han et al., 2020). For negative correlation of k and t_0 in

cured cases, it could also be explained as when the increasing rate (k) of the cured case

is larger, it will be faster to cure most patients.

Suggestion for governments around the world on the prevention and control of

COVID-19

More importantly, we found there is a strong correlation between the timing of

governmental control measures for COVID-19 in 31 countries and the semi-saturation

period estimated (Fig 6 BE). This indicates the early implementation of the

government's prevention and control policy effectively shortened the turning point of

the epidemic. Taken China as an example, when the COVID-19 outbreak in Wuhan, the

strict isolation policy was announcement on Jan. 23, and the inflection point there was

around Feb 9, 2020, the time lag was 17 days. Another result showed that the time of

governmental policy is significantly shorter than the inflection time point of infection,

which is also a strong reference for governments of all countries to take control

measures as early as possible.

Reference

Reference

Sina, https://news.sina.cn/zt_d/yiqing0121

WHO$_a$, https://www.who.int/docs/default-source/coronaviruse/situation-reports

CCDC$_a$ Prevention, C. C. f. D. C. a. Prevention and control measures of COVID-19,

<http://www.nhc.gov.cn/jkj/s3577/202002/a5d6f7b8c48c451c87dba14889b30147/files/

3514cb996ae24e2faf65953b4ecd0df4.pdf> (2020).

CCDC$_b$ http://www.nhc.gov.cn/xcs/yqfkdt/gzbd_index.shtml

Li R, Pei S, Chen B, Song Y, Zhang T, Yang W, Shaman J (2020) Substantial
undocumented infection facilitates the rapid dissemination of novel coronavirus
(SARS-CoV2). Science.

Zhou P et al. (2020) A pneumonia outbreak associated with a new coronavirus of
probable bat origin. Nature.

Wang D, Hu B, Hu C, Zhu F, Liu X, Zhang J, Wang B, Xiang H, Cheng Z, Xiong Y,
Zhao Y, Li Y, Wang X, Peng Z (2020) Clinical Characteristics of 138 Hospitalized
Patients With 2019 Novel Coronavirus-Infected Pneumonia in Wuhan, China. JAMA.

Li Q et al. (2020) Early Transmission Dynamics in Wuhan, China, of Novel
Coronavirus-Infected Pneumonia. N Engl J Med.

Zhu N, Zhang D, Wang W, Li X, Yang B, Song J, Zhao X, Huang B, Shi W, Lu R, Niu P,
Zhan F, Ma X, Wang D, Xu W, Wu G, Gao GF, Tan W, China Novel Coronavirus I,
Research T (2020) A Novel Coronavirus from Patients with Pneumonia in China, 2019.
N Engl J Med.

Hui DS, E IA, Madani TA, Ntoumi F, Kock R, Dar O, Ippolito G, McHugh TD, Memish
ZA, Drosten C, Zumla A, Petersen E (2020) The continuing 2019-nCoV epidemic threat
of novel coronaviruses to global health - The latest 2019 novel coronavirus outbreak in
Wuhan, China. Int J Infect Dis 91:264-266.

Choi SC, Ki M (2020) Estimating the reproductive number and the outbreak size of
Novel Coronavirus disease (COVID-19) using mathematical model in Republic of
Korea. Epidemiol Health:e2020011.

Ki M, nCo VT (2020) Epidemiologic characteristics of early cases with 2019 novel
coronavirus (2019-nCoV) disease in Republic of Korea. Epidemiol Health:e2020007.

Shim E, Tariq A, Choi W, Lee Y, Chowell G (2020) Transmission potential and severity
of COVID-19 in South Korea. Int J Infect Dis.

Abdi M (2020) Coronavirus disease 2019 (COVID-19) outbreak in Iran; actions and
problems. Infect Control Hosp Epidemiol:1-5.

Tuite AR, Bogoch, II, Sherbo R, Watts A, Fisman D, Khan K (2020) Estimation of
Coronavirus Disease 2019 (COVID-19) Burden and Potential for International
Dissemination of Infection From Iran. Ann Intern Med.

Zhuang Z, Zhao S, Lin Q, Cao P, Lou Y, Yang L, He D (2020) Preliminary estimation of
the novel coronavirus disease (COVID-19) cases in Iran: A modelling analysis based on

overseas cases and air travel data. Int J Infect Dis.

Livingston E, Bucher K (2020) Coronavirus Disease 2019 (COVID-19) in Italy. JAMA.

Rosenbaum L (2020) Facing Covid-19 in Italy - Ethics, Logistics, and Therapeutics on
the Epidemic's Front Line. N Engl J Med.

Spina S, Marrazzo F, Migliari M, Stucchi R, Sforza A, Fumagalli R (2020) The response
of Milan's Emergency Medical System to the COVID-19 outbreak in Italy. Lancet
395:e49-e50.

Han C, Liu Y, Yang S (2020) Intrinsic growth rules of patients infected, dead and cured
with 2019 novel coronavirus in mainland China. medRxiv:2020.2002.2023.20024802.

Zhou P et al. (2020) A pneumonia outbreak associated with a new coronavirus of
probable bat origin. Nature.

Huang C et al. (2020) Clinical features of patients infected with 2019 novel coronavirus
in Wuhan, China. Lancet.

Li R, Pei S, Chen B, Song Y, Zhang T, Yang W, Shaman J (2020) Substantial
undocumented infection facilitates the rapid dissemination of novel coronavirus
(SARS-CoV2). Science.

Wang H, Wang Z, Dong Y, Chang R, Xu C, Yu X, Zhang S, Tsamlag L, Shang M,
Huang J, Wang Y, Xu G, Shen T, Zhang X, Cai Y (2020) Phase-adjusted estimation of
the number of Coronavirus Disease 2019 cases in Wuhan, China. Cell Discov 6:10.

H. Tian et al., Science 10.1126/science.abb6105 (2020).

Azman AS, Luquero FJ (2020) From China: hope and lessons for COVID-19 control.
Lancet Infect Dis.

Figure 1 Example provinces for the time series of infected and cured cases with
corresponding fitted curve in mainland China

A-C shows three example cases in three provinces (Anhui, Heilongjiang and Hebei). The horizontal axis
in each panel denotes the x^{th} day after January 1, 2020. The vertical axis in each panel is the cumulative
number of infected (red circles) and cured case (blue circles). The red and blue line are the fitting curves
by the sigmoid model, and the dashed red and blue curves mark the 95% confidence interval of the fitting
curves.

Figure 2 Relationship between intrinsic rules for infected and cured case in mainland
China

Panel A shows the scatter plot of increasing rate (k) and semi-saturation period (t_0) for infected (red circles)
and cured case (blue circles).

Panel B is the scatter plot of increasing rate (k) of infected and cured case (black circles), the dashed line
is the diagonal line of the coordinate system (the same for Panel C). The relation between k of infected
and cured cases is modeled with a linear function, which is shown as the black curve. The bar graph
shown in northwest corner demonstrates the difference between *k* of infected and cured cases.

Panel C shows the scatter plot of semi-saturation period (t_0). The black curve is the linear function
estimated based on the relation between infected and cured case. And the bar graph shown in southeast
corner demonstrates difference between t_0 of infected and cured cases.

Figure 3 Time stability of logistic model in infected case of Chinese mainland

Panel A is an example case in Fujian province. The horizontal axis in each panel denotes the x th day after

January 1, 2020. The vertical axis in each panel is the cumulative number of infected (black dots).

The lines with different colors are the fitting curve by the logistic model using data with different time

length as shown in color bar in the southeast corner. Panel B and C illustrates how estimated parameters (k

and t_0) changes with increase of the data's time length. The black curve shows the average of increasing

rate (k) across 30 provinces in mainland China changing with time length of data and the dashed line

marks its standard error.

Figure 4 Example countries for the time series of infected cases with corresponding
fitted curve around world and its temporal stability

Panel A-C show infected cases and their fitted curves of Australia, Italy and Iran. The horizontal axis is
the x^{th} day after Jan 1, 2020. The vertical axis denotes the number of infected cases in corresponding
country. The black dots in each panel are raw data of the cumulative number of infected cases. The red
line is the fitted curves by our descriptive model. The dashed red lines mark the 95% confidence interval
of the fitted curves. Panel D-F show the difference of parameters estimated from relative longer (by the
end of Jun 13) and shorter (by the end of Apr 12) time series data, each dot denotes for one country.

Figure 5 Prediction for intrinsic rule of cured case around world based on Chinese
enlightenment

The scatter plot of increasing rate (k) and semi-saturation period (t_0) for infected cases is showed in red
circles. Based on the linear model shown in Fig 2BC, a prediction of k and t_0 for cured case is plotted as
blue circles.

Figure 6 Relationship between time of government policy from each countries and parameters estimated from the model

Panel A-C show the scatter plot of the time of government measures and the model parameters of infected case (k, t_0 and A, respectively). In the southeast corner of Panel B, a bar graph was shown to indicate the difference between timing of government measures (grey bar) and semi-saturation period (SAP, black bar).

D-F shows the same data of A-C but in the order form. The red circle in six panels denotes China.

Supplement Figure S1 Intrinsic growth rules of patients infected with 2019 novel coronavirus in all provinces of mainland China

The deep red and blue dots in each panel are raw data of the cumulative infected and cured cases. The red and blue line is the corresponding fitted curves by our descriptive model.

407 **Supplement Figure S2 Intrinsic growth rules of patients infected with 2019 novel**

408 **coronavirus in 31 countries around world**

409 The black dots in each panel are raw data of the cumulative infected cases. The red line is the fitted curve

410 by our descriptive model.

413